

Technical notes: Kinetic data for MISTRA

The Potential Importance of Frost Flowers, Recycling on Snow, and Open Leads for Ozone Depletion Events

M. Piot and R. von Glasow

Atmos. Chem. Phys., 2008

May 7, 2008

1 Tables of reaction rates

This collection comprises a complete listing of all gas and aqueous phase species (Table 1), gas phase (Table 2) and aqueous phase (Table 3) reaction rates, as well as rates for the heterogeneous (particle surface) reactions (Table 4), aqueous phase equilibrium constants (Table 5), Henry constants and accommodations coefficients (Table 6).

Table 1: Species in MISTRA

| |
|---|
| Gas phase |
| O ¹ D, O ₂ , O ₃ , OH, HO ₂ , H ₂ O ₂ , H ₂ O |
| NO, NO ₂ , NO ₃ , N ₂ O ₅ , HONO, HNO ₃ , HNO ₄ , PAN, NH ₃ , RONO ₂ |
| CO, CO ₂ , CH ₄ , C ₂ H ₆ , C ₂ H ₄ , HCHO, HCOOH, ALD (i.e., CH ₃ CHO), CH ₂ O ₂ , HOCH ₂ O ₂ , CH ₃ CO ₃ , CH ₃ O ₂ , C ₂ H ₅ O ₂ , H ₃ CO ₂ , EO ₂ (i.e., H ₂ C(OH)CH ₂ OO), CH ₂ O ₂ , ROOH (i.e., alkylhydroperoxides), DOM |
| SO ₂ , SO ₃ , HOSO ₂ , H ₂ SO ₄ , DMS, CH ₃ SCH ₂ OO, DMSO, DMSO ₂ , CH ₃ S, CH ₃ SO, CH ₃ SO ₂ , CH ₃ SO ₃ , CH ₃ SO ₂ H, CH ₃ SO ₃ H |
| Cl, ClO, OClO, HCl, HOCl, Cl ₂ , Cl ₂ O ₂ , ClNO ₂ , ClNO ₃ |
| Br, BrO, HBr, HOBr, Br ₂ , BrNO ₂ , BrNO ₃ , BrCl, CHBr ₃ , CH ₃ Br |
| Liquid phase (neutral) |
| O ₂ , O ₃ , OH, HO ₂ , H ₂ O ₂ , H ₂ O |
| NO, NO ₂ , NO ₃ , HONO, HNO ₃ , HNO ₄ , NH ₃ |
| CO ₂ , HCHO, HCOOH, CH ₃ OH, CH ₃ OO, CH ₃ OOH |
| SO ₂ , H ₂ SO ₄ , DMSO, DMSO ₂ , CH ₃ SO ₂ H, CH ₃ SO ₃ H |
| Cl, HCl, HOCl, Cl ₂ |
| Br, HBr, HOBr, Br ₂ , BrCl |
| Liquid phase (ions) |
| H ⁺ , OH ⁻ , O ₂ ⁻ |
| NO ₂ ⁻ , NO ₃ ⁻ , NO ₄ ⁻ , NH ₄ ⁺ |
| HCO ₃ ⁻ , CO ₃ ⁻ , HCOO ⁻ |
| HSO ₃ ⁻ , SO ₃ ²⁻ , HSO ₄ ⁻ , SO ₄ ²⁻ , HSO ₅ ⁻ , SO ₃ ⁻ , SO ₄ ⁻ , SO ₅ ⁻ , CH ₃ SO ₃ ⁻ , CH ₂ OHSO ₂ ⁻ , CH ₂ OHSO ₃ ⁻ |
| Cl ⁻ , Cl ₂ ⁻ , ClO ⁻ , ClOH ⁻ |
| Br ⁻ , Br ₂ ⁻ , BrO ⁻ , BrCl ₂ ⁻ , Br ₂ Cl ⁻ , BrOH ⁻ |

Table 2: Gas phase reactions.

| no | reaction | n | $A [(\text{cm}^{-3})^{1-n} \text{s}^{-1}]$ | $-E_a / R$ [K] | reference |
|------|---|-----|--|----------------|---------------------------|
| O 1 | $\text{O}^1\text{D} + \text{O}_2 \longrightarrow \text{O}_3$ | 2 | 3.2×10^{-11} | 70 | Atkinson et al. (2006) |
| O 2 | $\text{O}^1\text{D} + \text{N}_2 \longrightarrow \text{O}_3$ | 2 | 1.8×10^{-11} | 110 | Atkinson et al. (2006) |
| O 3 | $\text{O}^1\text{D} + \text{H}_2\text{O} \longrightarrow 2 \text{OH}$ | 2 | 2.2×10^{-10} | | Atkinson et al. (2006) |
| O 4 | $\text{OH} + \text{O}_3 \longrightarrow \text{HO}_2 + \text{O}_2$ | 2 | 1.7×10^{-12} | -940 | Atkinson et al. (2006) |
| O 5 | $\text{OH} + \text{HO}_2 \longrightarrow \text{H}_2\text{O} + \text{O}_2$ | 2 | 4.8×10^{-11} | 250 | Atkinson et al. (2006) |
| O 6 | $\text{OH} + \text{H}_2\text{O}_2 \longrightarrow \text{HO}_2 + \text{H}_2\text{O}$ | 2 | 2.9×10^{-12} | -160 | Atkinson et al. (2006) |
| O 7 | $\text{HO}_2 + \text{O}_3 \longrightarrow \text{OH} + 2\text{O}_2$ | 2 | 1.0×10^{-14} | -490 | Atkinson et al. (2004) |
| O 8 | $\text{HO}_2 + \text{HO}_2 \longrightarrow \text{H}_2\text{O}_2 + \text{O}_2$ | 2 | 2.3×10^{-13} | 600 | Atkinson et al. (2006) |
| O 9 | $\text{O}_3 + h\nu \longrightarrow \text{O}_2 + \text{O}^1\text{D}$ | 1 | 1 | | DeMore et al. (1997) |
| O 10 | $\text{H}_2\text{O}_2 + h\nu \longrightarrow 2\text{OH}$ | 1 | 1 | | DeMore et al. (1997) |
| N 1 | $\text{NO} + \text{OH} \xrightarrow{M} \text{HONO}$ | 3 | 2 | | Sander et al. (2003) |
| N 2 | $\text{NO} + \text{HO}_2 \longrightarrow \text{NO}_2 + \text{OH}$ | 2 | 3.5×10^{-12} | 250 | Atkinson et al. (2004) |
| N 3 | $\text{NO} + \text{O}_3 \longrightarrow \text{NO}_2 + \text{O}_2$ | 2 | 3.0×10^{-12} | -1500 | Sander et al. (2003) |
| N 4 | $\text{NO} + \text{NO}_3 \longrightarrow 2\text{NO}_2$ | 2 | 1.5×10^{-11} | 170 | Sander et al. (2003) |
| N 5 | $\text{NO}_2 + \text{OH} \xrightarrow{M} \text{HNO}_3$ | 3 | 2 | | Sander et al. (2003) |
| N 6 | $\text{NO}_2 + \text{HO}_2 \xrightarrow{M} \text{HNO}_4$ | 3 | 2 | | Atkinson et al. (2006) |
| N 7 | $\text{NO}_2 + \text{O}_3 \longrightarrow \text{NO}_3 + \text{O}_2$ | 2 | 1.2×10^{-13} | -2450 | Sander et al. (2003) |
| N 8 | $\text{NO}_2 + h\nu \longrightarrow \text{NO} + \text{O}_3$ | 1 | 1 | | DeMore et al. (1997) |
| N 9 | $\text{NO}_2 + \text{NO}_3 \xrightarrow{M} \text{N}_2\text{O}_5$ | 3 | 2 | | Sander et al. (2003) |
| N 10 | $\text{NO}_3 + h\nu \longrightarrow \text{NO} + \text{O}_2$ | 1 | 1 | | Wayne et al. (1991) |
| N 11 | $\text{NO}_3 + \text{HO}_2 \longrightarrow 0.3 \text{HNO}_3 + 0.7 \text{OH} + 0.7 \text{NO}_2 + \text{O}_2$ | 2 | 4.0×10^{-12} | | Atkinson et al. (2006) |
| N 12 | $\text{NO}_3 + \text{NO}_3 \longrightarrow \text{NO}_2 + \text{NO}_2 + \text{O}_2$ | 2 | 8.5×10^{-13} | -2450 | Sander et al. (2003) |
| N 13 | $\text{NO}_3 + h\nu \longrightarrow \text{NO}_2 + \text{O}_3$ | 1 | 1 | | Wayne et al. (1991) |
| N 14 | $\text{N}_2\text{O}_5 \xrightarrow{M} \text{NO}_2 + \text{NO}_3$ | 2 | 2 | | Sander et al. (2003) |
| N 15 | $\text{N}_2\text{O}_5 + \text{H}_2\text{O} \longrightarrow 2\text{HNO}_3$ | 2 | 2.6×10^{-22} | | Atkinson et al. (2006) |
| N 16 | $\text{N}_2\text{O}_5 + h\nu \longrightarrow \text{NO}_2 + \text{NO}_3$ | 1 | 1 | | DeMore et al. (1997) |
| N 17 | $\text{HONO} + \text{OH} \longrightarrow \text{NO}_2$ | 2 | 1.8×10^{-11} | -390 | Sander et al. (2003) |
| N 18 | $\text{HONO} + h\nu \longrightarrow \text{NO} + \text{OH}$ | 1 | 1 | | DeMore et al. (1997) |
| N 19 | $\text{HNO}_3 + h\nu \longrightarrow \text{NO}_2 + \text{OH}$ | 1 | 1 | | DeMore et al. (1997) |
| N 20 | $\text{HNO}_3 + \text{OH} \longrightarrow \text{NO}_3 + \text{H}_2\text{O}$ | 2 | 2 | | Atkinson et al. (2006) |
| N 21 | $\text{HNO}_4 \xrightarrow{M} \text{NO}_2 + \text{HO}_2$ | 2 | 2 | | Sander et al. (2003) |
| N 22 | $\text{HNO}_4 + \text{OH} \longrightarrow \text{NO}_2 + \text{H}_2\text{O} + \text{O}_2$ | 2 | 1.3×10^{-12} | 380 | Haggerstone et al. (2005) |
| N 23 | $\text{HNO}_4 + h\nu \longrightarrow \text{NO}_2 + \text{HO}_2$ | 1 | 1 | | DeMore et al. (1997) |
| N 24 | $\text{HNO}_4 + h\nu \longrightarrow \text{OH} + \text{NO}_3$ | 1 | 1 | | DeMore et al. (1997) |
| N 25 | $\text{RONO}_2 + \text{OH} \longrightarrow \text{R} + \text{H}_2\text{O} + \text{NO}_2$ | 2 | 1.3×10^{-12} | | Sander et al. (1997) |
| N 26 | $\text{RONO}_2 + h\nu \longrightarrow \text{RO} + \text{NO}_2$ | 1 | assumed similar as N 19 | | |

Table 2: Continued.

| no | reaction | n | A [(cm ⁻³) ¹⁻ⁿ s ⁻¹] | $-E_a / R$ [K] | reference |
|------|--|-----|---|----------------|---------------------------------|
| C 1 | CO + OH $\xrightarrow{O_2}$ HO ₂ + CO ₂ | 2 | 2 | | Sander et al. (2003) |
| C 2 | CH ₄ + OH $\xrightarrow{O_2}$ CH ₃ OO + H ₂ O | 2 | 2.4×10^{-12} | -1775 | Sander et al. (2003) |
| C 3 | C ₂ H ₆ + OH \rightarrow C ₂ H ₅ O ₂ + H ₂ O | 2 | 1.7×10^{-11} | -1232 | Lurmann et al. (1986) |
| C 4 | C ₂ H ₄ + OH \rightarrow EO ₂ | 2 | 1.66×10^{-12} | 474 | Sander et al. (1997), see note |
| C 5 | C ₂ H ₄ + O ₃ \rightarrow HCHO + 0.4CH ₂ O ₂ + 0.12HO ₂ + 0.42CO + 0.06CH ₄ | 2 | 1.2×10^{-14} | -2633 | Lurmann et al. (1986), see note |
| C 6 | HO ₂ + CH ₃ OO \rightarrow ROOH + O ₂ | 2 | 4.1×10^{-13} | 750 | Sander et al. (2003) |
| C 7 | HO ₂ + C ₂ H ₅ O ₂ \rightarrow ROOH + O ₂ | 2 | 7.5×10^{-13} | 700 | Sander et al. (2003) |
| C 8 | HO ₂ + CH ₃ CO ₃ \rightarrow ROOH + O ₂ | 2 | 4.5×10^{-13} | 1000 | DeMore et al. (1997) |
| C 9 | CH ₃ OO + CH ₃ OO \rightarrow 1.4HCHO + 0.8HO ₂ + O ₂ | 2 | 1.5×10^{-13} | 220 | Lurmann et al. (1986) |
| C 10 | C ₂ H ₅ O ₂ + NO \rightarrow ALD + HO ₂ + NO ₂ | 2 | 4.2×10^{-12} | 180 | Lurmann et al. (1986) |
| C 11 | 2C ₂ H ₅ O ₂ \rightarrow 1.6ALD + 1.2HO ₂ | 2 | 5.00×10^{-14} | | Lurmann et al. (1986) |
| C 12 | EO ₂ + NO \rightarrow NO ₂ + 2.0HCHO + HO ₂ | 2 | 4.2×10^{-12} | 180 | Lurmann et al. (1986) |
| C 13 | EO ₂ + EO ₂ \rightarrow 2.4HCHO + 1.2HO ₂ + 0.4ALD | 2 | 5.00×10^{-14} | | Lurmann et al. (1986) |
| C 14 | HO ₂ + EO ₂ \rightarrow ROOH + O ₂ | 2 | 3.00×10^{-12} | | Lurmann et al. (1986) |
| C 15 | HCHO + hν \rightarrow 2HO ₂ + CO | 1 | 1 | | DeMore et al. (1997) |
| C 16 | HCHO + hν \rightarrow CO + H ₂ | 1 | 1 | | DeMore et al. (1997) |
| C 17 | HCHO + OH $\xrightarrow{O_2}$ HO ₂ + CO + H ₂ O | 2 | 1.00×10^{-11} | | DeMore et al. (1997) |
| C 18 | HCHO + HO ₂ \rightarrow HOCH ₂ O ₂ | 2 | 6.7×10^{-15} | 600 | Sander et al. (2003) |
| C 19 | HCHO + NO ₃ $\xrightarrow{O_2}$ HNO ₃ + HO ₂ + CO | 2 | 5.8×10^{-16} | | DeMore et al. (1997) |
| C 20 | ALD + OH \rightarrow CH ₃ CO ₃ + H ₂ O | 2 | 6.9×10^{-12} | 250 | Lurmann et al. (1986) |
| C 21 | ALD + NO ₃ \rightarrow HNO ₃ + CH ₃ CO ₃ | 2 | 1.40×10^{-15} | | DeMore et al. (1997) |
| C 22 | ALD + hν \rightarrow CH ₃ OO + HO ₂ + CO | 1 | 1 | | Lurmann et al. (1986) |
| C 23 | ALD + hν \rightarrow CH ₄ + CO | 1 | 1 | | Lurmann et al. (1986) |
| C 24 | HOCH ₂ O ₂ + NO \rightarrow HCOOH + HO ₂ + NO ₂ | 2 | 4.2×10^{-12} | 180 | Lurmann et al. (1986) |
| C 25 | HOCH ₂ O ₂ + HO ₂ \rightarrow HCOOH + H ₂ O + O ₂ | 2 | 2.00×10^{-12} | | Lurmann et al. (1986) |
| C 26 | 2 HOCH ₂ O ₂ \rightarrow 2HCOOH + 2HO ₂ + 2O ₂ | 2 | 1.00×10^{-13} | | Lurmann et al. (1986) |
| C 27 | HCOOH + OH $\xrightarrow{O_2}$ HO ₂ + H ₂ O + CO ₂ | 2 | 4.0×10^{-13} | | DeMore et al. (1997) |
| C 28 | CH ₃ CO ₃ + NO ₂ \rightarrow PAN | 2 | 4.70×10^{-12} | | Lurmann et al. (1986) |
| C 29 | PAN \rightarrow CH ₃ CO ₃ + NO ₂ | 1 | 1.9×10^{16} | -13543 | DeMore et al. (1997) |
| C 30 | CH ₃ CO ₃ + NO \rightarrow CH ₃ OO + NO ₂ + CO ₂ | 2 | 4.2×10^{-12} | 180 | Lurmann et al. (1986) |
| C 31 | CH ₃ OO + NO $\xrightarrow{O_2}$ HCHO + NO ₂ + HO ₂ | 2 | 3.0×10^{-12} | 280 | DeMore et al. (1997) |
| C 32 | ROOH + OH \rightarrow 0.7 CH ₃ OO + 0.3 HCHO + 0.3 OH | 2 | 3.8×10^{-12} | 200 | DeMore et al. (1997), see note |
| C 33 | ROOH + hν \rightarrow HCHO + OH + HO ₂ | 1 | 1 | | DeMore et al. (1997), see note |

Table 2: Continued.

| no | reaction | n | A [(cm ⁻³) ¹⁻ⁿ s ⁻¹] | $-E_a / R$ [K] | reference |
|------|--|-----|---|----------------|---|
| S 1 | $\text{SO}_2 + \text{OH} \xrightarrow{M} \text{HOSO}_2$ | 3 | 2 | | Atkinson et al. (2006) |
| S 2 | $\text{HOSO}_2 + \text{O}_2 \rightarrow \text{HO}_2 + \text{SO}_3$ | 2 | 1.3×10^{-12} | 330 | Atkinson et al. (2006) |
| S 3 | $\text{SO}_3 \xrightarrow{\text{H}_2\text{O}} \text{H}_2\text{SO}_4$ | 1 | 2 | | Jayne et al. (1997) |
| S 4 | $\text{CH}_3\text{SCH}_3 + \text{OH} \rightarrow \text{CH}_3\text{SCH}_2\text{OO} + \text{H}_2\text{O}$ | 2 | 2 | | Atkinson et al. (1997) |
| S 5 | $\text{CH}_3\text{SCH}_3 + \text{OH} \xrightarrow{\text{O}_2} \text{CH}_3\text{SOCH}_3 + \text{HO}_2$ | 2 | 2 | | Atkinson et al. (1997) |
| S 6 | $\text{CH}_3\text{SCH}_3 + \text{NO}_3 \xrightarrow{\text{O}_2} \text{CH}_3\text{SCH}_2\text{OO} + \text{HNO}_3$ | 2 | 1.9×10^{-13} | 520 | Atkinson et al. (1999) |
| S 7 | $\text{CH}_3\text{SCH}_3 + \text{Cl} \xrightarrow{\text{O}_2} \text{CH}_3\text{SCH}_2\text{OO} + \text{HCl}$ | 2 | 3.3×10^{-10} | | Atkinson et al. (1999) |
| S 8 | $\text{CH}_3\text{SCH}_3 + \text{Br} \xrightarrow{\text{O}_2} \text{CH}_3\text{SCH}_2\text{OO} + \text{HBr}$ | 2 | 9.0×10^{-11} | -2386 | Jefferson et al. (1994) |
| S 9 | $\text{CH}_3\text{SCH}_3 + \text{BrO} \rightarrow \text{CH}_3\text{SOCH}_3 + \text{Br}$ | 2 | 2.54×10^{-14} | 850 | Ingham et al. (1999) |
| S 10 | $\text{CH}_3\text{SCH}_3 + \text{ClO} \rightarrow \text{CH}_3\text{SOCH}_3 + \text{Cl}$ | 2 | 9.5×10^{-15} | | Barnes et al. (1991) |
| S 11 | $\text{CH}_3\text{SCH}_3 + \text{IO} \rightarrow \text{CH}_3\text{SOCH}_3 + \text{I}$ | 2 | 1.4×10^{-14} | | THALOZ (2005) |
| S 12 | $\text{CH}_3\text{SCH}_2\text{OO} + \text{NO} \rightarrow \text{HCHO} + \text{CH}_3\text{S} + \text{NO}_2$ | 2 | 4.9×10^{-12} | 263 | Urbanski et al. (1997) |
| S 13 | $\text{CH}_3\text{SCH}_2\text{OO} + \text{CH}_3\text{SCH}_2\text{OO} \xrightarrow{\text{O}_2} 2 \text{HCHO} + 2 \text{CH}_3\text{S}$ | 2 | 1.0×10^{-11} | | Urbanski et al. (1997); Atkinson et al. (2006) |
| S 14 | $\text{CH}_3\text{S} + \text{O}_3 \rightarrow \text{CH}_3\text{SO} + \text{O}_2$ | 2 | 1.15×10^{-12} | 432 | Atkinson et al. (2006) |
| S 15 | $\text{CH}_3\text{S} + \text{NO}_2 \rightarrow \text{CH}_3\text{SO} + \text{NO}$ | 2 | 3.0×10^{-11} | 210 | Atkinson et al. (2006) |
| S 16 | $\text{CH}_3\text{SO} + \text{NO}_2 \xrightarrow{\text{O}_2} 0.82 \text{CH}_3\text{SO}_2 + 0.18 \text{SO}_2 + 0.18 \text{H}_3\text{CO}_2 + \text{NO}$ | 2 | 1.2×10^{-11} | | Atkinson et al. (2006); Kukui et al. (2000), product ratios from van Dingenen et al. (1994) |
| S 17 | $\text{CH}_3\text{SO} + \text{O}_3 \xrightarrow{\text{O}_2} \text{CH}_3\text{SO}_2$ | 2 | 6.0×10^{-13} | | Atkinson et al. (2006) |
| S 18 | $\text{CH}_3\text{SO}_2 \rightarrow \text{SO}_2 + \text{CH}_3\text{OO}$ | 1 | 1.9×10^{13} | -8661 | Barone et al. (1995) |
| S 19 | $\text{CH}_3\text{SO}_2 + \text{NO}_2 \rightarrow \text{CH}_3\text{SO}_3 + \text{NO}$ | 2 | 2.2×10^{-12} | | Ray et al. (1996) |
| S 20 | $\text{CH}_3\text{SO}_2 + \text{O}_3 \rightarrow \text{CH}_3\text{SO}_3$ | 2 | $3. \times 10^{-13}$ | | Barone et al. (1995) |
| S 21 | $\text{CH}_3\text{SO}_3 + \text{HO}_2 \rightarrow \text{CH}_3\text{SO}_3\text{H}$ | 2 | $5. \times 10^{-11}$ | | Barone et al. (1995) |
| S 22 | $\text{CH}_3\text{SO}_3 \xrightarrow{\text{H}_2\text{O}, \text{O}_2} \text{CH}_3\text{OO} + \text{H}_2\text{SO}_4$ | 1 | 1.36×10^{14} | -11071 | Barone et al. (1995) |
| S 23 | $\text{CH}_3\text{SOCH}_3 + \text{OH} \rightarrow 0.95 \text{CH}_3\text{SO}_2\text{H} + 0.95 \text{CH}_3\text{OO} + 0.05 \text{DMSO}_2$ | 2 | 8.7×10^{-11} | | Urbanski et al. (1998) |
| S 24 | $\text{CH}_3\text{SO}_2\text{H} + \text{OH} \rightarrow 0.95 \text{CH}_3\text{SO}_2 + 0.05 \text{CH}_3\text{SO}_3\text{H} + 0.05 \text{HO}_2 + \text{H}_2\text{O}$ | 2 | $9. \times 10^{-11}$ | | Kukui et al. (2003) |
| S 25 | $\text{CH}_3\text{SO}_2\text{H} + \text{NO}_3 \rightarrow \text{CH}_3\text{SO}_2 + \text{HNO}_3$ | 2 | 1.0×10^{-13} | | Yin et al. (1990) |

Table 2: Continued.

| no | reaction | n | A [(cm ³) ¹⁻ⁿ s ⁻¹] | $-E_a / R$ [K] | reference |
|-------|--|-----|--|----------------|--|
| Cl 1 | Cl + O ₃ → ClO + O ₂ | 2 | 2.8 × 10 ⁻¹¹ | -250 | Atkinson et al. (2006) |
| Cl 2 | Cl + HO ₂ → HCl + O ₂ | 2 | 1.8 × 10 ⁻¹¹ | 170 | Sander et al. (2003) |
| Cl 3 | Cl + HO ₂ → ClO + OH | 2 | 4.1 × 10 ⁻¹¹ | -450 | Sander et al. (2003) |
| Cl 4 | Cl + H ₂ O ₂ → HCl + HO ₂ | 2 | 1.1 × 10 ⁻¹¹ | -980 | Atkinson et al. (2006) |
| Cl 5 | Cl + CH ₃ OO → 0.5 ClO + 0.5 HCHO + 0.5 HO ₂ + 0.5 HCl + 0.5 CO + 0.5 H ₂ O | 2 | 1.6 × 10 ⁻¹⁰ | | Sander et al. (2003) |
| Cl 6 | Cl + NO ₃ → ClO + NO ₂ | 2 | 2.4 × 10 ⁻¹¹ | | Mellouki et al. (1987) |
| Cl 7 | Cl + CH ₄ $\xrightarrow{O_2}$ HCl + CH ₃ OO | 2 | 9.6 × 10 ⁻¹² | -1360 | Sander et al. (2003) |
| Cl 8 | Cl + C ₂ H ₆ $\xrightarrow{O_2}$ HCl + C ₂ H ₅ O ₂ | 2 | 7.7 × 10 ⁻¹¹ | -90 | Sander et al. (2003) |
| Cl 9 | Cl + C ₂ H ₄ $\xrightarrow{O_2}$ HCl + C ₂ H ₅ O ₂ | 2 | 1. × 10 ⁻¹⁰ | | Sander et al. (1997), see note |
| Cl 10 | Cl + HCHO $\xrightarrow{O_2}$ HCl + HO ₂ + CO | 2 | 8.1 × 10 ⁻¹¹ | -30 | Sander et al. (2003) |
| Cl 11 | Cl + ROOH → CH ₃ OO + HCl | 2 | 5.7 × 10 ⁻¹¹ | | Wallington et al. (1990), see note |
| Cl 12 | Cl + OClO → ClO + ClO | 2 | 3.2 × 10 ⁻¹¹ | 170 | Atkinson et al. (2006) |
| Cl 13 | Cl + ClNO ₃ → Cl ₂ + NO ₃ | 2 | 6.5 × 10 ⁻¹² | 135 | Sander et al. (2003) |
| Cl 14 | Cl + PAN → HCl + HCHO + NO ₃ | 2 | 1.0 × 10 ⁻¹⁴ | | Tsalkani et al. (1988) |
| Cl 15 | Cl + HNO ₃ → HCl + NO ₂ | 2 | 1.0 × 10 ⁻¹⁶ | | Wine et al. (1988) |
| Cl 16 | Cl + RONO ₂ → HCl + NO ₂ | 2 | 7.7 × 10 ⁻¹¹ | | estimated from Muthuramu et al. (1994) |
| Cl 17 | ClO + OH → Cl + HO ₂ | 2 | 7.4 × 10 ⁻¹² | -270 | Sander et al. (2003) |
| Cl 18 | ClO + OH → HCl + O ₂ | 2 | 6.0 × 10 ⁻¹³ | -230 | Sander et al. (2003) |
| Cl 19 | ClO + HO ₂ → HOCl + O ₂ | 2 | 2.2 × 10 ⁻¹² | 340 | Atkinson et al. (2006) |
| Cl 20 | ClO + CH ₃ OO → Cl + HCHO + HO ₂ | 2 | 3.3 × 10 ⁻¹² | -115 | Sander et al. (2003) |
| Cl 21 | ClO + NO → Cl + NO ₂ | 2 | 6.2 × 10 ⁻¹² | 295 | Atkinson et al. (2006) |
| Cl 22 | ClO + NO ₂ \xrightarrow{M} ClNO ₃ | 3 | 2 | | Atkinson et al. (2006) |
| Cl 23 | ClO + ClO → Cl ₂ O ₂ | 2 | 2 | | Atkinson et al. (2006) |
| Cl 24 | ClO + ClO → Cl ₂ + O ₂ | 2 | 1.0 × 10 ⁻¹² | -1590 | Atkinson et al. (2006) |
| Cl 25 | ClO + ClO → Cl ₂ O ₂ | 2 | 3.0 × 10 ⁻¹¹ | -2450 | Atkinson et al. (2006) |
| Cl 26 | ClO + ClO → Cl + OClO | 2 | 3.5 × 10 ⁻¹³ | -1370 | Atkinson et al. (2006) |
| Cl 27 | OClO + OH → HOCl + O ₂ | 2 | 4.5 × 10 ⁻¹³ | 800 | Atkinson et al. (2006) |
| Cl 28 | OClO + NO → ClO + NO ₂ | 2 | 1.1 × 10 ⁻¹³ | 350 | Atkinson et al. (2006) |
| Cl 29 | Cl ₂ O ₂ → ClO + ClO | 1 | 2 | | Atkinson et al. (2006) |
| Cl 30 | HOCl + OH → ClO + H ₂ O | 2 | 3.0 × 10 ⁻¹² | -500 | Sander et al. (2003) |
| Cl 31 | HCl + OH → H ₂ O + Cl | 2 | 1.8 × 10 ⁻¹² | -240 | Atkinson et al. (2006) |
| Cl 32 | ClNO ₂ + OH → HOCl + NO ₂ | 2 | 2.4 × 10 ⁻¹² | -1250 | Atkinson et al. (2006) |

Table 2: Continued.

| no | reaction | n | $A [(\text{cm}^{-3})^{1-n} \text{s}^{-1}]$ | $-E_a / R [\text{K}]$ | reference |
|-------|---|-----|--|-----------------------|---------------------------------|
| Cl 33 | $\text{ClNO}_3 + \text{OH} \longrightarrow 0.5 \text{ClO} + 0.5 \text{HNO}_3 + 0.5 \text{HOCl} + 0.5 \text{NO}_3$ | 2 | 1.2×10^{-12} | -330 | Atkinson et al. (2006) |
| Cl 34 | $\text{ClNO}_3 \longrightarrow \text{ClO} + \text{NO}_2$ | 1 | 2 | | Anderson and Fahey (1990) |
| Cl 35 | $\text{OCIO} + h\nu \xrightarrow{\text{O}_2, \text{O}_3} \text{O}_3 + \text{ClO}$ | 1 | 1 | | DeMore et al. (1997) |
| Cl 36 | $\text{Cl}_2\text{O}_2 + h\nu \longrightarrow \text{Cl} + \text{Cl} + \text{O}_2$ | 1 | 1 | | DeMore et al. (1997) |
| Cl 37 | $\text{Cl}_2 + h\nu \longrightarrow 2 \text{Cl}$ | 1 | 1 | | DeMore et al. (1997) |
| Cl 38 | $\text{HOCl} + h\nu \longrightarrow \text{Cl} + \text{OH}$ | 1 | 1 | | DeMore et al. (1997) |
| Cl 39 | $\text{ClNO}_2 + h\nu \longrightarrow \text{Cl} + \text{NO}_2$ | 1 | 1 | | DeMore et al. (1997) |
| Cl 40 | $\text{ClNO}_3 + h\nu \longrightarrow \text{Cl} + \text{NO}_3$ | 1 | 1 | | DeMore et al. (1997) |
| Br 1 | $\text{Br} + \text{O}_3 \longrightarrow \text{BrO} + \text{O}_2$ | 2 | 1.7×10^{-11} | -800 | Atkinson et al. (2006) |
| Br 2 | $\text{Br} + \text{HO}_2 \longrightarrow \text{HBr} + \text{O}_2$ | 2 | 7.7×10^{-12} | -450 | Atkinson et al. (2006) |
| Br 3 | $\text{Br} + \text{C}_2\text{H}_4 \xrightarrow{\text{O}_2} \text{HBr} + \text{C}_2\text{H}_5\text{O}_2$ | 2 | $5. \times 10^{-14}$ | | Sander et al. (1997), see note |
| Br 4 | $\text{Br} + \text{HCHO} \xrightarrow{\text{O}_2} \text{HBr} + \text{CO} + \text{HO}_2$ | 2 | 1.7×10^{-11} | -800 | Sander et al. (2003) |
| Br 5 | $\text{Br} + \text{ROOH} \longrightarrow \text{CH}_3\text{OO} + \text{HBr}$ | 2 | 2.66×10^{-12} | -1610 | Mallard et al. (1993), see note |
| Br 6 | $\text{Br} + \text{NO}_2 \longrightarrow \text{BrNO}_2$ | 2 | 2 | | Sander et al. (2003) |
| Br 7 | $\text{Br} + \text{BrNO}_3 \longrightarrow \text{Br}_2 + \text{NO}_3$ | 2 | 4.9×10^{-11} | | Orlando and Tyndall (1996) |
| Br 8 | $\text{BrO} + \text{OH} \longrightarrow \text{Br} + \text{HO}_2$ | 2 | 1.8×10^{-11} | 250 | Atkinson et al. (2006) |
| Br 9 | $\text{BrO} + \text{HO}_2 \longrightarrow \text{HOBr} + \text{O}_2$ | 2 | 4.5×10^{-12} | 500 | Atkinson et al. (2006) |
| Br 10 | $\text{BrO} + \text{CH}_3\text{OO} \longrightarrow \text{HOBr} + \text{HCHO}$ | 2 | 4.1×10^{-12} | | Aranda et al. (1997) |
| Br 11 | $\text{BrO} + \text{CH}_3\text{OO} \longrightarrow \text{Br} + \text{HCHO} + \text{HO}_2$ | 2 | 1.6×10^{-12} | | Aranda et al. (1997) |
| Br 12 | $\text{BrO} + \text{HCHO} \xrightarrow{\text{O}_2} \text{HOBr} + \text{CO} + \text{HO}_2$ | 2 | 1.5×10^{-14} | | Hansen et al. (1999) |
| Br 13 | $\text{BrO} + \text{NO} \longrightarrow \text{Br} + \text{NO}_2$ | 2 | 8.7×10^{-12} | 260 | Atkinson et al. (2006) |
| Br 14 | $\text{BrO} + \text{NO}_2 \xrightarrow{M} \text{BrNO}_3$ | 3 | 2 | | Atkinson et al. (2006) |
| Br 15 | $\text{BrO} + \text{BrO} \longrightarrow 2 \text{Br} + \text{O}_2$ | 2 | 2.4×10^{-12} | 40 | Sander et al. (2003) |
| Br 16 | $\text{BrO} + \text{BrO} \longrightarrow \text{Br}_2 + \text{O}_2$ | 2 | 2.9×10^{-14} | 860 | Sander et al. (2003) |
| Br 17 | $\text{HBr} + \text{OH} \longrightarrow \text{Br} + \text{H}_2\text{O}$ | 2 | 5.5×10^{-12} | 205 | Atkinson et al. (2006) |
| Br 18 | $\text{BrNO}_3 \longrightarrow \text{BrO} + \text{NO}_2$ | 1 | 2 | | Orlando and Tyndall (1996) |
| Br 19 | $\text{BrO} + h\nu \xrightarrow{\text{O}_2} \text{Br} + \text{O}_3$ | 1 | 1 | | DeMore et al. (1997) |
| Br 20 | $\text{Br}_2 + h\nu \longrightarrow 2 \text{Br}$ | 1 | 1 | | Hubinger and Nee (1995) |
| Br 21 | $\text{HOBr} + h\nu \longrightarrow \text{Br} + \text{OH}$ | 1 | 1 | | Ingham et al. (1999) |
| Br 22 | $\text{BrNO}_2 + h\nu \longrightarrow \text{Br} + \text{NO}_2$ | 1 | 1 | | Scheffer et al. (1997) |
| Br 23 | $\text{BrNO}_3 + h\nu \longrightarrow \text{Br} + \text{NO}_3$ | 1 | 1 | | DeMore et al. (1997) |
| Br 24 | $\text{Br}_2 + \text{OH} \longrightarrow \text{HOBr} + \text{Br}$ | 1 | 1 | | Atkinson et al. (2006) |
| Br 25 | $\text{CH}_3\text{Br} + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{Br}$ | 2 | 2.0×10^{-11} | 240 | Atkinson et al. (2006) |
| Br 26 | $\text{CHBr}_3 + \text{OH} \longrightarrow \text{H}_2\text{O} + \text{Br}$ | 2 | 1.7×10^{-12} | -1215 | Atkinson et al. (2006) |
| | | 2 | 1.35×10^{-12} | -600 | Atkinson et al. (2004) |

Table 2: Continued.

| no | reaction | n | $A [(\text{cm}^{-3})^{1-n} \text{s}^{-1}]$ | $-E_a / R [\text{K}]$ | reference |
|------|--|-----|--|-----------------------|------------------------|
| Hx 1 | $\text{Cl} + \text{BrCl} \rightarrow \text{Br} + \text{Cl}_2$ | 2 | 1.5×10^{-11} | | Mallard et al. (1993) |
| Hx 2 | $\text{Cl} + \text{Br}_2 \rightarrow \text{BrCl} + \text{Br}$ | 2 | 1.2×10^{-10} | | Mallard et al. (1993) |
| Hx 3 | $\text{Br} + \text{OCIO} \rightarrow \text{BrO} + \text{ClO}$ | 2 | 2.6×10^{-11} | -1300 | Atkinson et al. (2006) |
| Hx 4 | $\text{Br} + \text{Cl}_2 \rightarrow \text{BrCl} + \text{Cl}$ | 2 | 1.1×10^{-15} | | Mallard et al. (1993) |
| Hx 5 | $\text{Br} + \text{BrCl} \rightarrow \text{Br}_2 + \text{Cl}$ | 2 | 3.3×10^{-15} | | Mallard et al. (1993) |
| Hx 6 | $\text{BrO} + \text{ClO} \rightarrow \text{Br} + \text{OCIO}$ | 2 | 1.6×10^{-12} | 430 | Atkinson et al. (2006) |
| Hx 7 | $\text{BrO} + \text{ClO} \rightarrow \text{Br} + \text{Cl} + \text{O}_2$ | 2 | 2.9×10^{-12} | 220 | Atkinson et al. (2006) |
| Hx 8 | $\text{BrO} + \text{ClO} \rightarrow \text{BrCl} + \text{O}_2$ | 2 | 5.8×10^{-13} | 170 | Atkinson et al. (2006) |
| Hx 9 | $\text{BrCl} + h\nu \rightarrow \text{Br} + \text{Cl}$ | 1 | 1 | | DeMore et al. (1997) |

n is the order of the reaction. ¹ photolysis rates calculated online, ² special rate functions (pressure dependent and/or humidity dependent). Notes: The rates for ROOH were assumed as that of CH_3OOH ; C_2H_4 is used as generic alkene as in the Lurmann et al. (1986) mechanism. The rate coefficients are calculated with $k = A \times \exp(-\frac{E_a}{RT})$.

Table 3: Aqueous phase reactions.

| no | reaction | n | k_0 [(M ¹⁻ⁿ)s ⁻¹] | $-E_a / R$ [K] | reference |
|------|--|---|---|----------------|--|
| O 1 | $O_3 + OH \rightarrow HO_2$ | 2 | 1.1×10^8 | | Sehested et al. (1984) |
| O 2 | $O_3 + O_2^- \rightarrow OH + OH^-$ | 2 | 1.5×10^9 | | Sehested et al. (1983) |
| O 3 | $OH + OH \rightarrow H_2O_2$ | 2 | 5.5×10^9 | | Buxton et al. (1988) |
| O 4 | $OH + HO_2 \rightarrow H_2O$ | 2 | 7.1×10^9 | | Sehested et al. (1968) |
| O 5 | $OH + O_2^- \rightarrow OH^-$ | 2 | 1.0×10^{10} | | Sehested et al. (1968) |
| O 6 | $OH + H_2O_2 \rightarrow HO_2$ | 2 | 2.7×10^7 | -1684 | Christensen et al. (1982) |
| O 7 | $HO_2 + HO_2 \rightarrow H_2O_2$ | 2 | 9.7×10^5 | -2500 | Christensen and Sehested (1988) |
| O 8 | $HO_2 + O_2^- \xrightarrow{H^+} H_2O_2$ | 2 | 1.0×10^8 | -900 | Christensen and Sehested (1988) |
| N 1 | $HONO + OH \rightarrow NO_2$ | 2 | 1.0×10^{10} | | assumed =N7 Barker et al. (1970) |
| N 2 | $HONO + H_2O_2 \xrightarrow{H^+} HNO_3$ | 3 | 4.6×10^3 | -6800 | Damschen and Martin (1983) |
| N 3 | $NO_3 + OH^- \rightarrow NO_3^- + OH$ | 2 | 8.2×10^7 | -2700 | Exner et al. (1992) |
| N 4 | $NO_2 + NO_2 \rightarrow HNO_3 + HONO$ | 2 | 1.0×10^8 | | Lee and Schwartz (1981) |
| N 5 | $NO_2 + HO_2 \rightarrow HNO_4$ | 2 | 1.8×10^9 | | Warneck (1999) |
| N 6 | $NO_2^- + O_3 \rightarrow NO_3^- + O_2$ | 2 | 5.0×10^5 | -6950 | Damschen and Martin (1983) |
| N 7 | $NO_2^- + OH \rightarrow NO_2 + OH^-$ | 2 | 1.0×10^{10} | | Barker et al. (1970) |
| N 8 | $NO_4^- \rightarrow NO_2^- + O_2$ | 1 | 8.0×10^{-1} | | Warneck (1999) |
| C 1 | $HCHO + OH \rightarrow HCOOH + HO_2$ | 2 | 7.7×10^8 | -1020 | Chin and Wine (1994) |
| C 2 | $HCOOH + OH \rightarrow HO_2 + CO_2$ | 2 | 1.1×10^8 | -991 | Chin and Wine (1994) |
| C 3 | $HCOO^- + OH \rightarrow OH^- + HO_2 + CO_2$ | 2 | 3.1×10^9 | -1240 | Chin and Wine (1994) |
| C 4 | $CH_3OO + HO_2 \rightarrow CH_3OOH$ | 2 | 4.3×10^5 | | estimated by Jacob (1986) |
| C 5 | $CH_3OO + O_2^- \rightarrow CH_3OOH + OH^-$ | 2 | 5.0×10^7 | | estimated by Jacob (1986) |
| C 6 | $CH_3OH + OH \rightarrow HCHO + HO_2$ | 2 | 9.7×10^8 | | Buxton et al. (1988) |
| C 7 | $CH_3OOH + OH \rightarrow CH_3OO$ | 2 | 2.7×10^7 | -1715 | estimated by Jacob (1986) |
| C 8 | $CH_3OOH + OH \rightarrow HCHO + OH$ | 2 | 1.1×10^7 | -1715 | estimated by Jacob (1986) |
| C 9 | $CO_3^- + O_2^- \rightarrow HCO_3^- + OH^-$ | 2 | 6.5×10^8 | | Ross et al. (1992) |
| C 10 | $CO_3^- + H_2O_2 \rightarrow HCO_3^- + HO_2$ | 2 | 4.3×10^5 | | Ross et al. (1992) |
| C 11 | $CO_3^- + HCOO^- \rightarrow HCO_3^- + HCO_3^- + HO_2$ | 2 | 1.5×10^5 | | Ross et al. (1992) |
| C 12 | $HCO_3^- + OH \rightarrow CO_3^-$ | 2 | 8.5×10^6 | | Ross et al. (1992) |
| C 13 | $DOM + OH \rightarrow HO_2$ | 2 | 5.0×10^9 | | estimated by (C. Anastasio, pers. comm.) from Ross et al. (1998) |

Table 3: Continued.

| no | reaction | n | k_0 [$(M^{1-n})s^{-1}$] | $-E_a / R$ [K] | reference |
|------|---|---|---|----------------|---|
| S 1 | $SO_3^- + O_2 \rightarrow SO_5^-$ | 2 | 1.5×10^9 | | Huie and Neta (1987) |
| S 2 | $HSO_3^- + O_3 \rightarrow SO_4^{2-} + H^+ + O_2$ | 2 | 3.7×10^5 | -5500 | Hoffmann (1986) |
| S 3 | $SO_3^{2-} + O_3 \rightarrow SO_4^{2-} + O_2$ | 2 | 1.5×10^9 | -5300 | Hoffmann (1986) |
| S 4 | $HSO_3^- + OH \rightarrow SO_3^-$ | 2 | 4.5×10^9 | | Buxton et al. (1988) |
| S 5 | $SO_3^{2-} + OH \rightarrow SO_3^- + OH^-$ | 2 | 5.5×10^9 | | Buxton et al. (1988) |
| S 6 | $HSO_3^- + HO_2 \rightarrow SO_4^{2-} + OH + H^+$ | 2 | 3.0×10^3 | | upper limit D. Sedlak pers. comm. with R. Sander |
| S 7 | $HSO_3^- + O_2^- \rightarrow SO_4^{2-} + OH$ | 2 | 3.0×10^3 | | upper limit D. Sedlak pers. comm. with R. Sander |
| S 8 | $HSO_3^- + H_2O_2 \rightarrow SO_4^{2-} + H^+$ | 2 | $5.2 \times 10^6 \times \frac{[H^+]}{[H^+] + 0.1M}$ | -3650 | Damschen and Martin (1983) |
| S 9 | $HSO_3^- + NO_2 \xrightarrow{NO_2} HSO_4^- + HONO + HONO$ | 2 | 2.0×10^7 | | Clifton et al. (1988) |
| S 10 | $SO_3^{2-} + NO_2 \xrightarrow{NO_2} SO_4^{2-} + HONO + HONO$ | 2 | 2.0×10^7 | | Clifton et al. (1988) |
| S 11 | $HSO_3^- + NO_3 \rightarrow SO_3^- + NO_3^- + H^+$ | 2 | 1.4×10^9 | -2000 | Exner et al. (1992) |
| S 12 | $HSO_3^- + HNO_4 \rightarrow HSO_4^- + NO_3^- + H^+$ | 2 | 3.1×10^5 | | Warneck (1999) |
| S 13 | $HSO_3^- + CH_3OOH \xrightarrow{H^+} SO_4^{2-} + H^+ + CH_3OH$ | 3 | 1.6×10^7 | -3800 | Lind et al. (1987) |
| S 14 | $SO_3^{2-} + CH_3OOH \xrightarrow{H^+} SO_4^{2-} + CH_3OH$ | 3 | 1.6×10^7 | -3800 | Lind et al. (1987) |
| S 15 | $HSO_3^- + HCHO \rightarrow CH_2OHSO_3^-$ | 2 | 4.3×10^{-1} | | Boyce and Hoffmann (1984) |
| S 16 | $SO_3^{2-} + HCHO \xrightarrow{H^+} CH_2OHSO_3^-$ | 2 | 1.4×10^4 | | Boyce and Hoffmann (1984) |
| S 17 | $CH_2OHSO_3^- + OH^- \rightarrow SO_3^{2-} + HCHO$ | 2 | 3.6×10^3 | | Seinfeld and Pandis (1998) |
| S 18 | $HSO_3^- + HSO_5^- \xrightarrow{H^+} SO_4^{2-} + SO_4^{2-} + H^+ + H^+$ | 3 | 7.1×10^6 | | Betterton and Hoffmann (1988) |
| S 19 | $SO_4^- + OH \rightarrow HSO_5^-$ | 2 | 1.0×10^9 | | Jiang et al. (1992) |

Table 3: Continued.

| no | reaction | n | k_0 [$(M^{1-n})s^{-1}$] | $-E_a / R$ [K] | reference |
|------|---|---|-----------------------------|----------------|--|
| S 20 | $SO_4^- + HO_2 \rightarrow SO_4^{2-} + H^+$ | 2 | 3.5×10^9 | | Jiang et al. (1992) |
| S 21 | $SO_4^- + O_2^- \rightarrow SO_4^{2-}$ | 2 | 3.5×10^9 | | assumed =S20 |
| S 22 | $SO_4^- + H_2O \rightarrow SO_4^{2-} + H^+ + OH$ | 2 | 1.1×10^{11} | -1110 | Herrmann et al. (1995) |
| S 23 | $SO_4^- + H_2O_2 \rightarrow SO_4^{2-} + H^+ + HO_2$ | 2 | 1.2×10^7 | | Wine et al. (1989) |
| S 24 | $SO_4^- + NO_3^- \rightarrow SO_4^{2-} + NO_3$ | 2 | 5.0×10^4 | | Exner et al. (1992) |
| S 25 | $SO_4^- + HSO_3^- \rightarrow SO_3^- + SO_4^{2-} + H^+$ | 2 | 8.0×10^8 | | Huie and Neta (1987) |
| S 26 | $SO_4^- + SO_3^{2-} \rightarrow SO_3^- + SO_4^{2-}$ | 2 | 4.6×10^8 | | Huie and Neta (1987) |
| S 27 | $SO_4^{2-} + NO_3 \rightarrow NO_3^- + SO_4^-$ | 2 | 1.0×10^5 | | Logager et al. (1993) |
| S 28 | $SO_5^- + HSO_3^- \rightarrow SO_4^- + SO_4^{2-} + H^+$ | 2 | 7.5×10^4 | | Huie and Neta (1987) |
| S 29 | $SO_5^- + SO_3^{2-} \rightarrow SO_4^- + SO_4^{2-}$ | 2 | 9.4×10^6 | | Huie and Neta (1987) |
| S 30 | $SO_5^- + HSO_3^- \rightarrow SO_3^- + HSO_5^-$ | 2 | 2.5×10^4 | | Huie and Neta (1987); Deister and Warneck (1990) |
| S 31 | $SO_5^- + SO_3^{2-} \xrightarrow{H^+} SO_3^- + HSO_5^-$ | 2 | 3.6×10^6 | | Huie and Neta (1987); Deister and Warneck (1990) |
| S 32 | $SO_5^- + O_2^- \xrightarrow{H^+} HSO_5^- + O_2$ | 2 | 2.3×10^8 | | Buxton et al. (1996) |
| S 33 | $SO_5^- + SO_5^- \rightarrow H_2O$ | 2 | 1.0×10^8 | | Ross et al. (1992) |
| S 34 | $DMS + O_3 \rightarrow O_2 + DMSO$ | 2 | 8.6×10^8 | -2600 | Gershenzon et al. (2001) |
| S 35 | $DMS + OH \rightarrow 0.5 CH_3SO_3^- + 0.5 CH_3OO + 0.5 HSO_4^- + HCHO + H^+$ | 2 | 1.9×10^{10} | | Ross et al. (1998) |
| S 36 | $DMSO + OH \rightarrow CH_3SO_2^- + CH_3OO + H^+$ | 2 | 4.5×10^9 | | Bardouki et al. (2002) |
| S 37 | $CH_3SO_2^- + OH \rightarrow CH_3SO_3^- + H_2O - O_2$ | 2 | 1.2×10^{10} | | Bardouki et al. (2002) |
| S 38 | $CH_3SO_3^- + OH \rightarrow SO_4^{2-} + H^+ + CH_3OO$ | 2 | 1.2×10^7 | | Bonsang et al. (1991) |

Table 3: Continued.

| no | reaction | n | k_0 [(M ¹⁻ⁿ)s ⁻¹] | $-E_a / R$ [K] | reference |
|-------|--|---|---|----------------|--|
| Cl 1 | $\text{Cl} + \text{H}_2\text{O}_2 \rightarrow \text{HO}_2 + \text{Cl}^- + \text{H}^+$ | 2 | 2.0×10^9 | | Yu (2001) |
| Cl 2 | $\text{Cl} + \text{H}_2\text{O} \rightarrow \text{H}^+ + \text{ClOH}^-$ | 2 | 1.8×10^5 | | Yu (2001) |
| Cl 3 | $\text{Cl} + \text{NO}_3^- \rightarrow \text{NO}_3 + \text{Cl}^-$ | 2 | 1.0×10^8 | | Buxton et al. (1999b) |
| Cl 4 | $\text{Cl} + \text{DOM} \rightarrow \text{Cl}^- + \text{HO}_2$ | 2 | 5.0×10^9 | | estimated (C. Anastasio, pers. comm.) from Ross et al. (1998) |
| Cl 5 | $\text{Cl} + \text{SO}_4^{2-} \rightarrow \text{SO}_4^- + \text{Cl}^-$ | 2 | 2.1×10^8 | | Buxton et al. (1999a) |
| Cl 6 | $\text{Cl} + \text{Cl} \rightarrow \text{Cl}_2$ | 2 | 8.8×10^7 | | Wu et al. (1980) |
| Cl 7 | $\text{Cl}^- + \text{OH} \rightarrow \text{ClOH}^-$ | 2 | 4.2×10^9 | | Yu (2001) |
| Cl 8 | $\text{Cl}^- + \text{O}_3 \rightarrow \text{ClO}^- + \text{O}_2$ | 2 | 3.0×10^{-3} | | Hoigné et al. (1985) |
| Cl 9 | $\text{Cl}^- + \text{NO}_3 \rightarrow \text{NO}_3^- + \text{Cl}$ | 2 | 9.3×10^6 | -4330 | Exner et al. (1992) |
| Cl 10 | $\text{Cl}^- + \text{SO}_4^- \rightarrow \text{SO}_4^{2-} + \text{Cl}$ | 2 | 2.5×10^8 | | Buxton et al. (1999a) |
| Cl 11 | $\text{Cl}^- + \text{HSO}_5^- \rightarrow \text{HOCl} + \text{SO}_4^{2-}$ | 2 | 1.8×10^{-3} | -7352 | Fortnum et al. (1960) |
| Cl 12 | $\text{Cl}^- + \text{HOCl} + \text{H}^+ \rightarrow \text{Cl}_2$ | 3 | 2.2×10^4 | -3508 | Ayers et al. (1996) |
| Cl 13 | $\text{Cl}_2 \rightarrow \text{Cl}^- + \text{HOCl} + \text{H}^+$ | 1 | 2.2×10^1 | -8012 | Ayers et al. (1996) |
| Cl 14 | $\text{Cl}_2^- + \text{OH} \rightarrow \text{HOCl} + \text{Cl}^-$ | 2 | 1.0×10^9 | | Ross et al. (1998) |
| Cl 15 | $\text{Cl}_2^- + \text{OH}^- \rightarrow \text{Cl}^- + \text{Cl}^- + \text{OH}$ | 2 | 4.0×10^6 | | Jacobi (1996) |
| Cl 16 | $\text{Cl}_2^- + \text{HO}_2 \rightarrow \text{Cl}^- + \text{Cl}^- + \text{H}^+ + \text{O}_2$ | 2 | 3.1×10^9 | | Yu (2001) |
| Cl 17 | $\text{Cl}_2^- + \text{O}_2^- \rightarrow \text{Cl}^- + \text{Cl}^- + \text{O}_2$ | 2 | 6.0×10^9 | | Jacobi (1996) |
| Cl 18 | $\text{Cl}_2^- + \text{H}_2\text{O}_2 \rightarrow \text{Cl}^- + \text{Cl}^- + \text{H}^+ + \text{HO}_2$ | 2 | 7.0×10^5 | -3340 | Jacobi (1996) |
| Cl 19 | $\text{Cl}_2^- + \text{NO}_2^- \rightarrow \text{Cl}^- + \text{Cl}^- + \text{NO}_2$ | 2 | 6.0×10^7 | | Jacobi (1996) |
| Cl 20 | $\text{Cl}_2^- + \text{CH}_3\text{OOH} \rightarrow \text{Cl}^- + \text{Cl}^- + \text{H}^+ + \text{CH}_3\text{OO}$ | 2 | 7.0×10^5 | -3340 | Jacobi (1996) |
| Cl 21 | $\text{Cl}_2^- + \text{DOM} \rightarrow \text{Cl}^- + \text{Cl}^- + \text{HO}_2$ | 2 | 1.0×10^6 | | assumed by Jacobi (1996) estimated (C. Anastasio, pers. comm.) from Ross et al. (1998) |
| Cl 22 | $\text{Cl}_2^- + \text{HSO}_3^- \rightarrow \text{SO}_3^- + \text{Cl}^- + \text{Cl}^- + \text{H}^+$ | 2 | 4.7×10^8 | -1082 | Shoute et al. (1991) |
| Cl 23 | $\text{Cl}_2^- + \text{SO}_3^{2-} \rightarrow \text{SO}_3^- + \text{Cl}^- + \text{Cl}^-$ | 2 | 6.2×10^7 | | Jacobi et al. (1996) |
| Cl 24 | $\text{Cl}_2^- + \text{Cl}_2 \rightarrow \text{Cl}_2 + 2\text{Cl}^-$ | 2 | 6.2×10^9 | | Yu (2001) |
| Cl 25 | $\text{Cl}_2^- + \text{Cl}^- \rightarrow \text{Cl}^- + \text{Cl}_2$ | 2 | 2.7×10^9 | | Yu (2001) |
| Cl 26 | $\text{Cl}_2^- + \text{DMS} \rightarrow 0.5 \text{CH}_3\text{SO}_3^- + 0.5 \text{CH}_3\text{OO} + 0.5 \text{HSO}_4^- + \text{HCHO} + 2 \text{Cl}^- + 2 \text{H}^+$ | 2 | 3.0×10^9 | | rate from Ross et al. (1998) |
| Cl 27 | $\text{ClOH}^- \rightarrow \text{Cl}^- + \text{OH}$ | 1 | 6.0×10^9 | | Yu (2001) |
| Cl 28 | $\text{ClOH}^- + \text{H}^+ \rightarrow \text{Cl}$ | 2 | 4.0×10^{10} | | Yu (2001) |
| Cl 29 | $\text{HOCl} + \text{HO}_2 \rightarrow \text{Cl} + \text{O}_2$ | 2 | 7.5×10^6 | | assumed = Cl30 Long and Bielski (1980) |
| Cl 30 | $\text{HOCl} + \text{O}_2^- \rightarrow \text{Cl} + \text{OH}^- + \text{O}_2$ | 2 | 7.5×10^6 | | Long and Bielski (1980) |
| Cl 31 | $\text{HOCl} + \text{SO}_3^{2-} \rightarrow \text{Cl}^- + \text{HSO}_4^-$ | 2 | 7.6×10^8 | | Fogelman et al. (1989) |
| Cl 32 | $\text{HOCl} + \text{HSO}_3^- \rightarrow \text{Cl}^- + \text{HSO}_4^- + \text{H}^+$ | 2 | 7.6×10^8 | | assumed = Cl31 Fogelman et al. (1989) |
| Cl 33 | $\text{Cl}_2 + \text{HO}_2 \rightarrow \text{Cl}_2^- + \text{H}^+ + \text{O}_2$ | 2 | 1.0×10^9 | | Bjergbakke et al. (1981) |
| Cl 34 | $\text{Cl}_2 + \text{O}_2^- \rightarrow \text{Cl}_2^- + \text{O}_2$ | 2 | 1.0×10^9 | | assumed = Cl33 Bjergbakke et al. (1981) |
| Cl 35 | $\text{Cl}^- + \text{HNO}_4 \rightarrow \text{HOCl} + \text{NO}_3^-$ | 2 | 1.4×10^{-2} | | Régimbal and Mozurkewich (1997) |

Table 3: Continued.

| no | reaction | n | k_0 [(M ¹⁻ⁿ)s ⁻¹] | $-E_a / R$ [K] | reference |
|-------|--|---|---|----------------|--|
| Br 1 | $\text{Br} + \text{OH}^- \rightarrow \text{BrOH}^-$ | 2 | 1.3×10^{10} | | Zehavi and Rabani (1972) |
| Br 2 | $\text{Br} + \text{DOM} \rightarrow \text{Br}^- + \text{HO}_2$ | 2 | 2.0×10^8 | | estimated (C. Anastasio, pers. comm.) from Ross et al. (1998) |
| Br 3 | $\text{Br}^- + \text{OH} \rightarrow \text{BrOH}^-$ | 2 | 1.1×10^{10} | | Zehavi and Rabani (1972) |
| Br 4 | $\text{Br}^- + \text{O}_3 \rightarrow \text{BrO}^-$ | 2 | 2.1×10^2 | -4450 | Haag and Hoigné (1983) |
| Br 5 | $\text{Br}^- + \text{NO}_3 \rightarrow \text{Br} + \text{NO}_3^-$ | 2 | 3.8×10^9 | | Zellner et al. 1996 in Herrmann et al. (2000) |
| Br 6 | $\text{Br}^- + \text{SO}_4^- \rightarrow \text{Br} + \text{SO}_4^{2-}$ | 2 | 2.1×10^9 | | Jacobi (1996) |
| Br 7 | $\text{Br}^- + \text{HSO}_5^- \rightarrow \text{HOBr} + \text{SO}_4^{2-}$ | 2 | 1.0 | -5338 | Fortnum et al. (1960) |
| Br 8 | $\text{Br}^- + \text{HOBr} + \text{H}^+ \rightarrow \text{Br}_2$ | 3 | 1.6×10^{10} | | Liu and Margerum (2001) |
| Br 9 | $\text{Br}_2 \rightarrow \text{Br}^- + \text{HOBr} + \text{H}^+$ | 1 | 9.7×10^1 | 7457 | Liu and Margerum (2001) |
| Br 10 | $\text{Br}_2^- + \text{O}_2^- \rightarrow \text{Br}^- + \text{Br}^-$ | 2 | 1.7×10^8 | | Wagner and Strehlow (1987) |
| Br 11 | $\text{Br}_2^- + \text{HO}_2 \rightarrow \text{Br}_2 + \text{H}_2\text{O}_2 - \text{H}^+$ | 2 | 4.4×10^9 | | Matthew et al. (2003) |
| Br 12 | $\text{Br}_2^- + \text{H}_2\text{O}_2 \rightarrow \text{Br}^- + \text{Br}^- + \text{H}^+ + \text{HO}_2$ | 2 | 5.0×10^2 | | Chameides and Stelson (1992) |
| Br 13 | $\text{Br}_2^- + \text{Br}_2^- \rightarrow \text{Br}^- + \text{Br}^- + \text{Br}_2$ | 2 | 1.9×10^9 | | Ross et al. (1992) |
| Br 14 | $\text{Br}_2^- + \text{CH}_3\text{OOH} \rightarrow \text{Br}^- + \text{Br}^- + \text{H}^+ + \text{CH}_3\text{OO}$ | 2 | 1.0×10^5 | | assumed by Jacobi (1996) |
| Br 15 | $\text{Br}_2^- + \text{DOM} \rightarrow \text{Br}^- + \text{Br}^- + \text{HO}_2$ | 2 | 1.0×10^5 | | estimated (C. Anastasio, pers. comm.) from Ross et al. (1998) |
| Br 16 | $\text{Br}_2^- + \text{NO}_2^- \rightarrow \text{Br}^- + \text{Br}^- + \text{NO}_2$ | 2 | 1.7×10^7 | -1720 | Shoute et al. (1991) |
| Br 17 | $\text{Br}_2^- + \text{HSO}_3^- \rightarrow \text{Br}^- + \text{Br}^- + \text{H}^+ + \text{SO}_3^-$ | 2 | 6.3×10^7 | -782 | Shoute et al. (1991) |
| Br 18 | $\text{Br}_2^- + \text{SO}_3^{2-} \rightarrow \text{Br}^- + \text{Br}^- + \text{SO}_3^-$ | 2 | 2.2×10^8 | -650 | Shoute et al. (1991) |
| Br 19 | $\text{Br}_2^- + \text{DMS} \rightarrow 0.5 \text{CH}_3\text{SO}_3^- + 0.5 \text{CH}_3\text{OO} + 0.5 \text{HSO}_4^- + \text{HCHO} + 2 \text{H}^+$ | 2 | 3.2×10^9 | | rate from Ross et al. (1998) |
| Br 20 | $\text{BrOH}^- \rightarrow \text{Br}^- + \text{OH}$ | 1 | 3.3×10^7 | | Zehavi and Rabani (1972) |
| Br 21 | $\text{BrOH}^- \rightarrow \text{Br} + \text{OH}^-$ | 1 | 4.2×10^6 | | Zehavi and Rabani (1972) |
| Br 22 | $\text{BrOH}^- + \text{H}^+ \rightarrow \text{Br}$ | 2 | 4.4×10^{10} | | Zehavi and Rabani (1972) |
| Br 23 | $\text{BrOH}^- + \text{Br}^- \rightarrow \text{Br}_2^- + \text{OH}^-$ | 2 | 1.9×10^8 | | Zehavi and Rabani (1972) |
| Br 24 | $\text{BrO}^- + \text{SO}_3^{2-} \rightarrow \text{Br}^- + \text{SO}_4^{2-}$ | 2 | 1.0×10^8 | | Troy and Margerum (1991) |
| Br 25 | $\text{HOBr} + \text{HO}_2 \rightarrow \text{Br} + \text{O}_2$ | 2 | 1.0×10^9 | | Herrmann et al. (1999) |
| Br 26 | $\text{HOBr} + \text{O}_2^- \rightarrow \text{Br} + \text{OH}^- + \text{O}_2$ | 2 | 3.5×10^9 | | Schwarz and Bielski (1986) |
| Br 27 | $\text{HOBr} + \text{H}_2\text{O}_2 \rightarrow \text{Br}^- + \text{H}^+ + \text{O}_2$ | 2 | 1.2×10^6 | | von Gunten and Oliveras (1998) |
| Br 28 | $\text{HOBr} + \text{SO}_3^{2-} \rightarrow \text{Br}^- + \text{HSO}_4^-$ | 2 | 5.0×10^9 | | Troy and Margerum (1991) |
| Br 29 | $\text{HOBr} + \text{HSO}_3^- \rightarrow \text{Br}^- + \text{HSO}_4^- + \text{H}^+$ | 2 | 5.0×10^9 | | assumed = Br28 |
| Br 30 | $\text{Br}_2 + \text{HO}_2 \rightarrow \text{Br}_2^- + \text{H}^+ + \text{O}_2$ | 2 | 1.1×10^8 | | Ross et al. (1998) |
| Br 31 | $\text{Br}_2 + \text{O}_2^- \rightarrow \text{Br}_2^- + \text{O}_2$ | 2 | 5.6×10^9 | | Ross et al. (1998) |
| Br 32 | $\text{Br}^- + \text{HNO}_4 \rightarrow \text{HOBr} + \text{NO}_3^-$ | 2 | 5.4×10^{-1} | | Régimbal and Mozurkewich (1997) |
| Br 33 | $\text{Br}^- + \text{O}_3 + \text{H}^+ \rightarrow \text{HOBr} + \text{O}_2$ | 3 | 11.7 | | Haag and Hoigné (1983) |

Table 3: Continued.

| no | reaction | n | k_0 [(M ¹⁻ⁿ)s ⁻¹] | $-E_a / R$ [K] | reference |
|------|--|---|---|----------------|---|
| Hx 1 | Br ⁻ + HOCl + H ⁺ → BrCl | 3 | 1.3 x 10 ⁶ | | Liu and Margerum (2001) |
| Hx 2 | Cl ⁻ + HOBr + H ⁺ → BrCl | 3 | 2.3 x 10 ¹⁰ | | Liu and Margerum (2001) |
| Hx 3 | BrCl → Cl ⁻ + HOBr + H ⁺ | 1 | 3.0 x 10 ⁶ | | Liu and Margerum (2001) |
| Hx 4 | Br ⁻ + ClO ⁻ + H ⁺ → BrCl + OH ⁻ | 3 | 3.7 x 10 ¹⁰ | | Kumar and Margerum (1987) |
| Hx 5 | Cl ₂ + Br ⁻ → BrCl ₂ ⁻ | 2 | 7.7 x 10 ⁹ | | Liu and Margerum (2001) |
| Hx 6 | BrCl ₂ ⁻ → Cl ₂ + Br ⁻ | 1 | 1.83 x 10 ³ | | Liu and Margerum (2001) |
| hv 1 | O ₃ + hv → OH + OH + O ₂ | 1 | 1 | | assumed 2x gas phase |
| hv 2 | H ₂ O ₂ + hv → OH + OH | 1 | 1 | | assumed 2x gas phase |
| hv 3 | NO ₃ ⁻ + hv $\xrightarrow{H^+}$ NO ₂ + OH | 1 | 1 | | Zellner et al. (1990) |
| hv 4 | NO ₂ ⁻ + hv $\xrightarrow{H^+}$ NO + OH | 1 | 1 | | Zellner et al. (1990); Burley and Johnston (1992) |
| hv 5 | HOCl + hv → OH + Cl | 1 | 1 | | assumed 2x gas phase |
| hv 6 | Cl ₂ + hv → Cl + Cl | 1 | 1 | | assumed 2x gas phase |
| hv 7 | HOBr + hv → OH + Br | 1 | 1 | | assumed 2x gas phase |
| hv 8 | Br ₂ + hv → Br + Br | 1 | 1 | | assumed 2x gas phase |
| hv 9 | BrCl + hv → Cl + Br | 1 | 1 | | assumed 2x gas phase |

n is the order of the reaction. ¹ photolysis rates calculated online. The temperature dependence is $k = k_0 \times \exp(\frac{-E_a}{R}(\frac{1}{T} - \frac{1}{T_0}))$; $T_0 = 298$ K.

Table 4: Heterogeneous reactions.

| no | reaction | k | reference |
|-----|---|--|--|
| H 1 | $\text{N}_2\text{O}_5 \xrightarrow{\text{H}_2\text{O}} \text{HNO}_{3aq} + \text{HNO}_{3aq}$ | $\bar{k}_t(\text{N}_2\text{O}_5)w_{l,i}[\text{H}_2\text{O}]/\text{Het}_T$ | Behnke et al. (1994), Behnke et al. (1997) |
| H 2 | $\text{N}_2\text{O}_5 \xrightarrow{\text{Cl}^-} \text{ClNO}_2 + \text{NO}_3^-$ | $\bar{k}_t(\text{N}_2\text{O}_5)w_{l,i}f(\text{Cl}^-)[\text{Cl}^-]/\text{Het}_T$ | Behnke et al. (1994), Behnke et al. (1997) |
| H 3 | $\text{N}_2\text{O}_5 \xrightarrow{\text{Br}^-} \text{BrNO}_2 + \text{NO}_3^-$ | $\bar{k}_t(\text{N}_2\text{O}_5)w_{l,i}f(\text{Br}^-)[\text{Br}^-]/\text{Het}_T$ | Behnke et al. (1994), Behnke et al. (1997) |
| H 4 | $\text{ClNO}_3 \xrightarrow{\text{H}_2\text{O}} \text{HOCl}_{aq} + \text{HNO}_{3aq}$ | $\bar{k}_t(\text{ClNO}_3)w_{l,i}[\text{H}_2\text{O}]/\text{Het}_T$ | see note |
| H 5 | $\text{ClNO}_3 \xrightarrow{\text{Cl}^-} \text{Cl}_{2aq} + \text{NO}_3^-$ | $\bar{k}_t(\text{ClNO}_3)w_{l,i}f(\text{Cl}^-)[\text{Cl}^-]/\text{Het}_T$ | see note |
| H 6 | $\text{ClNO}_3 \xrightarrow{\text{Br}^-} \text{BrCl}_{aq} + \text{NO}_3^-$ | $\bar{k}_t(\text{ClNO}_3)w_{l,i}f(\text{Br}^-)[\text{Br}^-]/\text{Het}_T$ | see note |
| H 7 | $\text{BrNO}_3 \xrightarrow{\text{H}_2\text{O}} \text{HOBr}_{aq} + \text{HNO}_{3aq}$ | $\bar{k}_t(\text{BrNO}_3)w_{l,i}[\text{H}_2\text{O}]/\text{Het}_T$ | see note |
| H 8 | $\text{BrNO}_3 \xrightarrow{\text{Cl}^-} \text{BrCl}_{aq} + \text{NO}_3^-$ | $\bar{k}_t(\text{BrNO}_3)w_{l,i}f(\text{Cl}^-)[\text{Cl}^-]/\text{Het}_T$ | see note |
| H 9 | $\text{BrNO}_3 \xrightarrow{\text{Br}^-} \text{Br}_{2aq} + \text{NO}_3^-$ | $\bar{k}_t(\text{BrNO}_3)w_{l,i}f(\text{Br}^-)[\text{Br}^-]/\text{Het}_T$ | see note |

For a definition of \bar{k}_t and $w_{l,i}$ see von Glasow et al. (2002) or von Glasow (2000). $\text{Het}_T = [\text{H}_2\text{O} + f(\text{Cl}^-)[\text{Cl}^-] + f(\text{Br}^-)[\text{Br}^-]]$, with $f(\text{Cl}^-) = 5.0 \times 10^2$ and $f(\text{Br}^-) = 3.0 \times 10^5$. H4 - H9: the total rate is determined by \bar{k}_t , the distribution among the different reaction paths was assumed to be the same as for reactions H1 - H3.

Table 5: Aqueous phase equilibrium constants.

| no | reaction | m | n | K_0 [M^{n-m}] | $-\Delta H/R$ [K] | reference |
|-------|---|-----|-----|-----------------------|-------------------|-------------------------------------|
| EQ 1 | $\text{CO}_{2aq} \leftrightarrow \text{H}^+ + \text{HCO}_3^-$ | 1 | 2 | 4.3×10^{-7} | -913 | Chameides (1984) |
| EQ 2 | $\text{NH}_{3aq} \leftrightarrow \text{OH}^- + \text{NH}_4^+$ | 1 | 2 | 1.7×10^{-5} | -4325 | Chameides (1984) |
| EQ 3 | $\text{H}_2\text{O}_{aq} \leftrightarrow \text{H}^+ + \text{OH}^-$ | 1 | 2 | 1.0×10^{-14} | -6716 | Chameides (1984) |
| EQ 4 | $\text{HCOOH}_{aq} \leftrightarrow \text{H}^+ + \text{HCOO}^-$ | 1 | 2 | 1.8×10^{-4} | | Weast (1980) |
| EQ 5 | $\text{HSO}_3^- \leftrightarrow \text{H}^+ + \text{SO}_3^{2-}$ | 1 | 2 | 6.0×10^{-8} | 1120 | Chameides (1984) |
| EQ 6 | $\text{H}_2\text{SO}_{4aq} \leftrightarrow \text{H}^+ + \text{HSO}_4^-$ | 1 | 2 | 1.0×10^3 | | Seinfeld and Pandis (1998) |
| EQ 7 | $\text{HSO}_4^- \leftrightarrow \text{H}^+ + \text{SO}_4^{2-}$ | 1 | 2 | 1.2×10^{-2} | 1120 | Weast (1980) |
| EQ 8 | $\text{HO}_{2aq} \leftrightarrow \text{O}_2 + \text{H}^+$ | 1 | 2 | 1.6×10^{-5} | | Weinstein-Lloyd and Schwartz (1991) |
| EQ 9 | $\text{SO}_{2aq} \leftrightarrow \text{H}^+ + \text{HSO}_3^-$ | 1 | 2 | 1.7×10^{-2} | 2090 | Chameides (1984) |
| EQ 10 | $\text{Cl}_2^- \leftrightarrow \text{Cl}_{aq} + \text{Cl}^-$ | 1 | 2 | 5.2×10^{-6} | | Jayson et al. (1973) |
| EQ 11 | $\text{HOCl}_{aq} \leftrightarrow \text{H}^+ + \text{ClO}^-$ | 1 | 2 | 3.2×10^{-8} | | Lax (1969) |
| EQ 12 | $\text{HBr}_{aq} \leftrightarrow \text{H}^+ + \text{Br}^-$ | 1 | 2 | 1.0×10^9 | | Lax (1969) |
| EQ 13 | $\text{Br}_2^- \leftrightarrow \text{Br}_{aq} + \text{Br}^-$ | 1 | 2 | 9.1×10^{-6} | | Mamou et al. (1977) |
| EQ 14 | $\text{HOBr}_{aq} \leftrightarrow \text{H}^+ + \text{BrO}^-$ | 1 | 2 | 2.3×10^{-9} | -3091 | Kelley and Tartar (1956) |
| EQ 15 | $\text{BrCl}_{aq} + \text{Cl}^- \leftrightarrow \text{BrCl}_2^-$ | 2 | 1 | 3.8 | 1143 | Wang et al. (1994) |
| EQ 16 | $\text{BrCl}_{aq} + \text{Br}^- \leftrightarrow \text{Br}_2\text{Cl}^-$ | 2 | 1 | 1.8×10^4 | | Wang et al. (1994) |
| EQ 17 | $\text{Br}_{2aq} + \text{Cl}^- \leftrightarrow \text{Br}_2\text{Cl}^-$ | 2 | 1 | 1.3 | | Wang et al. (1994) |
| EQ 18 | $\text{HNO}_{3aq} \leftrightarrow \text{H}^+ + \text{NO}_3^-$ | 1 | 2 | 1.5×10^1 | | Davis and de Bruin (1964) |
| EQ 19 | $\text{HCl}_{aq} \leftrightarrow \text{H}^+ + \text{Cl}^-$ | 1 | 2 | 1.7×10^6 | | Marsh and McElroy (1985) |
| EQ 20 | $\text{HONO}_{aq} \leftrightarrow \text{H}^+ + \text{NO}_2^-$ | 1 | 2 | 5.1×10^{-4} | -1260 | Schwartz and White (1981) |
| EQ 21 | $\text{HNO}_{4aq} \leftrightarrow \text{NO}_4^- + \text{H}^+$ | 1 | 2 | 1.0×10^{-5} | 8700 | Warneck (1999) |

The temperature dependence is $K = K_0 \times \exp\left(\frac{-\Delta H}{R} \left(\frac{1}{T} - \frac{1}{T_0}\right)\right)$, $T_0 = 298$ K.

Table 6: Henry constants and accommodation coefficients.

| specie | K_H^0 [M/atm] | $-\Delta_{soln}H/R$ [K] | reference | α^0 | $-\Delta_{obs}H/R$ [K] | reference |
|-----------------------------------|----------------------|-------------------------|----------------------------------|------------------|------------------------|------------------------------|
| O ₃ | 1.2×10^{-2} | 2560 | Chameides (1984) | 0.002 | (at 292 K) | DeMore et al. (1997) |
| O ₂ | 1.3×10^{-3} | 1500 | Wilhelm et al. (1977) | 0.01 | 2000 | estimated |
| OH | 3.0×10^1 | 4300 | Hanson et al. (1992) | 0.01 | (at 293 K) | Takami et al. (1998) |
| HO ₂ | 3.9×10^3 | 5900 | Hanson et al. (1992) | 0.2 | (at 293 K) | DeMore et al. (1997) |
| H ₂ O ₂ | 1.0×10^5 | 6338 | Lind and Kok (1994) | 0.077 | 2769 | Worsnop et al. (1989) |
| NO ₂ | 6.4×10^{-3} | 2500 | Lelieveld and Crutzen (1991) | 0.0015 | (at 298 K) | Ponche et al. (1993) |
| NO ₃ | 2.0 | 2000 | Thomas et al. (1993) | 0.04 | (at 273? K) | Rudich et al. (1996) |
| HONO | 4.9×10^1 | 4780 | Schwartz and White (1981) | 0.04 | (at 247-297 K) | DeMore et al. (1997) |
| HNO ₃ | 1.7×10^5 | 8694 | Lelieveld and Crutzen (1991) | 0.5 | (at RT) | Abbatt and Waschewsky (1998) |
| HNO ₄ | 1.2×10^4 | 6900 | Régimbald and Mozurkewich (1997) | 0.1 | (at 200 K) | DeMore et al. (1997) |
| NH ₃ | 5.8×10^1 | 4085 | Chameides (1984) | 0.06 | (at 295 K) | DeMore et al. (1997) |
| CH ₃ OO | 6.0 | =HO ₂ | Pandis and Seinfeld (1989) | 0.01 | 2000 | estimated |
| ROOH | 3.0×10^2 | 5322 | Lind and Kok (1994) | 0.0046 | 3273 | Magi et al. (1997) |
| HCHO | 7.0×10^3 | 6425 | Chameides (1984) | 0.04 | (at 260-270 K) | DeMore et al. (1997) |
| HCOOH | 3.7×10^3 | 5700 | Chameides (1984) | 0.014 | 3978 | DeMore et al. (1997) |
| CO ₂ | 3.1×10^{-2} | 2423 | Chameides (1984) | 0.01 | 2000 | estimated |
| HCl | 1.2 | 9001 | Brimblecombe and Clegg (1989) | 0.074 | 3072 | Schweitzer et al. (2000) |
| HOCl | 6.7×10^2 | 5862 | Huthwelker et al. (1995) | =HOBr | =HOBr | estimated |
| ClNO ₃ | ∞ | — | — | 0.1 | (at RT) | Koch and Rossi (1998) |
| Cl ₂ | 9.1×10^{-2} | 2500 | Wilhelm et al. (1977) | 0.038 | 6546 | Hu et al. (1995) |
| HBr | 1.3 | 10239 | Brimblecombe and Clegg (1989) | 0.031 | 3940 | Schweitzer et al. (2000) |
| HOBr | 9.3×10^1 | =HOCl | Vogt et al. (1996) | 0.5 | (at RT) | Wachsmuth et al. (2002) |
| BrNO ₃ | ∞ | — | — | 0.8 | 0 | Hanson et al. (1996) |
| Br ₂ | 7.6×10^{-1} | 4094 | Dean (1992) | 0.038 | 6546 | Hu et al. (1995) |
| BrCl | 9.4×10^{-1} | 5600 | Bartlett and Margerum (1999) | =Cl ₂ | =Cl ₂ | estimated |
| DMSO | 5.0×10^4 | =HCHO | De Bruyn et al. (1994) | 0.048 | 2578 | De Bruyn et al. (1994) |
| DMSO ₂ | ∞ | — | assumed | 0.03 | 5388 | DeMore et al. (1997) |
| SO ₂ | 1.2 | 3120 | Chameides (1984) | 0.11 | 0 | Pöschl et al. (1998) |
| H ₂ SO ₄ | ∞ | — | assumed | 0.65 | (at 303 K) | Lucas and Prinn (2002) |
| CH ₃ SO ₂ H | ∞ | — | assumed | 0.0002 | 0 | De Bruyn et al. (1994) |
| CH ₃ SO ₃ H | ∞ | — | assumed | 0.076 | 1762 | De Bruyn et al. (1994) |

For ROOH the values of CH₃OOH have been assumed. The temperature dependence is for the Henry constants is $K_H = K_H^0 \times \exp(-\frac{\Delta_{soln}H}{R}(\frac{1}{T} - \frac{1}{T_0}))$, $T_0 = 298$ K and for the accommodation coefficients $dl n(\frac{\alpha}{1-\alpha})/d(\frac{1}{T}) = -\frac{\Delta_{obs}H}{R}$. RT stands for “room temperature”.

References

- Abbatt, J. P. D. and Waschewsky, G. C. G.: Heterogeneous interactions of HOBr, HNO₃, O₃ and NO₂ with deliquescent NaCl aerosols at room temperature, *J. Phys. Chem. A*, 102, 3719 – 3725, 1998.
- Anderson, L. C. and Fahey, D. W.: Studies with ClONO₂: Thermal dissociation rate and catalytic conversion to NO using an NO/O₃ chemiluminescence detector, *J. Phys. Chem.*, 94, 644 – 652, 1990.
- Aranda, A., Le Bras, G., La Verdet, G., and Poulet, G.: The BrO + CH₃O₂ reaction: Kinetics and role in the atmospheric ozone budget, *Geophys. Res. Lett.*, 24, 2745 – 2748, 1997.
- Atkinson, R., Baulch, D. L., Cox, R. A., Hampson, Jr., R. F., Kerr, J. A., Rossi, M. J., and Troe, J.: Evaluated Kinetic and Photochemical Data for Atmospheric Chemistry: Supplement VI, *J. Phys. Chem. Ref. Data*, 26, 1329 – 1499, 1997.
- Atkinson, R., Baulch, D. L., Cox, R. A., Hampson, Jr., R. F., Kerr, J. A., Rossi, M. J., and Troe, J.: Summary of Evaluated Kinetic and Photochemical Data for Atmospheric Chemistry, Web Version, <http://www.iupac-kinetic.ch.cam.ac.uk>, 1999.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Jenkin, R. G. H. M. E., Kerr, J. A., Rossi, M. J., and Troe, J.: Summary of Evaluated Kinetic and Photochemical Data for Atmospheric Chemistry, Web Version, Jul. 2004, <http://www.iupac-kinetic.ch.cam.ac.uk>, 2004.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Kerr, J. A., Rossi, M. J., and Troe, J.: Summary of Evaluated Kinetic and Photochemical Data for Atmospheric Chemistry, Web Version, February 2006, <http://www.iupac-kinetic.ch.cam.ac.uk>, 2006.
- Ayers, G. P., Penkett, S. A., Gillett, R. W., Bandy, B., Galbally, I. E., Meyer, C. P., Elsworth, C. M., Bentle, S. T., and Forgan, B. W.: The Annual Cycle of Peroxides and Ozone in Marine Air at Cape Grim, Tasmania, *J. Atmos. Chem.*, 23, 221 – 252, 1996.
- Bardouki, H., da Rosa, M. B., Mihalopoulos, N., Palm, W.-U., and Zetzsch, C.: Kinetics and mechanism of the oxidation of dimethylsulfoxide (DMSO) and methanesulfinic acid (MSI⁻) in aqueous medium, *Atmos. Environ.*, 36, 4627 – 4634, 2002.
- Barker, G. C., Fowles, P., and Stringer, B.: Pulse radiolytic induced transient electrical conductance in liquid solutions, *Trans. Faraday Soc.*, 66, 1509–1519, 1970.
- Barnes, I., Bastian, V., Becker, K. H., and Overath, R. D.: Kinetic studies of the reactions of IO, BrO and ClO with DMS, *Int. J. Chem. Kinet.*, 23, 579 – 591, 1991.
- Barone, S. B., Turnipseed, A. A., and Ravishankara, A. R.: Role of adducts in the atmospheric oxidation of dimethyl sulfide, *Faraday Discuss.*, 100, 39 – 54, 1995.
- Bartlett, W. P. and Margerum, D. W.: Temperature dependencies of the Henry's law constant and the aqueous phase dissociation constant of bromine chloride, *Environ. Sci. Technol.*, 33, 3410–3414, 1999.
- Behnke, W., Scheer, V., and Zetzsch, C.: Production of BrNO₂, Br₂ and ClNO₂ from the Reaction between Sea Spray Aerosol and N₂O₅, *J. Aerosol Sci.*, 25, Suppl.1, S277 – S278, 1994.
- Behnke, W., George, C., Scheer, V., and Zetzsch, C.: Production and decay of ClNO₂ from the reaction of gaseous N₂O₅ with NaCl solution: Bulk and aerosol experiments, *J. Geophys. Res.*, 102, 3795 – 3804, 1997.

- Betterton, E. A. and Hoffmann, M. R.: Oxidation of aqueous SO₂ by peroxymonosulfate, *J. Phys. Chem.*, 92, 5962–5965, 1988.
- Bjergbakke, E., Navartnam, S., Parsons, B. J., and Swallow, A. J.: Reaction between HO₂· and chlorine in aqueous solution, *J. Am. Chem. Soc.*, 103, 5926–5928, 1981.
- Bonsang, B., Martin, D., Lambert, G., Kanakidou, M., de Roulley, J., and Sennequier, G.: Vertical distribution of nonmethane hydrocarbons in the remote marine boundary layer, *J. Geophys. Res.*, 96, 7313 – 7324, 1991.
- Boyce, S. D. and Hoffmann, M. R.: Kinetics and mechanism of the formation of hydroxymethanesulfonic acid at low pH, *J. Phys. Chem.*, 88, 4740–4746, 1984.
- Brimblecombe, P. and Clegg, S. L.: Erratum, *J. Atmos. Chem.*, 8, 95, 1989.
- Burley, J. D. and Johnston, H. S.: Ionic mechanisms for heterogeneous stratospheric reactions and ultraviolet photoabsorption cross-sections for NO₂⁺, HNO₃, and NO₃⁻ in sulfuric-acid, *Geophys. Res. Lett.*, 19, 1359 – 1362, 1992.
- Buxton, G. V., Greenstock, C. L., Helman, W. P., and Ross, A. B.: Critical review of rate constants for reactions of hydrated electrons, hydrogen atoms and hydroxyl radicals (·OH/·O⁻) in aqueous solution, *J. Phys. Chem. Ref. Data*, 17, 513–886, 1988.
- Buxton, G. V., McGowan, S., Salmon, G. S., Williams, J. E., and Wood, N. D.: A study of the spectra and reactivity of oxysulphur-radical anions involved in the chain oxidation of S(IV): A pulse and γ-radiolysis study, *Atmos. Environ.*, 30, 2483–2493, 1996.
- Buxton, G. V., Bydder, M., and Salmon, G. A.: The reactivity of chlorine atoms in aqueous solution Part II: The equilibrium SO₄⁻ + Cl⁻ ↔ Cl + SO₄²⁻, *Phys. Chem. Chem. Phys.*, 1, 269 – 273, 1999a.
- Buxton, G. V., Salmon, G. A., and Wang, J.: The equilibrium NO₃ + Cl⁻ ↔ NO₃⁻ + Cl: A laser flash photolysis and pulse radiolysis study of the reactivity of NO₃ with chloride ion in aqueous solution, *Phys. Chem. Chem. Phys.*, 1, 3589 – 3593, 1999b.
- Chameides, W. L.: The Photochemistry of a Remote Marine Stratiform Cloud, *J. Geophys. Res.*, 89, 4739 – 4755, 1984.
- Chameides, W. L. and Stelson, A. W.: Aqueous-Phase Chemical Processes in Deliquescent Sea-Salt Aerosols: A Mechanism That Couples the Atmospheric Cycles of S and Sea Salt, *J. Geophys. Res.*, 97, 20 565 – 20 580, 1992.
- Chin, M. and Wine, P. H.: A temperature-dependent competitive kinetics study of the aqueous-phase reactions of OH radicals with formate, formic acid, acetate, acetic acid, and hydrated formaldehyde, in *Aquatic and Surface Photochemistry*, edited by G. R. Helz, R. G. Zepp, and D. G. Crosby, pp. 85–96, A. F. Lewis, NY, 1994.
- Christensen, H. and Sehested, K.: HO₂ and O₂⁻ radicals at elevated temperatures, *J. Phys. Chem.*, 92, 3007–3011, 1988.
- Christensen, H., Sehested, K., and Corfitzen, H.: Reactions of hydroxyl radicals with hydrogen peroxide at ambient and elevated temperatures, *J. Phys. Chem.*, 86, 1588–1590, 1982.

- Clifton, C. L., Altstein, N., and Huie, R. E.: Rate constant for the reaction of NO_2 with sulfur(IV) over the pH range 5.3–13, *Environ. Sci. Technol.*, 22, 586–589, 1988.
- Damschen, D. E. and Martin, L. R.: Aqueous aerosol oxidation of nitrous acid by O_2 , O_3 and H_2O_2 , *Atmos. Environ.*, 17, 2005–2011, 1983.
- Davis, Jr., W. and de Bruin, H. J.: New activity coefficients of 0-100 per cent aqueous nitric acid, *J. Inorg. Nucl. Chem.*, 26, 1069 – 1083, 1964.
- De Bruyn, W. J., Shorter, J. A., Davidovits, P., Worsnop, D. R., Zahniser, M. S., and Kolb, C. E.: Uptake of gas phase sulfur species methanesulfonic acid, dimethylsulfoxide, and dimethyl sulfone by aqueous surface, *J. Geophys. Res.*, 99, 16 927 – 16 932, 1994.
- Dean, J. A.: *Lange's Handbook of Chemistry*, McGraw-Hill, Inc., 1992.
- Deister, U. and Warneck, P.: Photooxidation of SO_3^{2-} in aqueous solution, *J. Phys. Chem.*, 94, 2191–2198, 1990.
- DeMore, W. B., Sander, S. P., Golden, D. M., Hampson, R. F., Kurylo, M. J., Howard, C. J., Ravishankara, A. R., Kolb, C. E., and Molina, M. J.: *Chemical Kinetics and Photochemical Data for Use in Stratospheric Modeling*, Tech. Rep. JPL Publication 97-4, Jet Propulsion Laboratory, Pasadena, CA, 1997.
- Exner, M., Herrmann, H., and Zellner, R.: Laser-based studies of reactions of the nitrate radical in aqueous solution, *Ber. Bunsenges. Phys. Chem.*, 96, 470 – 477, 1992.
- Fogelman, K. D., Walker, D. M., and Margerum, D. W.: Non-metal redox kinetics: Hypochlorite and hypochlorous acid reactions with sulfite, *Inorg. Chem.*, 28, 986 – 993, 1989.
- Fortnum, D. H., Battaglia, C. J., Cohen, S. R., and Edwards, J. O.: The kinetics of the oxidation of halide ions by monosubstituted peroxides, *J. Am. Chem. Soc.*, 82, 778–782, 1960.
- Gershenson, M., Davidovits, P., Jayne, J. T., Kolb, C. E., and Worsnop, D. R.: Simultaneous Uptake of DMS and Ozone on Water, *J. Phys. Chem. A*, 105, 7031 – 7036, 2001.
- Haag, W. R. and Hoigné, J.: Ozonation of bromide-containing waters: Kinetics of formation of hypobromous acid and bromate, *Environ. Sci. Technol.*, 17, 261 – 267, 1983.
- Haggerstone, A.-L., Carpenter, L. J., Carslaw, N., and McFiggans, G.: Improved model predictions of HO_2 with gas to particle mass transfer rates calculated using aerosol number size distributions, *J. Geophys. Res.*, 110, D04 303, doi: 10.1029/2004JD005 282, 2005.
- Hansen, J. C., Li, Y., Francisco, J. S., and Li, Z.: On the Mechanism of the $\text{BrO} + \text{CH}_2\text{O}$ Reaction, *J. Phys. Chem. A*, 103, 8543 – 8546, 1999.
- Hanson, D. R., Burkholder, J. B., Howard, C. J., and Ravishankara, A. R.: Measurement of OH and HO_2 radical uptake coefficients on water and sulfuric acid surfaces, *J. Phys. Chem.*, 96, 4979–4985, 1992.
- Hanson, D. R., Ravishankara, A. R., and Lovejoy, E. R.: Reaction of BrONO_2 with H_2O on submicron sulfuric acid aerosol and the implications for the lower stratosphere, *J. Geophys. Res.*, 101D, 9063–9069, 1996.

- Herrmann, H., Reese, A., and Zellner, R.: Time resolved UV/VIS diode array absorption spectroscopy of SO_x^- ($x=3, 4, 5$) radical anions in aqueous solution, *J. Mol. Struct.*, 348, 183–186, 1995.
- Herrmann, H., Ervens, B., Nowacki, P., Wolke, R., and Zellner, R.: A chemical aqueous phase radical mechanism for tropospheric chemistry, *Chemosphere*, 38, 1223 – 1232, 1999.
- Herrmann, H., Ervens, B., Jacobi, H.-W., Wolke, R., Nowacki, P., and Zellner, R.: CAPRAM2.3: A Chemical Aqueous Phase Radical Mechanism for Tropospheric Chemistry, *J. Atmos. Chem.*, 36, 231 – 284, 2000.
- Hoffmann, M. R.: On the kinetics and mechanism of oxidation of aquated sulfur dioxide by ozone, *Atmos. Environ.*, 20, 1145 – 1154, 1986.
- Hoigné, J., Bader, H., Haag, W. R., and Staehelin, J.: Rate constants of reactions of ozone with organic and inorganic compounds in water — III Inorganic compounds and radicals, *Wat. Res.*, 19, 993–1004, 1985.
- Hu, J. H., Shi, Q., Davidovits, P., Worsnop, D. R., Zahniser, M. S., and Kolb, C. E.: Reactive uptake of Cl_2 and Br_2 by aqueous surfaces as a function of Br^- and I^- ion concentration: The effect of chemical reaction at the interface, *J. Phys. Chem.*, 99, 8768 – 8776, 1995.
- Hubinger, S. and Nee, J. B.: Absorption spectra of Cl_2 , Br_2 and BrCl between 190 and 600 nm, *J. Photochem. Photobiol. A: Chem.*, 86, 1–7, 1995.
- Huie, R. E. and Neta, P.: Rate constants for some oxidations of S(IV) by radicals in aqueous solutions, *Atmos. Environ.*, 21, 1743–1747, 1987.
- Huthwelker, T., Clegg, S. L., Peter, T., Carslaw, K., Luo, B. P., and Brimblecombe, P.: Solubility of HOCl in water and aqueous H_2SO_4 to stratospheric temperatures, *J. Atmos. Chem.*, 21, 81–95, 1995.
- Ingham, T., Bauer, D., Sander, R., Crutzen, P. J., and Crowley, J. N.: Kinetics and Products of the Reactions $\text{BrO} + \text{DMS}$ and $\text{Br} + \text{DMS}$ at 298 K, *J. Phys. Chem. A*, 103, 7199 – 7209, 1999.
- Jacob, D. J.: Chemistry of OH in Remote Clouds and Its Role in the Production of Formic Acid and Peroxymonosulfate, *J. Geophys. Res.*, 91, 9807 – 9826, 1986.
- Jacobi, H.-W.: Kinetische Untersuchungen und Modellrechnungen zur troposphärischen Chemie von Radikalanionen und Ozon in wässriger Phase, Ph.D. thesis, Universität Essen, Germany, 1996.
- Jacobi, H.-W., Herrmann, H., and Zellner, R.: Kinetic investigation of the Cl_2^- radical in the aqueous phase, in *Air Pollution Research Report 57: Homogenous and heterogenous chemical Processes in the Troposphere*, edited by P. Mirabel, pp. 172–176, Office for official Publications of the European Communities, Luxembourg, 1996.
- Jayne, J. T., Pöschl, U., Chen, Y., Dai, D., Molina, L. T., Worsnop, D. R., Kolb, C. E., and Molina, M. J.: Pressure and Temperature Dependence of the Gas-Phase Reaction of SO_3 with H_2O and the Heterogeneous Reaction of SO_3 with $\text{H}_2\text{O}/\text{H}_2\text{SO}_4$ Surfaces, *J. Phys. Chem. A*, 101, 10000 – 10011, 1997.
- Jayson, G. G., Parsons, B. J., and Swallow, A. J.: Some simple, highly reactive, inorganic chlorine derivatives in aqueous solution, *J. Chem. Soc. Faraday Trans.*, 69, 1597–1607, 1973.

- Jefferson, A., Nicovich, J. M., and Wine, P. H.: Temperature-dependent kinetics studies of the reactions $\text{Br}(^2\text{P}_{3/2}) + \text{CH}_3\text{SCH}_3 \leftrightarrow \text{CH}_3\text{SCH}_2 + \text{HBr}$. Heat of formation of the CH_3SCH_2 radical, *J. Phys. Chem.*, 98, 7128 – 7135, 1994.
- Jiang, P.-Y., Katsumura, Y., Nagaishi, R., Domae, M., Ishikawa, K., Ishigure, K., and Yoshida, Y.: Pulse radiolysis study of concentrated sulfuric acid solutions. Formation mechanism, yield and reactivity of sulfate radicals, *J. Chem. Soc. Faraday Trans.*, 88, 1653–1658, 1992.
- Kelley, C. M. and Tartar, H. V.: On the system: bromine-water, *J. Am. Chem. Soc.*, 78, 5752–5756, 1956.
- Koch, T. G. and Rossi, M. J.: Direct measurement of surface residence times: Nitryl chloride and chlorine nitrate on alkali halides at room temperature, *J. Phys. Chem. A*, 102, 9193–9201, 1998.
- Kukui, A., Bossoutrot, V., Laverdet, G., and Bras, G. L.: Mechanism of the Reaction of CH_3SO with NO_2 in Relation to Atmospheric Oxidation of Dimethyl Sulfide: Experimental and Theoretical Study, *J. Phys. Chem. A*, 104, 935 – 946, 2000.
- Kukui, A., Borissenko, D., Laverdet, G., and Bras, G. L.: Gas phase reactions of OH radicals with dimethyl sulfoxide and methane sulfonic acid using turbulent flow reactor and chemical ionization mass spectrometry, *J. Phys. Chem. A*, 107, 5732 – 5742, 2003.
- Kumar, K. and Margerum, D. W.: Kinetics and mechanism of general-acid-assisted oxidation of bromide by hypochlorite and hypochlorous acid, *Inorg. Chem.*, 26, 2706 – 2711, 1987.
- Lax, E.: Taschenbuch für Chemiker und Physiker, Springer Verlag, Berlin, 1969.
- Lee, Y.-N. and Schwartz, S. E.: Reaction kinetics of nitrogen dioxide with liquid water at low partial pressure, *J. Phys. Chem.*, 85, 840–848, 1981.
- Lelieveld, J. and Crutzen, P. J.: The Role of Clouds in Tropospheric Photochemistry, *J. Atmos. Chem.*, 12, 229 – 267, 1991.
- Lind, J. A. and Kok, G. L.: Correction to “Henry’s law determinations for aqueous solutions of hydrogen peroxide, methylhydroperoxide, and peroxyacetic acid” by John A. Lind and Gregory L. Kok, *J. Geophys. Res.*, 99D, 21 119, 1994.
- Lind, J. A., Lazrus, A. L., and Kok, G. L.: Aqueous phase oxidation of sulfur(IV) by hydrogen peroxide, methylhydroperoxide, and peroxyacetic acid, *J. Geophys. Res.*, 92D, 4171–4177, 1987.
- Liu, Q. and Margerum, D. W.: Equilibrium and Kinetics of Bromine Chloride Hydrolysis, *est*, 35, 1127 – 1133, 2001.
- Logager, T., Sehested, K., and Holcman, J.: Rate constants of the equilibrium reactions $\text{SO}_4 + \text{HNO}_3 \leftrightarrow \text{HSO}_4^- + \text{NO}_3$ and $\text{SO}_4 + \text{NO}_3 \leftrightarrow \text{SO}_4^{2-} + \text{NO}_3$., *Radiat. Phys. Chem.*, 41, 539 – 543, 1993.
- Long, C. A. and Bielski, B. H. J.: Rate of reaction of superoxide radical with chloride-containing species, *J. Phys. Chem.*, 84, 555–557, 1980.
- Lucas, D. D. and Prinn, R. G.: Mechanistic studies of dimethylsulfide oxidation products using an observationally constrained model, *J. Geophys. Res.*, 107, doi: 10.1029/2001JD000 843, 2002.

- Lurmann, F. W., Lloyd, A. C., and Atkinson, R.: A Chemical Mechanism for Use in Long-Range Transport/Acid Deposition Computer Modeling, *J. Geophys. Res.*, 91, 10 905 – 10 936, 1986.
- Magi, L., Schweitzer, F., Pallares, C., Cherif, S., Mirabel, P., and George, C.: Investigation of the uptake rate of ozone and methyl hydroperoxide by water surfaces, *J. Phys. Chem. A*, 101, 4943 – 4949, 1997.
- Mallard, W. G., Westley, F., Herron, J. T., Hampson, R. F., and Frizzel, D. H.: NIST Chemical Kinetics Database: Version 5.0, National Institute of Standards and Technology, Gaithersburg, MD, 1993.
- Mamou, A., Rabani, J., and Behar, D.: On the oxidation of aqueous Br^- by OH radicals, studied by pulse radiolysis, *J. Phys. Chem.*, 81, 1447–1448, 1977.
- Marsh, A. R. W. and McElroy, W. J.: The dissociation constant and Henry's law constant of HCl in aqueous solution, *Atmos. Environ.*, 19, 1075 – 1080, 1985.
- Matthew, B. M., George, I., and Anastasio, C.: Hydroperoxyl radical (HO_2) oxidizes dibromide radical anion (Br_2^-) to bromine (Br_2) in aqueous solution: Implications for the formation of Br_2 in the marine boundary layer, *Geophys. Res. Lett.*, 30 (24), 2297, doi: 10.1029/2003GL018 572, 2003.
- Mellouki, A., Bras, G. L., and Poulet, G.: Discharge Flow Kinetic Study of NO_3 Reactions with Free Radicals: The Reaction of NO_3 with Cl, *J. Phys. Chem.*, 91, 5760–5764, 1987.
- Muthuramu, K., Shepson, P. B., Bottenheim, J. W., Jobson, T., Niki, H., and Anlauf, K. G.: Relationships Between Organic Nitrates and Surface Ozone Destruction During Polar Sunrise Experiment 1992, *J. Geophys. Res.*, 99 (D12), 25 369–25 378, doi:10.1029/94JD01 309, 1994.
- Orlando, J. J. and Tyndall, G. S.: Rate coefficients for the thermal decomposition of BrONO_2 and the heat of formation of BrONO_2 , *J. Phys. Chem.*, 100, 19 398 – 19 405, 1996.
- Pandis, S. N. and Seinfeld, J. H.: Sensitivity Analysis of a Chemical Mechanism for Aqueous-Phase Atmospheric Chemistry, *J. Geophys. Res.*, 94, 1105 – 1126, 1989.
- Ponche, J. L., George, C., and Mirabel, P.: Mass transfer at the air/water interface: Mass accommodation coefficients of SO_2 , HNO_3 , NO_2 and NH_3 , *J. Atmos. Chem.*, 16, 1–21, 1993.
- Pöschl, U., Canagaratna, M., Jayne, J. T., Molina, L. T., Worsnop, D. R., Kolb, C. E., and Molina, M. J.: Mass accommodation coefficient of H_2SO_4 vapor on aqueous sulfuric acid surfaces and gaseous diffusion coefficient of H_2SO_4 in $\text{N}_2/\text{H}_2\text{O}$, *J. Phys. Chem. A*, 102, 10 082–10 089, 1998.
- Ray, A., Vassalli, I., G.Laverdet, and Bras, G. L.: Kinetics of the Thermal Decomposition of the CH_3SO_2 Radical and Its Reaction with NO_2 at 1 Torr and 298 K, *J. Phys. Chem.*, 100, 8895 – 8900, 1996.
- Régimbal, J.-M. and Mozurkewich, M.: Peroxynitric acid decay mechanisms and kinetics at low pH, *J. Phys. Chem. A*, 101, 8822–8829, 1997.
- Ross, A. B., Mallard, W. G., Helman, W. P., Bielski, B. H. J., Buxton, G. V., Cabelli, D. E., Greenstock, C. L., Huie, R. E., and Neta, P.: NDRL-NIST Solution Kinetics Database: - Ver. 1, National Institute of Standards and Technology, Gaithersburg, MD, 1992.
- Ross, A. B., Mallard, W. G., Helman, W. P., Buxton, G. V., Huie, R. E., and Neta, P.: NDRL-NIST Solution kinetics database: Version 3.0, Notre Dame Radiation Laboratory, Notre Dame, and National Institut of Standards and Technology, Gaithersburg, 1998.

- Rudich, Y., Talukdar, R. K., Imamura, T., Fox, R. W., and Ravishankara, A. R.: Uptake of NO_3 on KI solutions: Rate coefficient for the $\text{NO}_3 + \text{I}^-$ reaction and gas-phase diffusion coefficients for NO_3 , Chem. Phys. Lett., 261, 467–473, 1996.
- Sander, R., Vogt, R., Harris, G. W., and Crutzen, P. J.: Modeling the chemistry of ozone, halogen compounds, and hydrocarbons in the arctic troposphere during spring, Tellus, 49B (5), 522–532, 1997.
- Sander, S. P., Friedl, R. R., Golden, D. M., Kurylo, M. J., Huie, R. E., Orkin, V. L., Moortgat, G. K., Ravishankara, A. R., Kolb, C. E., Molina, M. J., and Finlayson-Pitts, B. J.: Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies, Tech. Rep. JPL Publication 02-25, Jet Propulsion Laboratory, Pasadena, CA, 2003.
- Scheffler, D., Grothe, H., Willner, H., Frenzel, A., and Zetzsch, C.: Properties of Pure Nitryl Bromide. Thermal Behaviour, UV/Vis and FTIR Spectra, and Photoisomerization to *trans*-BrONO in an Argon Matrix, Inorg. Chem., 36, 335 – 338, 1997.
- Schwartz, S. E. and White, W. H.: Solubility equilibria of the nitrogen oxides and oxyacids in dilute aqueous solution, in Advances in Environmental Science and Engineering, edited by J. R. Pfafflin and E. N. Ziegler, vol. 4, pp. 1–45, Gordon and Breach Science Publishers, NY, 1981.
- Schwarz, H. A. and Bielski, B. H. J.: Reactions of HO_2 and O_2^- with iodine and bromine and the I_2^- and I atom reduction potentials, J. Phys. Chem., 90, 1445–1448, 1986.
- Schweitzer, F., Mirabel, P., and George, C.: Uptake of hydrogen halides by water droplets, J. Phys. Chem. A, 104, 72–76, 2000.
- Sehested, K., Rasmussen, O. L., and Fricke, H.: Rate constants of OH with HO_2 , O_2^- , and H_2O_2^+ from hydrogen peroxide formation in pulse-irradiated oxygenated water, J. Phys. Chem., 72, 626–631, 1968.
- Sehested, K., Holcman, J., and Hart, E. J.: Rate constants and products of the reactions of e_{aq}^- , O_2^- and H with ozone in aqueous solutions, J. Phys. Chem., 87, 1951–1954, 1983.
- Sehested, K., Holcman, J., Bjergbakke, E., and Hart, E. J.: A pulse radiolytic study of the reaction $\text{OH} + \text{O}_3$ in aqueous medium, J. Phys. Chem., 88, 4144–4147, 1984.
- Seinfeld, J. H. and Pandis, S. N.: Atmospheric Chemistry and Physics, John Wiley & Sons, New York, Chichester, Weinheim, 1998.
- Shoute, L. C. T., Alfassi, Z. B., Neta, P., and Huie, R. E.: Temperature dependence of the rate constants for reaction of dihalide and azide radicals with inorganic reductants, J. Phys. Chem., 95, 3238 – 3242, 1991.
- Takami, A., Kato, S., Shimono, A., and Koda, S.: Uptake coefficient of OH radical on aqueous surface, Chem. Phys., 231, 215–227, 1998.
- THALOS: Final report of the EU project THALOS: Tropospheric Halogens - effect on ozone, coordinated by R. A. Cox, University of Cambridge, U.K., 2005.
- Thomas, K., Volz-Thomas, A., and Kley, D.: Zur Wechselwirkung von NO_3 -Radikalen mit wässrigen Lösungen: Bestimmung des Henry- und des Massenakkommodationskoeffizienten, Ph.D. thesis, Institut für Chemie und Dynamik der Geosphäre 2, Forschungszentrum Jülich GmbH, FRG, 1993.

- Troy, R. C. and Margerum, D. W.: Non-metal redox kinetics: Hypobromite and hypobromous acid reactions with iodide and with sulfite and the hydrolysis of bromosulfate, *Inorg. Chem.*, 30, 3538 – 3543, 1991.
- Tsalkani, N., Mellouki, A., Poulet, G., Toupance, G., and Bras, G. L.: Rate constant measurement for the reactions of OH and Cl with peroxyacetyl nitrate at 298 K, *J. Atmos. Chem.*, 7, 409–419, 1988.
- Urbanski, S., Stickel, R. E., Zhao, Z. Z., and Wine, P. H.: Mechanistic and kinetic study of formaldehyde production in the atmospheric oxidation of dimethyl sulfide, *J. Chem. Soc. Faraday Trans.*, 93, 2813 – 2819, 1997.
- Urbanski, S., Stickel, R. E., and Wine, P. H.: Mechanistic and kinetic study of the gas-phase reaction of hydroxyl radical with dimethyl sulfoxide, *J. Phys. Chem. A*, 102, 10 522 – 10 529, 1998.
- van Dingenen, R., Jensen, N. R., Hjorth, J., and Raes, F.: Peroxynitrate Formation During the Night-Time Oxidation of Dimethylsulfide: Its Role as a Reservoir Species for Aerosol Formation, *J. Atmos. Chem.*, 18, 211 – 237, 1994.
- Vogt, R., Crutzen, P. J., and Sander, R.: A mechanism for halogen release from sea-salt aerosol in the remote marine boundary layer, *Nature*, 383, 327 – 330, 1996.
- von Glasow, R.: Modeling the gas and aqueous phase chemistry of the marine boundary layer, Ph.D. thesis, Universität Mainz, Germany, <http://www.rolandvonglasow.de>, 2000.
- von Glasow, R., Sander, R., Bott, A., and Crutzen, P. J.: Modeling halogen chemistry in the marine boundary layer. 1. Cloud-free MBL, *J. Geophys. Res.*, 107, 4341, doi: 10.1029/2001JD000 942, 2002.
- von Gunten, U. and Oliveras, Y.: Advanced oxidation of bromide-containing waters: Bromate formation mechanisms, *Environ. Sci. Technol.*, 32, 63 – 70, 1998.
- Wachsmuth, M., Gäggeler, H. W., von Glasow, R., and Ammann, M.: Accommodation coefficient of HOBr on deliquescent sodium bromide aerosol particles, *Atmos. Chem. Phys.*, 2, 121 – 131, 2002.
- Wagner, I. and Strehlow, H.: On the flash photolysis of bromide ions in aqueous solution, *Ber. Bunsenges. Phys. Chem.*, 91, 1317 – 1321, 1987.
- Wallington, T. J., Andino, J. M., Ball, J. C., and Japar, S. M.: Fourier transform infrared studies of the reaction of Cl atoms with PAN, PPN, CH₃OOH, HCOOH, CH₃COCH₃ and CH₃COC₂H₅ at 295±2 K, *J. Atmos. Chem.*, 10, 301 – 313, 1990.
- Wang, T. X., Kelley, M. D., Cooper, J. N., Beckwith, R. C., and Margerum, D. W.: Equilibrium, kinetic, and UV-spectral characteristics of aqueous bromine chloride, bromine, and chlorine species, *Inorg. Chem.*, 33, 5872 – 5878, 1994.
- Warneck, P.: The relative importance of various pathways for the oxidation of sulfur dioxide and nitrogen dioxide in sunlit continental fair weather clouds, *Phys. Chem. Chem. Phys.*, 1, 5471 – 5483, 1999.
- Wayne, R. P., Barnes, I., Biggs, P., Burrows, J. P., Canosa-Mas, C. E., Hjorth, J., Le Bras, G., Moortgat, G. K., Perner, D., Poulet, G., Restelli, G., and Sidebottom, H.: The nitrate radical: Physics, chemistry, and the atmosphere, *Atmos. Environ.*, 25A, 1–203, 1991.
- Weast, R. C., ed.: *CRC Handbook of Chemistry and Physics*, 61st Edition, CRC Press, Inc., Boca Raton, FL, 1980.

- Weinstein-Lloyd, J. and Schwartz, S. E.: Low-intensity radiolysis study of free-radical reactions in cloud-water: H_2O_2 production and destruction, *Environ. Sci. Technol.*, 25, 791–800, 1991.
- Wilhelm, E., Battino, R., and Wilcock, R. J.: Low-pressure solubility of gases in liquid water, *Chem. Rev.*, 77, 219–262, 1977.
- Wine, P. H., Wells, J. R., and Nicovich, J. M.: Kinetics of the reactions of F^2P and Cl^2P with HNO_3 , *J. Chem. Phys.*, 92, 2223–2228, 1988.
- Wine, P. H., Tang, Y., Thorn, R. P., Wells, J. R., and Davis, D. D.: Kinetics of aqueous phase reactions of the SO_4^- radical with potential importance in cloud chemistry, *J. Geophys. Res.*, 94D, 1085–1094, 1989.
- Worsnop, D. R., Zahniser, M. S., Kolb, C. E., Gardner, J. A., Watson, L. R., van Doren, J. M., Jayne, J. T., and Davidovits, P.: The temperature dependence of mass accommodation of SO_2 and H_2O_2 on aqueous surfaces, *J. Phys. Chem.*, 93, 1159–1172, 1989.
- Wu, D., Wong, D., and Di Bartolo, B.: Evolution of Cl_2^- in aqueous NaCl solutions, *J. Photochem.*, 14, 303–310, 1980.
- Yin, F., Grosjean, D., and Seinfeld, J. H.: Photooxidation of Dimethyl Sulfide and Dimethyl Disulfide. I: Mechanism Development, *J. Atmos. Chem.*, 11, 309 – 364, 1990.
- Yu, X.-Y.: Kinetics of free radical reactions generated by laser flash photolysis of $\text{OH} + \text{Cl}^-$ and $\text{SO}_4^- + \text{Cl}^-$ in the aqueous phase – Chemical mechanism, kinetics data and their implications, Ph.D. thesis, University of Michigan, Ann Arbor, 2001.
- Zehavi, D. and Rabani, J.: Oxidation of aqueous bromide ions by hydroxyl radicals. Pulse radiolytic investigation, *J. Phys. Chem.*, 76, 312 – 319, 1972.
- Zellner, R., Exner, M., and Herrmann, H.: Absolute OH Quantum Yield in the Laser Photolysis of Nitrate, Nitrite and Dissolved H_2O_2 at 308 and 351 nm in the Temperature Range 278–353 K, *J. Atmos. Chem.*, 10, 411 – 425, 1990.