

Supplementary Material For: Loading-Dependent Elemental Composition of α -Pinene SOA Particles

by

John E. Shilling,^{1,†} Qi Chen,¹ Stephanie M. King,¹ Thomas Rosenoern,¹ Jesse H. Kroll,²
Douglas R. Worsnop,² Peter F. DeCarlo,^{3,4,‡} Allison C. Aiken,^{4,5} Donna Sueper,^{2,4,5}
Jose L. Jimenez,^{4,5} and Scot T. Martin^{1,6*}

¹ *School of Engineering and Applied Sciences, Harvard University, Cambridge, MA
02138*

² *Aerodyne Research, Inc., Billerica, MA 08121*

³ *Department of Atmospheric and Oceanic Sciences University of Colorado, Boulder, CO
80309*

⁴ *Cooperative Institute for Research in the Environmental Sciences (CIRES), University of
Colorado, Boulder, CO 80309*

⁵ *Department of Chemistry and Biochemistry, University of Colorado, Boulder, CO 80309*

⁶ *Department of Earth and Planetary Sciences, Harvard University, Cambridge, MA
02138*

[†] *Now at Pacific Northwest National Laboratory, Atmospheric Sciences and Global
Change Division, Richland, WA 99352*

[‡] *Now at the Paul Scherrer Institut, Laboratory of Atmospheric Chemistry, Villigen-PSI,
CH-5232*

E-mail: scot_martin@harvard.edu

<http://www.seas.harvard.edu/environmental-chemistry>

***To Whom Correspondence Should be Addressed**

Table S1. SOA particle mass loadings and the measured O/C and H/C atomic ratios used in the basis-set fitting (i.e., entries of Table 2).

Figure S1. Correlation plots between unit-mass-resolution signal intensity and high-resolution oxygen-to-carbon atomic ratio. Except for the bottom-left panel, vertical axes show the percent contribution of the signal intensity of the indicated m/z value to the total organic signal. For the bottom-left panel, the vertical axis shows the ratio of the signal intensity at m/z 44 to that at m/z 43.

Figure S2. Measured oxygen-to-carbon atomic ratio vs. the 44/org signal for SOA particles produced from the dark ozonolysis of α -pinene. The linear regression through the data is shown in the solid line. The data points and the regression of *Aiken et al.* (2008) for ambient ground and aircraft measurements during the MILAGRO campaign are also shown. Equations for the linear fits and correlation coefficients are given in the figure.

Figure S3. Example of the deconvolution of a high-resolution mass spectrum for fragments at m/z 43. Peaks representing high-resolution model fits of $C_2H_3O^+$ and $C_3H_7^+$ are shown in the figure (DeCarlo *et al.*, 2006). Panel i shows the residual between the recorded data and model fits for panel ii (gold), panel iii (grey), and panel iv (red). Panel ii, which was recorded with the AMS chopper open, shows the mass spectrum of the particles and the background. Panel iii, which was recorded with the AMS chopper closed, shows the mass spectrum of the background. Panel iv, which shows the difference of panel iii from panel ii, represents the mass spectrum of the particles. In each panel, the open circles show the recorded signal (Hz). The

solid lines show the model fits in gold, grey, or red for individual peaks and in black for the sum of the model fits. The solid green bars indicate the peak intensity of each fit.

Figure S4. Contribution of the signal intensity at m/z 60 to the total organic signal intensity (i.e., 60/org) for increasing SOA particle mass loading for the dark ozonolysis of α -pinene. The average 60/org across all loading is 0.23%. Increased signal at m/z 60 is interpreted as a marker of biomass burning particles when observed in the mass spectra of atmospheric particles (Schneider *et al.*, 2006; Docherty *et al.*, 2008). Figure S4, however, shows that some of the signal intensity at m/z 60 can also be attributed to SOA particles.

Literature Cited

DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T., Docherty, K. S., Worsnop, D. R. and Jimenez, J. L.: A field-deployable high-resolution time-of-flight aerosol mass spectrometer, *Anal. Chem.*, **78**, 8281-8289, 2006.

Aiken, A. C., DeCarlo, P. F., Kroll, J. H., Worsnop, D. R., Huffman, J. A., Cocherty, K., Ulbrich, I. M., Mohr, C., Kimmel, J. R., Sueper, D., Zhang, Q., Sun, Y., Trimborn, A., Northway, M., Ziemann, P. J., Canagaratna, M. R., Alfarra, R., Prevot, A. S. H., Dommen, J., Duplissy, J., Metzger, A., Baltensperger, U. and Jimenez, J. L.: O/C and OM/OC ratios of primary, secondary, and ambient organic aerosols with high resolution time-of-flight aerosol mass spectrometry, *Environ. Sci. Technol.*, **42**, 4478-4485, 2008.

Docherty, K. S., Stone, E. A., Ulbrich, I. M., DeCarlo, P. F., Snyder, D. C., Schauer, J. J., Peltier, R. E., Weber, R. J., Murphy, S. N., Seinfeld, J. H., Grover, B. D., Eatough, D. J. and Jimenez, J. L.: Apportionment of primary and secondary organic aerosols in southern California during the 2005 study of organic aerosols in Riverside (SOAR-1), *Environ. Sci. Technol.*, **42**, 7655-7662, 2008.

Schneider, J., Weimer, S., Drewnick, F., Borrmann, S., Helas, G., Gwaze, P., Schmid, O., Andreae, M. O. and Kirchner, U.: Mass spectrometric analysis and aerodynamic properties of various types of combustion-related aerosol particles, *Int. J. Mass Spectrom.*, **258**, 37-49, 2006.