

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.

Heterogeneous uptake of gaseous hydrogen peroxide by Gobi and Saharan dust aerosols: a potential missing sink for H₂O₂ in the troposphere

M. Pradhan, G. Kyriakou, A. T. Archibald, A. C. Papageorgiou, M. Kalberer, and R. M. Lambert

Department of Chemistry, University of Cambridge, Lensfield Road, Cambridge, CB2 1EW, UK

Received: 16 April 2010 – Accepted: 20 April 2010 – Published: 26 April 2010

Correspondence to: R. M. Lambert (rml1@cam.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

The first direct laboratory measurements of gaseous hydrogen peroxide uptake by authentic Gobi and Saharan dust aerosol particles as a function of relative humidity (RH) have been carried out in an entrained aerosol flow tube coupled to a chemical ionization mass spectrometer. Gobi dust shows uptake coefficients, $\gamma_{\text{H}_2\text{O}_2} = (3.33 \pm 0.26) \times 10^{-4}$ at 15% RH rising to $\gamma_{\text{H}_2\text{O}_2} = (6.03 \pm 0.42) \times 10^{-4}$ at 70% RH; the corresponding values for Saharan dust are systematically higher ($\gamma_{\text{H}_2\text{O}_2} = (6.20 \pm 0.22) \times 10^{-4}$ at 15% RH rising to $\gamma_{\text{H}_2\text{O}_2} = (9.42 \pm 0.41) \times 10^{-4}$ at 70% RH). High resolution X-ray photoelectron spectroscopy (XPS) measurements of the surface chemical composition of the two mineral dust samples together with published water adsorption isotherms of their principal constituents enables rationalization of these observations, which are relevant to nighttime tropospheric chemistry. A box model study performed by incorporating the experimentally determined data set reveals that uptake of H_2O_2 onto dust can be an important loss process for this species which has been, until now, poorly constrained.

1 Introduction

The Gobi desert in northwest China and the Sahara desert in northern Africa are the two most important global sources of mineral dust aerosol with current estimates of between 1000–3000 Tg/yr being advected into the atmosphere (Tegen and Fung, 1994; Dentezer et al., 1996; Luo et al., 2003). Dust aerosols produced from windblown soils in arid and semi-arid regions can be viewed as a mixture of inorganic oxides and carbonates, possibly with some adsorbed organics and/or sulfates present (Usher et al., 2003; Hatch and Grassian, 2008). The atmospheric residence time of mineral dust aerosol depends on several factors such as metrological conditions and the particle size distribution. Large dust particles with diameters of 100 μm are only found in the vicinity of desert regions whereas small particles with diameters typically <4 μm have

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

an atmospheric lifetime of up to several weeks (Prospero, 1999; Bauer et al., 2004). They can be transported over thousands of kilometers by individual dust storms from Africa across the Atlantic to the east coast of the United States or from Asia across the Pacific Ocean (Duce et al., 1980; Savoie and Prospero, 1982; Ott et al., 1991).

5 During long-range transport, particles interact with various organic and inorganic atmospheric trace gas-species, and can consequently impact numerous atmospheric processes, potentially changing the chemical balance of the atmosphere through heterogeneous reactions by providing reactive surface sites (Cwiertny et al., 2008; Hatch and Grassian, 2008). Other well-known processes include direct and indirect climate
10 forcing (due to scattering and absorbing of incoming solar radiation) or effects on biogeochemical cycles of trace elements such as iron (Ravishankara, 1997; Kaufman et al., 2002; Usher et al., 2003).

Mineral dust particles can act as a sink or reactive surface for a variety of atmospheric chemical species (Usher et al., 2003). Accordingly, heterogeneous reactions
15 occurring on the surface of mineral dust aerosol particles may provide additional chemical pathways, which may account for the discrepancies between field and laboratory measurements. During the last decade, the heterogeneous chemical kinetics of gas-phase species such as SO_2 , O_3 , NO_3 , N_2O_5 , and HNO_3 interacting with mineral dust aerosol particles has attracted intensive study: both laboratory studies and field ex-
20 periments have demonstrated the importance of this chemistry (Hanisch and Crowley, 2003; Adams et al., 2005; Karagulian and Rossi, 2005; de Reus et al., 2005; Vlasenko et al., 2006; Wagner et al., 2008; Crowley et al., 2010).

In contrast, laboratory measurements of the heterogeneous chemical kinetics of gaseous hydrogen peroxide (H_2O_2) under realistic atmospheric conditions have not
25 been explored thus far. H_2O_2 is an important atmospheric oxidant with a relatively short atmospheric lifetime (~ 1 day) and serves as a sensitive indicator for HO_x ($\text{HO}_2 + \text{OH}$) chemistry (Kuhlmann et al., 2003; Rinsland et al., 2007; Hua et al., 2008). H_2O_2 is highly water soluble, acting as an important oxidant of sulfur compounds in the aqueous phase where it may be involved in the formation of secondary organic aerosol (Hua

et al., 2008). Detailed quantitative understanding of the removal of atmospheric H₂O₂ by mineral dust aerosol is therefore of importance for an accurate assessment of its lifetime, its potential impact on the tropospheric HO_x family chemistry, OH budgets, and accordingly, ozone balance. Moreover, it is important to understand how tropospheric photochemical oxidant cycles and the behaviour of other atmospheric pollutants are affected as a consequence of heterogeneous reactions of H₂O₂.

To our knowledge, there is almost no information involving direct measurement of H₂O₂ uptake by authentic mineral dust aerosol particles. Due to the absence of experimental values for the heterogeneous H₂O₂ uptake coefficient ($\gamma_{\text{H}_2\text{O}_2}$) on mineral dust aerosol, previous work on combined aerosol/gas-phase chemical box models and on global and regional chemistry-transport models was necessarily based on assumptions and estimates. Inevitably, substantial uncertainties arise from the lack of reliable values for uptake coefficients used in such studies. For example, in their modeling studies Zhang et al. (1994) and Phadnis and Carmichael (2000), used a value of $\gamma_{\text{H}_2\text{O}_2}=0.1$, whereas in similar studies by Dentener et al. (1996) and Martin et al. (2003), the uptake of H₂O₂ on dust was simply neglected. Other modeling studies have used widely divergent values in the range $8.0 \times 10^{-4} \leq \gamma_{\text{H}_2\text{O}_2} \leq 0.18$, corresponding to an upper limit and lower limit (Zhang and Carmichael, 1999; Zhu et al., 2009).

Here, we present a detailed investigation of the heterogeneous chemical kinetics of authentic mineral dust aerosol with gas-phase H₂O₂. To our knowledge, this is the first laboratory study of the heterogeneous uptake kinetics of gaseous H₂O₂ by authentic Gobi and Saharan desert dust aerosol particles over a wide range of relative humidity pertinent to the troposphere. Additionally, we explore the potential impact of heterogeneous losses of H₂O₂ on tropospheric chemistry by means of a box model study that incorporates our own experimentally determined data set.

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2 Experimental

Experiments were carried out in an aerosol flow tube (AFT) coupled to a chemical ionization mass spectrometer (CIMS) for the detection of H_2O_2 . The experimental set-up and measurement procedure have been described in detail in our recent publication (Pradhan et al., 2010) and are summarized below briefly in Sect. 2.2.

2.1 Composition of dust aerosols: X-ray photoelectron spectroscopy

Two types of authentic desert dust aerosols were used: the Saharan dust originates from Cape Verde Islands and is characteristic of mineral dust aerosol transported from the western Sahara, whereas Gobi sand was collected from the Khongoryn Els in Mongolia (central China) and was mechanically ground and sieved to $\leq 1 \mu\text{m}$ diameter. The chemical composition of both dusts was determined by high resolution X-ray photoelectron spectroscopy (XPS) (Fig. 1) a technique that provides information about surface as opposed to bulk chemical composition.

Details of sample preparation and handling, instrumentation, data acquisition, quantification and representative spectra are given in the supporting information (see <http://www.atmos-chem-phys-discuss.net/10/11081/2010/acpd-10-11081-2010-supplement.pdf>). Figure 1 summarizes the quantified results obtained for both types of sample. It is evident from Fig. 1 that silica and alumina are the most abundant components in both samples, as would be expected, and that certain minor components show large differences in relative concentrations. Note that Saharan dust contains a small amount of Ti present as TiO_2 whereas this material is undetectable in the Gobi sample. Moreover, both dusts contain Fe as Fe_2O_3 : the significance of these findings will be discussed below.

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2 Generation and characterization of dust aerosol

Aerosol of either Gobi or Saharan dust was generated from an aqueous suspension using a commercial atomizer (TSI 3076). The aerosol relative humidity (RH) was controlled using a silica gel drying unit in series with a humidifier system and monitored with an accuracy of $\pm 2\%$ at the inlet to the AFT with a commercial hygrometer (Vaisala HMP 143A). The size distributions of the aerosol particles were characterized by a differential mobility analyzer (DMA, Hauke model EMS VIE-08) with a Faraday cup electrometer as the detector. Scans over mobility diameters ranging from 0.01–1 μm produced a size distribution from which total surface areas were determined. Prior to entering the DMA or AFT, aerosol particles passed through a cut-off Berner cascade impactor to remove particles of diameters greater than 1 μm . The particle number density and aerosol surface area both remained constant for the two different dusts to within 5% over the time of a kinetic run (≤ 1 h). Furthermore, the particle size distributions for both dusts remained constant with increasing the RH in the range 15–70%.

Typical size distributions of the dust particles are depicted in Fig. 2. Gobi dust exhibits a bimodal distribution whereas that of the Saharan dust is monomodal. The origin of the bimodal distribution of the Gobi dust is not clear. We may speculate that the bimodal distribution is a consequence of agglomeration of the smaller particles in the atomizer suspension. The insets in Fig. 2 show the distributions in terms of surface area ($\mu\text{m}^2 \text{cm}^{-3}$) given as the function $\pi D^2 \frac{dN}{d \ln D}$ obtained for each of the 34 size fractions measured in the DMA. The particle diameters at the maximum in the surface area distribution are 0.40 μm and 0.17 μm for the Gobi and Saharan dusts, respectively.

2.3 Aerosol Flow Tube reactor for kinetic experiments

The aerosol flow tube (AFT) reactor has been described in detail previously (Pradhan et al., 2010). The halocarbon wax coated AFT (length 60 cm, ID 3 cm) was maintained at ambient temperature (295 ± 2 K) and pressure. Under typical measurement conditions the constant volumetric flow rate was 1.25 slpm corresponding to an average linear

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



flow velocity of $\sim 2.4 \text{ cm s}^{-1}$; the Reynold's number and time for diffusional mixing were estimated to be ~ 107 and 2.4 s , respectively. Contact times were varied by means of a sliding injector and all heterogeneous kinetic measurements were performed at contact times of 6–15 s (see Fig. 3) to ensure that data were acquired under well-mixed and fully developed laminar flow conditions.

2.4 Generation of gas-phase H_2O_2 and detection system

A detailed description of the generation with delivery system of gaseous H_2O_2 has been reported previously (Pradhan et al., 2010) and only operational parameters are given here. A concentrated H_2O_2 sample of ($\geq 95 \text{ wt} \%$) was prepared from an aqueous solutions of H_2O_2 (Aldrich, $50 \text{ wt} \%$) for the kinetic measurements. To generate the concentrated sample, a flow of dry N_2 was bubbled through the diluted aqueous solution for two days; then the H_2O_2 finger (pyrex U-tube), kept in an ice bath, was transferred to a vacuum line where the vapour pressure of H_2O_2 was monitored. The vapour pressure of pure H_2O_2 at 273.2 K is 0.352 Torr as measured by Manatt and Manatt (2004). Measurement of the vapour pressure in the present case indicated that purity was $\geq 95 \text{ wt} \%$ H_2O_2 . The H_2O_2 vapour was delivered to the experimental gas line by flowing 20 sccm dry N_2 through the cold finger at 273.2 K and the resulting H_2O_2 saturated flow was then mixed with a 1200 sccm dry N_2 carrier flow. A fraction of this flow (25 sccm) was introduced into the AFT via the injector. When this flow was mixed with the main gas flow (1000 sccm), it resulted in a typical initial flow tube concentration of $\text{H}_2\text{O}_2 \sim 4.2 \times 10^{12} \text{ molecules cm}^{-3}$ (169 ppbv).

H_2O_2 was detected with the CIMS (Extrel Model 150-QC) using CF_3O^- ($m/z=85$) as a reagent ion. CF_3O^- was generated by flowing trace amounts of CF_3OOCF_3 in 2.0 slpm N_2 over a radioactive ^{210}Po source into the chemical ionization region. The use of CF_3O^- for sensitive and selective CIMS detection of H_2O_2 was first demonstrated by Crouse et al. (2006) and subsequently used by ourselves for kinetic measurements (Pradhan et al., 2010). H_2O_2 was monitored as the cluster ion

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



CF₃O⁻·H₂O₂ detected at $m/z=119$. The mass spectrometer was modified by incorporation of a collisional dissociation chamber (CDC) after the ion source region: this allowed sensitive detection of the target cluster ion of CF₃O⁻·H₂O₂, free from any higher cluster ions of the type CF₃O⁻·(H₂O)_{*n*}. The typical CIMS detection limit for H₂O₂ was estimated to be $\sim 1.6 \times 10^{11}$ molecules cm⁻³ (6 ppbv) at the 1σ standard deviation noise level with respect to the baseline under an initial flow tube concentration of H₂O₂ $\sim 4.2 \times 10^{12}$ molecules cm⁻³.

3 Results and discussion

3.1 Heterogeneous kinetic measurements on Gobi dust aerosol

The kinetic experiments were performed in the AFT in which the contact or reaction time between the aerosol and H₂O₂ were controlled by a sliding injector. The first order heterogeneous rate coefficient for loss of H₂O₂ solely to the aerosol particles, k_{aerosol} is defined by

$$k_{\text{aerosol}} = \frac{d[\text{H}_2\text{O}_2]}{dt[\text{H}_2\text{O}_2]} = \gamma_{\text{H}_2\text{O}_2} \frac{\langle c \rangle}{4} S_a \quad (1)$$

where S_a is the total surface area of the aerosol particles per unit volume (cm² cm⁻³) as calculated from the DMA measurements, $\langle c \rangle$ is the mean molecular velocity of H₂O₂ gas molecules (cm s⁻¹) and $\gamma_{\text{H}_2\text{O}_2}$ is the uptake coefficient. Figure 3 illustrates typical results from kinetic run of H₂O₂ uptake by Gobi dust aerosol particles [$S_a = (1.09 \pm 0.05) \times 10^{-3}$ cm² cm⁻³] at 35% RH and initial [H₂O₂] $\approx 4.2 \times 10^{12}$ molecules cm⁻³ in the AFT with a H₂O₂ sensitivity of ~ 5 cps/ppbv. In converting mobility diameters to surface areas it is assumed that the particles have spherical shape. The kinetics of H₂O₂ loss to the walls was measured both prior to and following the aerosol kinetic run. First-order rate constants, k_{wall} for loss in the absence

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of aerosol and k_{total} for losses to both aerosol and wall (i.e. $k_{\text{total}} = k_{\text{aerosol}} + k_{\text{wall}}$), are determined from the gradients of weighted linear least square fits to the data sets. The resulting first-order rate constants were then corrected for non-plug flow conditions using the iterative method of Brown (1978). In this correction the gas-phase diffusion constant for H_2O_2 was taken as $0.184 \text{ cm}^2 \text{ s}^{-1}$ as calculated by McMurtrie and Keyes (1978). The H_2O_2 uptake coefficient, $\gamma_{\text{H}_2\text{O}_2}$ was then determined from the rate constant, $k_{\text{aerosol}} = (k_{\text{total}} - k_{\text{wall}})$ by using Eq. (1).

The results shown in Fig. 3 for the Gobi dust aerosol yield a value of $\gamma_{\text{H}_2\text{O}_2} = (3.51 \pm 0.28) \times 10^{-4}$. However, it should be mentioned that the value of the experimentally determined uptake coefficient (γ_{exp}) obtained in this way can lead to an underestimation of the true uptake coefficient (γ_{true}). This effect resulting from gas phase diffusion can be corrected for, as described by the method of Fuchs and Sutugin (1970) using:

$$\frac{1}{\gamma_{\text{true}}} = \frac{1}{\gamma_{\text{exp}}} - \frac{0.75 + 0.283K_n}{K_n(K_n + 1)} \quad (2)$$

$$K_n = \frac{3D_g}{\langle c \rangle r_{\text{sw}}} \quad (3)$$

where D_g is the gas-phase diffusion coefficient of H_2O_2 , r_{sw} is the radius of the particle at the maximum of the surface area weighted size distribution and K_n is the Knudsen number. We estimated that this correction is negligible ($< 1\%$) over the entire aerosol size range ($0.01\text{--}1 \mu\text{m}$) and for the relatively small reaction probabilities we observed in all kinetic runs. We therefore neglect the diffusion correction and all uptake coefficients reported here are derived solely from the Eq. (1). Additionally, the effect of varying H_2O_2 concentration was investigated by performing measurements over the range of $\sim (3.5\text{--}8.2) \times 10^{12} \text{ molecules cm}^{-3}$; no significant dependence of H_2O_2 uptake on the H_2O_2 concentration was observed within the experimental error.

3.2 Surface area dependence of heterogeneous rate constants

To check the validity of the observed first order kinetics, we performed kinetic measurements in which the aerosol surface area was varied at a constant RH. This was achieved by varying the atomizer flow mixed into the humidified bulk N₂ flow.

5 Figure 4 displays a plot of the retrieved heterogeneous rate constant (k_{aerosol}) versus aerosol surface area (S_a) at 35% RH and $[\text{H}_2\text{O}_2] \sim 4.2 \times 10^{12}$ molecules cm^{-3} in the AFT. A linear relationship with a gradient of $\gamma_{\text{H}_2\text{O}_2} \frac{(c)}{4}$ was found for both dust aerosols. This demonstrates that the observed kinetics are first order in surface area and free from systematic errors. The gradient of the linear fits yield uptake coefficients

10 $\gamma_{\text{H}_2\text{O}_2} = (3.63 \pm 0.24) \times 10^{-4}$ for the Gobi dust and $\gamma_{\text{H}_2\text{O}_2} = (6.76 \pm 0.22) \times 10^{-4}$ for the Saharan dust at 35% RH in excellent agreement with the measurements made using a single surface area as a function of RH shown in Fig. 5.

3.3 Humidity dependence of H₂O₂ uptake on Gobi and Saharan dusts

Uptake coefficients were measured over a wide range of relative humidity (15%–70%) relevant to the troposphere. It is apparent from Fig. 5 that the uptake coefficients, $\gamma_{\text{H}_2\text{O}_2}$, of both dusts initially increases slowly with RH and then more rapidly for $\text{RH} \geq 30\%$. Such positive correlations of γ with RH have been reported previously, for example in the case of HNO₃ on Arizona Test Dust (ATD) (Vlasenko et al., 2006). As expected, both dust samples are composed mainly of silica, alumina and magnesia (Fig. 1) accompanied by smaller amounts of other oxidic materials. Moreover, the water adsorption isotherms of SiO₂, Al₂O₃ and MgO have been accurately measured at 296 K over the RH range 0–98% (Usher et al., 2003). Interestingly, these data show initial Langmuir-like water uptake with monolayer completion occurring at ~30% RH in every case. Thereafter, multilayer formation commences, but the rate of accumulation of these higher layers is markedly dependent on the oxide in question: for example, it is much faster on SiO₂ than on MgO (Usher et al., 2003). This behavior is in striking qualitative accord with the results shown in Fig. 5. Keeping in mind the complexity of

15

20

25

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



our mineral surfaces, we may offer the following admittedly approximate rationalization of these results. At low RH the dust surface is relatively water-free so that impinging H_2O_2 molecules encounter either “bare” pristine surface sites or those on which dissociative adsorption of H_2O_2 has occurred. In these circumstances, the different chemical compositions of the two types of dust inevitably results in significantly different values of $\gamma_{\text{H}_2\text{O}_2}$. At $\text{RH} \sim 30\%$ water multilayers nucleate and start to spread to varying extents on all the solid phases present (Usher et al., 2003) and $\gamma_{\text{H}_2\text{O}_2}$ increases as hydrogen peroxide molecules now impinge on a patchy surface where H_2O_2 uptake into the growing islands of liquid-like water acts to enhance the net uptake coefficient. The approximately constant difference in γ values for the two dusts that persists up to the highest RH accessible in our experiments (70%) is understandable in that at 70% RH all the principal dust components are still only partially covered by multilayers. Presumably the two curves would converge at $\sim 100\%$ RH as MgO and Al_2O_3 multilayer uptake “catches up” with SiO_2 multilayer uptake: the respective isotherms do indeed show large differences in water uptake between 70% and 100% RH.

Finally, we draw attention to a potentially important issue. Saharan dust contains a significant amount of TiO_2 and it is known that this material catalyzes the reduction of NO_2 to nitrous acid when present as an aerosol and illuminated with near UV light under conditions pertinent to the troposphere (Gustafsson et al., 2006). Moreover, both dusts contain Fe_2O_3 (bandgap 2.0 eV) which also exhibits photocatalytic activity under near UV irradiation (Leland et al., 1987). Therefore it is possible that the daytime heterogeneous chemistry of both dusts could be significantly different from their nighttime chemistry.

3.4 Implications for modeling studies

The present results provided the first data for the uptake of H_2O_2 on authentic desert dust aerosol particles: to our knowledge, there are no other experimental findings with which to compare our data. Accordingly, in order to explore the potential impact of the heterogeneous uptake of H_2O_2 by mineral aerosol dust on tropospheric chemistry, we

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

performed a simple box model study (see below) in which our experimentally determined RH-dependent uptake coefficients were incorporated in the calculations. Before describing these findings, we note in passing that in a recent combined field and modeling study by de Reus et al. (2005), the H_2O_2 uptake on Saharan dust plume was estimated to be in the order of $\gamma_{\text{H}_2\text{O}_2} = 5 \times 10^{-4}$ in order to obtain agreement between calculated and observed values of H_2O_2 concentration: this estimate is in very good agreement with our experimental measurements.

The box model was written using the FACSIMILE chemical kinetics program (Curtis and Sweetenham, 1987) and was constructed using inputs from the MINATROC campaign (de Reus et al., 2005). No specific meteorological conditions were simulated; however, a parameterization was included to allow both the model temperature and RH to vary throughout the day (min RH=22%, max RH=70%). A simple photolysis procedure relating the rate of photolysis to the solar zenith angle was used in line with previous box model studies using the Master Chemical Mechanism (MCM) (Saunders et al., 2003). The model was run for a four day period using campaign average mixing ratios of a number of trace gas species as initial conditions. The model included a fairly detailed chemical mechanism based on the MCM (Saunders et al., 2003) comprising 304 species and 1072 reactions, describing the oxidation of the following compounds; CH_4 , CO, HCHO, CH_3OH , CH_3COCH_3 , C_3H_8 , C_4H_{10} and C_5H_8 . The initial conditions for the box model runs are shown in Table 1. A reference simulation was run in which no heterogeneous chemistry was included. This run was then used to derive the effects of inclusion of heterogeneous loss of H_2O_2 on to mineral dust. In all runs dry deposition was included following the results of de Reus (2005) who highlight its importance for, in particular, H_2O_2 . The following species were dry deposited with the maximum loss rates indicated in parenthesis; HNO_3 and N_2O_5 ($2.5 \times 10^{-5} \text{ s}^{-1}$), NO_2 ($1.8 \times 10^{-6} \text{ s}^{-1}$) and O_3 and H_2O_2 ($6.2 \times 10^{-6} \text{ s}^{-1}$).

As we have highlighted, several previous studies have used a large range of values for the uptake coefficient of H_2O_2 onto mineral dust (Zhang et al., 1994; Phadnis and Carmichael, 2000; Zhu et al., 2009). The choice of uptake coefficient in these previous

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



studies was unconstrained due to the lack of previous experimental data. In our box model study, heterogeneous chemistry was simply accounted for by including loss of H_2O_2 by dust using Eq. (1). As with the initialization of chemical species, we have used real world measurement data as inputs of S_a (the dust aerosol surface area) from de Reus et al. (2005). In the model $\gamma_{\text{H}_2\text{O}_2}$ was calculated as a function of the observed RH dependent data for Saharan dust (Fig. 5) as it was Saharan dust that was encountered in the study of de Reus et al. (2005). Thus our calculated values of k_{aerosol} vary with the changing RH throughout time in the model. The complex interplay of dust with radiation is not currently considered in our simple model, but it is duly noted that several previous studies have highlighted that dust has a dampening effect on the photolysis rates of many labile trace gases. In light of this we acknowledge that our model formulation is not entirely representative of real world conditions. However, we note that the aims of this work are to assess the first order sensitivity of the chemical system to the new measurements.

The chemistry of H_2O_2 in the reference simulation can be expressed analytically as:

$$\frac{d[\text{H}_2\text{O}_2]}{dt} = k_1 [\text{HO}_2]^2 - j_2 [\text{H}_2\text{O}_2] - k_3 [\text{OH}][\text{H}_2\text{O}_2] - v_d [\text{H}_2\text{O}_2] \quad (4)$$

where the first term corresponds to the production of H_2O_2 from the $\text{HO}_2 + \text{HO}_2$ self reaction. The rate constant for this reaction (k_1) shows both bi-molecular and ter-molecular components and most notably shows significant enhancement in the presence of hydrogen bonding gases (e.g. H_2O , Stone and Rowley, 2005). The loss of H_2O_2 in the reference run is then accounted for by photolysis (j_2), the reaction with OH (k_3) and dry deposition (v_d) with kinetic data taken from the MCM (apart from v_d , see above). Equation (4) is then modified in the sensitivity tests by inclusion of the additional loss process of H_2O_2 to dust aerosol.

Dust was accounted for in the model using a single parameter, S_a . This value was maintained as a constant throughout the model runs and thus these runs highlight the effects of time integrated exposure of the chemical system to dust. Hence these runs mimic a dust laden air parcel undergoing photochemical aging. Several sensitivity tests

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

were performed in which the input S_a was varied from a minimum of $15 \mu\text{m}^2 \text{cm}^{-3}$ to a maximum of $200 \mu\text{m}^2 \text{cm}^{-3}$ representing the range of measured values quoted by de Reus et al. (2005). The results of these tests are shown in Figs. 6 and 7, relative to the reference run in which no heterogeneous loss of H_2O_2 was included. The results of the simulations highlight that heterogeneous uptake of H_2O_2 on dust can act as an important loss process of H_2O_2 , currently not considered in many models. In the base run the loss of H_2O_2 is dominated by dry deposition to the surface (assuming an average $v_d=0.5 \text{cm s}^{-1}$) with the contributions from photolysis and the reaction with OH being roughly equal. Figure 6 shows that with reasonably high levels of dust loading (daily average $S_a \leq 100 \mu\text{m}^2 \text{cm}^{-3}$), there are increases in the loss of H_2O_2 of $\sim 10\%$ and greater. This increase in H_2O_2 removal leads to decreases in the H_2O_2 lifetime with respect to dust of a factor of ~ 4 . In Fig. 7 the effects of inclusion of H_2O_2 uptake are shown as a function of S_a for a range of important trace gas species. The results are presented as the relative difference (%) after the second day of model integration. From Fig. 7 it is clear that the single biggest effect of inclusion of this extra loss term for H_2O_2 is that the model simulated H_2O_2 concentrations are substantially reduced (up to $\sim 40\%$ after two days). The effects on total organic peroxy radicals (ΣRO_2), OH, O_3 and NO_x are all smaller than $\pm 5\%$.

Finally, a further run was performed whereby the uptake of H_2O_2 into the aqueous phase was included. The role of H_2O_2 as an in cloud oxidant has been known for several decades and it is widely acknowledged that H_2O_2 is one of the principal oxidants involved in conversion of SO_2 to $\text{SO}_4^{(-2)}$. In this run again only perturbations to the chemistry of H_2O_2 are considered with the aqueous uptake acting as an additional loss process for Eq. (4). Data for the aqueous uptake were based on the work of Redington et al. (2009). Considering a cloud fraction of 30% and a cloud liquid water content of 0.3g m^{-3} (typical of cumulus clouds) the aqueous phase uptake outweighs the other loss processes by a further 26%. The model simulated H_2O_2 mixing ratios are thus substantially reduced, being $\sim 60\%$ lower than the reference run after two days. Thus under these conditions the effects of aqueous removal H_2O_2 are a factor of ~ 2

greater than for heterogeneous loss onto dust. However, we note that in many of the regions where severe dust events occur the local meteorology may dictate that cloud formation is weak and so this aqueous phase loss shall be much less important.

4 Conclusions

5 The uptake coefficients of hydrogen peroxide on Saharan and Gobi mineral aerosol dusts are significantly different and vary markedly with relative humidity. They are very different from either the much bigger or the negligible values that have been assumed in earlier modeling studies. Using these experimental values in a simple box model indicates that the loss of H_2O_2 to dust aerosol may present an important additional
10 loss process for H_2O_2 . This process is dependent on the simulated dust loading (in our case the dust aerosol surface area) and increases strongly with increasing loading. Under typically very dusty conditions we have calculated that dust accounts for $\sim 75\%$ of the loss of H_2O_2 and this term leads to an $\sim 15\%$ increase in the total rate of loss of H_2O_2 compared to a reference simulation. This quickly leads to significant reductions
15 in the modeled mixing ratios of H_2O_2 after only a few days.

In the light of the measured surface chemical composition of the two mineral dusts and given the water adsorption isotherms of their principal constituents, it is possible to rationalize our observations, which are characteristic of their nighttime tropospheric chemistry. The presence of relatively minor but strongly UV-photoactive components
20 points to the possibility that the daytime chemistry may be significantly different.

Acknowledgements. We are very grateful to Helge Willner, Bergische Universität, Wuppertal, Germany for kindly providing the CF_3OOCF_3 used in this study. We would like to thank Tony Cox for his kind provision of the Saharan dust sample and Mette High for collecting Gobi sample used for measurements. M. P. acknowledges a post-doctoral research funding by the
25 Isaac Newton Trust, Trinity College, Cambridge. A. T. A. wishes to thank NCAS Climate for funding. G. K. and A. C. P. acknowledge financial support from the UK Engineering and Physical Sciences Research Council.

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

- Adams, J. W., Rodriguez, D., and Cox, R. A.: The uptake of SO₂ on Saharan dust: a flow tube study, *Atmos. Chem. Phys.*, 5, 2679–2689, 2005, <http://www.atmos-chem-phys.net/5/2679/2005/>.
- 5 Bauer, S. E, Balkanski, Y., Schulz, M., Hauglustaine, D. A., and Dentener, F.: Global modeling of heterogeneous chemistry on mineral aerosol surfaces: Influences on tropospheric ozone chemistry and composition to observations, *J. Geophys. Res.*, 109, D02304, doi:10.1029/2003JD003868, 2004.
- Brown, R. L.: Tubular flow reactors with first-order kinetics, *J. Res. Nat. Bur. Standards*, 83, 1–8, 1978.
- 10 Crounse, J. D., Mckinney, K. A., Kwan, A. J., and Wennberg, P. O.: Measurement of gas-phase hydroperoxides by chemical ionization mass spectrometry, *Anal. Chem.*, 78, 6726–6732, 2006.
- Crowley, J. N., Ammann, M., Cox, R. A., Hynes, R. G., Jenkin, M. E., Mellouki, A., Rossi, M. J., Troe, J., and Wallington, T. J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume V heterogeneous reactions on solid substrates, *Atmos. Chem. Phys. Discuss.*, 10, 5233–5564, 2010, <http://www.atmos-chem-phys-discuss.net/10/5233/2010/>.
- 15 Curtis A. R. and Sweetenham, W. P. Facsimile Users Manual; Computer Science and Systems Division, H. L., Ed.; United Kingdom Atomic Energy Authority: Oxford, England, 1987.
- Cwiertny, D. M., Young, M. A., and Grassian, V. H.: Chemistry and photochemistry of mineral dust aerosol, *Annu. Rev. Phys. Chem.*, 59, 27–51, 2008.
- Dentener, F. J., Carmichael, G. R., Zhang, Y., Lelieveld, J., and Crutzen, P. J.: Role of mineral aerosol as a reactive surface in the global troposphere, *J. Geophys. Res.*, 101, 22869–22889, 1996.
- 25 de Reus, M., Fischer, H., Sander, R., Gros, V., Kormann, R., Salisbury, G., Van Dingenen, R., Williams, J., Zöllner, M., and Lelieveld, J.: Observations and model calculations of trace gas scavenging in a dense Saharan dust plume during MINATROC, *Atmos. Chem. Phys.*, 5, 1787–1803, 2005, <http://www.atmos-chem-phys.net/5/1787/2005/>.
- 30 Duce, R. A., Unni, C. K., Ray, B. J., Prospero, J. M., and Merrill, J. T.: Long-range atmospheric transport of soil dust from Asia to the Tropical North Pacific: Temporal Variability, *Science*,

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



209, 1522–1524, 1980.

- Fuchs, N. A. and Sutugin, A. G.: Highly dispersed aerosols, Ann Arbor Sci., Ann Arbor, 1970.
- Gustafsson, R. J., Orlov, A., Griffiths, P. T., Cox, R. A., and Lambert, R. M.: Reduction of NO₂ to nitrous acid on illuminated titanium dioxide aerosol surfaces: implications for photocatalysis and atmospheric chemistry, Chem. Commun., 37, 3936–3938, 2006.
- Hanisch, F. and Crowley, J. N.: Ozone decomposition on Saharan dust: an experimental investigation, Atmos. Chem. Phys., 3, 119–130, 2003, <http://www.atmos-chem-phys.net/3/119/2003/>.
- Hatch, C. D. and Grassian, V. H.: 10th Anniversary Review: Applications of analytical techniques in laboratory studies of the chemical and climatic impacts of mineral dust aerosol in the Earth's atmosphere, J. Environ. Monit., 10, 919–934, 2008.
- Hua, W., Chen, Z. M., Jie, C. Y., Kondo, Y., Hofzumahaus, A., Takegawa, N., Chang, C. C., Lu, K. D., Miyazaki, Y., Kita, K., Wang, H. L., Zhang, Y. H., and Hu, M.: Atmospheric hydrogen peroxide and organic hydroperoxides during PRIDE-PRD'06, China: their concentration, formation mechanism and contribution to secondary aerosols, Atmos. Chem. Phys., 8, 6755–6773, 2008, <http://www.atmos-chem-phys.net/8/6755/2008/>.
- Karagulian, F. and Rossi, M. J.: The heterogeneous chemical kinetics of NO₃ on atmospheric mineral dust surrogates, Phys. Chem. Chem. Phys., 7, 3150–3162, 2005
- Kaufman, Y. J., Tanre, D., and Boucher, O.: A satellite view of aerosols in the climate system, Nature, 419, 215–223, 2002.
- Kuhlmann, R. V., Lawrence, M. G., and Crutzen, P. J.: A model for studies of tropospheric ozone and nonmethane hydrocarbons: Model evaluation of ozone-related species. J. Geophys. Res., 108(D23), 4729, doi:10.1029/2002JD00, 2003.
- Leland, J. K. and Bard, A. J.: Photochemistry of colloidal semiconducting iron oxide polymorphs, J. Phys. Chem., 91, 5076–5083, 1987.
- Luo, C., Mahowald, N. M., and del Corral, J.: Sensitivity study of meteorological parameters on mineral aerosol mobilization, transport and distribution, J. Geophys. Res., 108, 4447, doi:10.1029/2003JD003483, 2003.
- Manatt, S. L. and Manatt, M. R. R.: On the analyses of mixture vapor pressure data: The hydrogen peroxide/ water system and its excess thermodynamic functions, Chem. Eur. J., 10, 6540–6557, 2004.
- Martin, R. V., Jacob, D. J., Yantosca, R. M., Chin, M., and Ginoux, P.: Global and regional

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

decrease in tropospheric oxidants from photochemical effects of aerosols, *J. Geophys. Res.*, 108(D3), 4097, doi:10.1029/2002JD002622, 2003.

McMurtrie, R. L. and Keyes, F. G.: A measurement of the diffusion coefficient of hydrogen peroxide vapor into air, *J. Am. Chem. Soc.*, 70(11), 3755–3758, 1948.

5 Ott, S. T., Ott, A., Martin, D. W., and Young, J. A.: Analysis of trans-Atlantic Saharan dust outbreak based on satellite and GATE data, *Mon. Weather Rev.*, 119, 1832–1850, 1991.

Phadnis, M. J. and Carmichael, G. R.: Numerical investigation of the influence of mineral dust on the tropospheric chemistry of East Asia, *J. Atmos. Chem.*, 36, 285–323, 2000.

10 Pradhan, M., Kalberer, M., Griffiths, P. T., Braban, C. F., Pope, F. D., Cox, R. A., and Lambert, R. M.: Uptake of gaseous hydrogen peroxide by submicrometer titanium dioxide aerosol as a function of relative humidity, *Environ. Sci. Technol.*, 44, 1360–1365, 2010.

Prospero, J. M.: Long range transport of mineral dust in the global atmosphere: Impact of African dust on the environment of the southeastern United States, *P. Natl. Acad. Sci. USA*, 96, 3396–3403, 1999.

15 Ravishankara, A. R.: Heterogeneous and multiphase chemistry in the troposphere, *Science*, 276, 1058–1065, 1997.

Redington, A. L., Derwent, R. G., Witham, C. S., and Manning, A. J.: Sensitivity of modelled sulphate and nitrate aerosol to cloud, pH and ammonia emissions, *Atmos. Environ.*, 43, 3227–3234, 2009.

20 Rinsland, C. P., Coheur, P. F., Herbin, H., Clerbaux, C., Boone, C., Bernath, P., and Chiou, L. S.: Detection of elevated tropospheric hydrogen peroxide (H_2O_2) mixing ratios in atmospheric chemistry experiment (ACE) subtropical infrared solar occultation spectra, *J. Quant. Spectrosc. Ra.*, 107, 340–348, 2007.

25 Saunders, S. M., Jenkin, M. E., Derwent, R. G., and Pilling, M. J.: Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part A): tropospheric degradation of non-aromatic volatile organic compounds, *Atmos. Chem. Phys.*, 3, 161–180, 2003, <http://www.atmos-chem-phys.net/3/161/2003/>.

Savoie, D. L. and Prospero, J. M.: Particle-size distribution of nitrate and sulfate in the marine atmosphere, *Geophys. Res. Lett.* 9, 1207–1210, 1982.

30 Stone, D. and Rowley, D. M.: Kinetics of the gas phase HO_2 self-reaction: Effects of temperature, pressure, water and methanol vapours, *Phys. Chem. Chem. Phys.*, 7, 2156–2163, 2005.

Tegen, I. and I. Fung.: Modelling of Mineral dust in the Atmosphere-Sources, Transport and



Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.

- Optical-Thickness, *J. Geophys. Res.-Atmos.*, 99, 22897–22914, 1994.
- Underwood, G. M., Li, P., Miller, Al-Abadleh, H., and Grassian, V. H.: A Knudsen cell study of the heterogeneous reactivity of nitric acid on oxide and mineral dust particles, *J. Phys. Chem. A*, 105, 6609–6620, 2001.
- 5 Usher, C. R., Michel, A. E., and Grassian, V. H.: Reactions on mineral dust, *Chem. Rev.*, 103, 4883–4939, 2003.
- Vlasenko, A., Sjogren, S., Weingartner, E., Gaggeler, H.W., and Ammann, M.: Generation of submicron Arizona Test Dust aerosol: chemical and hygroscopic properties, *Aerosol Sci. Technol.*, 39, 452–460, 2005.
- 10 Vlasenko, A., Sjogren, S., Weingartner, E., Stemmler, K., Gäggeler, H. W., and Ammann, M.: Effect of humidity on nitric acid uptake to mineral dust aerosol particles, *Atmos. Chem. Phys.*, 6, 2147–2160, 2006, <http://www.atmos-chem-phys.net/6/2147/2006/>.
- Wagner, C., Hanisch, F., Holmes, N., de Coninck, H., Schuster, G., and Crowley, J. N.: The interaction of N_2O_5 with mineral dust: aerosol flow tube and Knudsen reactor studies, *Atmos. Chem. Phys.*, 8, 91–109, 2008, <http://www.atmos-chem-phys.net/8/91/2008/>.
- 15 Zhang, Y. and Carmichael, G. R.: The role of mineral aerosol in tropospheric chemistry in East Asia—a model study, *J. Appl. Meteorol.*, 38, 353–366, 1999.
- Zhang, Y., Sunwoo, Y., Kotamarthi, V., and Carmichael, G. R.: Photochemical oxidant processes in the presence of dust: An evaluation of the impact of dust on particulate nitrate and ozone formation, *J. Appl. Meteorol.*, 33, 813–824, 1994.
- 20 Zhang, Y., Sunwoo, Y., Kotamarthi, V., and Carmichael, G. R.: Photochemical oxidant processes in the presence of dust: An evaluation of the impact of dust on particulate nitrate and ozone formation, *J. Appl. Meteorol.*, 33, 813–824, 1994.
- Zhu, S., Butler, T., Sander, R., Ma, J., and Lawrence, M. G.: Impact of dust on tropospheric photochemistry over polluted regions: a case study of the Beijing megacity, *Atmos. Chem. Phys. Discuss.*, 9, 20145–20194, 2009, <http://www.atmos-chem-phys-discuss.net/9/20145/2009/>.
- 25

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Initial chemical and physical conditions for the box model runs.

Parameter	Value
Temperature*	292.0 K
Relative Humidity (RH)*	45.0 %
Pressure	1000 HPa
[H ₂ O]	1.5×10 ¹⁷ molecules cm ⁻³
[CH ₄]	1.8 ppbv
[NO _x]	0.25 ppbv
[O ₃]	45.0 ppbv
[CO]	100.0 ppbv
[HCHO]	0.5 ppbv
[H ₂ O ₂]	1.0 ppbv
[CH ₃ OOH]	1.0 ppbv
[HNO ₃]	0.8 ppbv
[CH ₃ OH]	2.0 ppbv
[CH ₃ COCH ₃]	1.3 ppbv
[C ₃ H ₈]	0.05 ppbv
[C ₄ H ₁₀]	0.03 ppbv
[C ₅ H ₈]	0.15 ppbv

* vary throughout the model run.

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.




Fig. 1. Composition of Saharan and Gobi dust aerosol, as determined by high resolution XPS.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.




Fig. 2. Typical size distributions of dust aerosol particles used to study the heterogeneous reaction with gaseous hydrogen peroxide. The distributions shown result from atomizing a 4.0 wt% suspension of dust in water. Particle diameters at the maximum in the surface area distribution are $0.08 \mu\text{m}$ and $0.40 \mu\text{m}$ for the Gobi (bimodal) and $0.17 \mu\text{m}$ for the Saharan dusts (monomodal).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.




Fig. 3. An example of a heterogeneous kinetic run of H₂O₂ uptake by Gobi dust aerosol at 35% RH in the aerosol flow tube. The lines show 2 σ error-weighted least-squares fits to the data from which first-order rate constants are determined. Error bars on the H₂O₂ signal are 2 σ standard deviation of the noise level in the measured signal. Error bars on contact times result from $\pm 2\%$ uncertainties in the flow rate measurements.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.




Fig. 4. Aerosol surface area, S_a , versus the first-order rate constants for loss of H_2O_2 to the aerosol k_{aerosol} at 35% RH. The calculated uptake coefficients from the gradients of this plot are $\gamma_{\text{H}_2\text{O}_2} = (3.63 \pm 0.24) \times 10^{-4}$ for the Gobi dust and $\gamma_{\text{H}_2\text{O}_2} = (6.76 \pm 0.22) \times 10^{-4}$ for the Saharan dust. The lines represent 2σ error-weighted least-squares fits forced through the origin.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.




Fig. 5. Variation of uptake coefficients for H_2O_2 with relative humidity for the Gobi and Saharan dust samples. Error bars are 2σ statistical. The additional data points at 35% RH for both dusts are calculated from the data shown in Fig. 4.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.




Fig. 6. The increase in H_2O_2 loss as a function of increasing dust exposure relative to the control run with no heterogeneous loss (filled circles). The e -folding lifetime of H_2O_2 is also shown with respect to dust exposure (filled triangles).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Heterogeneous uptake of gaseous hydrogen peroxide

M. Pradhan et al.




Fig. 7. The effects of including reactive uptake of H_2O_2 as a function of dust loading for a range of trace gases.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

