

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Unraveling the complex local-scale flows influencing ozone patterns in the southern Great Lakes of North America

I. Levy¹, P. A. Makar¹, D. Sills², J. Zhang¹, K. L. Hayden¹, C. Mihele¹, J. Narayan¹, M. D. Moran¹, S. Sjostedt¹, and J. Brook¹

¹Air Quality Research Division, Science and Technology Branch, Environment Canada, 4905 Dufferin Street, Toronto, Ontario, Canada

²National Laboratory for Nowcasting and Remote Sensing Meteorology, Environment Canada, 4905 Dufferin Street, Toronto, Ontario, Canada

Received: 9 July 2010 – Accepted: 13 August 2010 – Published: 23 August 2010

Correspondence to: J. Brook (jeff.brook@ec.gc.ca)

Published by Copernicus Publications on behalf of the European Geosciences Union.

ACPD

10, 19763–19810, 2010

**Unraveling the
complex local-scale
flows influencing
ozone patterns**

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

This study examines the complexity of various processes influencing summertime ozone levels in the southern Great Lakes region of North America. Results from the Border Air Quality and Meteorology (BAQS-Met) field campaign in the summer of 2007 are examined with respect to land-lake differences and local meteorology using a large array of ground-based measurements, aircraft data and simulation results from a high resolution (2.5 km) regional air-quality model, AURAMS.

Analyses of average ozone mixing ratio from the entire BAQS-Met intensive campaign period support previous findings that ozone levels are higher over the southern Great Lakes than over the adjacent land. However, there is great heterogeneity in the spatial distribution of surface ozone over the lakes, particularly over Lake Erie during the day, with higher levels located over the southwestern end of the lake. Results suggest that some of these increased ozone levels are due to local emission sources in large nearby urban centers. The land-lake differences in ozone mixing ratios are most pronounced during the night in a shallow inversion layer of about 200 m above the surface. After sunrise, these differences have a limited effect on the total mass of ozone over the lakes during the day time, though they may cause elevated ozone levels in the lake breeze air.

A large reservoir layer of ozone is predicted by the AURAMS model over Lake Erie at night, centered between 600–1000 m above ground and extending into the land over Cleveland. The model also predicts a vertical circulation during the day with an updraft over Detroit-Windsor and downdraft over Lake St. Clair, which transports ozone up to 1500 m above ground and results in high ozone over the lake.

Oscillations in ground level ozone mixing ratios were observed on several nights and several ground monitoring sites, with amplitudes of up to 40 ppbv and time periods of 15–40 min.

ACPD

10, 19763–19810, 2010

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

1 Introduction

Land and sea/lake breeze winds play an important role in determining many aspects of coastal environments, and air quality in particular. Air pollutants levels in coastal regions, that hold a large part of the world's population along with emission sources, are the outcome of the combined effect of several factors, such as long range transport, short range transport, local emissions, photochemical activity and surface properties (Banta et al., 2005). While the main driving force of the breeze winds is the temperature difference between land and sea/lake, other factors have been shown to interact with these winds, such as large scale synoptic flow (e.g., Oh et al., 2006), the urban heat island (e.g., Ohashi and Kida, 2002), vicinity to a mountain ridge (e.g., Lu and Turco, 1994), curvature of the shoreline (e.g., Alpert and Getenio, 1988; Levy et al., 2008b) and other topographical features such as vegetation, land use and land cover.

The effect of sea/lake breeze winds on air pollution has been shown in many locations around the world, such as Athens (e.g., Kallos et al., 1993; Kambezidis et al., 1998); the Iberian peninsula (Millan et al., 2000); southern France (e.g., Lasry et al., 2005); Israel (e.g., Alper-Siman Tov et al., 1997; Levy et al., 2008a); Taiwan (e.g., Cheng, 2002; Liu et al., 2002); Korea (Oh et al., 2006), southwestern British Columbia (Brook et al., 2004) and Los Angeles (Lu and Turco, 1995). The large populations living in these coastal cities are often exposed to higher pollution levels due to higher emissions and may experience greater health impacts due to higher chronic exposures accentuated or even induced by the complex local meteorology. Thus, this meteorology can lead to more complex yet potentially repeatable exposure patterns.

The impact of the North American Great Lakes on pollution levels have been studied since the 1960's (e.g., Anlauf et al., 1975; Cooper et al., 2001; Dye et al., 1995; Fast and Heilman, 2003, 2005; Hanna and Chang, 1995; Keeler et al., 1990; Lyons and Cole, 1976, 1973; Mukammal, 1965; Sillman et al., 1993). These multi-year observations combined with models output suggest that ozone is systematically higher over the lakes. Dye et al. (1995) analyzed observations from the Lake Michigan Ozone

ACPD

10, 19763–19810, 2010

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Study in 1991, and found the highest ozone concentrations occur in a shallow and cool conduction layer over the lake. Ozone precursors emitted during the night and early morning were trapped in this shallow layer of cool and stable air over the water to react and produce high concentrations of ozone during the day. Hastie et al. (1999) reported increases of 30 ppbv in ozone levels measured in air masses arriving with the Lake Ontario lake-breeze front, measured at two ground stations north of the lake and from an aircraft. The authors postulate that the origin of these polluted air masses is emissions from source areas that are entrained over the lake by the land breeze at night. Fast and Heilman (2005) used the PEGASUS air-quality model to evaluate ozone levels over the Great Lakes region for two summers (May–September 1999 and 2001). Ozone exceedances above 60 and 80 ppbv were found to be higher on average over the southern Great Lakes than over land. Similar findings were published by Capps et al. (2010) for a 14-day period in August 2002 using the CMAQ air-quality model.

The magnitude of the land-lake differences in ozone are not well characterized due to a lack of routinely available pollutant data of fine spatiotemporal resolution. For example, Hastie et al. (1999) reported ozone levels to change between 60–100 ppbv over Lake Ontario during measurements with an instrumented aircraft on 26 August 1993 around noon, while NO_x changed between 1.5–6 ppbv. In a subsequent flight about one hour later, ozone levels were within the same range, but the location of the maximum ozone had changed, suggesting that temporal changes, perhaps related to advection patterns, also play an important part.

During the Border Air Quality and Meteorology (BAQS-Met) study, which took place in summer 2007 in southwestern Ontario, Canada, a comprehensive suite of air-quality and meteorological measurements were made from both fixed and mobile platforms in order to study air pollutant transport and transformation in relation to lake and land breezes and to gain more insight into the nature of the high ozone over and near the lake in particular. The study region is situated between two of the Great Lakes, Lake Erie and Lake Huron, and a third, smaller lake, Lake St. Clair (see Fig. 1). This paper describes some of the results from this study that are related to ozone levels over

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Lake Erie and Lake St. Clair. First, average ozone mixing ratios from ground-level measurements and from the AURAMS regional air-quality model are examined in time and space. Then, a case study is presented to illustrate the influence of local meteorological circulations driven by the southern Great Lakes on air quality. Included in the analysis are measurements from a meso-network, a chemistry supersite, an instrumented aircraft, an instrumented ferry, an instrumented buoy, and a tethersonde. The analyses, which are unique in the extent to which detailed measurements and model applications have been combined and interpreted, provide insight regarding the behaviour and significance of the elevated ozone levels over the complex coastal environment.

2 Datasets

The BAQS-Met field campaign was conducted during the summer of 2007 to study the effect of mesoscale meteorology on air pollution in the southwestern Ontario. Figure 1 provides an overview of the study area and measurement station locations used in this paper. In addition to routine air-quality monitoring stations operated by the Province of Ontario (Thermo Scientific instrumentation with ozone logged at 1 min resolution) and the State of Michigan (1 h resolution for ozone), spatial detail was enhanced by deploying a meso-network of ozone and meteorological instruments. This was in operation before, during and after the intensive observing period of the study (June 20 to July 10) to obtain 1–5 min time-resolved measurements for the late spring to late summer period. Ozone mixing ratios at the meso-network sites were measured using low-power monitors from 2B Technologies (Model #202). Ozone instruments were also installed on a buoy in Lake Erie and on a ferry operating in Lake Erie between Leamington and Pelee Island (Thermo Scientific instrumentation with ozone logged at 1 min resolution).

Additional detailed chemical measurements were made with research-grade instruments at three chemistry supersites (Harrow, Bear Creek, and Ridgetown) which operated for the intensive period (e.g., Stroud et al., 2010). These sites used instruments

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



from Thermo Scientific to measure O₃ (Model 49C), SO₂ (Model 43C), CO (Model 48 TLE) and NO/NO_x (Model 42C). A standard chemiluminescence-based NO/NO_x instrument (Thermo Scientific Model 42C) was modified in-house to conduct continuous measurements of NO, NO₂ and NO_y with 1 min time resolution. Gas phase volatile organic compounds were measured at Harrow with 5 min resolution by an Ionicon Proton Transfer Mass Spectrometer (PTR-MS) (Gouw and Warneke, 2007). An instrumented Twin Otter aircraft was also deployed for approximately 30 h of flight time measuring gas and particle phase pollutants as well as meteorology at a 1 s time resolution. These measurements are described in detail in Hayden et al. (2010). At the Ridgetown supersite, a Vaisala Tethersonde system consisting of three ECC ozonesondes from Science Pump (model 6A) and tethered meteorological sondes (model # TTS111/RSS911) made measurements from the surface up to about 1000 m above ground level (a.g.l.) on multiple days during the period.

To better understand the BAQS-Met observations, output from the GEM/AURAMS regional air-quality modeling system (Cho et al., 2009; Côté et al., 1998; Makar et al., 2009), which was run with three nested grids with 42, 15 and 2.5 km horizontal grid spacing (Fig. 1), are also analyzed in this paper. A more detailed description of the model set up and a comprehensive evaluation of the model's performance is given by Makar et al. (2010). The main purpose of the model applications presented here was for a more detailed interpretation of the pollutant behavior in the study region and to assess the prevalence and/or validity of the commonly observed and modeled high ozone levels over the lakes. Model output was analyzed with the open-source statistical language R (RDCT, 2009) and visualized with the Unidata Integrated Data Viewer (IDV) (Murray et al., 2003).

Diurnal averages of measured ozone were calculated at 11 rural sites and 6 urban sites, as well as for the Buoy and Pelee Island sites in Lake Erie (shown in Fig. 1, listed in Table 1). For consistency between different sites, averages were calculated for the Buoy operational period only (26 June–10 July 2007). Diurnal averages were created by calculating 2-h running averages of ozone on a 1-min basis (1-h for Michigan

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

sites) for each site during the entire period, then averaging by time of day for each site. The 2-h period was selected for consistency with Michigan sites. Composite diurnal averages were then obtained by averaging over sites grouped by type, i.e., rural and urban, as described in Table 1. Diurnal change rates in ozone concentration were calculated by hour as the difference between the averages of two consecutive hours. Also, the AURAMS predicted ozone levels for the near-surface layer during the buoy operational period were extracted over a relevant analysis sub-domain around southwestern Ontario, Lake St. Clair, Detroit, and the northwestern quadrant of Lake Erie (Fig. 1), and model grid cells were grouped by different surface type (i.e., rural, lake, and urban).

3 Results and discussion

3.1 Diurnal pattern of surface ozone

To examine the diurnal behavior of ground-level ozone over the lakes as compared to land, 2-h averages of ozone and their hourly change rates were calculated for the Buoy (representing the lake), Pelee Island (representing lake influenced conditions), and the groups of 11 rural and six urban sites (Fig. 2). Ozone measurements over Lake Erie were also made by a monitor installed for BAQS-Met on the Leamington–Pelee Island ferry, but due to its routine movement it was not possible to compute relevant diurnal averages. The ferry measurements were therefore averaged by trip times (approximately 90 min for each trip), for four sailings that were scheduled for the same departure times on all weekdays (Fig. 2a). Thus only its over-lake measurements were combined for comparison with and to further support the buoy and Pelee Island measurements. Although differences in ozone concentrations between land and lake were suggested in previous studies, this is the first time a detailed diurnal pattern and change rate are available at high temporal resolution and compared between lake, rural and urban sites.

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Figure 2 shows the averaged ozone diurnal pattern (2a) and change rates (2b) measured at the different site types, as well as the average ozone diurnal pattern (2c) and change rates (2d) predicted for corresponding surface types for the same period by the AURAMS model. The nighttime (00:00–06:00 Eastern Daylight Time [EDT]) ozone mixing ratio over the lake (buoy) is ~5 ppbv higher than the Pelee Island site, ~15 ppbv higher than the rural-site average, and close to 25 ppbv higher than the urban-site average. At 10:00 EDT, average ozone levels at the rural and urban sites match those of the buoy. Higher ozone concentrations over the lake at night are due to a lack of fresh NO emissions, particularly in comparison to urban areas, and lower deposition rates to water surface than land surfaces (e.g., Dye et al., 1995; Sillman et al., 1993). Examination of the morning (09:00 EDT) rate of increase in ozone at the buoy, shown in Fig. 2b, indicates that it is slower over water than over land (e.g., 1.5 vs. 3.4, 5.8, and 7.5 ppbv h⁻¹, for the buoy, Pelee Island, rural, and urban sites, respectively). These differences are due to greater vertical mixing over land which quickly replenishes the ozone at the surface in the morning, particularly over the urban areas where ozone experiences the greatest depletion at night. This behavior suggests that the nighttime differences between lake and land are limited to a shallow layer above the surface since within a few hours after sunrise mixing from aloft results in a minimum in spatial variability in surface ozone from 10:00–11:00 EDT.

Previous studies have shown that the break up of the surface inversion and vertical mixing with the ozone residual layer aloft has a significant impact on morning build up of ground-level ozone, whereas photochemical production processes augment surface ozone levels later in the day (Zhang and Rao, 1999). Zhang and Rao (1999) found comparable summer mean ozone build-up rates at a rural site, with the highest rates (~6.5 ppbv h⁻¹) at 10:00 EDT, compared to 10 ppbv h⁻¹ at 11:00 EDT at an urban site in New York City. Kleinman et al. (2002) used a chemical box model to estimate ozone build-up rates at five large cities in the United States. The authors found the highest rates of 11.3 ppbv h⁻¹ in Philadelphia, PA and Houston TX, compared to 6, 4.3, and 3.5 ppbv h⁻¹ in Nashville, TN, New York City, NY, and Phoenix, AZ, respectively. In this

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



study over southwestern Ontario, measured rates of change of ozone mixing ratio at the urban sites (Fig. 2b) are between 8 ppbv h^{-1} (at 09:00 EDT) and -6 ppbv h^{-1} (at 21:00 EDT).

To compare the measured diurnal cycles of ozone to those predicted by AURAMS, an analysis sub-domain of the 2.5-km model grid was selected over the study region (Fig. 1). The AURAMS mean ozone diurnal pattern for the lakes versus land surfaces in the region (Figs. 2c, d) is similar to the measurements, with a $\sim 15 \text{ ppbv}$ difference at night and build-up rates of 2.7 and 6 ppbv h^{-1} over lake and land surfaces, respectively, at 09:00 EDT. Urban areas (i.e., mainly Detroit) over the modeled region show a higher build-up rate of 10.8 ppbv h^{-1} in the morning (10:00 EDT) compared to the maximum rate of 7.5 ppbv h^{-1} measured at the urban sites an hour earlier, as well as faster loss rates at night (-13 ppbv h^{-1} at 20:30 EDT) compared to measurements (-6 ppbv h^{-1} at 21:00 EDT). These differences might be due, at least in part, to the six urban sites being more influenced by traffic NO_x emissions than the part of the model domain that is assigned to the “urban” group (cf. Fig. 4b).

From 23:00 to 06:00 EDT Fig. 2b shows a general trend of decreasing measured ozone loss rates. In contrast, the model predicts an almost constant loss rate with a magnitude of about 2 ppbv h^{-1} . The result of these differences in night time change rates is lower predicted ozone mixing ratios in the morning (06:00 EDT) compared to measurements. These lower model values at sunrise are then compensated by higher predicted versus measured change rates in the morning, particularly over rural (8 vs. 6 ppbv h^{-1} , respectively) and urban (11 vs. 8 ppbv h^{-1} , respectively) areas. As a result, the predicted daily maxima in the afternoon are higher than measured for urban areas (63 vs. 50 ppbv , respectively), but are in good agreement for rural areas ($\sim 58 \text{ ppbv}$).

Given the limited ability to obtain systematic three-dimensional measurements, particularly over the lakes, and after showing reasonable agreement between measured and predicted average ozone levels near the surface in the study region in Fig. 2, the AURAMS output was examined to gain more insight into the vertical structure of ozone over the different surface types of rural, lakes and urban and to examine the suggestion

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

that higher ozone levels over the lakes are limited to a shallow layer above the surface. Time-height cross sections extracted from the AURAMS mean hourly ozone fields for the buoy operational period (26 June–10 July), averaged further over the three surface types in the sub-domain are displayed in Fig. 3a–c. While overall relatively small differences are predicted throughout much of the vertical domain between the rural and lake surface types (Figs. 3a and 3b, respectively), not surprisingly, the model produces lower ozone levels over the urban areas within about 200 m of the surface at night. In contrast, according to the model there are considerably higher levels within the lower 1500 m over the urban areas during the day (Fig. 3c).

Quantitative model estimates of the differences in the vertical structure of ozone between different surface types is shown in Fig. 3d–f, together with a closer examination of the lowest vertical levels (outlined with the red frames in Fig. 3d–f) shown in Fig. 3g–i. Confirming the interpretation of Fig. 2 above that the significant land-lake differences are limited to a shallow layer, Fig. 3d shows small differences (less than ± 2 ppbv) in ozone levels over land vs. lake at the 200–1500 m levels in the early morning. Near the surface, however, differences are as high as 10–12 ppbv, but rapidly decrease to 4 ppbv at 100 m (Fig. 3g). A detailed case study presented in Sect. 3.3 presents observations confirming this vertical structure and further detailing its impact on observed surface ozone. The largest contrast in the time-height cross sections is between the urban and lake surface types (Fig. 3f and i) near the surface at night (21:00–04:00), with the urban area being 14–20 ppbv lower than the lakes, in good agreement with observations shown in Fig. 2a, of about 25 ppbv difference between the buoy and urban sites at night. These night time differences are the result of two loss mechanisms near the surface that act at different intensities over land and over the water. First, greater NO_x emissions over land (and over urban areas in particular) result in stronger ozone titration. Second, ozone has higher deposition velocities over land than over water.

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3.2 Spatial pattern of mean ozone

The scarcity of continuous (in time and space) air pollution observations over the Great Lakes generally reduces the examination of their spatial distribution to short term field campaigns where multiple monitors are used around the lakes (e.g., Fast and Heilman, 2005), and/or from aircraft measurements are taken above the region (e.g., Hastie et al., 1999). In an attempt to bridge this gap and examine the spatial distribution of ozone over the study region, ozone mixing ratios from AURAMS for the entire 23-day BAQS-Met intensive observing campaign (June 20 – July 10) were calculated. The spatial distributions of mean surface ozone mixing ratios for different times of day are shown in Fig. 4a. Modeled surface ozone mixing ratios are higher over the lakes as compared to their surroundings at all times of day (Fig. 4a1), but the difference is more pronounced at night with 10–15 ppbv difference between 21:00 and 06:00 EDT. This result is consistent with the actual measurements shown in Fig. 2a. However, the output from AURAMS indicates that ozone levels are not homogeneous over Lake Erie. This difference is most pronounced over the south-central part of Lake Erie (between Cleveland and Wheatley), where during the study period daily mean ozone levels reached a high of about 50 ppbv, compared to 41 ppbv in southwestern Ontario. Note that particularly low ozone mixing ratios of 20–30 ppbv are predicted over the urban regions of Cleveland and Detroit, where emission levels are typically higher. It should also be noted that mean ozone mixing ratios over Lake Huron are lower than over Lake Erie. With the southwestern end of Lake Erie being ringed by the urban centers of Detroit, Toledo, and Cleveland (Fig. 1), this suggests that proximity to pollution emission sources is also an important factor. Fast and Heilman (2005) presented a similar finding for Lakes Erie and Huron in their modeling results.

Figure 4 also shows a vertical cross section between the surface and 3000 m a.g.l. of ozone (4b) and vertical wind (4c) along a transect across Lake St. Clair and Lake Erie, marked A-F in Fig. 4. The simulated mean ozone cross section shows a reservoir layer persisting after 21:00 EDT (Fig. 4b5), centered between 600 and 1000 m a.g.l.,

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

over Lake Erie and Cleveland. At the start of the overnight period (i.e., 00:00 EDT) the model suggests that higher ozone levels over Lake Erie are not limited to the lowest layers, but extend to about 1500 m a.g.l. By early morning (06:00 EDT) the differences over land and over the lakes are much reduced and are limited to a shallow layer of less than 200 m a.g.l. In contrast, above about 500 m at this time the horizontal variation in ozone is small, with no apparent lake-related features. The main differences in the early morning are over the large emission areas (Cleveland and west of Lake St. Clair) due to ozone titration by NO that is emitted into a shallow nocturnal surface layer.

After noon, when photochemical production is peaking, higher ozone over the lake (60–70 ppbv, ~10 ppbv higher than over land, Fig. 4b4) is limited to a shallower layer of about 100 m over Lake Erie. This is most evident over the US side of the lake, while on the Canadian side, ozone mixing ratios are lower (50–60 ppbv). This pattern is likely related to the assumed spatial distribution in emissions over the lake, where emissions on the US side are spread over the southern part of the lake while emissions on the Canadian side are allocated along the main shipping lane (see Makar et al., 2010). At the same time, over Lake St. Clair high ozone levels of about 70 ppbv extend up to 1500 m a.g.l. following the vertical extent of the coupled lake and urban circulation (Fig. 4c4). In the evening (21:00 EDT) while ground-level ozone is reduced over land, it is 10–15 ppbv higher over Lake Erie, up to about 2000 m a.g.l. This difference is noticeable through the night and until early morning, and is due to differences in the rate of surface deposition of ozone between land and water, and the increased impact of NO_x titration of ozone after sunset over the land, as mentioned earlier.

High ozone levels are also noticeable near the surface during the day over the larger metropolitan areas of Detroit and Cleveland (Fig. 4a4), with values going from 35 ppbv at the downtown areas (e.g., point E at the cross section in Fig. 4a4) to above 60 ppbv in the cities' outskirts. Unlike the high ozone predicted close to the surface of Lake Erie mentioned earlier, the high values over land extend up to about 1500 m a.g.l. (Fig. 4b4), corresponding to the vertical extent of the breeze circulation cell, with rising air over land and descending over the lakes (Fig. 4c4). However, the high values over the lake

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

are more widespread compared to the cities, particularly over the southwestern and central parts of Lake Erie, though emission sources over the water are not high.

It is interesting to note that during the night, low ozone mixing ratios over Detroit extend to hundreds of meters above the ground (at 00:00 EDT over point B in Fig. 4b2), while over Cleveland the titrated layer is shallower. This difference in the vertical extent of ozone titration could be explained in part by the vertical motion over Detroit and over the adjacent part of Lake St. Clair at this time (over point B in Fig. 4c2). However, similar vertical flow is also predicted by the model over Cleveland (Fig. 4c2 over point E), without a similar effect on ozone above the surface. Additional measurements are needed to be able to determine the extent to which these features and city-to-city differences occur and their implications on the local air quality.

3.3 Case study: 6–7 July 2007

The analyses in the two previous subsections have examined the average temporal and spatial patterns of ozone in the study area for the BAQS-Met intensive observing period. In order to fully understand the complex influence of local terrain-driven circulations in this area, however, it is necessary to consider a day-specific case study. Based on a review of the BAQS-Met air-quality and meteorological measurement data, the 6–7 July period was identified as being an interesting example of the impact of local circulations associated with lake and land breezes on the time evolution and spatial distribution of ozone and other pollutants. The following paragraphs describe this period focusing on conditions at the Harrow supersite, located 30 km southeast of the Detroit-Windsor urban complex. The Harrow area is affected by both Lake Erie and Lake St. Clair, as well as Detroit-Windsor. Measurements are also presented from additional surface sites, the Twin-Otter aircraft and the Ridgetown tetheredsonde to obtain a more complete understanding of this case.

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.3.1 Meteorology

A high-pressure system approached the BAQS-Met study area from the west on 6 July, and passed south of the study region from SW to S on 7 July. The ridge passage caused the light to moderate gradient flow to shift from northwesterly on 6 July to westerly on 7 July in the morning. In addition, as shown in Fig. 5, superimposed on these synoptic features during this period were lake-induced winds and their associated sequence of mesoscale lake-breeze fronts. The positions of these fronts, which were determined for every hour of each day during BAQS-Met (Sills et al., 2010), were based upon multiple information sources including the meso-network and routine meteorological observation stations, satellite and radar images.

The meso-analysis in Fig. 5a for 6 July 12:00 EDT, shows that a portion of the Lake Erie lake-breeze front (marked A and A1 in Fig. 5a) had just formed on the northwestern and western shores and was moving inland and northward south of Harrow. By 20:00 EDT (Fig. 5b) this lake-breeze front had passed over Harrow, reached the south shore of Lake St. Clair and stretched from Windsor towards the east and then southeast to Lake Erie. At this time a large part of extreme southwestern Ontario was experiencing southerly winds under an air mass that originated over Lake Erie, while northerly winds were prevalent over the reminder of the domain. The lake-breeze flow was moderately deformed by the gradient winds so that lake-breeze fronts did not form on the downwind side (south shores) of the lakes (Sills et al., 2010). At 20:00 EDT on July 6, the portion of the Lake Erie lake-breeze front that stretched from Windsor and to the south (marked A2 in Fig. 5b), was retreating eastward towards Harrow as the synoptic flow had swung around from northwesterly to westerly. A few hours later at 00:00 EDT on 7 July, a third lake-breeze front has almost reached Harrow from the northeast, this time generated earlier in the day at the southern end of Lake Huron (marked B in Fig. 5b and c).

Thus, during this period of less than a day, the detailed analysis of the meteorological conditions, enabled by deployment of the meso-network, revealed that air was transported to Harrow first from the northwest, then the south, then the west, and then

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the northeast (Fig. 6a). In addition, this analysis showed that size and location of the land area impacted by the “Lake Erie influenced air” changed continuously during the day and both Lakes St. Clair and Huron played a role in the mesoscale patterns as they evolved. These relatively dramatic meteorological changes provided a unique opportunity to examine in detail how the lakes influence ozone and other air pollutants. Insights gained through such analyses provide additional measures for evaluating the capabilities of the high resolution GEM and AURAMS models and for better assessing the true nature of the higher ozone occurring over and near the lakes. The discussion below will thus focus on how the chemical characteristics of the air masses changed with the lake breeze front transitions described above and in Sills et al. (2010).

3.3.2 Air pollution

Figure 6 shows concurrent time series from the Harrow supersite of wind speed and direction (Fig. 6a), O_3 , SO_2 and CO mixing ratios (Fig. 6b), nitrogen oxides (NO , NO_2 , NO_x and NO_y) and the toluene to benzene ratio (TBR, Fig. 6c) for 6–7 July. Also shown is O_3 at three BAQS-Met sites east of Harrow (Wheatley, Leamington, and the Lake Erie buoy, Fig. 6d, see Fig. 1 for locations) for 6–7 July. As NO_2 measurements from the supersite were not available for the morning of 6 July, measurements taken by the co-located instrument operated by the Ontario Ministry of Environment (OME) are presented in Fig. 6. Toluene and benzene measurements from the PTR-MS were used as one measure of the photochemical age of the air masses affecting the site (Roberts et al., 1984). Since both species are emitted from anthropogenic sources (mostly vehicle exhaust), and toluene is approximately four times more reactive with OH radicals, the lower the toluene/benzene ratio the more aged the air mass is. A ratio of about 4 indicates a freshly emitted air mass, while a ratio of about 1 marks an air mass emitted approximately 1–2 days earlier (Vlasenko et al., 2009). Below is a description of the different air masses identified during the 40 h period of 6 July 08:00 EDT to 8 July 00:00 EDT of the case study. For better clarity, certain time periods are marked in Fig. 6 in numbers (1–11) and referenced in the following text as different

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



stages.

Stage 1 (08:00–10:05 EDT): A NNW flow (Fig. 6a) brings relatively fresh pollution to Harrow in the early morning hours, with CO levels of 400 ppbv, NO of 5 ppbv and NO_y of 20 ppbv. The high TBR of almost 3 and a wind speed of about 3–5 m s⁻¹ (11–18 km h⁻¹) from the NNW support the assumption that these are polluted air masses arriving in an almost direct route from the Detroit-Windsor area.

Stage 2 (10:05–12:50 EDT): A sharp drop of about 200 ppbv in CO mixing ratios at 10:00 EDT, along with significant changes in ozone, SO₂, NO and NO_y mark a different air mass. This air is more photochemically aged, as seen by the lower TBR of 2.3–1.5, and with the winds having a more easterly component (wind direction of 0–45 deg, Fig. 6a), suggests an air mass that followed a longer path from Detroit to the east (i.e., north of Harrow) in the morning and now arrives at Harrow from the north, allowing more time for photochemical activity. Increases in mixing height due to the surface heating at this time, would also contribute to the above mentioned decrease in primary pollutants. However, the decrease in TBR at that time (from 2.3 to 1.5), being a ratio between ambient levels of two species and therefore not sensitive to changes in mixing height, suggests this air mass had also undergone more photochemical processing.

Stage 3 (12:50–13:25 EDT): The Lake Erie lake-breeze front passing at Harrow at 12:50 EDT (marked A1 in Fig. 6 between stages 2 and 3) causes an abrupt change in wind direction from northwesterly to southerly and an increase in wind speed from 5 to 7 m s⁻¹. The frontal passage coincides with a sharp drop in ozone, a dramatic increase in NO_y (from 7 to 21 ppbv), NO (from 1 to 3 ppbv), NO₂ (from 3 to 13 ppbv), and the TBR rises, which are all indicative of a fresher air mass. Indeed Fig. 6b shows that CO also experienced an abrupt jump, from 202 to 491 ppbv, and SO₂ increased from 0.2 to 5.6 ppbv. During Stage 3 there is also a rapid recovery in the ozone mixing ratio after the drop. This is either due to a change in its build-up rate from 18 ppb h⁻¹ in Stage 2 to 25 ppb h⁻¹ as a result of the availability of precursors and/or spatial heterogeneity in this new air mass with “pools” of higher NO locally destroying ozone, which would have been the cause of the drop at the start of Stage 3.

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Stages 4 (13:25–13:50 EDT) and 5 (13:50–15:10 EDT): During the short time period labeled as Stage 4 the wind continues to blow from the south, ozone increases more slowly and there are small decreases in CO, NO_y, NO₂ and NO. These variations, as well as the observations during Stage 5, tend to indicate that there was considerable spatial heterogeneity in the air mass behind the lake breeze front. The peak in SO₂ during Stage 5, which is about 10 ppbv higher than the peak in Stage 3, and the maximum CO mixing ratio near this time further suggest that this fresh air mass contained a complex, variable mixture of primary pollutants.

The second intrusion of relatively fresh emissions at Harrow starting Stage 3 is surprising given the apparent origin of the air over Lake Erie. The wind directions over the region during the morning hours, however, suggest that this air mass likely originated near Detroit-Windsor. With NNW winds in the early morning these urban/industrial emissions could have followed the Detroit River traveling to the area of Lake Erie south and west of Harrow. Then this “pool” of relatively fresh emissions would have been transported northward across Harrow (at 12:50 EDT) into extreme southwestern Ontario behind the advancing lake-breeze front. The amount of temporal variability in the pollutant levels after the frontal passage (Stages 3, 4 and 5), which were all during southerly wind off the lake, suggests that the pollutants were recently emitted and experienced limited horizontal and vertical mixing. Transport early in the morning from Detroit-Windsor and then accumulation in the stable air over the lake could have led to this scenario. It is interesting to note that, with the exception of SO₂, pollutant levels observed in Stages 3 and 5 were similar. This raises the possibility that the air mass in Stage 4 was a short “interruption” to a larger air mass containing ample ozone precursors and hence experiencing relatively rapid ozone formation.

Stage 6 (15:10–17:30 EDT): A sharp increase of 9 ppbv in ozone mixing ratios in two minutes (from 78 to 87 between 15:04 and 15:06) that is followed by a drop in CO, SO₂ and NO_y mixing ratios a few minutes later marks another dramatic change of air mass. Ozone levels during this stage maintain an almost constant and high value (85–90 ppbv), indicating an air mass that has been fully processed. This is supported

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

by the low NO (<1 ppbv), the large ratio of NO_z to NO_x and minimum values in the TBR of 0.8.

5 Stage 7 (17:30–19:10 EDT): A change in wind direction from S to SW (Fig. 6a) and a decrease in ozone from 90 to 70 ppbv at 17:30 that coincides with changes in CO, SO₂, NO_y, NO_z and NO₂ (Figs. 6b and c) mark another change in air mass. This air also maintains a near-constant ozone level for almost two hours that together with the low TBR and high NO_z/NO_y indicates this air mass was also photochemically aged. However, its origin and history were clearly different than in Stage 6 given that less ozone had been produced. Figure 5b shows that during the period of southerly flow
10 over the region (Stages 3-7) the Lake Erie air mass moves northward to the south shore of Lake St. Clair. The meso-network station at Lighthouse Cove (LIG in Fig. 1) experienced its peak ozone at ~20:00 EDT just after the front (A) passed over, thus confirming the existence of higher ozone in this air mass, which covered a distinct area of about 2000 km² at 20:00 EDT.

15 Stage 8 (19:10–20:35 EDT): As the retreating Lake Erie lake-breeze front passes Harrow for the second time from west to east (marked A2 on Fig. 5b and on Fig. 6) there was a large drop in ozone of 40 ppbv in 35 min (between 19:50–20:25 EDT, Fig. 6b). This is accompanied by changes in other species, as well as wind speed and direction and are hypothesized to be the result of complex three dimensional flows in this frontal zone. This complexity is further indicated by the dramatic oscillation in ozone, dropping first by about 10 ppbv then rising by 25 ppbv and then halving from 80 to 40 ppbv. A detailed description of the meteorological and pollutant changes across lake breeze frontal zones in this region is given in Hayden et al. (2010), based on BAQS-Met measurements taken at multiple heights on board the Twin Otter aircraft.
20

25 Stage 9 (20:35–07:40 EDT): After the front (A2) passes east of Harrow, a westerly synoptic flow develops over the SW Ontario region. Wind speeds during this overnight period are light to stagnant, but gradually push the air mass with maximum values in ozone, east and south back towards Lake Erie. A fresher air mass moves over Harrow, as indicated by the increasing TBR after 20:00 EDT, a decrease in NO_z and an increase

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

in NO₂ from 5 ppbv at 21:00 to 14 ppbv at 04:00 EDT (Fig. 6c). The westerly winds in the first few hours after the frontal passage and the analysis of the local meteorology in Section 3.3.1 suggest the Detroit-Windsor region as the source for the pollution at this time.

The light wind flow pushes the ozone-rich air eastward, where it is marked by a 40 ppbv drop in ozone mixing ratio at Leamington (LEA in Fig. 1) between 21:00–22:00 EDT. This is shown in Fig. 6d along with ozone levels over the lake measured by the buoy and at the Wheatley meso-network site located further east (marked ECO and WHE in Fig. 1, respectively). Throughout the evening there is also continued south and eastward movement of a third lake-breeze front (Lake Huron front), which eventually affects the region. At 20:00 EDT this front (marked B on Fig. 5b) stretches from rural areas in Michigan north of Detroit, through Lake St. Clair and then towards Lake Erie. Eventually, it pushes southwestward and passes Leamington at about midnight with ozone levels dropping further and stabilizing at about 10–20 ppbv. According to the ozone in Fig. 6b this front appears to have passed over Harrow at around 04:00 EDT on 7 July.

The result of these two later lake-breeze frontal passages (A2 and B) is a reduction in ozone mixing ratios over land in southwestern Ontario. However, high mixing ratios remain over Lake Erie since the Lake Huron front does not penetrate far enough south. The difference between the two air masses is clearly seen by comparing Leamington ozone to the observations at the buoy (Fig. 6d). At 00:00 EDT (7 July) the mixing ratio was 70 ppbv at the buoy and 20 ppbv in Leamington, with the large difference between these two locations persisting until 05:30 EDT. The location of the edge of the ozone-rich air mass is evident at Wheatley, where ozone mixing ratios fluctuate during the night between the Leamington and Buoy levels. This boundary between the air masses was also observed by the Twin Otter aircraft as it flew over Lake Erie during the early morning hours of 7 July. The aircraft passed approximately 5 km NE of the buoy at 150 m.a.g.l. at 05:20 EDT (Fig. 7a) and detected a sudden steep drop in ozone from 70 ppbv to 20 ppbv and a simultaneous increase in NO₂ (Fig. 7b). The

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

vertical differences in ozone over land and over Lake Erie at this time are described in the following Sect. 3.3.3 and the oscillations in ozone observed at several sites are discussed in Sect. 3.3.4.

5 Stage 10 (07:40–18:50 EDT, 7 July): Low wind speeds of $0\text{--}4\text{ m s}^{-1}$ (Fig. 6a) in the morning, an increase in NO from 2 to 4 ppbv (Fig. 6c), a sharp increase from 300 to 800 ppbv in CO and from almost 0 to 5 ppbv in SO₂ (Fig. 6b) suggest a different and more polluted air mass arriving at Harrow. These mark the beginning of another stage, with a typical morning build-up of ozone from 19 ppbv at 08:00 to 55 ppbv at 13:00 (i.e., 7.2 ppbv h⁻¹, comparable with the average 5.8 ppbv h⁻¹ for 11 rural sites in the morning shown in Fig. 2). This stage continues through the day with a high of 60 ppbv in ozone persisting through the evening. During this period wind speed increases to 6 m s^{-1} and gradually shifts from NW to W and southerly flow (Fig. 6a). SO₂ peaks of 12 ppbv at 15:45 EDT and 27 ppbv at 17:35 EDT, both associated with high NO_y and low (<1) TBR, are observed during the afternoon southwesterly flow. These suggest a local to regional transport of polluted air crossing over Lake Erie, potentially related to a large power plant in the west shore of the lake.

20 Stage 11 (18:50–00:00 EDT, 7 July): Uncharacteristically high ozone after sunset at Harrow (Fig. 6b) and other sites in the region (Fig. 6d) is observed at this stage. There was also a higher TBR ratio compared to the previous stage, reaching 2, and an increase in CO, NO₂ and NO_y after 22:00 EDT. The regional spread of the high ozone, being measured at multiple sites in the study region (Fig. 6d), and the persistent and moderate southwesterly winds (Fig. 6a) suggest that a large regional airmass with elevated ozone was being transported from southwest of Lake Erie. At the same time, some more local sources were likely adding to this airmass, as evidenced by the higher TBR and primary pollutant levels. Ozone remained high across the region after sunset due to turbulent/mechanical mixing induced by moderate winds, which continued to bring the higher ozone aloft to the surface and tended to homogenize the levels throughout the region.

25 The detailed description above demonstrates the complexity of the factors control-

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ling ozone levels in southwestern Ontario observed in a short period of 40 h. Pollution sources affecting the region can be from long range transport (Stage 11) or relatively local emissions (e.g., Detroit-Windsor) that are transported in complex pathways by land-lake breeze winds (e.g., stages 6 & 7), while undergoing photochemical processing, and by larger scale flows (e.g., stage 9). Furthermore, the air arriving with the lake breeze is often heterogeneous in its composition and photochemical age, as demonstrated in stages 3–7.

3.3.3 Vertical structure

Given the complexity described above in surface ozone for the case study, the vertical structure of ozone in the lower troposphere is examined in the following paragraphs. Measurements supporting the limited vertical extent of enhanced ozone levels over Lake Erie discussed in Sects. 3.1 and 3.2, are found in the early morning of 7 July. At this time vertical profiles of ozone were obtained by the aircraft (Fig. 8) and by tethered sonde launches at Ridgeway (Fig. 9). A temperature profile taken by the aircraft over Lake Erie at 05:40 EDT ascending from Pelee Island (Fig. 8b and marked P1 in Figs. 7a and 8a; timing corresponds to Stage 9 described above) shows a thermal inversion layer between 200–235 m a.g.l. The inversion separates a low-ozone layer near the surface with about 25 ppbv that is also NO₂-rich (10 ppbv) from the regional ozone levels of 40–45 ppbv observed above the inversion along with lower NO₂ levels. Over land, however, the thermal inversion starts at the surface due to radiative cooling overnight (profiles P2–P7 and P9 in Figs. 7, 8c–h and 8j). This results in higher NO₂ (e.g., 20 ppbv over Windsor in P4 and over Detroit in P7, Fig. 8e and h, respectively) and consequently much lower ozone levels near the ground (e.g., about 5 ppbv near Windsor, P4, Fig. 8e). However, ozone is rapidly increasing with height within the surface inversion layer of about 150 m a.g.l., and above that layer it is over 40 ppbv, the same as over the lake at this altitude.

Tethered sonde launches at Ridgeway on the same morning support these findings and show a surface inversion up to about 100 m a.g.l. at 06:00 EDT (Fig. 9b), with

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ozone levels increasing from zero to 50 ppbv with altitude, and stabilizing at 40 ppbv above 300 m. The air mass sampled by the early morning launches P1-P6 (Figs. 9b-g, respectively) is the same as the one described in Stage 9 at Harrow, dominated by a westerly large scale flow. Thus, the uniqueness of the conditions over the lake (i.e., higher ozone) is limited to a shallow surface layer (<200 m), while the reservoir of ozone aloft is similar across this larger region with no evidence of the land-lake difference.

Of special note is the difference in the ozone vertical profile below the inversion layer between the lake and land before sunrise. While ozone mixing ratios are almost constant over the lake in this layer (about 25 ppbv, Fig. 8b), over the adjacent land (Fig. 8c) the observed mixing ratios are much lower near the surface (17 ppbv at 47 m) but rapidly increase (34 ppbv at 73 m) with altitude. These vertical changes result from the different thermal structure over the two surfaces during the early morning hours. The surface inversion over land suppresses any vertical mixing and allows much of the ozone near the ground to be titrated by nitrogen oxides emissions or be deposited to the surface.

Subsequent tethered sonde launches in that morning show that ozone levels are rapidly increasing after sunrise at the lowest surface level from 0 to 16 ppbv between 06:00 to 07:34 (Fig. 9b to e), while decreasing at the 50 m a.g.l. level from 33 to 20 ppbv. These changes correspond to changes in the temperature profile with a breakup of the surface inversion occurring between 06:51 and 07:34 EDT (marked P3 and P4 in Fig. 9d and e, respectively). More evidence of this process is found in the morning change rates in surface ozone at the monitoring sites over lake and over land. While differences of ~10 ppbv in surface ozone levels between lake (Buoy) and land (Wheatley) are observed in the early morning hours (07:00 EDT, Fig. 6d), ozone change rates in the morning are faster over land than over the lake, so that by 09:00 EDT both sites reach 35 ppbv of ozone. The early morning change rate in surface ozone over land is thus shown to be the result of vertical mixing with an ozone-rich layer at a higher altitude.

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

An interesting feature observed at Ridgetown is a decrease in ozone mixing ratios in a shallow layer above the surface. This layer starts at 250 m a.g.l. (06:09 EDT) and descends to 78 m a.g.l. by 07:32 EDT (Figs. 9b and 9e, respectively). A similar feature observed over the Chatham-Kent airport at 200 m a.g.l. by the Twin Otter at 07:16 EDT (Fig. 8j and marked P9 in Fig. 7a), and the NNW wind over the region suggest the presence of a fan shaped NO_x plume, possibly from the Lambton power plant south of Sarnia, ON (Fig. 1). Cases of NO₂ aloft over Ridgetown are discussed by Halla et al. (2010).

A second Twin Otter flight made later on 7 July from 13:30 to 16:20 EDT examined the vertical cross section over Lake St. Clair up to 2600 m a.g.l. A moderate westerly flow dominated the region in the late morning (also observed at Harrow, Fig. 6a), transporting ozone and its precursors from the Detroit region over Lake St. Clair. By noon a stationary Lake St. Clair lake-breeze front is established over the southern and western shores of the lake and stalls for several hours until evening (Fig. 5d), while the regional flow changes to WSW (Fig. 6a). A plan view of the flight (Fig. 10a) and a time series (Fig. 11a) show an increase in in ozone mixing ratios from about 55 ppbv to 65 ppbv as the aircraft crosses the shore at the southern side of the lake, both at 500 m and 700 m a.g.l. at 14:40 and 14:59 EDT, respectively. The stationary Lake St. Clair lake-breeze front located on the shore at that time marks the boundary between two air masses. On the northeastern side of the lake, high ozone values extend some 20 km out of the lake at 300 m, 500 m, and 740 m altitudes a.g.l. (Fig. 10b). At the 1400 m level, high ozone is more limited in its horizontal extent and does not go beyond the lake. The vertical extent of the high ozone values reaches 1900 m a.g.l., where a subsidence inversion layer is encountered, and ozone mixing ratios decrease at a steady rate (Fig. 11). These measurements suggest the presence of a dome-shaped high ozone region over Lake St. Clair at noon, that is shifted and stretched at lower levels to the northeast, with peak values of about 85 ppbv at the lake surface. Other model simulations (Makar et al., 2010) examine the formation mechanism of this feature in detail: the coupling of emissions from Detroit, the city's heat island circulation with the

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Lake St. Clair subsidence flow, and possible recirculation of pollutants over Lake St. Clair.

3.3.4 Nocturnal ozone oscillations

Night-time oscillations in ozone mixing ratios were observed at several monitoring sites during the BAQS-Met campaign. These were characterized by amplitudes of 20–40 ppbv and time periods of 15–40 min, changing from site to site and continuing for several hours. The oscillations occurred at multiple sites and different nights (e.g., 24–25 June, 29–30 June and 4–5 July), but had the highest amplitude and longest duration on the night of 6–7 July. Figure 12 shows meteorological measurements (wind speed and direction, temperature, dew point temperature and relative humidity) as well as ozone at four ground sites in SW Ontario where oscillations were observed on the night of 6–7 July (sites locations are indicated in Fig. 1): Bear Creek (mostly between 02:00–03:00 EDT, Fig. 12a), Lighthouse Cove (between 21:00–01:30 EDT, Fig. 12b), Leamington (between 21:00–01:00 EDT, Fig. 12c) and Wheatley (between 21:00–04:00 EDT, Fig. 12d). The oscillations occurred during periods with calm winds at the surface ($0\text{--}3\text{ m s}^{-1}$, red line in top panels in Fig. 12) and are associated with oscillations in temperature and relative humidity (black and blue lines, respectively, in bottom panels of Fig. 12), so that peaks in ozone correspond to peaks in temperature and lows in relative humidity. Dew point temperature (shown in green in Fig. 12) shows very little change related to the oscillations, with the exception of Leamington at certain times (e.g., 22:30 and 23:30 EDT in Fig. 12c). It is not clear whether these changes are related to the oscillations in ozone or to condensation of water vapor to the surface.

Increased night-time ozone levels were identified in other studies at different locations, and explained by nocturnal low level jets (e.g., Corsmeier et al., 1997) and stratospheric intrusions (e.g., Mavrakis et al., 2010). However, these studies observed either a secondary night-time maxima or an abrupt increase in ozone mixing ratios, not oscillatory behavior. The difference in observations between the BAQS-Met study described here and previous studies could be the result of different physical mechanisms

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



producing high nocturnal ozone, or in part simply due to the finer temporal resolution of the ozone measurements taken during BAQS-Met, of 1 min (e.g. at Bear Creek) and 5 min (at Leamington, Lighthouse Cove and Wheatley) and of 1 min for meteorological parameters in all sites. In spite of the strong amplitude of the observed oscillations (e.g., 40 ppbv drop in 25 min at Wheatley), they would not necessarily appear in hourly averaged ozone data that is typically recorded by air quality monitoring networks because of the smaller time periods of 15–40 min.

A possible explanation for the oscillations could be that a vertical stratification occurs near the surface either due to Lake Huron breeze air mass presence over the region (as described in Sect. 3.3.1), that is confined to a very shallow layer above the surface, or by radiative cooling of the surface. This vertical structure is supported by the vertical profiles taken by the aircraft in the early morning over land between Leamington and Wheatley (Fig. 8c, location marked as P2 in Fig. 7), showing a surface inversion layer up to 200 m a.g.l. and by the tether sonde launch at Ridgeway at 06:00 EDT (Fig. 9b) showing a surface inversion layer up to 100 m a.g.l. Such a vertical structure would result in low ozone (due to titration and deposition) and cooler air near the surface and higher ozone and warmer air just above it. Stronger winds in the layer aloft could cause a wind shear between the two layers, resulting in turbulence and injection of air from aloft to reach the surface. This mechanism is similar to the nocturnal low level jets as described by Reitebuch et al. (2000) and Corsmeier et al. (1997). The caveat with this explanation is that an intrusion of a different air mass from aloft is expected to show different water content, resulting in a more significant change in dew point temperature than is observed in Fig. 12. GEM/AURAMS model results support the low level jets explanation, showing high wind speeds just above the surface on several occasions during that night. Figure 13 shows plan views at the surface (Fig. 13a, b) and cross sections of wind speeds above SW Ontario (Figs. 13c, d) at two times, 22:00 and 03:20 EDT. The model predicts high wind speeds just above the surface either as extensions of stronger winds above 1000 m a.g.l. at 22:00 EDT (Fig. 13c, marked i and j) or isolated from the stronger winds aloft such as at 03:20 EDT in Fig. 13d (marked k,

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

l and m). Strong gradients in wind speed from the surface to 100 m a.g.l. are predicted at these locations, with speeds increasing from 2 m s^{-1} near the surface to 6 m s^{-1} (Fig. 13c, near i and j) and even 8 m s^{-1} (Fig. 13d, near k, l and m). Furthermore, the heterogeneity in the horizontal gradients of the wind speed predicted by the model near the surface (e.g., near point D in Fig. 13a, b) and just above the surface at the same location (e.g., Fig. 13c, d at 200 m a.g.l.) would result in higher potential for the vertical mixing to occur over certain locations.

Other explanations are suggested for this behavior. Assuming that the oscillations are related to the Lake Huron breeze front described above in Stage 9, it is possible that this night-time front has a similar motion to that noted by Drobinski et al. (2007), with oscillatory behavior in its advance, and a buildup of pollutants (NO_x , resulting in ozone titration at night), just behind the front. However, the observations at SW Ontario occurred during the night, making it less likely for the breeze front to propagate by this mechanism. It is also possible that differential radiative cooling of the surface would cause parcels of warmer air near the surface to ascend, thus forcing air from higher levels to reach the surface, where it is observed as warmer and more ozone-rich air. The caveats with this explanation are that the surface layer would have to be very shallow to allow air from aloft to reach the ground in such a mechanism, and that only small oscillations are observed in the dew point temperature in Fig. 12 that correlate with the oscillations in ozone, suggesting that the water content at the two layers is the same, which is unlikely for air masses from different sources. Last, given the presence of a lake breeze front over the region on that night, as described in Sect. 3.3.1, it is possible that density currents induced by the front inject air from higher elevations to the surface, as described by Sun et al. (2002). However, as with the first explanation, this mechanism would also result in short bursts of strong winds measured at the surface, which are not observed in the measurements.

As none of the above conceptual models given for the oscillations is adequate and explains it in a satisfactory way, further study is needed to find their cause, extent and impact on surface ozone.

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4 Conclusions

The average ozone from the entire BAQS-Met intensive campaign period (20 June to 10 July 2007), both from ground-level observations and from AURAMS model output, support previous findings that near-surface ozone levels are higher over the southern Great Lakes than over the surrounding land. During the night and early morning, average ozone mixing ratios over the lake are ~15 ppbv higher than over rural sites and ~25 ppbv higher than urban sites. Vertical profiles taken over land and over the lakes, as well as differences in the average diurnal change rates of ozone over lakes, rural and urban surfaces and model simulations suggest that the higher ozone levels observed over the larger lakes are limited to a shallow layer of about 200 m deep. Differences in NO_x emissions/titration and ozone deposition velocities between land and water have a significant impact on ozone concentration contrasts between the two surfaces at night, and result in the higher ozone levels observed over water in the early morning. As soon as vertical mixing begins in the morning, ozone levels over land increase rapidly, and match the lake levels by 10:00 a.m. local time, so that in late morning until just after noon very little difference is observed. Just after noon, however, ozone levels over the lake are higher again by about 5 and 15 ppbv compared to rural and urban sites, respectively, as seen in both observations and model results.

Model simulations over the region show great spatial heterogeneity in ozone levels over the lakes, particularly over Lake Erie with higher levels over the southwestern end of the lake. Some of this heterogeneity is due to multiple emission sources in the region that are concentrated around southwestern Lake Erie, such as large urban centers (e.g., Windsor, Detroit, Toledo, Cleveland). An important cause of the heterogeneity in ozone over Lake Erie as well as the land areas in the region is the complex meteorology. Land and lake breezes from the different lakes, which form on the majority of days, interact with the larger-scale synoptic flow, as demonstrated for the 6–7 July 2007 case study. These flows transport ozone and its precursors over the region, including from emission sources to the lakes. The pollutants that are stored in the more stable at-

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

mosphere over the water at night experience limited fresh emissions and little removal by dry deposition. Comparison of modeled and measured ozone diurnal changes over the lake at night indicate that the model is simulating these processes reasonably well with no evidence that it is over-predicting nocturnal ozone over the lake. When the lake breeze starts blowing towards the land in the morning, a significant increase in ozone may then be observed over land. Values as high as 90 ppbv were observed during BAQS-Met over extreme southwestern Ontario. When primary air pollutants (e.g., CO, NO_x) also accumulate within shallow layers and in distinct pockets over the lakes at night (e.g., northwest Lake Erie) enhanced ozone formation on the subsequent day results. This process can lead to local emissions having a greater local impact (i.e., within 50 km) on surface ozone mixing ratios. The case study presented for 6–7 July 2007, shows that these situations, with enhanced levels of locally-emitted precursors, are distinct from other cases of higher ozone over the lake when the accumulation of ozone has occurred over multiple days so that the available precursors have been depleted and the mixing ratio in ozone ceases to increase.

Considerable differences in temperature and ozone levels are observed in the vertical over land and over water in the early morning hours. While a surface inversion over land causes a very low ozone layer close to the surface (particularly over urban areas with mixing ratios reaching 0 ppbv), there is a rapid increase to background levels above the inversion layer. In contrast, over the lake, ozone mixing ratios are fairly constant below the inversion up to 200 m a.g.l. and rapidly increase to background levels above the inversion layer. These differences in the vertical profile result in higher ozone levels over the lake at the lowest layers, but lower below the inversion layer.

The mean ozone mixing ratios of the AURAMS model simulations for the entire study period show a large ozone reservoir layer over Lake Erie, extending to the land over Cleveland, which is centered between 600–1000 m a.g.l. Furthermore, a vertical wind ascending over Detroit-Windsor and descending over Lake St. Clair is predicted during the day. The vertical circulation transports ozone and its precursors up to 1500 m a.g.l., resulting in high ozone levels in the entire column.

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The high temporal resolution of the measurements examined in this study allowed the observation of nocturnal oscillations in ozone. These were recorded on several nights, and covered a relatively large spatial domain. Possible explanations to this behavior include low level jets that inject ozone-rich air from higher levels to the surface, the dynamics related to a lake front progression, heterogeneity in surface cooling that result in vertical mixing of air masses and density currents. Further study is needed to establish the mechanism causing this behavior and determine the impact of such oscillations on ground level ozone, examining their spatial extent and frequency.

Acknowledgements. The authors acknowledge with thanks the many BAQS-Met data providers, including the Ontario Ministry of the Environment for the measurements conducted at Ridgetown and on the Peele Island, the researchers from the University of Toronto and York University for the measurements at Harrow and Ridgetown, the Owen Sound Transportation Company for use of their MV *Jiimaan* ferry and the National Research Council of Canada for the instrumented Twin Otter aircraft. Special thanks are due to Frank Froude for the tethersondes data, Andrew Sheppard for the IADN buoy data, Tomasz Stapf for operating the mesonet sites, Kurt Anlauf and Jason O'Brien for their effort with the Twin Otter O₃ and NO₂ measurements. Ilan Levy gratefully acknowledges support by the post-doctoral fellowship from the Environment and Health Fund, Jerusalem, Israel.

References

- Alper-Siman Tov, D., Peleg, M., Matveev, V., Mahrer, Y., Seter, I., and Luria, M.: Recirculation of polluted air masses over the East Mediterranean coast, *Atmos. Environ.*, 31(10), 1441–1448, 1997.
- Alpert, P. and Getenio, B.: One-Level Diagnostic Modeling of Mesoscale Surface Winds in Complex Terrain .1. Comparison with 3-Dimensional Modeling in Israel, *Mon. Weather Rev.*, 116(10), 2025-2046, 1988.
- Anlauf, K., Lusi, M., Wiebe, H. and Stevens, R.: High ozone concentrations measured in the vicinity of Toronto, Canada, *Atmos. Environ.*, 9(12), 1137–1139, doi:10.1016/0004-6981(75)90193-6, 1975.

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



▶

►

[Back](#)

Close

Full Screen / Esc

[Printer-friendly Version](#)

Interactive Discussion



- Banta, R. M., Seniff, C. J., Nielsen-Gammon, J., Darby, L. S., Ryerson, T. B., Alvarez, R. J., Sandberg, S. R., Williams, E. J., and Trainer, M.: A bad air day in Houston, *Bull. Am. Meteorol. Soc.*, 86(5), 657–669, 2005.
- Brook, J. R., Strawbridge, K. B., Snyder, B. J., Boudries, H., Worsnop, D., Sharma, S., Anlauf, K., Lu, G., and Hayden, K.: Towards an understanding of the fine particle variations in the LFV: integration of chemical, physical and meteorological observations, *Atmos. Environ.* 38(34), 5775–5788, 2004.
- Capps, S. L., Hu, Y., and Russell, A. G.: Assessing near-field and downwind impacts of reactivity-based substitutions, *J. Air Waste Manage. Assoc.*, 60(3), 316–327, 2010.
- Cheng, W.: Ozone distribution in coastal central Taiwan under sea-breeze conditions, *Atmos. Environ.*, 36(21), 3445–3459, doi:10.1016/S1352-2310(02)00307-2, 2002.
- Cho, S., Makar, P., Lee, W., Herage, T., Liggio, J., Li, S., Wiens, B., and Graham, L.: Evaluation of a unified regional air-quality modeling system (AURAMS) using PrAIRie2005 field study data: The effects of emissions data accuracy on particle sulphate predictions, *Atmos. Environ.*, 43(11), 1864–1877, doi:10.1016/j.atmosenv.2008.12.048, 2009.
- Cooper, O. R., Moody, J. L., Thornberry, T. D., Town, M. S., and Carroll, M. A.: PROPHET 1998 meteorological overview and air-mass classification, *J. Geophys. Res.*, 106, 24289–24300, 2001.
- Corsmeier, U., Kalthoff, N., Kolle, O., Kotzian, M., and Fiedler, F.: Ozone concentration jump in the stable nocturnal boundary layer during a LLJ-event, *Atmos. Environ.*, 31(13), 1977–1989, doi:10.1016/S1352-2310(96)00358-5, 1997.
- Côté, J., Gravel, S., Méthot, A., Patoine, A., Roch, M., and Staniforth, A.: The operational CMC-MRB global environmental multiscale (GEM) model. Part I: Design considerations and formulation, *Mon. Weather Rev.*, 126(6), 1373–1395, 1998.
- Drobinski, P., Saïd, F., Ancellet, G., Arteta, J., Augustin, P., Bastin, S., Brut, A., Caccia, J. L., Campistron, B., Cautenet, S., Colette, A., et al.: Regional transport and dilution during high-pollution episodes in southern France: Summary of findings from the Field Experiment to Constraint Models of Atmospheric Pollution and Emissions Transport (ESCOMPTE), *J. Geophys. Res.*, 112, 26 pp., doi:200710.1029/2006JD007494, 2007.
- Dye, T. S., Roberts, P. T. and Korc, M. E.: Observations of Transport Processes for Ozone and Ozone Precursors during the 1991 Lake Michigan Ozone Study, *J. Appl. Meteorol.*, 34(8), 1877–1889, 1995.
- Fast, J. D. and Heilman, W. E.: The Effect of Lake Temperatures and Emissions on Ozone

- Exposure in the Western Great Lakes Region, *J. Appl. Meteorol.*, 42(9), 1197–1217, 2003.
- Fast, J. D. and Heilman, W. E.: Simulated sensitivity of seasonal ozone exposure in the Great Lakes region to changes in anthropogenic emissions in the presence of interannual variability, *Atmos. Environ.*, 39(29), 5291–5306, doi:10.1016/j.atmosenv.2005.05.032, 2005.
- 5 Gouw, J. D. and Warneke, C.: Measurements of volatile organic compounds in the earth's atmosphere using proton-transfer-reaction mass spectrometry, *Mass Spectrom. Rev.*, 26(2), 223–257, doi:10.1002/mas.20119, 2007.
- Halla, J., Wagner, T., Beirle, S., Ng, A., Strawbridge, K., Firanski, B., Brook, J., Makar, P. A., and McLaren, R.: Determination of Vertical Columns of NO₂ and Aerosol Optical Properties
- 10 in a Rural Setting Using MAX-DOAS, *Atmos. Chem. Phys. Discuss.*, in preparation, 2010.
- Hanna, S. R. and Chang, J. C.: Relations between Meteorology and Ozone in the Lake Michigan Region, *J. Appl. Meteorol.*, 34(3), 670–678, 1995.
- Hastie, D. R., Narayan, J., Schiller, C., Niki, H., Shepson, P. B., Sills, D. M. L., Taylor, P. A., Moroz, W. J., Drummond, J. W., Reid, N., Taylor, R., et al.: Observational evidence for the
- 15 impact of the lake breeze circulation on ozone concentrations in Southern Ontario, *Atmos. Environ.*, 33(2), 323–335, doi:10.1016/S1352-2310(98)00199-X, 1999.
- Hayden, K., Sills, D. M. L., Li, S., Brook, J., Anlauf, K. G., O'Brien, J., and Sharma, S.: The impact of lake breezes on trace gases and particles during the Border Air Quality and Meteorology Study (BAQS-Met), *Atmos. Chem. Phys. Discuss.*, in preparation, 2010.
- 20 Kallos, G., Kassomenos, P., and Pielke, R. A.: Synoptic and Mesoscale Weather Conditions During Air-Pollution Episodes in Athens, Greece, *Bound.-Lay. Meteorol.*, 62(1-4), 163–184, 1993.
- Kambezidis, H. D., Weidauer, D., Melas, D., and Ulbricht, M.: Air quality in the Athens basin during sea breeze and non-sea breeze days using laser-remote-sensing technique, *Atmos. Environ.*, 32(12), 2173–2182, doi:10.1016/S1352-2310(97)00409-3, 1998.
- 25 Keeler, G. J., Spengler, J. D., Koutrakis, P., Allen, G. A., Raizenne, M., and Stern, B.: Transported acid aerosols measured in southern Ontario, *Atmos. Environ. Part A. General Topics*, 24(12), 2935–2950, doi:10.1016/0960-1686(90)90474-2, 1990.
- Kleinman, L. I., Daum, P. H., Imre, D., Lee, Y., Nunnermacker, L. J., Springston, S. R., Weinstein-Lloyd, J. and Rudolph, J.: Ozone production rate and hydrocarbon reactivity in
- 30 5 urban areas: A cause of high ozone concentration in Houston, *Geophys. Res. Lett.*, 29, 105-1, 2002.
- Lasry, F., Coll, I. and Buisson, E.: An insight into the formation of severe ozone episodes:

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



modeling the 21/03/01 event in the ESCOMPTE region, Atmos. Res., 74(1–4), 191–215, 2005.

Levy, I., Dayan, U. and Mahrer, Y.: A five-year study of coastal recirculation and its effect on air pollutants over the East Mediterranean region, J. Geophys. Res.-Atmospheres, 113(D16), doi:10.1029/2007JD009529, 2008a.

Levy, I., Dayan, U., and Mahrer, Y.: Studying coastal recirculation with a simplified analytical land-sea breeze model, J. Geophys. Res.-Atmospheres, 113, D3104, doi:10.1029/2007JD008628, 2008b.

Liu, K., Wang, Z., and Hsiao, L.: A modeling of the sea breeze and its impacts on ozone distribution in northern Taiwan, Environ. Model. Softw., 17(1), 21–27, doi:10.1016/S1364-8152(01)00049-4, 2002.

Lu, R. and Turco, R. P.: Air Pollutant Transport in a Coastal Environment .1. 2-Dimensional Simulations of Sea-Breeze and Mountain Effects, J. Atmos. Sci., 51(15), 2285–2308, 1994.

Lu, R. and Turco, R. P.: Air Pollutant Transport in a Coastal Environment .2. 3-Dimensional Simulations over Los-Angeles Basin, Atmos. Environ., 29(13), 1499–1518, 1995.

Lyons, W. A. and Cole, H. S.: Photochemical Oxidant Transport – Mesoscale Lake Breeze and Synoptic-Scale Aspects, J. Appl. Meteorol., 15(7), 733–743, 1976.

Lyons, W. A. and Cole, H. S.: Fumigation and Plume Trapping on the Shores of Lake Michigan During Stable Onshore Flow, J. Appl. Meteorol., 12(3), 494–510, 1973.

Makar, P., Moran, M., Zheng, Q., Cousineau, S., Sassi, M., Duhamel, A., Besner, M., Davignon, D., Crevier, L. and Bouchet, V.: Modelling the impacts of ammonia emissions reductions on North American air quality, Atmos. Chem. Phys., 9(18), 7183–7212, doi:10.5194/acp-9-7183-2009, 2009.

Makar, P. A., Zhang, J., Gong, W., Stroud, C., Sills, D., Hayden, K. L., Brook, J., Levy, I., Mihele, C., Moran, M. D., Tarasick, D. W., and He, H.: Mass tracking for chemical analysis: the causes of ozone formation in southern Ontario during BAQS-Met 2007, Atmos. Chem. Phys. Discuss., 10, 14241–14312, doi:10.5194/acpd-10-14241-2010, 2010.

Mavrakakis, A., Flocas, H. A., Mavromatidis, E., Kallos, G., Theoharatos, G., and Christides, A.: A case of nighttime high ozone concentration over the greater Athens area, Meteorologische Zeitschrift, 19, 35–45, doi:10.1127/0941-2948/2010/0428, 2010.

Millan, M. M., Mantilla, E., Salvador, R., Carratala, A., Sanz, M. J., Alonso, L., Gangoiti, G., and Navazo, M.: Ozone cycles in the western Mediterranean basin: Interpretation of monitoring data in complex coastal terrain, J. Appl. Meteorol., 39(4), 487–508, 2000.

ACPD

10, 19763–19810, 2010

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Mukammal, E. I.: Ozone as a cause of tobacco injury, *Agr. Meteorol.*, 2(3), 145–165, doi:10.1016/0002-1571(65)90016-6, 1965.
- Murray, D., McWhirter, J., Wier, S., and Emmerson, S.: The Integrated Data Viewer: a web-enabled application for scientific analysis and visualization, in *Preprints, 19th Intl Conf. on IIPS for Meteorology, Ocean. Hydrol.*, 2003.
- Oh, I., Kim, Y., Lee, H. W., and Kim, C.: An observational and numerical study of the effects of the late sea breeze on ozone distributions in the Busan metropolitan area, Korea, *Atmos. Environ.*, 40(7), 1284–1298, 2006.
- Ohashi, Y. and Kida, H.: Local circulations developed in the vicinity of both coastal and inland urban areas: A numerical study with a mesoscale atmospheric model, *J. Appl. Meteorol.*, 41(1), 30–45, 2002.
- RDCT: (R Development Core Team). R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria. [online] Available from: <http://www.R-project.org>, 2009.
- Reitebuch, O., Strassburger, A., Emeis, S., and Kuttler, W.: Nocturnal secondary ozone concentration maxima analysed by sodar observations and surface measurements, *Atmos. Environ.*, 34(25), 4315–4329, doi:10.1016/S1352-2310(00)00185-0, 2000.
- Roberts, J. M., Fehsenfeld, F. C., Liu, S. C., Bollinger, M. J., Hahn, C., Albritton, D. L., and Sievers, R. E.: Measurements of aromatic hydrocarbon ratios and NO_x concentrations in the rural troposphere: Observation of air mass photochemical aging and NO_x removal, *Atmos. Environ.*, 18(11), 2421–2432, doi:10.1016/0004-6981(84)90012-X, 1984.
- Sillman, S., Samson, P. J., and Masters, J. M.: Ozone Production in Urban Plumes Transported Over Water: Photochemical Model and Case Studies in the Northeastern and Midwestern United States, *J. Geophys. Res.*, 98(D7), 12687–12699, doi:10.1029/93JD00159, 1993.
- Sills, D. M. L., Brook, J., Taylor, P., Junhua, Z., Levy, I., Makar, P., and Hayden, K.: Lake Breezes in the Southern Great Lakes and Their Influence During BAQS-Met 2007, *Atmos. Chem. Phys. Discuss.*, in preparation, 2010.
- Stroud, C., Makar, P., Moran, M., Gong, W. M., Gong, S., Zhang, J., Hayden, K., Mihele, C., Brook, J., Abbatt, J., and Slowik, J.: Impact of model grid spacing on regional air quality predictions of organic aerosol, *Atmos. Chem. Phys. Discuss.*, in preparation, 2010.
- Sun, J., Burns, S. P., Lenschow, D. H., Banta, R., Newsom, R., Coulter, R., Frasier, S., Ince, T., Nappo, C., Cuxart, J., Blumen, W., et al.: Intermittent Turbulence Associated with a Density

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Current Passage in the Stable Boundary Layer, Bound.-Lay. Meteorol., 105(2), 199–219, doi:10.1023/A:1019969131774, 2002.

Vlasenko, A., Slowik, J. G., Bottenheim, J. W., Brickell, P. C., Chang, R. Y., Macdonald, A. M., Shantz, N. C., Sjostedt, S. J., Wiebe, H. A., Leaitch, W. R., and Abbatt, J. P. D.: Measurements of VOCs by proton transfer reaction mass spectrometry at a rural Ontario site: Sources and correlation to aerosol composition, J. Geophys. Res., 114(D21), doi:10.1029/2009JD012025, 2009.

Zhang, J. and Rao, S. T.: The Role of Vertical Mixing in the Temporal Evolution of Ground-Level Ozone Concentrations, J. Appl. Meteorol., 38(12), 1674–1691, 1999.

ACPD

10, 19763–19810, 2010

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. List of sites used for ozone analysis.

Name	Abbreviation	Type	Latitude	Longitude	Network
Palmyra	PAL	Rural	42°26.525' N	81°44.370' W	MESONet
Croton	CRO	Rural	42°36.571' N	82°04.812' W	MESONet
Sombra	SOM	Rural	42°41.780' N	82°25.781' W	MESONet
Paquette Corners	PAQ	Rural	42°11.717' N	82°57.922' W	MESONet
Leamington	LEA	Rural	42°04.121' N	82°36.826' W	MESONet
Wheatley	WHE	Rural	42°07.978' N	82°23.731' W	MESONet
Lighthouse Cove	LIG	Rural	42°17.499' N	82°31.343' W	MESONet
Essex	ESS	Rural	42°09.600' N	82°50.000' W	OME ¹
Ridgetown (Supersite)	XRG	Rural	42°27.200' N	81°53.268' W	OME
Harrow (Supersite)	HAR	Rural	42°01.978' N	82°53.603' W	EC ² , OME
Bear Creek (Supersite)	BEC	Rural	42°32.153' N	82°23.352' W	OME
Pelee Island	PEI	Lake	41°46.767' N	82°40.220' W	OME
Lake Erie Buoy	ECO	Lake	42°02.005' N	82°59.006' W	IADN ³
Windsor Downtown	WND	Urban	42°18.983' N	83°02.664' W	OME
Windsor West	WNW	Urban	42°17.567' N	83°04.400' W	OME
Allen Park	ALP	Urban	42°13.700' N	83°12.000' W	MI ⁴
Detroit	DET	Urban	42°18.250' N	83°06.000' W	MI
Oak Park	OAK	Urban	42°27.784' N	83°10.998' W	MI
Warren	WAR	Urban	42°30.800' N	83°00.000' W	MI

¹ OME – Ontario Ministry of the Environment

² EC – Environment Canada

³ IADN – Integrated Atmospheric Deposition Network

⁴ MI – Michigan State

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.




Fig. 1. Map of study region, indicating (left) model analysis sub-domain and the locations of chemistry supersites (BEC – Bear Creek, XRG – Ridgetown, HAR – Harrow), monitoring stations, buoy (ECO), ferry route, and main urban areas (gray shaded regions), and (right) the 2.5-km AURAMS model domain and larger setting. LSC marks Lake St. Clair; LH marks Lake Huron, and LO marks Lake Ontario.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.




Fig. 2. (a) 2-h running averages of surface ozone for the Buoy, Pelee Island, rural-site composite, urban-site composite, and the average of ferry measurements for trips scheduled on all weekdays; (b) running change rate between two consecutive hours of the measured data; (c) AURAMS diurnal averages at surface level over the 2.5-km grid sub-domain in Fig. 1 grouped by lakes, land and urban areas; (d) running change rates between two consecutive hours of the model data.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.




Fig. 3. Time-height cross-sections of the AURMAS predicted ozone for the sub-domain shown in Fig. 2, sorted by surface type: **(a)** rural; **(b)** lake; and **(c)** urban. Differences are shown between **(d)** rural and lake, **(e)** rural and urban, and **(f)** urban and lake surface types, with the red frames in the middle panels marking a higher-resolution cross-section for midnight-noon up to 240 m a.g.l. shown in panels **(g)**, **(h)** and **(i)**, respectively.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.




Fig. 4. (a) Mean ground-level ozone, (b) mean ozone cross sections (ppbv) and (c) mean vertical velocity (m s^{-1}) from AURAMS for all 24 h (a1, b1 & c1) and selected times (00:00, 06:00, 14:00 and 21:00 EDT, a2-5, b2-5 and c2-5, respectively), calculated for the entire 23 days of the intensive campaign (18 June–10 July 2007). Cross sections are along transect A–F marked in (a), with coloring intervals of 2 ppbv for ozone and 0.01 m s^{-1} for vertical velocity.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.




Fig. 5. Breeze fronts locations on 6 July at **(a)** 12:00 EDT and **(b)** 20:00 EDT, and 7 July at **(c)** 00:00 EDT and **(d)** 14:00 EDT.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.




Fig. 6. Measurements from 6–7 July including: **(a)** wind speed and direction at Harrow; **(b)** O₃, CO and SO₂ mixing ratios at Harrow; **(c)** NO, NO₂, NO_x and NO_y, as well as Toluene to Benzene ratio (TBR) at Harrow; and **(d)** O₃ at three additional sites (Wheatley, Leamington and the buoy). Thick gray vertical lines mark the times of the first two lake-breeze frontal passages at Harrow, marked A1 and A2. Numbering (1–11) refers to different stages detailed in the text.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.




Fig. 7. Ozone (left) and NO₂ (right) measured at early morning hours (04:30–07:20 EDT) of 7 July by the Twin Otter aircraft. Notation P1–P9 refers to vertical profiles shown in Fig. 8.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.




Fig. 8. (a): Twin Otter time series for 7 July showing O₃, NO₂, and elevation; (b–j): selected vertical profiles of O₃, NO₂, and temperature. Profile locations are indicated on Fig. 7.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.




Fig. 9. (a): time series for 7 July showing tethersonde elevation (black) and ozone mixing ratio (red) and ground-level ozone measured at the Ridgetown site (blue). (b–k): tethersonde vertical profiles of O₃ (red) and temperature (green) at Ridgetown. Profiles are numbered from P1 to P10.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.




Fig. 10. (a) Plan view and (b) side view from southeast of the aircraft flight on 7 July 2007 between 13:40 and 16:20 EDT, with colors indicating ozone mixing ratios (ppbv).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.




Fig. 11. (a): Twin Otter time series of O_3 , NO_2 , and elevation; (b): vertical profiles of O_3 , NO_2 , and temperature over Lake St. Clair. Profile period is indicated as shaded area. Top notation refers to periods when the aircraft is located over, to the north, or to the south of Lake St. Clair (LSC).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.




Fig. 12. Meteorological (wind speed (black) and direction (red), temperature (black), dew point temperature (green) and relative humidity (blue)) and ozone (red) measurements during the night of 6–7 July 2007 at four ground stations ((a) – Leamington, (b) – Lighthouse Cove, (c) – Bear Creek and (d) – Wheatley).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Unraveling the complex local-scale flows influencing ozone patterns

I. Levy et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 13. Plan view at the surface (a, b) and vertical cross sections (c, d) of wind speed from the AURAMS model on the night of 6–7 July 2007 at 22:00 EDT and 03:20 EDT.