

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

**Greenhouse gases in  
the Indian summer  
monsoon plume**

T. J. Schuck et al.

# Greenhouse gas relationships in the Indian summer monsoon plume measured by the CARIBIC passenger aircraft

T. J. Schuck<sup>1</sup>, C. A. M. Brenninkmeijer<sup>1</sup>, A. K. Baker<sup>1</sup>, F. Slemr<sup>1</sup>,  
P. F. J. v. Velthoven<sup>2</sup>, and A. Zahn<sup>3</sup>

<sup>1</sup>Atmospheric Chemistry Division, Max Planck Institute for Chemistry, Mainz, Germany

<sup>2</sup>Royal Netherlands Meteorological Institute, de Bilt, The Netherlands

<sup>3</sup>Institute for Meteorology and Climate Research, Karlsruhe Institute of Technology, Karlsruhe, Germany

Received: 7 January 2010 – Accepted: 19 January 2010 – Published: 1 February 2010

Correspondence to: T. J. Schuck (tanja.schuck@mpic.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

During the Indian summer monsoon period 2008, regular measurement flights were performed by the CARIBIC aircraft between Germany and India. Measurements included whole air sampling and subsequent analysis for greenhouse gases (CH<sub>4</sub>, CO<sub>2</sub>, N<sub>2</sub>O, SF<sub>6</sub>) and in-situ measurements of CO, O<sub>3</sub>, and H<sub>2</sub>O. For all these tracers a distinct monsoon signature was observed in the longitude range 50–80° E south of 40° N at flight altitudes 8–12.5 km. The formation of a monsoon plume with enhanced mixing ratios was observed for CH<sub>4</sub>, N<sub>2</sub>O, SF<sub>6</sub>, CO and H<sub>2</sub>O. The plume began to develop in May and persisted through September, and maximum mixing ratios and maximum spatial extension of the plume were observed in August. For CO<sub>2</sub> and O<sub>3</sub> a minimum was observed. The amplitude of the CO<sub>2</sub> seasonal cycle was larger than at comparable latitudes outside the monsoon region, and the CO<sub>2</sub> spring maximum was observed with a delay of one month. Different tracer relationships show strong spatial variations across the plume. The comparison with NOAA ground station data shows a clear enhancement for CH<sub>4</sub>, N<sub>2</sub>O, SF<sub>6</sub> and CO and a decrease in CO<sub>2</sub> mixing ratios. Assuming seasonally invariant SF<sub>6</sub> emissions based on the EDGAR 4.0 inventory, monthly emissions were estimated for the period June–September for CH<sub>4</sub>, N<sub>2</sub>O, and CO, and the CO<sub>2</sub> uptake was estimated. While for N<sub>2</sub>O an emission decrease during the monsoon period was derived, emissions of CH<sub>4</sub> were highest in September indicating a different seasonal cycle of emissions. Net CO<sub>2</sub> uptake by the vegetation was largest in September.

## 1 Introduction

The Asian monsoon is a major meteorological phenomenon during Northern Hemisphere summer. The temperature differences between the Asian landmass, especially the Tibetan Plateau, and the Indian Ocean, drive a cyclonic circulation over the Indian Ocean at low altitudes (Hoskins and Rodwell, 1995; Hsu et al., 1999). At the same time,

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**Greenhouse gases in  
the Indian summer  
monsoon plume**T. J. Schuck et al.

---

5 during the summer monsoon period an anticyclone develops in the upper troposphere (Krishnamurti and Bhalme, 1976). The summer monsoon, and thus the rainy season, typically reaches Southern India in early June, and it usually starts to retreat at the beginning of September (IMD, 2008; Joseph et al., 2006; Wang et al., 2009). The monsoon circulation not only profoundly influences the large scale dynamics of the lower and upper troposphere, but it also is accompanied by enhanced troposphere – stratosphere exchange (Chen, 1995; Traub and Lelieveld, 2003; Fu et al., 2006; Krishnamurti et al., 2008). Model calculations combined with satellite measurements suggest that the monsoon convection and subsequent transport through the upper-level anticyclone moistens the stratosphere (Lelieveld et al., 2007). This mechanism may also be responsible for transport of pollution into the stratosphere.

10 During the summer monsoon, strong and widespread convection occurs over India, the Bay of Bengal and over the Tibetan Plateau, partly over densely populated regions where strong emission sources are located. Surface air masses with enhanced levels of H<sub>2</sub>O, CO, and many other trace gases are lifted and subsequently mixed into the large-scale circulation. The rapid convective vertical transport of polluted boundary layer air thus leads to enhanced mixing ratios of many trace gases in the upper-level anticyclone.

15 In addition, the intense precipitation associated with the summer monsoon affects the activity of several sources. For instance, CH<sub>4</sub> emissions from wetlands and rice paddies increase, CO<sub>2</sub> uptake by plants increases, while CO emissions from wildfires and agricultural waste burning are expected to decrease. The monsoon meteorology therefore not only influences the dynamics of the upper troposphere, it also affects its chemical composition. Air masses carrying a distinct monsoon signature such as enrichment of CH<sub>4</sub> can accumulate in the anticyclone, leading to the formation of a localized maximum that persists throughout the summer. In the following, the word “plume” denotes this large, quasi-stationary feature.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Greenhouse gases in  
the Indian summer  
monsoon plume**

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Satellite observations reveal strong seasonal variations of various tracers in the anticyclone. High summer time mixing ratios over South Asia were reported for CO (Fu et al., 2006; Park et al., 2008), for H<sub>2</sub>O (Park et al., 2004; Fu et al., 2006; Randel and Park, 2006), and for CH<sub>4</sub> (Park et al., 2004; Schneising et al., 2009; Xiong et al., 2009). Other trace gases that were observed to be enhanced in the monsoon region included NO<sub>x</sub> (Park et al., 2004), HCN, C<sub>2</sub>H<sub>6</sub>, and C<sub>2</sub>H<sub>2</sub> (Park et al., 2008). In addition, a concomitant decrease of O<sub>3</sub> was observed (Randel and Park, 2006; Liu et al., 2009). These observations were explained to be due to convective uplifting of boundary layer air in the monsoon region (Randel and Park, 2006) and were shown to be in agreement with model results, for example from the MOZART model. Simulations predicted high mixing ratios of CH<sub>4</sub> in the anticyclone between 60° E and 120° E around 30° N at pressure altitudes between 300 and 100 hPa, as well as elevated levels of H<sub>2</sub>O and NO<sub>x</sub>, in accordance with observations from the HALOE instrument (Park et al., 2004). The model also calculated a CO maximum at ~15 km altitude, in agreement with MLS and ACE-FTS measurements (Park et al., 2009). Moreover, calculations using the TM3 model generally agree with the observation of elevated CH<sub>4</sub> mixing ratios, although the model cannot accurately reproduce the position of the plume (Xiong et al., 2009). Satellite instruments, despite their limited vertical resolution, can provide information on the altitude range of the observed tracer enhancements. The observed monsoon plume in CH<sub>4</sub> is for example reported by the AIRS instrument to extend from 500 hPa to 150 hPa (Xiong et al., 2009), the MOZART model predicts a vertical extension from ~300–150 hPa (Park et al., 2004).

Strong variations during the summer monsoon period are also expected for CO<sub>2</sub> with its pronounced seasonal cycle related to photosynthesis and respiration. The drop in boundary layer concentrations for this greenhouse gas should be reflected by upper tropospheric levels due to rapid and widespread convective transport. Signatures of the Asian monsoon in upper tropospheric CO<sub>2</sub> have indeed been observed, e.g. by the IASI satellite instrument (Crevoisier et al., 2009). However, most studies based on CO<sub>2</sub> measurements by satellites concentrated on different aspects of the CO<sub>2</sub> distribution,

mainly taking a more global approach or focusing on highly industrialized regions rather than the Asian monsoon region (e.g., Chédin et al., 2003; Buchwitz et al., 2007a,b; Schneising et al., 2008).

Although an increasing number of satellite remote sensing observations of various trace gases is becoming available, systematic in-situ measurements of the composition of the upper troposphere are scarce in the areas most influenced by the Asian summer monsoon. Most studies investigating the influence of the monsoon circulation, such as the Arabian Sea Monsoon Experiment or the Bay of Bengal Monsoon Experiment (Sanjeeva Rao, 2005; Bhat and Narashima, 2007), involved ground and ship based measurements but no in-situ measurements in the upper troposphere. The Indian Ocean Experiment (INDOEX) did include aircraft measurements, but investigated the transport of pollution over the Indian Ocean in winter (Lelieveld et al., 2001). The Mediterranean Intensive Oxidant Study (MINOS) captured the outflow of pollutants from the monsoon anticyclone and their westward transport. It was shown that the upper troposphere over the Mediterranean was strongly influenced by long-range transport from the monsoon region (Lelieveld et al., 2002; Scheeren et al., 2003).

To study the influence of the Indian summer monsoon on the composition of the upper troposphere, not only over the Indian Ocean but also over land, 16 flights with the CARIBIC (Civil Aircraft for the Regular Investigation of the Atmosphere Based on an Instrument Container, [www.caribic-atmospheric.com](http://www.caribic-atmospheric.com)) aircraft took place between Frankfurt, Germany, and Chennai, India, from April to December 2008. Next to CH<sub>4</sub> and CO<sub>2</sub>, two other greenhouse gases, N<sub>2</sub>O and SF<sub>6</sub> will be discussed. SF<sub>6</sub>, an entirely anthropogenic gas, can be used as a tracer to estimate the emissions of the other greenhouse gases during the monsoon time. Further tracers measured are CO which, compared to the aforementioned greenhouse gases, has a short lifetime and H<sub>2</sub>O and O<sub>3</sub>, which show a very different behavior in the monsoon plume.

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2 Technical and analytical details

CARIBIC is a long-term atmospheric measurement program. It is based on the use of a fully automated scientific instrument package in a 1.5 t container aboard a passenger aircraft which is equipped with an advanced multi-probe inlet system (Brenninkmeijer et al., 2007). The present CARIBIC system came into service in December 2004, and since then it is operated monthly aboard a Lufthansa Airbus A340-600 passenger aircraft. With typical cruising altitudes ranging from 8.5 to 12 km (corresponding to 300–180 hPa), the CARIBIC aircraft frequently crosses the tropopause at mid-latitudes. At low latitudes the free tropical troposphere is probed. Figure 1 shows the CARIBIC flight tracks between Frankfurt (50.03° N, 8.54° E) and Chennai (12.99° N, 80.18° E) in 2008. Flights were performed monthly April through December. Usually one set of flights consisted of four consecutive flights, i.e. two round trips from Frankfurt to Chennai within three days, with exception of July and October, when only one round trip was performed.

The CARIBIC instrument container houses a range of in-situ trace gas and aerosol analyzers, complemented by an aerosol sampler and a whole air sample collection system (Brenninkmeijer et al., 2007). The latter comprises a pumping unit and two separate sampling units, each containing 14 glass cylinders with a volume of 2.67 l. Depending on the flight altitude, the total sample collection times range from 30 to 90 s, corresponding to a spatial resolution of about 7 to 21 km. Depending on the flight altitude, the air sample volume ranges from 9 to 11 l (STP). In order to get a higher spatial coverage along the flight track, given the number of samples carried aboard being limited to 28, sampling took always place during only the first round trip Frankfurt-Chennai, with 14 samples being filled on the first outbound flight and 14 samples on the corresponding return flight. Samples were taken at even time steps with on average one sample per half hour. In total, 251 whole air samples were collected during the 16 long-distance flights. After returning from the final flight leg, the container is de-installed from the aircraft. Subsequently, the sample units are removed and the pressurized

### Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



samples are analyzed in several laboratories. The standard greenhouse gas analysis is performed for all samples as soon as possible, usually within two days after the last flight. Briefly, it comprises simultaneous measurements using a HP 6890 gas chromatograph of CH<sub>4</sub> and CO<sub>2</sub> with a flame ionization detector (FID) and of N<sub>2</sub>O and SF<sub>6</sub> with an electron capture detector (ECD) (Schuck et al., 2009). The average precision obtained for the flights between Frankfurt and Chennai in 2008 is 0.12% for CH<sub>4</sub>, 0.07% for CO<sub>2</sub>, 0.11% for N<sub>2</sub>O, and 0.9% for SF<sub>6</sub>. Calibration is based on seven standard gas mixtures acquired from NOAA-ESRL. To calculate mixing ratios in dry air, the continuously measured total water mixing ratio is used for correction (see below).

A description of all instruments of the CARIBIC container and the inlet system is given in Brenninkmeijer et al. (2007). The package comprises several in-situ instruments for on-line measurements of various trace gases, among them CO, O<sub>3</sub>, and H<sub>2</sub>O. The air flow for measurements of CO and O<sub>3</sub> is through the same sampling line used for the collection of whole air samples with sampling perpendicular to the flight direction. CO is measured using a vacuum ultraviolet fluorescence instrument that is operated with a time resolution of 1 s and is calibrated in flight every 25 min. The uncertainty estimated from the propagation of the uncertainties of calibration gas and the standard deviations of the measurement, zero, and calibration signals is typically 1.9% or 1 ppbv, whichever is greater. O<sub>3</sub> is measured using a dual-beam UV-photometer with a time resolution of 4 s. Comparisons with a laboratory standard showed that at a time resolution of 4 s a precision of 0.3 ppbv (or 0.3%, whatever is highest) and a total uncertainty of ~1.5% can be achieved.

For measurements of H<sub>2</sub>O two instruments are used, a commercial chilled mirror frost point hygrometer (CR2, Buck Research, total uncertainty 0.5 K, time resolution 30–300 s) and a fast two-channel photo-acoustic laser spectrometer (precision 0.5 ppmV, time resolution 5–10 s), using laser light to excite water molecules and measuring their thermal relaxation. Two separate sampling lines are used to allow for measurements of gaseous and condensed water. Both instruments are supplied with air from an inlet facing flight direction for the detection of both, gaseous and condensed

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



water including ice particles (total water). The total water mixing ratio is used for the calculation of greenhouse gas mixing ratios in dry air. In addition, the second channel of the photo acoustic detector is connected to an orifice perpendicular to the flight direction for the detection of water vapor only.

5 The position and selected meteorological data recorded by the aircraft systems are provided to the CARIBIC master computer with a time resolution of 1 s. The master computer also records signals from different pressure and temperature sensors in the container and the inlet probe, and it communicates with the instruments via an Ethernet network, activating and de-activating them and continuously monitoring their status  
10 during a flight. To avoid sampling of polluted air in the vicinity of airports, measurements only take place at outside pressures below 650 mbar. Eventually, all data are reported in terms of the time signal from the master computer.

The routinely produced meteorological plots and trajectory calculations for each flight are based on meteorological analysis data from the European Centre for Medium range  
15 Weather Forecasts (ECMWF) (van Velthoven, 2009). Trajectories are calculated 5 days backward with the KNMI trajectory model TRAJKS (Scheele et al., 1996) for every 3 min of flight. For the trajectory calculations global gridded ECMWF wind fields with a horizontal resolution of  $1^\circ \times 1^\circ$  at time intervals of 6 h were used. Additionally, 8-day trajectories are calculated for every 6 s of flight during the collection time of each whole  
20 air sample. The ECMWF data themselves were also interpolated to the 3-D-location and time of the aircraft for every minute of flight for comparison with the measurements. They were then slightly further interpolated to meet the 10 s-averages of the merged observational data, and they were integrated in the same way as the continuous measurements over the sampling period of the flask samples.

### 25 3 Meteorology

In the upper troposphere, the Asian summer monsoon is associated with an anticyclone that forms in April over Southeast Asia at equatorial latitudes. In summer, this

---

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





---

**Greenhouse gases in  
the Indian summer  
monsoon plume**T. J. Schuck et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

anticyclone drifts northwestward, eventually reaching the Tibetan Plateau around 20–30° N (Krishnamurti and Bhalme, 1976; Hsu et al., 1999). In parallel, the westerly subtropical jet weakens in spring and shifts to more northern latitudes while a weak easterly jet develops at latitudes around 10° N (Krishnamurti et al., 2008). This easterly jet is a possible pathway for transport of pollution uplifted by convection from South Asia towards Northern Africa and the Mediterranean through the subsequent transport by the Arabian anticyclone (Lelieveld et al., 2002). Most summer rainfall occurs on the northern side of this easterly jet (Krishnamurti et al., 2008). During the dry season, rainfall occurs mainly south of 10° N (Krishnamurti et al., 1997).

The monsoon anticyclone is bordered by the westerly jet on its northern flank and in the south by the weaker easterly jet. Measurements of the zonal wind speed can be used as an indicator for the location and drift of the anticyclone. Therefore, of the observables that are recorded by the CARIBIC system, the horizontal wind speed is the one best suited to characterize the meteorological situation during a flight and to follow the seasonal changes. Figure 2 shows the westerly component of the horizontal wind speed averaged over 10 s intervals as a function of latitude for the flights to and from Chennai in 2008. The latitudinal variation of the zonal wind speed shows distinct changes throughout the year and can be used to identify the summer monsoon months. Panel a shows the distribution of the wind speed in summer from June through September. Maximum westerly wind speeds (positive numbers) are recorded north of ~35° N with values around 40 m/s. There is no localized maximum of the wind speed, but rather a plateau in the latitudinal distribution. South of 25° N, easterly winds (negative numbers) with speeds up to 30 m/s are observed. Maximum wind speeds of the tropical easterly jet are around 35 m/s at pressure altitudes around 175 hPa (Krishnamurti et al., 2008). The CARIBIC aircraft cruises at altitudes around 300–200 hPa, thus crossing the lower part this easterly jet, where maximum wind speeds are lower. Figure 3 shows a pressure level map of the wind speed at 250 hPa based on ECMWF data for the flight from Frankfurt to Chennai on 13 August. The cruise altitude during this flight varied between ~240 hPa and ~215 hPa. Clearly discernible are the large scale

anticyclonic circulation, the easterly jet south of 20° N, and the westerly jet around 40° N.

Panel b of Fig. 2 shows the wind speed for November and December 2008. The westerly jet now extends further south and a pronounced maximum of the wind speed is observed around 25–30° N, with maximum values of 60 m/s in December. Similar latitudinal distributions were observed during flights between Germany and the Philippines via China in winter. Not shown are the flights that took place in April, May and October. During these transition months, wind speeds are lower than in winter, comparable to the maximum levels in summer. The latitudinal distribution does not show a clear maximum and no tropical easterly jet was present in the latitude range 10–40° N. Studying the behavior of the horizontal wind speed as a function of latitude for all flights to India and to the Philippines for three years (2006–2008) shows that the common seasonal classification scheme according to month may not be appropriate. This is especially true since the transition between the summer and the winter mode does not happen at the same time each year. For example, in 2008, May was still a transition month, while in May 2007 during flights between Germany and the Philippines a summer-type profile was observed. These differences reflect the interannual variations of the upper tropospheric circulation and the varying onset times of the summer monsoon.

Generally, the rainfall season in India is expected to last from 1 June to 30 September (Joseph et al., 2006; Wang et al., 2009). In 2008, the onset of the monsoon rains over Southern India was on 31 May, and they progressed north slightly faster than usual, with total coverage of India about 5 days earlier than the long-term average from 1941–1990. Overall, average amounts of rain were reported by the India Meteorological Department for the whole season, but with large variations in time, even causing severe flooding in some areas. After a rain surplus in June, a rain deficiency was observed in July followed by average rainfall in August and September, and large spatial variations were observed. In 2008, the monsoon started to retreat in September, somewhat later than the long term average (IMD, 2008). Consequently, the features of the summer monsoon meteorology, namely the easterly winds at latitudes around

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



15–25° N are still observed during the CARIBIC flights on 10 and 11 September, and in the present analysis September is considered a summer month. According to the latitudinal variation of the horizontal wind speed, October was a transition month, while Fig. 2 shows a winter signature for November and December.

## 5 4 Measurements in the monsoon plume

### 4.1 The CH<sub>4</sub> plume

The most prominent feature observed in trace gas mixing ratios during the flights between Germany and India is the enhancement in upper tropospheric methane mixing ratios during summer. Figure 4 shows the latitudinal distribution of CH<sub>4</sub> for spring (April and May), summer (June to September) and autumn (October). The seasons were classified according to the zonal wind speed as described in the previous section. Shown are all tropospheric samples that were collected south of 40° N. Whereas the CARIBIC aircraft is always in the troposphere at tropical latitudes, it frequently crosses the tropopause in the mid-latitudes. Here, the selection of tropospheric samples is based on potential vorticity by excluding PV values >1.3 PVU (1 PVU ≐ 10<sup>-6</sup> K m<sup>2</sup> kg<sup>-1</sup> s<sup>-1</sup>). Excluded are 19 samples, all collected north of 25° N in spring and autumn and north of 30° N in summer. The total number of tropospheric samples collected south of 40° N is 96. Included for comparison is the monthly average CH<sub>4</sub> mixing ratio observed in the corresponding period at the Mauna Loa Observatory (MLO, 19° N, 155° W, 3397 m a.s.l.) (Dlugokencky et al., 2009). This station has been chosen because it is a high altitude station, situated in the latitude band where the monsoon plume is observed, but not in a region influenced by direct emissions and monsoon circulations.

In April and May (upper panel), little latitudinal structure is observed, and the values on average agree with those at MLO. Nonetheless, a small relative enhancement is visible around 20° N, whereas around 30° N mixing ratios are slightly lower. In summer

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Greenhouse gases in  
the Indian summer  
monsoon plume**

T. J. Schuck et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

(middle panel), the development of a pronounced  $\text{CH}_4$  plume can be seen. In June, values north of  $30^\circ \text{N}$  are still at the background level, whereas south of  $30^\circ \text{N}$  an enhancement of up to  $\sim 35$  ppb occurs. This plume grows in July and August when it extends further north and reaches a maximum value of  $1907.4 \pm 2.2$  ppb in August around  $28^\circ \text{N}$ , 112.8 ppb above the mean value of 1794.6 ppb observed at MLO in August. This amplitude agrees with that of  $\sim 100$  ppb observed by the AIRS satellite instrument (cf. Fig. 4 in Xiong et al., 2009). In September, the maximum  $\text{CH}_4$  mixing ratios remain high, but are observed at lower latitudes than in August as the plume weakens at its northern end. This southward shift reflects the retreat of the monsoon.

In October (lower panel), the latitudinal distribution of the  $\text{CH}_4$  mixing ratio has returned to the background level with a weak latitudinal gradient except for the first sample collected after the departure from Chennai on 15 October. This sample was collected during the ascent at an altitude of 8.6 km, while all other samples on the two flights in October were collected at cruise altitudes above 10 km. Backward trajectories indicate that for this flight section the air mass had crossed Eastern India at pressure altitudes below 400 hPa implying a high likelihood of urban pollution in the sample. The trajectories for the neighboring samples had been above 400 hPa for at least 24 h prior to sampling. The low altitude sample also had an exceptionally high water vapor content of  $\sim 2300$  ppmv compared to 120–160 ppmv measured for the neighboring samples and the corresponding sample on the inbound flight.

Figure 5 compares the latitudinal distribution of  $\text{CH}_4$  for the summer months with that of  $\text{N}_2\text{O}$ ,  $\text{SF}_6$ ,  $\text{CO}_2$ , and  $\text{CO}$ .  $\text{N}_2\text{O}$  and  $\text{CO}$  show a similar behavior, and their latitudinal distribution closely resembles that of  $\text{CH}_4$ . For both gases the monsoon plume is observed between  $15^\circ \text{N}$  and  $40^\circ \text{N}$  in June, and the latitude at which their maximum values are observed shifts northward between June and August, and southward again in September. In parallel, a broadening of the plume can be seen.  $\text{SF}_6$  also shows an increase throughout summer. Due to the larger analytical uncertainty, its latitudinal variation is less pronounced than for  $\text{CH}_4$  and  $\text{N}_2\text{O}$ , but the southward shift from August to September is also observed (cf. Figs. 7 and 8). The distributions of  $\text{N}_2\text{O}$ ,  $\text{SF}_6$ , and

CO that are similar to that of CH<sub>4</sub> will be discussed in more detail in Sect. 5.

## 4.2 Formation of a CO<sub>2</sub> minimum

For CO<sub>2</sub>, instead of the formation of a plume, the development of a minimum is observed. In June, the distribution looks very similar to that of CH<sub>4</sub>, with a plume-like structure south of 30° N. Around 20–25° N, the CARIBIC measurement is more than 1 ppm higher than the corresponding monthly mean value of 388.35 ppm observed at MLO. This indicates that surface level values of CO<sub>2</sub> are still high when the strong convection related to the onset of the monsoon sets in at lower latitudes, and upper tropospheric levels of CO<sub>2</sub> increase due to convective transport. In July, this plume disappears and a small decrease in the CO<sub>2</sub> mixing ratios at latitudes north of 30° N and a larger decrease at southern latitudes are observed. This trend continues in August and September and the formation of a “negative” plume is seen with lowest mixing ratios around 20° N. In September, this feature extends north to 40° N. No latitudinal variation is seen for CO<sub>2</sub> in spring and autumn (not shown).

In the monsoon region, the maximum mixing ratio of CO<sub>2</sub> is observed in June, the minimum mixing ratio in September. In contrast, for all CARIBIC flights to Asia from 2006 to 2008, including flights between Germany and India and between Germany and the Philippines, the maximum value is observed in May. Since the onset of the monsoon rains has a strong influence on the timing of the growth period, the monsoon activity could influence the seasonal cycle of CO<sub>2</sub> and the time when maximum mixing ratios occur. In addition, it may also influence the amplitude of the seasonal cycle in that region by enhanced vertical transport of low-CO<sub>2</sub> boundary layer air in late summer. In June, a mean CO<sub>2</sub> mixing ratio of 388.57±0.39 ppm is observed, in September it is 381.21±0.33 ppm. Compared with monthly mean values measured at MLO (Conway et al., 2009), a similar maximum value of 388.62 ppm is measured at the station. But while at MLO the maximum is observed in May, in the monsoon region it is observed in June, with a delay of one month. The minimum value of 382.70 ppm at MLO is observed in September, concomitant with the minimum in the CARIBIC data, but it is

### Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



higher than the CARIBIC value. Correcting both datasets for an average linear increase of  $0.14 \text{ ppm month}^{-1}$ , derived from the time series of monthly mean  $\text{CO}_2$  mixing ratios at MLO from 1969–2007, yields an amplitude of the seasonal cycle in 2008 of 6.48 ppm at MLO and 7.79 ppm for CARIBIC.

5 For all CARIBIC flights between Germany and India and between Germany and the Philippines a mean amplitude of only  $\sim 6$  ppm was derived for latitudes  $10\text{--}40^\circ \text{ N}$ . This result is in agreement with a value of about 6 ppm derived from flask measurements at  $10\text{--}30^\circ \text{ N}$  by the JAL/CONTRAIL collaboration during flights between Japan and Australia (Matsueda et al., 2008). These flights took place east of the monsoon plume  
10 at longitudes  $153\text{--}140^\circ \text{ E}$  in 1993–2007. In this data set, the maximum  $\text{CO}_2$  mixing ratio is typically observed in early May with a shift to the end of May at latitudes  $10\text{--}20^\circ \text{ N}$ , and the minimum is observed in late September. The fact that the amplitude of the seasonal cycle decreases with altitude throughout the free troposphere (Gurk et al., 2008), explains the somewhat smaller amplitude derived from aircraft measurements  
15 in comparison with the value observed at MLO. However, despite the higher altitude of the aircraft, the amplitude observed in the monsoon region in 2008 is markedly larger than that observed at MLO, and it is also higher than the average amplitude observed at similar altitudes further east.

At ground level, the monsoon circulation is associated with a cyclonic flow of air  
20 and southwesterly winds. These low-level winds transport air from more southern latitudes towards South Asia. In the beginning of the rainy season,  $\text{CO}_2$  in the continental boundary is still relatively high, and the strong convection leads to the formation of a  $\text{CO}_2$  plume at higher altitudes. With the growing season's progress, additional  $\text{CO}_2$  is taken up by the vegetation. Because convection is still strong in late summer, low-  
25  $\text{CO}_2$  air is effectively transported into the middle troposphere where then lower  $\text{CO}_2$  values are observed in the monsoon region than at corresponding latitudes further east, and the amplitude of the seasonal cycle increases.

---

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

### 4.3 Latitudinal distributions of O<sub>3</sub> and H<sub>2</sub>O

Ozone (O<sub>3</sub>) and water vapor (H<sub>2</sub>O) are important constituents of the atmosphere discussed in many publications. The CARIBIC measurements also document their changes during the summer monsoon. Figure 6 shows H<sub>2</sub>O (top panels) and O<sub>3</sub> (bottom panels) measurements integrated over the sampling period of each sample. In the latitudinal distribution of H<sub>2</sub>O, a systematic offset depending on the direction of the flight route is noticeable for July, August and September. The reason is that samples collected at identical latitudes do vary in altitude, because of systematic altitude differences between outbound and inbound flights. In the beginning of a flight, the cruise altitude is usually lower due to the greater weight of the fully fueled aircraft. Therefore, flights from India back to Germany tend to show higher H<sub>2</sub>O mixing ratios for the southernmost samples taken closest to Chennai. Touch-down and take-off in Chennai took place within two hours during the night, and the southernmost sample was collected about one flight hour away from Chennai. This means that no major difference in convective activity due to different times of day should occur. Such an asymmetry is not observed for any of the other tracers with exception of one low-level sample in October showing an exceptionally high CH<sub>4</sub> mixing ratio (cf. Fig. 4).

During all months from April through December, higher levels of H<sub>2</sub>O were observed towards southern latitudes with maximum values south of 20° N. In June, July, and August, mixing ratios were always above 100 ppmv south of 25° N, with a steep decrease towards 30° N. In September, this step-like decrease was seen a little further south, at 23–25° N. In autumn and winter (not shown) the increase towards southern latitudes was less steep, and it was observed as an increase over a wider latitude range with values above 100 ppmv only south of 20° N.

O<sub>3</sub> and H<sub>2</sub>O mixing ratios are generally seen to be anti-correlated throughout the observation period. During the summer months, a distinct increase of the O<sub>3</sub> mixing ratio from ~60 ppbv to ~100 ppbv can be seen between 25 and 30° N in Fig. 6. Moreover, the latitudinal position of this increase is shifted slightly southward in September.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Greenhouse gases in  
the Indian summer  
monsoon plume**

T. J. Schuck et al.

High mixing ratios of  $\text{N}_2\text{O}$  and  $\text{SF}_6$  and PV values  $<0.6$  PVU at latitudes south of  $30^\circ\text{N}$  clearly indicate tropospheric air masses. The increase in  $\text{O}_3$  mixing ratios towards northern latitudes between  $25^\circ\text{N}$  and  $30^\circ\text{N}$  is accompanied by an increase in PV reaching maximum values of 1.3 PVU in summer. During the monsoon period, values of  $\text{PV} > 1.5$  PVU were only found north of  $35^\circ\text{N}$ . The position of the steep increase in  $\text{O}_3$  at latitudes around  $25^\circ\text{N}$  corresponds to longitudes around  $65^\circ\text{E}$ . The observation of elevated  $\text{O}_3$  mixing ratios in this region is in agreement with a summertime tropospheric  $\text{O}_3$  maximum over the Middle East predicted by model calculations (Li et al., 2001). This maximum has been observed at pressures between 450 hPa and 300 hPa with the Tropospheric Emission Spectrometer, in agreement with observations from the MOZAIC project (Liu et al., 2009; Worden et al., 2009). Model simulations showed that the southern part of the  $\text{O}_3$  maximum is dominated by the easterly outflow of the Asian monsoon anticyclone, and it was proposed that the main contribution is from tropospheric  $\text{O}_3$  production (Li et al., 2001; Liu et al., 2009). In addition to the measurements of greenhouse gases, the CARIBIC air samples are also analyzed for a suite of non-methane hydrocarbons (Baker et al., 2009). These data indicate that there are different regimes in the monsoon plume with variations in the  $\text{O}_3$  production potential (Baker et al., 2010).

In the region of the monsoon anticyclone itself, satellite observations show an  $\text{O}_3$  minimum in the middle troposphere south of  $25^\circ\text{N}$  (Randel and Park, 2006; Liu et al., 2009), in agreement with what is observed during the CARIBIC flights. Data from the Global Ozone Monitoring Experiment (GOME) for 1997–2000 show considerable interannual variations in the strength and the extension of this  $\text{O}_3$  minimum (Liu et al., 2009). In the region where the steep increase in  $\text{O}_3$  is observed in the CARIBIC data ( $25^\circ\text{N}$ ,  $65^\circ\text{E}$ ), a similar increase from values below 70 ppbv to more than 90 ppbv was observed by the satellite instrument at 200 hPa in 1997, 1998, and 2000. In 1999, it was observed further north in the same longitude band. It has been suggested that the formation of the  $\text{O}_3$  minimum is caused by tropical marine boundary layer air with a low  $\text{O}_3$  content that is carried by the low level southwesterlies towards the monsoon

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



region where the O<sub>3</sub> depleted air is then transported into the free troposphere. It was argued that the well defined O<sub>3</sub> minimum is related to a blocking of northward flows by the Himalayas and the subtropical jet, both representing a barrier to transport (Liu et al., 2009). The steep cut-off observed in the H<sub>2</sub>O mixing ratio between 25° N and 30° N indeed indicates that convective uplifting of humid air is limited to this latitude band south of the Himalayas. Because the lifetime of gaseous H<sub>2</sub>O in the atmosphere is only of the order of days, it is effectively not transported horizontally, and the H<sub>2</sub>O plume is rather stationary, while tracers with longer lifetimes, such as the discussed greenhouse gases, may experience advection leading to a larger latitudinal extension of the plume in July, August and September. The atmospheric lifetime of O<sub>3</sub> depends on photochemistry and is also influenced by convection (Lawrence et al., 2003), but usually is of the order of weeks and thus shorter than the lifetime of the greenhouse gases shown in Fig. 5. (Randel and Park, 2006) showed, using trajectory calculations, that the monsoon anticyclone can trap air masses for up to several weeks depending on altitude, with more effective confinement occurring at higher altitudes. Depending on a tracer's lifetime, advection in the free troposphere and accumulation in the monsoon plume may thus play an important role. The observation that for all discussed long-lived tracers atmospheric mixing ratios decrease towards 40° N is in agreement with the concept of air being confined in the anticyclone by a northern barrier hindering transport to latitudes further north.

#### 4.4 Plume shape and position

In a simplifying approach, the spatial distributions of trace gas mixing ratios in the monsoon plume can be approximated by a Gaussian. Gaussian fits were applied to the observed latitudinal distributions, and from these mean latitudinal positions of the respective trace gas plumes and their maximum mixing ratios at these latitudes were determined. For CO, the continuous on-line measurements have been used. The inferred maximum mixing ratios are shown in Fig. 7a for May through September. For April and October the mean values over the latitude range 12–40° N are shown. For

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



all four gases an increase throughout summer and a subsequent steep decrease at the end of the summer monsoon season is observed. CH<sub>4</sub> and SF<sub>6</sub> show a smooth and monotonous increase from April through August. CO and N<sub>2</sub>O behave similar, but show a local maximum in June.

Figure 7b shows the latitudes at which the maximum mixing ratios are reached from May through September. Again, for all gases a similar systematic behavior is observed with the northernmost position being reached in August followed by a southward shift in September. The plume position as determined from the fits for these gases varies only within 2–7° during one month. This scatter is mainly due to the values determined for SF<sub>6</sub> for which the plume is slightly less well defined. For SF<sub>6</sub> the mixing ratios are lowest, and the analytic uncertainty is, with a value of 0.9%, larger than for the other compounds. For SF<sub>6</sub> the uncertainty of the plume position varies from 1.1° to 3.5° (1- $\sigma$ -uncertainty from fit). For CH<sub>4</sub> and N<sub>2</sub>O the uncertainty of the plume latitude varies between 0.4° and 1.6°, for CO, for which the fit is performed using high resolution data, it is always below 0.1°. The relative error of the maximum mixing ratio is up to 14%. For CH<sub>4</sub>, N<sub>2</sub>O, and SF<sub>6</sub>, the resulting functions are included in the latitudinal distributions shown in Fig. 8.

Figure 7c shows the concomitant changes in wind speed and H<sub>2</sub>O mixing ratios. The latitude at which maximum values of the zonal wind speed were observed indicates the latitudinal shift of the westerly jet, related to the migration of the monsoon anticyclone. From May until July, the maximum wind speed is observed at increasingly northern latitudes. In August, September and October a southward movement of the jet is observed. In addition, the latitude at which a zonal wind speed of ~0 m/s is observed, marking the position of the center of the anticyclonic circulation, is shown. It is located around 25° N in June, July and August, but further south in April, May and October. As an indicator of convective activity the latitude at which a H<sub>2</sub>O mixing ratio of 100 ppm is observed is also shown. At more southern latitudes H<sub>2</sub>O mixing ratios exceed 100 ppm, at more northern latitudes they are lower, as can also be seen in Fig. 6. A clear northward shift of this latitude is observed from May to June. In parallel to the position of

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the anticyclonic center, indicated by zonal wind speeds of 0 m/s, the convective region does not progress north from June to August. Both observables show a southward shift from August to September, following the monsoon retreat. While the measurements of wind speed and H<sub>2</sub>O indicate little change in convective activity from June to August the maximum greenhouse gas mixing ratios are observed in August. This again shows that trapping of air masses in the circulation and accumulation of trace gases may play an important role, mostly relevant for longer-lived tracers.

The monsoon activity has strong variations on different timescales throughout one season which can cause short-time variations in the chemical composition of the atmosphere (Fujinami and Yasunari, 2004; Randel and Park, 2006). Maximum CO values can also be influenced by single pollution events. The CARIBIC flights took place once per month during three consecutive days yielding the first systematic in-situ observations of trace gases inside the monsoon plume. Although the observations do not represent monthly averages, they give a consistent picture of the highly systematic evolution of the plume reflecting the progression of the Indian summer monsoon.

## 5 Trace gas enhancement in the monsoon plume

### 5.1 Enhancements of CH<sub>4</sub>, SF<sub>6</sub>, and N<sub>2</sub>O

In the Northern Hemisphere, ground station measurements show a pronounced seasonal cycle for CH<sub>4</sub> with a smaller amplitude at high altitude locations and a strong north-south gradient. Usually, a summer minimum is observed, predominantly due to reaction of CH<sub>4</sub> with OH (Dlugokencky et al., 1994). In contrast, satellite observations and model simulations show a summer maximum in the free troposphere at 10–40° N. Relative maxima of CH<sub>4</sub> are seen over the Asian and the North American monsoon regions and are associated with enhanced vertical transport (Park et al., 2004). Figure 8a again shows the latitudinal distributions of CH<sub>4</sub> observed by CARIBIC. The measurements in April and October show an increase in mixing ratios with increasing latitude as

observed by ground based measurements (Dlugokencky et al., 1994). But rather than a summer minimum, from June through August increasing mixing ratios are observed, in agreement with satellite observations.

To estimate the increase of the atmospheric methane load in the monsoon region, both, the seasonal and latitudinal variations of the atmospheric background need to be considered first. A background concentration is derived from the monthly mean values obtained from flask measurements at MLO in 2008 (Dlugokencky et al., 2009) and the latitudinal trend calculated from the CARIBIC October data. The resulting background estimates are shown as solid lines in Fig. 8a. Open symbols indicate samples with CH<sub>4</sub> mixing below this background level. The latitudinal extension of the monsoon plume along the CARIBIC flight path indicated by the vertical dashed lines is defined by all samples with CH<sub>4</sub> mixing ratios above the background level. Although there is not yet a distinct plume in April, five samples south of 25° N show enhanced mixing ratios relative to the background. From May through September, CH<sub>4</sub> mixing ratios in the plume are significantly higher than the background values with a maximum enhancement of ~123 ppb observed at 28° N in August.

For SF<sub>6</sub>, shown in Fig. 8b, no plume structure is evident in the latitudinal distributions for April and October. In April, a mean value of 6.41±0.01 ppt is observed, in October it is 6.54±0.01 ppt, resulting from an average monthly increase of 0.022 ppt. This is consistent with a value of 0.021 ppt month<sup>-1</sup> derived from all available CARIBIC data north of 10° N covering the time 1999–2001 and 2007–2009. Assuming the absence of a seasonal cycle and a linear trend for tropospheric SF<sub>6</sub> allows an estimate to be made how large the background values would have been during the summer months if vertical transport would not have been enhanced in the monsoon region (solid horizontal lines in Fig. 8b). The CARIBIC observations in October do not show a latitudinal variation. In April, three samples collected north of 35° N show higher SF<sub>6</sub> mixing ratios than the more southerly samples. These three samples also have lower N<sub>2</sub>O mixing ratios, but do not show deviations in CH<sub>4</sub> or CO. Although the corresponding PV integrals are low (0.51–0.61 PVU), the three samples are separated from the monsoon

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Greenhouse gases in  
the Indian summer  
monsoon plume**

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



region by a high PV intrusion ( $PV=1.6\text{--}3.1$  PVU) extending through the CARIBIC flight track at  $\sim 35\text{--}28^\circ$  N (van Velthoven, 2009).  $O_3$  mixing ratios in this filament exceeded 200 ppbv with maximum values of 460–490 ppb. This feature persisted for the three days during which the four successive CARIBIC flights took place, including the final two flights during which no sampling was performed. The samples collected in this high PV area were excluded in the present analysis, as only tropospheric samples are considered. Across the high PV area, the  $SF_6$  mixing ratio exhibits a gradient with values below 6.45 ppt in the south and higher values in the north. Northwest of this feature, where the samples with high  $SF_6$  mixing ratios were collected, a small anticyclone was located over the Arabian peninsula embedded in the westerly jet, clearly separated from the monsoon anticyclone that was centered around  $15^\circ$  N. For this flight, backward trajectories for all aircraft positions north of  $20^\circ$  N are influenced by the westerly jet and do not originate from the monsoon region. The  $SF_6$  enhancement is thus not related to convection in the monsoon region. Hence, the assumption of a constant  $SF_6$  background in the monsoon regions seems justified. Throughout summer not only the maximum mixing ratios increase in the plume, as shown in Fig. 7, but also the elevation relative to the background shows a clear seasonality. The maximum enhancement of 0.26 ppt in an individual sample is observed in August at  $30^\circ$  N.

$N_2O$  has a weak seasonal cycle in the NH troposphere, with a minimum in summer. Morgan et al. (2004) pointed out that the observed summer minimum in the  $N_2O$  seasonal cycle is caused by the downward flux of  $N_2O$  depleted stratospheric air at high latitudes. Since rapid vertical transport occurs in the monsoon region, and the upper level anticyclonic circulation is associated with enhanced transport of tropospheric air into the stratosphere rather than vice versa (Chen, 1995; Traub and Lelieveld, 2003; Fu et al., 2006; Krishnamurti et al., 2008), influx of stratospheric air is unlikely to influence  $N_2O$  mixing ratios in the monsoon plume, and thus a summer minimum is not expected in the monsoon region. Figure 8c shows the latitudinal distributions of  $N_2O$  measured during the CARIBIC flights. The solid lines indicate background mixing ratios derived from  $N_2O$  data from the NOAA/ESRL halocarbons in situ program

(Dutton et al., 2009). Given are monthly median values calculated from hourly collected air samples analyzed with a gas chromatograph at the Mauna Loa Observatory. In Fig. 8c, the seasonal cycle of  $N_2O$  is not obvious in the MLO data, because it is masked by the continuous increase, and highest mixing ratios at MLO are observed in October. This is not the case for the CARIBIC data. In the monsoon region, higher mixing ratios are observed in summer than in October, with mixing ratios in the plume clearly exceeding the background levels at MLO.

## 5.2 Quantification of tracer enhancements

Considering all samples with  $CH_4$  mixing ratios above the Mauna Loa reference line for the respective month defines the spatial extension of the monsoon plume along the CARIBIC flight track. In Fig. 8, the latitude range that is covered by the plume samples is indicated by the vertical dashed lines. The area between the data points and the background then yields the excess amount of  $CH_4$  in the atmosphere that is contained in the monsoon plume. From the integration of the observed trace gas mixing ratios over the latitude range defined by the samples with elevated  $CH_4$ , the mean mixing ratio increase in the plume can be calculated for each gas. Note that the Gaussian fits in Fig. 8 are included for illustration only, and all integrations have been performed numerically using the measured mixing ratios and sampling positions. Table 1 lists the average increase of  $CH_4$ ,  $SF_6$ ,  $N_2O$ ,  $CO_2$  and  $CO$  for April through September. The numbers for  $CO$  have been calculated from the continuous measurements using mean values observed at MLO as reference values (Novelli and Masarie, 2009).

In April, no significant increase in the mixing ratios at flight altitude is observed for all gases except for  $N_2O$ . The observed  $SF_6$  increase is close to 0 ppt, indicating little convective activity, and the  $CH_4$  increase is only 12.03 ppb, close to the standard deviation of the data which is 10.31 ppb. For  $CH_4$  and  $CO$ , initially strongly increasing enhancements are observed with maxima in August and lower values in September. This pattern is similar for  $SF_6$ , except that in May the increase is almost as strong as in

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



August. This is plausible if air masses would have mainly originated from industrialized regions with strong local SF<sub>6</sub> sources.

For N<sub>2</sub>O, a higher value is observed in August compared to July and September, but the maximum enhancement is observed in June. In April, the increase is similar to that in September. Figure 8 shows that in April the background level from MLO corresponds to the mixing ratios measured in the samples that were collected north of the high PV intrusion, while the samples collected further south have higher values. Throughout the summer, the CARIBIC observations approach the MLO background at the edges of the plume, and in October both data sets agree over the whole latitude range. Various studies on N<sub>2</sub>O emissions from rice paddies show strong variations depending on fertilization and water management (e.g., Cai et al., 1997; Akiyama et al., 2005, and references therein). Especially flooding practices such as intermittent drainages of rice paddies may influence N<sub>2</sub>O emissions. It was found that emissions of CH<sub>4</sub> and N<sub>2</sub>O do not occur simultaneously. CH<sub>4</sub> is mainly emitted during the growing season, when rice paddies are flooded, while the monsoon convection is active. In contrast, N<sub>2</sub>O emissions occur mainly during the fallow period (Cai et al., 1997; Nishimura et al., 2004) after the growing season, and over a short time interval at the beginning of the irrigation period, earlier than the CH<sub>4</sub> emissions. The different seasonality of emissions could thus explain the relatively large increase observed in N<sub>2</sub>O mixing ratios in June.

CO<sub>2</sub> shows an insignificant increase in April and May and a somewhat larger enhancement in June, indicating that at the beginning of the monsoon period, when the strong convection sets in, ground level mixing ratios of CO<sub>2</sub> are still relatively high. This indicates that, although the monsoon onset with strong rainfall happens in late May/early June, there is a delay in the uptake of CO<sub>2</sub> by the vegetation that is reflected in a delay in the upper troposphere seasonal cycle. From July through August, a decrease in the CO<sub>2</sub> mixing ratio is observed in the plume. The minimum in the seasonal cycle occurs in September when also the decrease relative to MLO is largest.

---

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 5.3 Structure of the plume

During its seasonal evolution the plume is changing in size, and it is moving northwest during the growing phase and southeast during the monsoon retreat. With summer progressing, the CARIBIC transect therefore probes different parts of the upper level anticyclone. Comparing outbound and return flights which, as mentioned above, show a systematic difference in flight altitude, higher water vapor mixing ratios were found for the southernmost samples upon leaving Chennai. These samples are collected about 1.5–3 km lower than the corresponding samples from the outbound flights. In the greenhouse gas distributions no influence of the altitude can be seen, and this is also the case for CO and O<sub>3</sub>. This indicates that in the altitude range 8–12 km there is little vertical structure in the plume for the longer-lived gases.

The CARIBIC aircraft crosses the plume at 55–80° E. The maximum greenhouse gas mixing ratios are observed around 28° N 65° E, while the center of the anticyclone, where maximum mixing ratios of CH<sub>4</sub> occur, is located further east, around 28° N 80° E. There, CH<sub>4</sub> mixing ratios are higher but currently nothing is known about the SF<sub>6</sub> concentrations. Comparing the enhancement of both, CH<sub>4</sub> and N<sub>2</sub>O, to that of SF<sub>6</sub>, reveals systematic differences for the samples collected near the plume center and those collected close to the plume edge. Here, plume center denotes the region where maximum mixing ratios were observed along the CARIBIC flight track west of the center of the anticyclonic circulation.

Calculating  $\frac{\Delta C_{\text{CH}_4}}{\Delta C_{\text{SF}_6}}$  for each individual sample that has a CH<sub>4</sub> mixing ratio above the background (filled symbols in Fig. 8, 62 samples in total), indeed shows variations across the plume with maximum values around 24° N and the lowest values at the northern plume edge. A similar distribution is found for  $\frac{\Delta C_{\text{N}_2\text{O}}}{\Delta C_{\text{SF}_6}}$ . For both ratios there is a weak correlation with horizontal wind speed. An explanation for this pattern could be transport of air from the plume center towards its edge and the mixing with air that has been transported upwards earlier. The ratio  $\frac{\Delta C_{\text{CO}_2}}{\Delta C_{\text{SF}_6}}$  also shows a gradient across the

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Greenhouse gases in  
the Indian summer  
monsoon plume**

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

plume with higher values in the center in June and lower values during the subsequent months. In June, a relatively high CO<sub>2</sub> mixing ratio is observed around 24° N because ground level mixing ratios are still high in the beginning of the growing season. In the following months, convection becomes stronger, uplifting more air with a high SF<sub>6</sub> content but with less CO<sub>2</sub>. Air from the plume center diverges with the large-scale circulation, thereby experiencing mixing with older plume air.

Backward trajectories based on ECMWF data show systematic patterns depending on the sampling location. Figure 9 (top panel) shows two examples of 10-day backward trajectories from a flight from Frankfurt to Chennai on 13 August 2008, one collected at 38.3° N 42.2° E, the second collected at 25.3° N 66.6° E. Blue colors indicate high altitudes, red and yellow colors indicate low altitudes. The trajectories are calculated over the sampling period for every 6 s of flight. For the northern sample the trajectory bundle is very compact with little dispersion. The air is moving rapidly within the outer parts of the anticyclone at altitudes above 250 hPa, without ground contact over the period of ten days. Similar trajectories are found for all samples collected in the monsoon plume north of 30° N. This air has already been part of the anticyclonic circulation for more than one week. A detailed analysis of chemical ages of air masses in the plume performed using non-methane hydrocarbon measurements will be published separately (Baker et al., 2010).

In contrast, trajectories for samples from the plume center and from the southern part of the circulation are much more diverse, usually with ground contact one to four days prior to sampling, so that the air had only recently been uplifted. The trajectory bundle is less compact and usually spreads over a larger area at altitudes below 300 hPa. The example in Fig. 9 illustrates the transport of air over the Arabian Sea towards the Indian subcontinent at altitudes below 850 hPa. Between 80° E and 95° E the flow turns around, and the air masses ascend, ultimately entering the upper tropospheric circulation. Also the trajectory patterns for samples collected south of 30° N are less homogeneous than for the northern samples. For all samples collected south of 20° N the turning point of the air flow is located over the Indian landmass or over the Bay of

---

**Greenhouse gases in  
the Indian summer  
monsoon plume**T. J. Schuck et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Bengal, close to the Indian east coast. For these samples, emissions from the densely populated subcontinent will mainly influence free tropospheric trace gas mixing ratios. For samples collected between 20° N and 30° N the position of the turning point varies. Often it lies over the Bay of Bengal, or the turn may occur over the western part of Mainland Southeast Asia. Here, greenhouse emissions are generally lower than from the densely populated northern Indian plains. Emission values quoted for India have therefore been used as a reference. Of the 62 plume samples only five trajectories indicate an influence of emissions east of 100° E. These regions are thus unlikely to influence the CARIBIC measurements in the western part of the anticyclone.

Figure 9 (bottom panel) shows a calculation of five day forward trajectories originating along the flight path of the CARIBIC aircraft for the flight section between 12:00 UTC and 18:00 UTC, for every 3 min of flight. The forward calculation also shows the air flow in the westerly jet north of the monsoon anticyclone that is crossed between ~42° N and ~32° N. Air masses encountered south of ~32° N are part of the anticyclonic circulation. Air masses around ~32° N that are transported with the monsoon circulation may eventually enter the jet stream to be transported east across the Pacific.

## 6 Estimates of emissions from the monsoon region

### 6.1 Scheme for emission estimates

Relating the measurements in the monsoon plume for the different tracers allows an estimate of the emissions during the monsoon period if the emissions of one of the measured tracers are known. In addition, it is required that the compared tracers behave similarly during atmospheric transport. Due to their long lifetimes, the greenhouse gases CH<sub>4</sub>, SF<sub>6</sub>, N<sub>2</sub>O, and CO<sub>2</sub> can be considered as chemically inert tracers during the four months duration of the monsoon period. For CO, with a lifetime of the order of weeks to months, this may still be an acceptable approximation. From the suite of tracers measured on the CARIBIC air samples, SF<sub>6</sub> has been chosen as reference gas

because it is a chemically inert, purely anthropogenic tracer with sources in the South Asian monsoon region that can be assumed to have no seasonal variation.

Frequently, the source-tracer-ratio (STR) method is used to derive emission estimates for a tracer that is measured under the same conditions as a second tracer for which emission rates are known. Emissions are estimated from the correlations of tracer mixing ratios, and for example  $\text{CH}_4$  emissions  $E_{\text{CH}_4}$  can be calculated from emissions  $E_{\text{SF}_6}$  of  $\text{SF}_6$  as

$$E_{\text{CH}_4} = E_{\text{SF}_6} \cdot \text{ER}_{\text{CH}_4/\text{SF}_6}, \quad (1)$$

$\text{ER}_{\text{CH}_4/\text{SF}_6}$  being the emission ratio of  $\text{CH}_4$  to  $\text{SF}_6$  calculated from the correlation of both tracers in the plume (Andreae and Merlet, 2001). The successful usage of  $\text{SF}_6$  as a reference tracer using this source-tracer-ratio method in tracer release experiments has for example been reported in Lamb et al. (1995); Möllmann-Coers et al. (2002). This method assumes the two gases to be emitted from similarly spatially distributed sources and to exhibit identical behavior during atmospheric transport. In the case of the monsoon plume, emission sources are situated in the boundary layer and polluted air is transported into the free troposphere by convection. Because the source region can be considered as a mosaic of point and extended sources, mixing is inherent, apart from large scale systematic differences. Equation (1) does not demand a priori assumptions about the tracer background concentrations in the free troposphere.

Figure 10 shows as an example the correlations of  $\text{CH}_4$ ,  $\text{N}_2\text{O}$ , and  $\text{CO}$  for September. The correlation coefficients  $r$  are 0.49 ( $\text{CH}_4$ ), 0.70 ( $\text{N}_2\text{O}$ ), and 0.58 ( $\text{CO}$ ). Also shown are the respective regression lines and the reference values that were used as background values previously. For all tracers, including  $\text{CO}_2$  which is not shown here, the chosen background values fall within the confidence band of the regression line. The correlation slopes are determined using weighted orthogonal regression, errors are calculated considering the analytical uncertainty of each data point. Because the range of mixing ratios in the plume is rather small, and the reference gas  $\text{SF}_6$  has the largest measurement uncertainty of the discussed tracers, this results in large uncertainties in the estimated emission ratios. The tracer correlations are only statistically

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



significant (on the 95% level) in July and September, and no significant correlation was found for CO<sub>2</sub>, which limits the applicability of the STR method for the current dataset.

A second way to estimate emissions is to compare the enhancement of the tracers relative to the atmospheric background (Andreae and Merlet, 2001). This will be referred to as background method. In this case, assumptions need to be made about tracer background mixing ratios. A mixing ratio increase  $\Delta c$  in the plume relative to the background value  $c$  indicates the existence of an additional net flux  $\varphi$  in summer, driven by convection. Assuming this flux to be proportional to the observed increase of mixing ratios,  $\varphi \sim \Delta c$ , allows to estimate the summer increase of CH<sub>4</sub> emissions. Surface emissions  $E$  represent a flux into the boundary layer of which a fraction  $f < 1$  is transported rapidly up to the plume altitude. This fraction is determined by the local strength of the convective activity. Assuming rapid convective transport without dilution and sinks,  $\varphi = f \cdot E$ , and the increase of mixing ratios  $\Delta c$  is proportional to the emissions  $E$ . The same ratio applies for the other fluxes into the monsoon plume and for the corresponding fluxes out of the boundary layer

$$\frac{\Delta c_{\text{CH}_4}}{\Delta c_{\text{SF}_6}} = \frac{\varphi_{\text{CH}_4}}{\varphi_{\text{SF}_6}} = \frac{f \cdot E_{\text{CH}_4}}{f \cdot E_{\text{SF}_6}} = \frac{E_{\text{CH}_4}}{E_{\text{SF}_6}}, \quad (2)$$

and the summer time emission of CH<sub>4</sub> can be estimated as

$$E_{\text{CH}_4} = E_{\text{SF}_6} \cdot \frac{\Delta c_{\text{CH}_4}}{\Delta c_{\text{SF}_6}}. \quad (3)$$

Supposed that SF<sub>6</sub> emissions remain constant during the monsoon period, it follows that the additional SF<sub>6</sub> in the plume is entirely due to more effective vertical transport during the monsoon period. Emissions of CH<sub>4</sub>, N<sub>2</sub>O and CO can then be calculated for the monsoon months relative to the emissions of SF<sub>6</sub>, and if these are known, absolute fluxes can be derived. Thereby it is again assumed that emissions of SF<sub>6</sub> and the other gases occur from similarly distributed sources, and the same fraction  $f$  of the emissions is transported by convection into the free troposphere, not taking into

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



account regional differences in convection strength. In addition, the assumption of proportionality between the fraction of the emissions that is transported into the free troposphere and  $\Delta c$  is based on the prerequisite that tropospheric sinks are the same for both gases.  $\text{SF}_6$  has no tropospheric sink, and due to the rapid vertical transport and the long lifetime this assumption is certainly justified for  $\text{CH}_4$ ,  $\text{CO}_2$ , and  $\text{N}_2\text{O}$  and to first approximation also for  $\text{CO}$ .

While  $\text{SF}_6$  is an industrial tracer, the most important  $\text{CH}_4$  sources in the monsoon region are rice paddies, wetlands, landfills, and ruminants.  $\text{N}_2\text{O}$  is emitted mainly from soils (Mosier et al., 1998) and estimations of the contribution of rice farming have large uncertainties (Akiyama et al., 2005). These different sources will most likely not be strictly co-located. However, for all gases, strong emissions originate in the densely populated Ganges-Indus plains in Northern India from industrial and agricultural sources. Different types of emission sources are distributed over this region, and it can be assumed that transport into the free troposphere occurs under similar conditions for all gases.

In the following, the EDGAR database (V4.0, Olivier et al., 2005, data obtained from <http://edgar.jrc.ec.europa.eu/index.php>) is used as a reference for greenhouse gas emission rates. For most source categories listed in the emission database, India is the country with the highest emissions in the monsoon region. In particular, emissions of  $\text{SF}_6$  in South Asia are only quoted for India (contributing 85%) and Pakistan (15%). As  $\text{SF}_6$  emissions are closely related to production and transmission of electricity, other countries in this region will likely emit  $\text{SF}_6$ , too. For India,  $\text{SF}_6$  emissions  $E_{\text{SF}_6}$  of 0.192 Gg are reported for 2005, corresponding to a mean monthly flux of 0.016 Gg or  $0.11 \times 10^6$  mol/month. In contrast, strong  $\text{CH}_4$  sources are located around the Bay of Bengal and highest yearly emissions are quoted for India and Bangladesh. Trajectories indicate that the CARIBIC measurements are strongly influenced by emissions from the Indian subcontinent (see also Sect. 5.3), therefore the most recent emission rates quoted for India will be used for comparison. But it has to be reiterated that the footprint is not completely the same for all tracers due to the distribution of sources.

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Greenhouse gases in  
the Indian summer  
monsoon plume**

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Based on the value of  $0.11 \times 10^6$  mol/month for  $\text{SF}_6$  emissions, both, Eqs. (3) and (1) have been used to derive emission estimates for each month during the monsoon period. To determine the value of  $\Delta c$ , for each gas a numerical integration of trace gas mixing ratios over latitude was performed for the samples that were collected in the plume. This procedure yields an average mixing ratio increase in the plume (cf. Table 1). The errors of the obtained emission rates were estimated from the mixing ratio difference observed in samples collected at the same latitude within one day during consecutive flights and from the standard deviation observed in October and April. For the determination of  $\frac{c_{\text{CH}_4}}{c_{\text{SF}_6}}$  a linear fit was applied to the correlation of mixing ratios for the samples collected in the monsoon plume using weighted orthogonal distance regression (ODRPACK, Boggs and Rogers, 1990). The error of the slope is calculated from the measurement uncertainty of each individual data point (Press, 2007). As mentioned previously, it has to be noted that according to a t-test the correlations of  $\text{CH}_4$ ,  $\text{N}_2\text{O}$ , and  $\text{CO}$  with  $\text{SF}_6$  were not significant in June and August, and for  $\text{N}_2\text{O}$  also in July. Thus, large uncertainties are associated with this method. Table 2 lists the emissions of  $\text{CH}_4$ ,  $\text{N}_2\text{O}$ , and  $\text{CO}$  obtained from Eq. (3) for June through September based on an annual  $\text{SF}_6$  flux of 0.192 Gg. Finally, Table 3 shows the resulting emission estimates from the STR-method. Results derived from non-significant correlations are set in parentheses. The results from both methods agree within  $\sim 30\%$  for the months with significant correlations, differences are largest for  $\text{CH}_4$  in June and August, for  $\text{N}_2\text{O}$  in August, and for  $\text{CO}$  in June.

## 6.2 $\text{CH}_4$ emissions

In India,  $\text{CH}_4$  emissions are dominated by enteric fermentation (ruminants) for which no seasonal dependence is expected. In the surrounding countries, this is not a relevant source of  $\text{CH}_4$ . The most important source type with seasonal variation of emissions in the monsoon region is rice cultivation, whose contribution to the total  $\text{CH}_4$  emissions varies strongly for the different countries from 8% (Bhutan) up to 51% (Sri Lanka) with

a regional mean value of 28% in 2005. In India, in 2005, 3.9 Tg out of 28.1 Tg total emissions are attributed to rice cultivation (13%).

The above integration-based estimate results in an average CH<sub>4</sub> emission rate of 1.5 Tg/month during the monsoon months. This is lower than the monthly average of 2.34 Tg/month derived from India's total estimated emissions in 2005 which points to a possible overestimation of CH<sub>4</sub> emissions or an underestimation of SF<sub>6</sub> emissions in South Asia, or both. Another reason for this relatively low value may be the position of the flight track. Along the flight track, slightly lower values for  $\frac{\Delta C_{CH_4}}{\Delta C_{SF_6}}$  were observed towards the plume edge. It is therefore likely that this fraction reaches its maximum in the center of the anticyclone, where, according to satellite observations, also CH<sub>4</sub> mixing ratios are highest. Emission estimates based on measurements off the center of the anticyclone may thus be too low, because the relative increase of CH<sub>4</sub> and SF<sub>6</sub> in the plume may be different. Then again, maximum mixing ratios of ~1900 ppb derived from satellite observations during 2003–2007 are similar to the maximum value measured by CARIBIC in 2008. From comparisons with the TM3 model it was concluded from this data that CH<sub>4</sub> emissions from rice farming are possibly overestimated in the model (Xiong et al., 2009).

For all three gases, CH<sub>4</sub>, N<sub>2</sub>O, and CO, the CARIBIC estimates show variations from month to month. Emissions of CH<sub>4</sub> increase strongly during the summer. This tendency can be explained by a strong increase in emissions from sources such as rice paddies, wetlands and landfills during the rainy season. Indeed, the largest sources in India for which a seasonal variation related to the monsoon rains is conceivable are waste water treatment (4.24 Tg), rice agriculture (3.9 Tg), manure management (1.1 Tg), and waste management (1.1 Tg), in sum accounting for ~1/3 of the total emissions of 28.1 Tg. During the monsoon period, CH<sub>4</sub> emissions strongly increase due to emissions from rice paddies, landfills, and wetlands over most of Asia (Bogner and Spokas, 1993; Chen and Prinn, 2006; Khalil and Butenhoff, 2008). Recent estimates of the total contribution of CH<sub>4</sub> emissions from rice farming to the global methane budget vary between 39 Tg/a (12%) (Olivier et al., 2005) and 112 Tg/a (26%) (Chen and

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Prinn, 2006). Although numerous field studies on methane fluxes from rice paddies are available, the global budget remains difficult to quantify, mainly because emissions have strong temporal and spatial variations, even in one location (Jain et al., 2000; Wassmann et al., 2000; Khalil and Butenhoff, 2008). Methane emissions also depend on agricultural practices such as irrigation patterns, that are varying in different locations and may even vary from season to season (Cai et al., 1997; Huang et al., 2004). Although absolute CH<sub>4</sub> fluxes are not very well known, the seasonality derived from the CARIBIC observations is consistent with the expected strong influence of rice agriculture and seasonally important sources such as wetlands.

### 6.3 Emissions of N<sub>2</sub>O and CO

Emissions of N<sub>2</sub>O are particularly difficult to assess because individual sources such as soil emissions or indirect emissions from rice farming may undergo very different seasonal variations. N<sub>2</sub>O emissions derived from the CARIBIC data show a decrease from 114.5 Gg to 43.4 Gg throughout the summer. This is consistent with results from observations in rice paddies showing an anti-correlation between CH<sub>4</sub> and N<sub>2</sub>O emissions (Nishimura et al., 2004). In the EDGAR inventory, emissions of 632 Gg were quoted for 2005, corresponding to average monthly emissions of 52.7 Gg/month. This is somewhat smaller than the average value of 72.0 Gg/month derived from the observations. However, the monthly mean value from the data archive does not consider the seasonality of emissions from different sources.

CO has a much shorter lifetime than CH<sub>4</sub>, therefore, its sink is more important, and the assumption of insignificant losses is not valid. While losses during the convective vertical transport are small because it happens rapidly, CO reacts with OH at higher altitudes while air masses are transported within the anticyclone. With a rate coefficient of  $k = 1.5 \times 10^{-13} \times (1 + 0.6p) \text{ cm}^3 \text{ s}^{-1}$ , at a pressure of  $p = 200 \text{ hPa}$  and an OH concentration  $[\text{OH}] = 1 \times 10^6 \text{ cm}^{-3}$  for 12 h per day (Spivakovsky et al., 2000; DeMore et al., 1997) about 80% of the initial CO will be left after one month. The above calculation is therefore expected to underestimate the monthly emissions accordingly. In addition, satellite

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





observations (Kar et al., 2004; Park et al., 2009) indicate that the maximum CO mixing ratios in the plume center exceed the maximum value of ~110 ppb measured by CARIBIC so that the position of the flight route west of the center may also result in an underestimation of emissions.

5 The estimated emissions of CO are lower at the beginning of the monsoon season. A plausible reason for this is a decrease in emissions from wildfires and agricultural biomass burning during the rainy season. Consequently, CO emissions will mainly originate from industrial sources, transportation and from domestic biofuel use, and their location will be similar to the distribution of SF<sub>6</sub> sources. On average, a value of  
10 1.2 Tg/month is obtained during the monsoon period. The most recent data available is for 2000 (EDGAR 3.2 Fast Track 2000). Total emissions of 63.8 Tg are quoted for India, including a contribution from biomass burning averaged for 1997–2001. This corresponds to a mean value of 5.3 Tg/month, considerably more than what is derived here from the CARIBIC flights. A seasonality of emissions is not resolved in the EDGAR  
15 data archive, and the monthly mean value derived from the yearly emissions from the data archive is most likely an overestimate during summer, because it does not consider an emission decrease in summer. Biomass burning emission data, based on GFED v2 data (Randerson et al., 2007), show a peak in CO emissions from biomass burning in spring in mainland Southeast Asia and minimum biomass burning activity  
20 during summer.

## 6.4 The CO<sub>2</sub> flux

The last column of Table 2 lists  $E_{\text{CO}_2}$  according to Eq. (3). CO<sub>2</sub> has strong emissions from fossil fuel, biofuel, and wood burning, but there is also a very large downward flux due to uptake of CO<sub>2</sub> by the vegetation during the growing season. Thus, this  
25 number  $E_{\text{CO}_2}$  does not quantify emissions into the free troposphere but rather a net flux of CO<sub>2</sub>. This flux is positive in June but changes sign in July reflecting the CO<sub>2</sub> uptake by the biosphere. This downward flux peaks in September when the lowest mixing ratios are observed at flight altitude. In August, the CO<sub>2</sub> uptake seems to be

---

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**Greenhouse gases in  
the Indian summer  
monsoon plume**T. J. Schuck et al.

---

lower, parallel to a lower CH<sub>4</sub> flux derived for this month, although plant growth and thus photosynthesis are not expected to weaken then. A possible explanation may be the variability of the monsoon activity. Randel and Park (2006) showed that variations in the convection strength, so-called active and break phases of the monsoon, are also reflected in upper tropospheric trace gas concentrations. In August, the upward vertical wind speed measured by the aircraft systems was lower around 20° N, while the highest values were encountered in September. Therefore, the flux decrease in August could reflect a decrease in convective activity masking increased emissions and uptake. The CARIBIC flights took place within three consecutive days during each month. Despite the highly systematic pattern, the representativeness of the data cannot be determined.

## 7 Summary and conclusions

Trace gas measurements from aboard the CARIBIC passenger aircraft during flights between Germany and India in 2008 show a clear monsoon signature in the free troposphere south of 40° N at altitudes between 8 and 12.5 km. The analytical precision of the measurements is good, but improvement for SF<sub>6</sub> is desirable. The most prominent feature is an enhancement in CH<sub>4</sub> mixing ratios, accompanied by elevated levels of SF<sub>6</sub>, N<sub>2</sub>O, and CO throughout the summer monsoon season. These tracers appear to be closely related, showing increasing mixing ratios from May through August followed by a decrease in September. Latitudinal distributions of mixing ratios show a similar plume structure for these four gases. In parallel to the change in maximum mixing ratios, a change in the plume position and extension occurs. From May through August, the plume is expanding and moving northwest, whereas in September during the retreat of the monsoon the plume shifts southeast. In October, after the monsoon season, no enhancement is observed. Elevated mixing ratios are also observed for water vapor. Because of its short lifetime, advection in the free troposphere plays a less important role for water vapor, and its plume does not move northward but is stationary, extending up to 30° N. Water vapor mixing ratios are anti-correlated with mixing

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ratios of  $O_3$ , consistent with the concept that convective mixing moistens the upper troposphere and reduces  $O_3$ .

The monsoon system over South Asia is associated with strong convective transport. At low altitudes, clean marine boundary layer air is transported towards the Asian landmass, and over the continent it is mixed with pollution. Trajectory calculations illustrate this transport of marine boundary layer air towards the Indian subcontinent followed by convective uplifting into the free troposphere. Air sampled in the northern part of the upper tropospheric anticyclone has generally traveled with the high altitude circulation for over one week, while trajectories for samples collected in the southern part of the anticyclone show recent ground contact, usually within the last four days prior to sampling. From the trajectory calculations it can be inferred that mainly emissions from India and the countries around the Bay of Bengal influence the measurements, while emissions from South-East Asia are less likely to be transported westward.

Based on tracer-tracer correlations and on numerical integration of mixing ratios across the plume, monthly emission rates were estimated for the monsoon period. Emissions from the EDGAR database for India have been used for comparison. Most trajectories showed ground contact of air masses over the subcontinent, being one of the most densely populated countries with the highest reported greenhouse gas emissions in the monsoon region.

$SF_6$  was used as a reference gas for emission estimates of  $CH_4$ ,  $N_2O$ , and  $CO$  and for an estimate of the net flux of  $CO_2$  in the monsoon region using measurements at the Mauna Loa observatory as background concentrations. Relating the observed enhancement of free tropospheric mixing ratios to surface fluxes requires precise measurements of a chemically inert tracer, that is emitted in the monsoon region without seasonal variations or for which the seasonality of emissions is well known. Also the absolute emission rates need to be known. From the present data set the compound that is therefore best suited for such emission estimates is  $SF_6$ .

On average, for  $CH_4$  and  $CO$ , integration based emission estimates from the background method were lower than emission rates from the EDGAR database. For  $N_2O$ ,

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Greenhouse gases in  
the Indian summer  
monsoon plume**

T. J. Schuck et al.

a higher value was found. Large uncertainties due to different regional emission patterns related to different agricultural practices make it difficult to quantify the contribution of emissions from rice paddies to the global methane budget. Considered that India is the main emitter of SF<sub>6</sub> in the monsoon region, but only part of the CH<sub>4</sub> emission originate there, higher inferred CH<sub>4</sub> fluxes would be expected. Keeping in mind the limitations of the approach, such as the usage of tracers from not co-located sources and that only a small part of the complete plume is probed, these results point to a possible underestimation of SF<sub>6</sub> emissions in South Asia, an overestimation of CH<sub>4</sub> emissions, or both. The difference found for N<sub>2</sub>O is insignificant and could indicate a seasonality in emissions not captured in the monthly mean emission rates derived from annual values in the database. The large difference found for CO remains puzzling. For CO<sub>2</sub>, a large net flux into the biosphere is found for which we have no corresponding literature value.

Another open question is the small-scale structure in the latitudinal distributions that may be related to the active and break phases of the monsoon. Besides the analysis for greenhouse gases, the CARIBIC samples are also analyzed for various non-methane hydrocarbons (Baker et al., 2009) and halocarbons (O'Sullivan, 2007). Many of these compounds are short-lived and due to their different reaction rates with the OH radical can serve as photochemical “clocks”. Extending the analysis to shorter-lived species may help to explain the structure of the plume structure in more detail. Of interest are for example the ages of air masses in different parts of the plume, and the O<sub>3</sub> production/destruction potential of non-methane hydrocarbons.

Hitherto, CARIBIC provides the only regular multi tracer in-situ measurements in the monsoon plume. We hope that these data may serve as input for model comparisons or comparisons with satellite data and may help to better constrain emission estimates of the discussed greenhouse gases. In addition, it is hoped that in the future more in situ measurements of trace gases in the monsoon plume can be performed, including higher precision measurements of SF<sub>6</sub>, preferably on a regular basis.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

---

**Greenhouse gases in  
the Indian summer  
monsoon plume**

---

T. J. Schuck et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

*Acknowledgements.* We wish to thank all CARIBIC partners as well as Lufthansa, especially T. Dauer and A. Waibel, and Lufthansa Technik for their ongoing support. We especially acknowledge D. Scharffe for the operation of this complex platform and C. Koeppel for the handling of the whole air sampler. The GC system has initially been set up by D. Worthy and P. Bergamaschi, A. Jordan gave valuable advice on calibration issues. We thank J. Lelieveld for his helpful comments on the manuscript and T. Rhee for many fruitful discussions. The Mauna Loa data was provided by E. Dlugokencky, G. Dutton, T. Conway, P. Novelli from NOAA/ESRL. We acknowledge the support of the European Commission through the GEOmon (Global Earth Observation and Monitoring) Integrated Project under the 6th Framework Program (contract number FP6-2005-Global-4-036677). T. J. S. acknowledges the Paul Crutzen Nobelpreisträger-Stipendium granted by the Max-Planck-Society.

The service charges for this open access publication have been covered by the Max Planck Society.

## References

- Akiyama, H., Yagi, K., and Yan, X.: Direct N<sub>2</sub>O emissions from rice paddy fields: summary of available data, *Global Biogeochem. Cy.*, 19, GB1005, doi:10.1029/2004GB002378, 2005. 2053, 2059
- Andreae, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, *Global Biogeochem. Cy.*, 15, 955–966, doi:10.1029/2000GB001382, 2001. 2057, 2058
- Baker, A. K., Slemr, F., and Brenninkmeijer, C. A. M.: Analysis of non-methane hydrocarbons in air samples collected aboard the CARIBIC passenger aircraft, *Atmos. Meas. Tech. Discuss.*, 2, 2377–2401, 2009, <http://www.atmos-meas-tech-discuss.net/2/2377/2009/>. 2046, 2066
- Baker, A. K., Brenninkmeijer, C. A. M., Slemr, F., et al., in preparation, 2010. 2046, 2055
- Bhat, G. and Narashima, N.: Indian summer monsoon experiments, *Curr. Sci. India*, 93, 153–164, 2007. 2035
- Boggs, P. and Rogers, J.: Orthogonal distance regression, *Contemp. Math.*, 112, 183–194, 1990. 2060

**Greenhouse gases in  
the Indian summer  
monsoon plume**

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Bogner, J. and Spokas, K.: Landfill CH<sub>4</sub>: rates, fates, and role in global carbon cycle, *Chemosphere*, 26, 369–386, doi:10.1016/0045-6535(93)90432-5, 1993. 2061
- Brenninkmeijer, C. A. M., Crutzen, P., Boumard, F., Dauer, T., Dix, B., Ebinghaus, R., Filippi, D., Fischer, H., Franke, H., Frieß, U., Heintzenberg, J., Helleis, F., Hermann, M., Kock, H. H., Koepfel, C., Lelieveld, J., Leuenberger, M., Martinsson, B. G., Miemczyk, S., Moret, H. P., Nguyen, H. N., Nyfeler, P., Oram, D., O'Sullivan, D., Penkett, S., Platt, U., Pucek, M., Ramonet, M., Randa, B., Reichelt, M., Rhee, T. S., Rohwer, J., Rosenfeld, K., Scharffe, D., Schlager, H., Schumann, U., Slemr, F., Sprung, D., Stock, P., Thaler, R., Valentino, F., van Velthoven, P., Waibel, A., Wandel, A., Waschitschek, K., Wiedensohler, A., Xueref-Remy, I., Zahn, A., Zech, U., and Ziereis, H.: Civil Aircraft for the regular investigation of the atmosphere based on an instrumented container: The new CARIBIC system, *Atmos. Chem. Phys.*, 7, 4953–4976, 2007, <http://www.atmos-chem-phys.net/7/4953/2007/>. 2036, 2037
- Buchwitz, M., Schneising, O., Burrows, J. P., Bovensmann, H., Reuter, M., and Notholt, J.: First direct observation of the atmospheric CO<sub>2</sub> year-to-year increase from space, *Atmos. Chem. Phys.*, 7, 4249–4256, 2007, <http://www.atmos-chem-phys.net/7/4249/2007/>. 2035
- Buchwitz, M., Schneising, O., Burrows, J. P., Bovensmann, H., Reuter, M., and Notholt, J.: Corrigendum to “First direct observation of the atmospheric CO<sub>2</sub> year-to-year increase from space” published in *Atmos. Chem. Phys.*, 7, 4249–4256, 2007, *Atmos. Chem. Phys.*, 7, 5341–5342, 2007, <http://www.atmos-chem-phys.net/7/5341/2007/>. 2035
- Cai, Z., Xing, G., Yan, X., Xu, H., Tsuruta, H., Yagi, K., and Minami, K.: Methane and nitrous oxide emissions from rice paddy fields as affected by nitrogen fertilisers and water management, *Plant Soil*, 196, 7–14, doi:10.1023/A:1004263405020, 1997. 2053, 2062
- Chédin, A., Serrar, S., Scott, N. A., Crevoisier, C., and Armante, R.: First global measurement of midtropospheric CO<sub>2</sub> from NOAA polar satellites: tropical zone, *J. Geophys. Res.*, 108, 4581, doi:10.1029/2003JD003439, 2003. 2035
- Chen, P.: Isentropic cross-tropopause mass exchange in the extratropics, *J. Geophys. Res.*, 100, 16661–16674, doi:10.1029/95JD01264, 1995. 2033, 2051
- Chen, Y.-H. and Prinn, R. G.: Estimation of atmospheric methane emissions between 1996 and 2001 using a three-dimensional global chemical transport model, *J. Geophys. Res.*, 111, 10307, doi:10.1029/2005JD006058, 2006. 2061
- Conway, T., Lang, P., and Masarie, K.: Carbon Dioxide Dry Air Mole Fractions from the NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network, 1968–2008, Version: 2009-07-15, <ftp://ftp.cmdl.noaa.gov/ccg/co2/flask/event/>, last access: 2 September, 2009. 2043



---

**Greenhouse gases in  
the Indian summer  
monsoon plume**T. J. Schuck et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Crevoisier, C., Chédin, A., Matsueda, H., Machida, T., Armante, R., and Scott, N. A.: First year of upper tropospheric integrated content of CO<sub>2</sub> from IASI hyperspectral infrared observations, *Atmos. Chem. Phys.*, 9, 4797–4810, 2009, <http://www.atmos-chem-phys.net/9/4797/2009/>. 2034

5 DeMore, W. B., Sander, S. P., Golden, D. M., Hampson, R. F., Kurylo, M. J., Howard, C. J., Ravishankara, A. R., Kolb, C. E., and Molina, M. J.: Chemical kinetics and photochemical data for use in stratospheric modeling, JPL Publication, 97–4, 1997. 2062

Dlugokencky, E., Lang, P., and Masarie, K.: Atmospheric Methane Dry Air Mole Fractions from the NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network, 1983–2008, Version: 2009-06-18, <ftp://ftp.cmdl.noaa.gov/ccg/ch4/flask/event/>, last access: 4 August, 2009. 2041, 2050

Dlugokencky, E. J., Steele, L. P., Lang, P. M., and Masarie, K. A.: The growth rate and distribution of atmospheric methane, *J. Geophys. Res.*, 99, 17021–17044, doi:10.1029/94JD01245, 1994. 2049, 2050

15 Dutton, G., Dutton, J. W., and Hall, B.: Nitrous Oxide N<sub>2</sub>O data from hourly in situ samples analyzed on a gas chromatograph located at Mauna Loa (MLO), Hawaii (19.539 N, 155.578 W, elevation: 3397 m), <ftp://ftp.cmdl.noaa.gov/hats/n2o/in situGCs/CATS/monthly/>, last access: 17 August, 2009. 2052

20 Fu, R., Hu, Y., Wrigh, J. S., Jiang, J. H., Dickinson, R. E., Chen, M., Filipiak, M., Read, W. G., Waters, J. W., and Wu, D. L.: Short circuit of water vapor and polluted air to the global stratosphere by convective transport over the Tibetan Plateau, *P. Natl. Acad. Sci. USA.*, 103, 5664–5669, doi:10.1073/pnas.0601584103, 2006. 2033, 2034, 2051

Fujinami, H. and Yasunari, T.: Submonthly variability of convection and circulation over and around the Tibetan Plateau during the boreal summer, *J. Meteorol. Soc. Jpn.*, 82, 1545–1564, 2004. 2049

25 Gurk, C., Fischer, H., Hoor, P., Lawrence, M. G., Lelieveld, J., and Wernli, H.: Airborne in situ measurements of vertical, seasonal and latitudinal distributions of carbon dioxide over Europe, *Atmos. Chem. Phys.*, 8, 6395–6403, 2008, <http://www.atmos-chem-phys.net/8/6395/2008/>. 2044

30 Hoskins, B. J. and Rodwell, M. J.: A model of the Asian summer monsoon. Part I: The global scale, *J. Atmos. Sci.*, 52, 1329–1340, doi:10.1175/1520-0469(1995)052, 1995. 2032

Hsu, H.-H., Terng, C.-T., and Chen, C.-T.: Evolution of large-scale circulation and heating during the first transition of Asian summer monsoon, *J. Climate*, 12, 793–810,

**Greenhouse gases in  
the Indian summer  
monsoon plume**

T. J. Schuck et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

doi:10.1175/1520-0442(1999)012, 1999. 2032, 2039

Huang, Y., Zhang, W., Zheng, X., Li, J., and Yu, Y.: Modeling methane emission from rice paddies with various agricultural practices, *J. Geophys. Res.*, 109, 8113, doi:10.1029/2003JD004401, 2004. 2062

5 IMD: Southwest Monsoon 2008: End of Season Report, India Meteorological Department, New-Delhi, 2008. 2033, 2040

Jain, M., Kumar, S., Wassmann, R., Mitra, S., Singh, S., Singh, J., Singh, R., Yadav, A. K., and Gupta, S.: Methane emissions from irrigated rice fields in Northern India (New Delhi), *Nutr. Cycl. Agroecosys.*, 58, 75–83, doi:10.1023/A:1009882216720, 2000. 2062

10 Joseph, P. V., Sooraj, K. P., and Rajan, C. K.: The summer monsoon onset process over South Asia and an objective method for the date of monsoon onset over Kerala, *Int. J. Climatol.*, 26, 1871–1893, doi:10.1002/joc.134, 2006. 2033, 2040

Kar, J., Bremer, H., Drummond, J. R., Rochon, Y. J., Jones, D. B. A., Nichitiu, F., Zou, J., Liu, J., Gille, J. C., Edwards, D. P., Deeter, M. N., Francis, G., Ziskin, D., and Warner, J.: Evidence of vertical transport of carbon monoxide from Measurements of Pollution in the Troposphere (MOPITT), *Geophys. Res. Lett.*, 31, 23105, doi:10.1029/2004GL021128, 2004. 2063

Khalil, M. A. K. and Butenhoff, C. L.: Spatial variability of methane emissions from rice fields and implications for experimental design, *J. Geophys. Res.*, 113, G00A09, doi:10.1029/2007JG000517, 2008. 2061, 2062

20 Krishnamurti, T. N. and Bhalme, H. N.: Oscillations of a Monsoon System. Part I. Observational Aspects., *J. Atmos. Sci.*, 33, 1937–1954, doi:10.1175/1520-0469(1976)033, 1976. 2033, 2039

Krishnamurti, T. N., Jha, B., Rasch, P. J., and Ramanathan, V.: A high resolution global reanalysis highlighting the winter monsoon, Part I: Reanalysis fields, *Meteorol. Atmos. Phys.*, 64, 123–150, 1997. 2039

25 Krishnamurti, T. N., Biswas, M. K., and Rao, D. V. B.: Vertical extension of the Tibetan high of the Asian summer monsoon, *Tellus A*, 60, 1038–1052, doi:10.1111/j.1600-0870.2008.00359.x, 2008. 2033, 2039, 2051

30 Lamb, B., McManus, J., Shorter, J., Kolb, C., Mosher, B., Harriss, R., Allwine, E., Blaha, D., Howard, T., Guenther, A., Lott, R., Siverson, R., Westburg, H., and Zimmerman, P.: Development of atmospheric tracer methods to measure methane emissions from natural gas facilities and urban areas, *Environ. Sci. Technol.*, 29, 1468–1479, doi:10.1021/es00006a007, 1995. 2057



**Greenhouse gases in  
the Indian summer  
monsoon plume**

T. J. Schuck et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Lawrence, M. G., Rasch, P. J., von Kuhlmann, R., Williams, J., Fischer, H., de Reus, M., Lelieveld, J., Crutzen, P. J., Schultz, M., Stier, P., Huntrieser, H., Heland, J., Stohl, A., Forster, C., Elbern, H., Jakobs, H., and Dickerson, R. R.: Global chemical weather forecasts for field campaign planning: predictions and observations of large-scale features during MINOS, CONTRACE, and INDOEX, *Atmos. Chem. Phys.*, 3, 267–289, 2003, <http://www.atmos-chem-phys.net/3/267/2003/>. 2047
- Lelieveld, J., Crutzen, P. J., Ramanathan, V., Andreae, M. O., Brenninkmeijer, C. A. M., Campos, T., Cass, G. R., Dickerson, R. R., Fischer, H., de Gouw, J. A., Hansel, A., Jefferson, A., Kley, D., de Laat, A. T. J., Lal, S., Lawrence, M. G., Lobert, J. M., Mayol-Bracero, O. L., Mitra, A. P., Novakov, T., Oltmans, S. J., Prather, K. A., Reiner, T., Rodhe, H., Scheeren, H. A., Sikka, D., and Williams, J.: The Indian Ocean Experiment: widespread air pollution from South and Southeast Asia, *Science*, 291, 1031–1036, doi:10.1126/science.1057103, 2001. 2035
- Lelieveld, J., Berresheim, H., Borrmann, S., Crutzen, P. J., Dentener, F. J., Fischer, H., Feichter, J., Flatau, P. J., Heland, J., Holzinger, R., Korrman, R., Lawrence, M. G., Levin, Z., Markowicz, K. M., Mihalopoulos, N., Minikin, A., Ramanathan, V., de Reus, M., Roelofs, G. J., Scheeren, H. A., Sciare, J., Schlager, H., Schultz, M., Siegmund, P., Steil, B., Stephanou, E. G., Stier, P., Traub, M., Warneke, C., Williams, J., and Ziereis, H.: Global air pollution crossroads over the mediterranean, *Science*, 298, 794–799, doi:10.1126/science.1075457, 2002. 2035, 2039
- Lelieveld, J., Brühl, C., Jöckel, P., Steil, B., Crutzen, P. J., Fischer, H., Giorgetta, M. A., Hoor, P., Lawrence, M. G., Sausen, R., and Tost, H.: Stratospheric dryness: model simulations and satellite observations, *Atmos. Chem. Phys.*, 7, 1313–1332, 2007, <http://www.atmos-chem-phys.net/7/1313/2007/>. 2033
- Li, Q., Jacob, D. J., Logan, J. A., Bey, I., Yantosca, R. M., Liu, H., Martin, R. V., Fiore, A. M., Field, B. D., Duncan, B. N., and Thouret, V.: A tropospheric ozone maximum over the Middle East, *Geophys. Res. Lett.*, 28, 3235–3238, 2001. 2046
- Liu, J. J., Jones, D. B. A., Worden, J. R., Noone, D., Parrington, M., and Kar, J.: Analysis of the summertime buildup of tropospheric ozone abundances over the Middle East and North Africa as observed by the Tropospheric Emission Spectrometer instrument, *J. Geophys. Res.*, 114, D05304, doi:10.1029/2008JD010993, 2009. 2046
- Liu, J. J., Jones, D. B. A., Worden, J. R., Noone, D., Parrington, M., and Kar, J.: Correction to “Analysis of the summertime buildup of tropospheric ozone abundances over the Middle

---

**Greenhouse gases in  
the Indian summer  
monsoon plume**T. J. Schuck et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

East and North Africa as observed by the Tropospheric Emission Spectrometer Instrument”,  
J. Geophys. Res., 114, 7399, doi:10.1029/2009JD012045, 2009. 2046

Liu, Y., Wang, Y., Liu, X., Cai, Z., and Chance, K.: Tibetan middle tropospheric ozone minimum in June discovered from GOME observations, Geophys. Res. Lett., 36, L05814, doi:10.1029/2008GL037056, 2009. 2034, 2046, 2047

Matsueda, H., Machida, T., Sawa, Y., Nakagawa, Y., Hirokuni, K., Ikeda, H., Kondo, N., and Goto, K.: Evaluation of atmospheric CO<sub>2</sub> measurements from new flask air sampling of JAL airliner observations, Pap. Meteorol. Geophys., 59, 1–17, 2008. 2044

Möllmann-Coers, M., Klemp, D., Mannschreck, K., and Slemr, F.: Determination of anthropogenic emissions in the Augsburg area by the source-tracer-ratio method, Atmos. Environ., 36, 95–107, doi:10.1016/S1352-2310(02)00212-1, 2002. 2057

Morgan, C. G., Allen, M., Liang, M. C., Shia, R. L., Blake, G. A., and Yung, Y. L.: Isotopic fractionation of nitrous oxide in the stratosphere: comparison between model and observations, J. Geophys. Res., 109, D04305, doi:10.1029/2003JD003402, 2004. 2051

Mosier, A., Kroeze, C., Nevison, C., Oenema, O., Seitzinger, S., and van Cleemput, O.: Closing the global N<sub>2</sub>O budget: nitrous oxide emissions through the agricultural nitrogen cycle, Nutr. Cycl. Agroecosys., 52, 225–248, doi:10.1023/A:1009740530221, 1998. 2059

Nishimura, S., Sawamoto, T., Akiyama, H., Sudo, S., and Yagi, K.: Methane and nitrous oxide emissions from a paddy field with Japanese conventional water management and fertilizer application, Global Biogeochem. Cy., 18, GB2017, doi:10.1029/2003GB002207, 2004. 2053, 2062

Novelli, P. and Masarie, K.: Carbon Monoxide Dry Air Mole Fractions from the NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network, 1988–2008, Version: 2009-07-28, ftp://ftp.cmdl.noaa.gov/ccg/co/flask/event/, last access: 2 September, 2009. 2052

Olivier, J., Van Aardenne, J., Dentener, F., Pagliari, V., Ganzeveld, L., and Peters, J.: Recent trends in global greenhouse gas emissions: regional trends 1970–2000 and spatial distribution of key sources in 2000, Environm. Sci., 2, 81–99, doi:10.1080/15693430500400345, 2005. 2061

O’Sullivan, D. A.: Temporal and spatial variability of halogenated compounds and other trace gases, Ph.D. thesis, University of East Anglia, 2007. 2066

Park, M., Randel, W. J., Kinnison, D. E., Garcia, R. R., and Choi, W.: Seasonal variation of methane, water vapor, and nitrogen oxides near the tropopause: satellite observations and model simulations, J. Geophys. Res., 109, 3302, doi:10.1029/2003JD003706, 2004. 2034,

2049

Park, M., Randel, W. J., Emmons, L. K., Bernath, P. F., Walker, K. A., and Boone, C. D.: Chemical isolation in the Asian monsoon anticyclone observed in Atmospheric Chemistry Experiment (ACE-FTS) data, *Atmos. Chem. Phys.*, 8, 757–764, 2008,

<http://www.atmos-chem-phys.net/8/757/2008/>. 2034

Park, M., Randel, W. J., Emmons, L. K., and Livesey, N. J.: Transport pathways of carbon monoxide in the Asian summer monsoon diagnosed from Model of Ozone and Related Tracers (MOZART), *J. Geophys. Res.*, 114, 08303, doi:10.1029/2008JD010621, 2009. 2034, 2063

Press, W.: *Numerical Recipes: The Art of Scientific Computing*, Cambridge University Press, 3 edition, 2007. 2060

Randel, W. J. and Park, M.: Deep convective influence on the Asian summer monsoon anticyclone and associated tracer variability observed with Atmospheric Infrared Sounder (AIRS), *J. Geophys. Res.*, 111, 12314, doi:10.1029/2005JD006490, 2006. 2034, 2046, 2047, 2049, 2064

Randerson, J. T., van der Werf, G. R., Giglio, L., Collatz, G. J., and Kasibhatla, P. S.: Global Fire Emissions Database, Version 2 (GFEDv2.1), Data set, available on-line <http://daac.ornl.gov/>, last access: 30 March 2007, from Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, USA, doi:10.3334/ORNLDAAAC/849, 2007 2063

Sanjeeva Rao, P.: Arabian Sea Monsoon Experiment: an overview, *Mausam*, 56, 1–6, 2005. 2035

Scheele, M., Siegmund, P., and Velthoven, P. V.: Sensitivity of trajectories to data resolution and its dependence on the starting point: in or outside a tropopause fold, *Meteorol. Appl.*, 3, 267–273, doi:10.1002/met.5060030308, 1996. 2038

Scheeren, H. A., Lelieveld, J., Roelofs, G. J., Williams, J., Fischer, H., de Reus, M., de Gouw, J. A., Warneke, C., Holzinger, R., Schlager, H., Klüpfel, T., Bolder, M., van der Veen, C., and Lawrence, M.: The impact of monsoon outflow from India and Southeast Asia in the upper troposphere over the eastern Mediterranean, *Atmos. Chem. Phys.*, 3, 1589–1608, 2003, <http://www.atmos-chem-phys.net/3/1589/2003/>. 2035

Schneising, O., Buchwitz, M., Burrows, J. P., Bovensmann, H., Reuter, M., Notholt, J., Macatangay, R., and Warneke, T.: Three years of greenhouse gas column-averaged dry air mole fractions retrieved from satellite - Part 1: Carbon dioxide, *Atmos. Chem. Phys.*, 8, 3827–3853, 2008,

ACPD

10, 2031–2087, 2010

---

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**Greenhouse gases in  
the Indian summer  
monsoon plume**

---

T. J. Schuck et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

<http://www.atmos-chem-phys.net/8/3827/2008/>. 2035

Schneising, O., Buchwitz, M., Burrows, J. P., Bovensmann, H., Bergamaschi, P., and Peters, W.: Three years of greenhouse gas column-averaged dry air mole fractions retrieved from satellite - Part 2: Methane, *Atmos. Chem. Phys.*, 9, 443–465, 2009,

<http://www.atmos-chem-phys.net/9/443/2009/>. 2034

Schuck, T. J., Brenninkmeijer, C. A. M., Slemr, F., Xueref-Remy, I., and Zahn, A.: Greenhouse gas analysis of air samples collected onboard the CARIBIC passenger aircraft, *Atmos. Meas. Tech.*, 2, 449–464, 2009,

<http://www.atmos-meas-tech.net/2/449/2009/>. 2037

Spivakovsky, C. M., Logan, J. A., Montzka, S. A., Balkanski, Y. J., Foreman-Fowler, M., Jones, D. B. A., Horowitz, L. W., Fusco, A. C., Brenninkmeijer, C. A. M., Prather, M. J., Wofsy, S. C., and McElroy, M. B.: Three-dimensional climatological distribution of tropospheric OH: update and evaluation, *J. Geophys. Res.*, 105, 8931–8980, doi:10.1029/1999JD901006, 2000. 2062

Traub, M. and Lelieveld, J.: Cross-tropopause transport over the eastern Mediterranean, *J. Geophys. Res.*, 108(D23), 4712, doi:10.1029/2003JD003754, 2003. 2033, 2051

van Velthoven, P. F. J.: Meteorological analysis of CARIBIC by KNMI, [http://www.knmi.nl/samenw/campaign\\_support/CARIBIC/#LH](http://www.knmi.nl/samenw/campaign_support/CARIBIC/#LH), last access: December, 2009. 2038, 2051

Wang, B., Ding, Q., and Joseph, P. V.: Objective definition of the Indian Summer Monsoon onset, *J. Climate*, 22, 3303–3316, doi:10.1175/2008JCLI2675.1, 2009. 2033, 2040

Wassmann, R., Neue, H.-U., Lantin, R., Buendia, L., and Rennenberg, H.: Characterization of methane emissions from rice fields in Asia. I. comparison among field sites in five countries, *Nutr. Cycl. Agroecosys.*, 58, 1–12, doi:10.1023/A:1009848813994, 2000. 2062

Worden, J., Jones, D. B. A., Liu, J., Parrington, M., Bowman, K., Stajner, I., Beer, R., Jiang, J., Thouret, V., Kulawik, S., Li, J.-L. F., Verma, S., and Worden, H.: Observed vertical distribution of tropospheric ozone during the Asian summertime monsoon, *J. Geophys. Res.*, 114, D13304, doi:10.1029/2008JD010560, 2009. 2046

Xiong, X., Houweling, S., Wei, J., Maddy, E., Sun, F., and Barnet, C.: Methane plume over south Asia during the monsoon season: satellite observation and model simulation, *Atmos. Chem. Phys.*, 9, 783–794, 2009,

<http://www.atmos-chem-phys.net/9/783/2009/>. 2034, 2042, 2061

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

**Table 1.** Enhancement of CH<sub>4</sub>, SF<sub>6</sub>, N<sub>2</sub>O, and CO<sub>2</sub> during the monsoon period, integrating along the CARIBIC flight track. No plume is observed in October.

	$\Delta\text{CH}_4$ [ppb]	$\Delta\text{SF}_6$ [ppt]	N <sub>2</sub> O [ppb]	$\Delta\text{CO}$ [ppbv]	$\Delta\text{CO}_2$ [ppm]
Error estimate	±10	±0.02	±0.1	±5	±0.3
April	12.03	−0.01	0.55	2.81	0.34
May	16.73	0.08	0.50	2.56	0.37
June	18.52	0.04	0.90	7.70	0.95
July	45.13	0.05	0.63	19.29	−1.10
August	65.76	0.09	0.81	29.36	−0.52
September	51.92	0.04	0.58	18.68	−1.65
October	–	–	–	–	–
June–September av.	45.33	0.06	0.73	18.75	−0.58

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

**Table 2.** Estimates of total net fluxes for CH<sub>4</sub>, N<sub>2</sub>O, CO, and CO<sub>2</sub> based on India's assumed SF<sub>6</sub> emissions in 2005. Error estimates are ~50% for all tracers. Emission data from the EDGAR inventory are for India in 2005 (v4.0, CH<sub>4</sub>, N<sub>2</sub>O) and 2000 (v3.2, CO).

	CH <sub>4</sub> [Tg/month]	N <sub>2</sub> O [Gg/month]	CO [Tg/month]	CO <sub>2</sub> [Tg/month]
June	0.9	114.5	0.7	121.2
July	1.6	60.6	1.3	-106.5
August	1.3	43.4	1.1	-28.1
September	2.3	69.6	1.6	-198.9
June–September av.	1.5	72.0	1.2	-53.1
EDGAR database	2.3	52.7	5.3	–

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.

**Table 3.** Estimates of total net fluxes for CH<sub>4</sub>, N<sub>2</sub>O, and CO based on India's assumed SF<sub>6</sub> emissions in 2005 using the STR-method. Emission data from the EDGAR inventory are for 2005 (CH<sub>4</sub>, N<sub>2</sub>O) and 2000 (CO). Values derived from non-significant correlations are set in parentheses.

	CH <sub>4</sub> [Tg/month]	N <sub>2</sub> O [Gg/month]	CO [Tg/month]
June	(1.3±1.5)	(109.1±146.5)	(1.9±2.3)
July	1.4±0.7	(58.4± 32.5)	1.0±0.5
August	(4.0±4.1)	(12.5± 7.0)	(1.0±0.6)
September	3.1±3.4	61.5± 46.6	1.9±1.8
June–September av.	2.5	60.4	1.5
EDGAR database	2.3	52.7	5.3

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Greenhouse gases in  
the Indian summer  
monsoon plume**

T. J. Schuck et al.




**Fig. 1.** CARIBIC flights between Germany and India in 2008. Shown are the flight routes for 16 round trips between Frankfurt and Chennai from April to December 2008. On 9 round trips air samples were taken at the positions indicated.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



**Greenhouse gases in  
the Indian summer  
monsoon plume**

T. J. Schuck et al.




**Fig. 2.** Zonal wind speed as a function of latitude observed during CARIBIC flights to India in summer and winter 2008. Spring and autumn months are not shown. Positive wind speeds indicate westerly flows, negative wind speeds easterly flows.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.



**Fig. 3.** Wind speed at 250 hPa on 13 August 2008 for a CARIBIC flight from Germany to India. Contours and coloring are in steps of 10 m/s for wind speeds between 20 m/s and 70 m/s. The CARIBIC flight path is indicated by the solid line, markers show the positions where air samples were collected.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Greenhouse gases in  
the Indian summer  
monsoon plume

T. J. Schuck et al.




**Fig. 4.** Mixing ratios of CH<sub>4</sub> as a function of latitude for flights between Frankfurt and Chennai from April to October 2008. The horizontal lines indicate the respective mean mixing ratio observed at Mauna Loa Observatory (19° N).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Greenhouse gases in  
the Indian summer  
monsoon plume

T. J. Schuck et al.




**Fig. 5.** Latitudinal distributions of  $\text{CH}_4$ ,  $\text{CO}_2$ ,  $\text{N}_2\text{O}$ , and  $\text{SF}_6$  from air samples and on-line measured  $\text{CO}$  mixing ratios integrated over the sampling period for flights between Frankfurt and Chennai in June, July, August, and September 2008. For  $\text{CO}_2$  the mean mixing ratio observed at Mauna Loa Observatory ( $19^\circ\text{N}$ ) is indicated by a solid line.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Greenhouse gases in  
the Indian summer  
monsoon plume

T. J. Schuck et al.




**Fig. 6.** Latitudinal distributions of H<sub>2</sub>O and O<sub>3</sub> mixing ratios integrated over the sampling periods of whole air samples for flights between Frankfurt and Chennai in June, July, August and September 2008.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Greenhouse gases in  
the Indian summer  
monsoon plume

T. J. Schuck et al.



**Fig. 7.** (a) Maximum mixing ratios of CH<sub>4</sub>, CO, N<sub>2</sub>O, and SF<sub>6</sub> observed in summer 2008 south of 40° N. For April and October, when no monsoon plume is apparent, the monthly mean values are plotted. (b) Latitudinal position at which the maximum mixing ratios of CH<sub>4</sub>, CO, N<sub>2</sub>O, and SF<sub>6</sub> were observed. (c) Latitudinal position of the maximum wind speed in the westerly jet and the latitude where the zonal wind speed is 0 m/s, both indicating the latitudinal position of the anticyclone. The latitude where a H<sub>2</sub>O mixing ratio of 100 ppm is observed indicates the northern extension of the strong convection.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Greenhouse gases in  
the Indian summer  
monsoon plume

T. J. Schuck et al.




**Fig. 8.** Latitudinal distributions of CH<sub>4</sub> (a), SF<sub>6</sub> (b), and N<sub>2</sub>O (c) for April–October 2008. For May–September a Gaussian has been fit to the data. Solid lines indicate background levels determined from the CARIBIC measurements in April and October for SF<sub>6</sub> and from the Mauna Loa Observatory for CH<sub>4</sub> and N<sub>2</sub>O. All samples for which CH<sub>4</sub> mixing ratios are above the reference background are indicated by closed symbols, open symbols denote samples with lower CH<sub>4</sub> mixing ratios. The vertical dashed lines mark the integration limits used to calculate the increase in the monsoon plume.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Greenhouse gases in the Indian summer monsoon plume

T. J. Schuck et al.



**Fig. 9.** Top: Ten day backward trajectories for two samples collected on 13 August 2008 during a flight from Frankfurt to Chennai. Along the trajectories, small markers indicate time steps of 24 h, the large symbols denote the sampling positions. Bottom: Five day forward trajectories calculated every 3 min along the flight track on 13 August 2008.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion





Greenhouse gases in  
the Indian summer  
monsoon plume

T. J. Schuck et al.



**Fig. 10.** Correlation of CH<sub>4</sub>, N<sub>2</sub>O, and CO with SF<sub>6</sub> in the monsoon plume in September. The black lines result from a weighted orthogonal distance regression, the black squares indicate the background mixing ratios (SF<sub>6</sub> interpolated from CARIBIC data, CH<sub>4</sub> and CO mean value, N<sub>2</sub>O median observed at MLO in September 2008).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)