

Supplementary information for:

Primary sources of PM_{2.5} particles in an industrial Mediterranean city, Marseille

Imad El Haddad^{1*}, Nicolas Marchand¹, Henri Wortham¹, Christine Piot^{2,3}, Jean-Luc Besombes², Julie Cozic³, Catherine Chauvel⁴, Alexandre Armengaud⁵, Dominique Robin⁵ and Jean-Luc Jaffrezo³

¹ Universités d'Aix-Marseille-CNRS, UMR 6264: Laboratoire Chimie Provence, Equipe Instrumentation et Réactivité Atmosphérique, Marseille, F-13331, France

² Laboratoire de Chimie Moléculaire et Environnement, Université Savoie-Polytech'Savoie, Chambéry, France

³ Université Joseph Fourier-Grenoble 1-CNRS, UMR 5183, Laboratoire de Glaciologie et Géophysique de l'Environnement, Saint Martin d'Hères, F-38402, France

⁴ Université Joseph Fourier-Grenoble 1-CNRS, UMR 5025, Laboratoire de Géodynamique des Chaînes Alpines, BP 53, Grenoble, F-38048, France

⁵ Regional Network for Air Quality Monitoring (ATMO-PACA), 146 rue Paradis 13006 Marseille, France

*Corresponding author:

Université Aix-Marseille I, II, III, CNRS UMR 6264

Laboratoire Chimie Provence (LCP)

Saint Charles Campus (Case 29)

3 Place Victor Hugo, 13331 Marseille

Tel: +33 4 91 10 65 90

Fax: +33 4 91 10 63 77

Email : IMAD.El-Haddad@etu.univ-provence.fr

Table S1: average, min and max concentrations [$\mu\text{g m}^{-3}$] of $\text{PM}_{2.5}$ components measured during the sampling period.

	Average	Min	Max
$\text{PM}_{2.5}$	15.565	6.917	33.167
OC	4.668	2.900	9.600
EC	1.342	0.700	3.400
WSOC	2.164	1.231	4.196
NH_4^+	1.108	0.289	2.197
K^+	0.093	0.042	0.193
Mg^{2+}	0.025	0.012	0.041
Na^+	0.048	<dl	0.175
Cl^-	0.072	0.010	0.263
NO_3^-	0.622	0.300	1.833
SO_4^{2-}	3.924	0.529	9.438

Table S2: average, min and max concentrations [ng m^{-3}] of elements measured during the sampling period.

Elements	Average	Min	Max	Elements	Average	Min	Max
Na	101.31	27.55	207.7	Sb	1.14	0.3	5.61
Mg	19.52	3.2	73.66	Cs	0.04	<dl	0.22
Al	34.72	8.47	115.45	Ba	0.86	0.38	1.58
Ca	62.1	14.26	113.53	La	0.12	0.03	0.43
V	7.18	0.77	22.72	Ce	0.09	0.02	0.27
Cr	5.4	0.70	13.41	Pr	0.005	<dl	0.01
Mn	1.41	0.27	5.14	Nd	0.01	<dl	0.05
Fe	52.77	14.55	131.42	Sm	<dl	<dl	<dl
Co	0.15	0.04	0.5	Eu	<dl	<dl	<dl
Ni	5.08	1.85	13.3	Tb	<dl	<dl	<dl
Cu	3.29	0.5	7.27	Gd	<dl	<dl	<dl
Zn	10.55	0.84	45.71	Dy	<dl	<dl	<dl
Ga	0.02	0.01	0.05	Ho	<dl	<dl	<dl
As	2.17	1.34	2.93	Er	<dl	<dl	<dl
Se	<dl	<dl	<dl	Yb	<dl	<dl	<dl
Rb	0.3	0.07	0.81	Lu	<dl	<dl	<dl
Sr	0.41	0.16	1.58	Hf	<dl	<dl	<dl
Y	0.01	<dl	0.04	Ta	0.01	<dl	0.04
Zr	0.21	0.03	2.42	W	0.13	0.03	0.44
Nb	<dl	<dl	<dl	Tl	0.03	<dl	0.11
Mo	1.63	0.11	9.08	Pb	2.4	0.57	8.85
Cd	0.05	0.01	0.26	Th	<dl	<dl	<dl
Sn	<dl	<dl	<dl	U	0.01	<dl	0.05

Table S3: average, min and max concentrations [ng m⁻³] of organic compounds measured during the sampling period.

Compounds	Average	Min	Max	Compounds	Average	Min	Max
n-nonadecane	0.08	<dl	0.28	n-octacosane	1.19	0.48	2.07
n-eicosane	0.60	0.10	2.00	n-nonacosane	4.44	1.48	10.10
n-heneicosane	0.45	<dl	0.99	n-triacontane	0.90	0.27	1.63
n-docosane	0.80	0.36	1.19	n-hentriacontane	3.79	1.35	7.93
n-tricosane	1.51	0.81	2.47	n-dotriacontane	0.72	0.13	1.32
n-tetracosane	1.51	0.78	2.31	n-tritriacontane	1.58	<dl	4.25
n-pentacosane	2.99	1.72	4.62	n-tetratriacontane	0.61	<dl	3.46
n-hexacosane	1.15	0.51	2.08	n-pentatriacontane	0.24	<dl	0.91
n-heptacosane	2.96	1.08	5.94	n-hexatriacontane	0.06	<dl	0.91
fluoranthene	0.26	0.01	1.12	benzo-e-pyrene	0.18	0.02	0.81
acephenanthrene	0.03	<dl	0.09	benzo-a-pyrene	0.14	0.01	0.85
pyrene	0.20	0.03	0.67	indeno[1,2,3-cd]fluoranthene	0.06	<dl	0.21
benzo[a]anthracene	0.16	0.02	1.19	indeno[1,2,3-cd]pyrene	0.17	0.02	0.84
chrysene/triphenylene	0.29	0.07	1.17	dibenzoanthracene	0.08	<dl	0.55
benzo[b,k]fluoranthene	0.34	0.05	1.69	benzo-ghi-perylene	0.18	0.02	0.66
benzo[j]fluoranthene	0.03	<dl	0.21				
20R+S,5 α (H),14 β (H),17 β (H)-cholestane	0.09	0.04	0.15	20R+S-5 α (H)-14 β (H)-17 β (H)-ergostane	0.10	0.04	0.29
20R,5 α (H),14 α (H),17 α (H)-cholestane	0.10	0.06	0.18	20R+S-5 α (H)-14 β (H)-17 β (H)-stigmastane	0.13	0.06	0.37
trisnorneohopane	0.04	0.01	0.08	17 α (H)-21 β (H)-22R-homohopane	0.09	0.03	0.18
17 α (H)-trisorhopane	0.04	0.01	0.10	17 α (H)-21 β (H)-22S-bishomohopane	0.09	<dl	0.19
17 α (H)-21 β (H)-norhopane	0.23	0.12	0.61	17 α (H)-21 β (H)-22R-bishomohopane	0.06	<dl	0.13
17 α (H)-21 β (H)-hopane	0.20	0.09	0.55	17 α (H)-21 β (H)-22S-trishomohopane	0.03	<dl	0.18
17 α (H)-21 β (H)-22S-homohopane	0.12	0.05	0.26	17 α (H)-21 β (H)-22R-trishomohopane	0.02	<dl	0.11
di-isobutyl phtalate	24.20	6.79	69.45	benzyl butyl phtalate	0.72	0.11	3.85
di-butyl phtalate	12.23	2.80	30.29	bis(2-ethyl hexyl) phtalate	10.85	1.79	25.58
cholesterol	0.41	0.13	3.32	stigmasterol	0.10	<dl	0.18
campesterol	0.09	<dl	0.13	β -sitosterol	0.25	<dl	0.98
palmitic acid	24.88	4.61	66.88	oleic acid	0.78	0.09	6.19
stearic acid	9.12	1.17	19.68	linoleic acid	1.77	0.06	12.70
1,2-phtalic acid	2.66	0.29	8.17	1,4-phtalic acid	0.68	0.09	4.16
1,3-phtalic acid	0.20	0.04	1.08				
D-fructose α	0.26	<dl	1.16	D-mannose α	<dl	<dl	<dl
D-fructose β	0.32	<dl	1.60	D-mannose β	<dl	<dl	<dl
D-galactose α	<dl	<dl	<dl	D-glucose α	2.40	<dl	15.54
D-galactose β	<dl	<dl	<dl	D-glucose β	2.38	<dl	11.98
D-arabitol	0.51	<dl	2.40	D-mannitol	0.45	<dl	2.39
sucrose	0.98	<dl	7.91	trehalose	0.10	<dl	0.57
galactosan	0.25	<dl	1.12	levoglucosan	5.02	0.26	18.70
mannosan	0.38	<dl	1.96				

Table S3: continues.

Compounds	Average	Min	Max	Compounds	Average	Min	Max
vanillin	0.26	<dl	1.28	3-guaiacylpropanol	0.01	<dl	0.04
acetovanillone	0.14	<dl	0.51	vanillic acid	0.16	<dl	0.40
coniferyl aldehyde	0.07	<dl	0.25				
syringaldehyde	0.14	<dl	0.51	3-syringylpropanol	<dl	<dl	<dl
acetosyringone	0.39	<dl	0.39	synapyl aldehyde	0.09	<dl	0.32
propionyl syringol	<dl	<dl	<dl	syringic acid	0.08	<dl	0.30
syringyl acetone	<dl	<dl	<dl	homosyringic acid	<dl	<dl	<dl
abietic acid	<dl	<dl	<dl				
glyceric acid	0.39	<dl	1.73				
2-methylglyceric acid	1.80	0.03	5.90	2-methylerythritol	1.34	0.02	9.20
2-methylthreitol	0.55	<dl	3.50		0.00	0.00	0.00
3-hydroxyglutaric acid	3.50	0.26	14.00	3-isopropylglutaric acid	1.73	0.16	6.15
3-(2-hydroxyethyl)-2,2-dimethylcyclobutane carboxylic acid	0.43	0.13	1.10	3-methyl-1,2,3-butanetricarboxylic	5.01	0.33	17.93
3-hydroxy-4,4-dimethylglutaric acid	0.95	<dl	2.20	pinonic acid	15.18	1.63	34.46
3-acetylglutaric acid	2.40	0.30	7.30	pinic acid	4.50	0.80	11.68
3-acetyl adipic acid	0.98	0.17	2.80				
β -caryophyllenic acid	0.27	<dl	1.10				
3-carboxyheptanedioic acid	0.83	0.16	2.20				
oxalic acid	82.20	35.70	322.00				

Table S4: Set of markers and source profiles injected in the CMB model

Markers	Sources	Profiles
EC, V, Ni, Pb, n-heptacosane, n-octacosane, n-nonacosane, n-triacontane, n-hentriacontane, n-dotriacontane, benzo[b,k]fluoranthene, benzo[e]pyrene, indeno[1,2,3-cd]pyrene, benzo-ghi-perylene, 17 α (H)-21 β (H)-norhopane, 17 α (H)-21 β (H)-hopane, 17 α (H)-21 β (H)-22S-homohopane and levoglucosan	Vehicular emissions Biomass burning Natural gas combustion Vegetative detritus Metallurgical coke production HFO combustion/Shipping Steel manufacturing	El Haddad et al. (2009) Fine et al., (2002) Rogge et al. (1993a) Rogge et al. (1993b) Weitkamp et al. (2005) Agrawal et al. (2008) Tsai et al. (2007)




Figure S1: Average contributions of primary sources to markers included in the CMB model. For the OC, the contributions of different primary sources are normalized to total primary fraction of OC. These results are in total agreement with the preliminary PCA (Figure 2) performed in section 3.2. EC, hopanes and even carbon number n-alkanes are dominated by vehicular emissions. Odd carbon number n-alkanes are dominated by vehicular emissions. PAH and heavy metals (Pb, Ni and V) are dominated by industrial processes.

Table S5: comparison between concentrations [ng m^{-3}] of sugars and sugar derivatives in Marseille and those obtained in other sites.

sugars and sugar alcohols	Potential sources ^c	This study ^d	K-puszta forest summer ^e	western Germany forest summer ^f	Norway urban background fall ^g	Eastern China forests summer ^h	Howland forest, USA ⁱ		
							spring	summer	fall
glucose	lichens, soil biota, pollen, fungi, bacteria, viruses, biomass burning.	4.78 (<0.05-27.5)	ND	16.4	7.20	20.4	28.7	9.75	26.2
fructose	lichens, soil biota, pollen.	0.57 (<0.05-2.71)	ND	10.9	3.30	6.17	4.33	0.37	0.93
arabitol	fungus spores	0.51 (<0.05-2.40)	4.80	15.2	1.00	6.36	2.35	2.65	4.97
mannitol	fungus spores	0.45 (<0.05-2.39)	5.30	13.5	1.60	13.2	3.35	3.65	7.43
sucrose	pollen, soil biota	0.98 (<0.05-7.91)	ND	ND	15.0	ND	2.43	<DL	<DL
trehalose	yeast, bacteria, fungi	0.11 (<0.05-0.57)	ND	ND	5.30	ND	6.88	6.27	14.7
n-alkanes (C23-C34)	anthropogenic and biogenic sources	25.8 (6.20-33.2)	ND	ND	ND	ND	2.27	3.15	2.1
biogenic n-alkanes (C23-C34) ^a	vegetative detritus	12.4 (4.41-27.5)	ND	ND	ND	ND	ND	ND	ND
total OC ^b		4.7 (2.9-9.6)	4.2	4.6	ND	5.5	ND	ND	ND
Modern OC		3.2 (2.1-6.2)	ND	ND	ND	ND	ND	ND	ND

ND: not determined

<DL: < detection limit. Galactose and mannose were not detected.

a Sum of biogenic linear alkanes determined according to Simoneit et al., 1991

b OC [$\mu\text{g.m}^{-3}$] measured by NIOSH method

c references: (Ion et al., 2005; Medeiros et al., 2006; Yttri et al., 2007; Bauer et al., 2008; Kourtchev et al., 2008; Wang et al., 2008)

d mean value (min-max)

e (Ion et al., 2005) ($\text{PM}_{2.5}$ median value)

f (Kourtchev et al., 2008) ($\text{PM}_{2.5}$ mean value)

g (Yttri et al., 2007) ($\text{PM}_{2.5}$ mean value)

h (Wang et al., 2008) ($\text{PM}_{2.5}$ mean value for 4 sites)

i (Medeiros et al., 2006) (Bulk aerosol mean value for each season; for the summer value the biomass burning episodes were not considered in the calculation of the mean).

References

- Agrawal, H., Welch, W. A., Miller, J. W., and Cocker, D. R.: Emission measurements from a crude oil tanker at sea; *Environ. Sci. Technol.*, 42, 7098-7103, 2008.
- Bauer, H.; Claeys, M.; Vermeylen, R.; Schueller, E.; Weinke, G.; Berger, A.; Puxbaum, H.: Arabitol and mannitol as tracers for the quantification of airborne fungal spores; *Atmosph. Environ.*, 3, 588-593, 2008.
- El Haddad, I., Marchand, N., Dron, J., Temime-Roussel, B., Quivet, E., Wortham, H., Jaffrezo, J. L., Baduel, C., Voisin, D., Besombes, J. L., and Gille, G.: Comprehensive primary particulate organic characterization of vehicular exhaust emissions in France; *Atmos. Environ.*, 43, 6190-6198, 2009.
- Fine, P. M., Cass, G. R., and Simoneit, B. R. T.: Chemical Characterization of Fine Particle Emissions from the Fireplace Combustion of Woods Grown in the Southern United States; *Environ. Sci. Technol.*, 36, 1442-1451, 10.1021/es0108988, 2002.
- Ion, A. C.; Vermeylen, R.; Kourtchev, I.; Cafmeyer, J.; Chi, X.; Gelencser, A.; Maenhaut, W.; Claeys, M.: Polar organic compounds in rural PM_{2.5} aerosols from K-puszta, Hungary, during a 2003 summer field campaign: Sources and diel variations; *Atmos. Chem. Phys.*, 1805-1814, 2005.
- Kourtchev, I.; Warnke, J.; Maenhaut, W.; Hoffmann, T.; Claeys, M.: Polar organic marker compounds in PM_{2.5} aerosol from a mixed forest site in western Germany; *Chemosphere*, 8, 1308-1314, 2008.
- Medeiros, P. M.; Conte, M. H.; Weber, J. C.; Simoneit, B. R. T.: Sugars as source indicators of biogenic organic carbon in aerosols collected above the Howland Experimental Forest, Maine; *Atmos. Environ.*, 9, 1694-1705, 2006.
- Rogge, W. F., Hildemann, L. M., Mazurek, M. A., Cass, G. R., and Simoneit, B. R. T.: Sources of Fine Organic Aerosol .4. Particulate Abrasion Products from Leaf Surfaces of Urban Plants; *Environ. Sci. Technol.*, 27, 2700-2711, 1993a.
- Rogge, W. F., Hildemann, L. M., Mazurek, M. A., Cass, G. R., and Simoneit, B. R. T.: Sources of Fine Organic Aerosol .5. Natural-Gas Home Appliances; *Environ. Sci. Technol.*, 27, 2736-2744, 1993b.
- Tsai, J.-H., Lin, K.-H., Chen, C.-Y., Ding, J.-Y., Choa, C.-G., and Chiang, H.-L.: Chemical constituents in particulate emissions from an integrated iron and steel facility; *J. Hazard Mater*, 147, 111-119, 2007.
- Wang, W.; Wu, M. H.; Li, L.; Zhang, T.; Liu, X. D.; Feng, J. L.; Li, H. J.; Wang, Y. J.; Sheng, G. Y.; Claeys, M.; Fu, J. M.: Polar organic tracers in PM_{2.5} aerosols from forests in eastern China; *Atmos. Chem. Phys.*, 24, 7507-7518, 2008.

Weitkamp, E. A., Lipsky, E. M., Pancras, P. J., Ondov, J. M., Polidori, A., Turpin, B. J., and Robinson, A. L.: Fine particle emission profile for a large coke production facility based on highly time-resolved fence line measurements; *Atmos. Environ.*, 39, 6719-6733, 2005.

Yttri, K. E.; Dye, C.; Kiss, G.: Ambient aerosol concentrations of sugars and sugar-alcohols at four different sites in Norway; *Atmos. Chem. Phys.*, 4267-4279, 2007.