

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

**Sensitivity of global
CCN to primary
sulfate emissions**

G. Luo and F. Yu

Sensitivity of global cloud condensation nuclei concentrations to primary sulfate emissions parameterizations

G. Luo and F. Yu

Atmospheric Sciences Research Center, State University of New York, 251 Fuller Road, Albany, New York 12203, USA

Received: 6 November 2010 – Accepted: 10 November 2010 – Published: 12 November 2010

Correspondence to: G. Luo (ganluo@asrc.cestm.albany.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The impact of primary sulfate emissions on cloud condensation nuclei (CCN) concentrations, one of the major uncertainties in global CCN predictions, depends on the fraction of sulfur mass emitted as primary sulfate particles (f_{sulfate}), the fraction of primary sulfate mass distributed into the nucleation mode particles (f_{nucl}), and the nucleation and growth process in the ambient atmosphere. Here, we use a global size-resolved aerosol microphysics model recently developed to study how the different parameterizations of primary sulfate emission affect particle properties and CCN abundance. Different from previous studies, we use the ion-mediated nucleation scheme to calculate tropospheric particle formation and explicitly simulate the condensation of low volatile secondary organic gas (in addition to H_2SO_4 gas) on nucleated particles. Our simulations show a compensation effect of nucleation to primary sulfate emission. We find that the change of f_{nucl} from 5% to 15% has a more significant impact on the simulated particle number budget than that of f_{sulfate} within the range of 2.5–5%. An increase of f_{sulfate} from 0% to 2.5% (with $f_{\text{nucl}} = 5\%$) does not improve the agreement between simulated and observed annual mean number concentrations of particles >10 nm at 21 stations but further increase of either f_{sulfate} from 2.5% to 5% (with $f_{\text{nucl}} = 5\%$) or f_{nucl} from 5% to 15% (with $f_{\text{sulfate}} = 2.5\%$) substantially deteriorates the agreement. For f_{sulfate} of 2.5%–5% and f_{nucl} of 5%, our simulations indicate that the global CCN at supersaturation of 0.2 increases by 8–11% in the boundary layer and 3–5% in the whole troposphere (compared to the case with $f_{\text{sulfate}} = 0$). Our derived impact of primary sulfate emission on global CCN abundance is about a factor of ~ 3 –7 smaller than those found in previous studies.

1 Introduction

Atmospheric aerosols significantly impact climate systems via direct and indirect radiative effects (IPCC, 2007). Atmospheric particles can affect climate indirectly by acting

ACPD

10, 27695–27723, 2010

Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



as cloud condensation nuclei (CCN), modifying cloud properties, altering precipitation, and changing global radiation and heat budgets (Twomey, 1977; Albrecht, 1989). The aerosol indirect effect is largely controlled by the CCN abundance which depends on particle size distributions and compositions. One of the uncertainties in global CCN simulations is associated with the parameterization of sub-grid sulfur oxidation and sulfate particle formation in anthropogenic SO₂ plumes which has been named “primary sulfate” emissions in the literature.

Uncertainties associated with primary sulfate emissions mainly arise from two parts: (1) fraction of sulfur mass emitted as sulfate particles (f_{sulfate}); and (2) parameterization of primary sulfate particle number size distribution. f_{sulfate} is employed to account for the sulfur oxidation and new particle formation occurring in the sub-grid scale anthropogenic SO₂ plumes. f_{sulfate} values ranging from 0% to 5% have been assumed in various global modeling studies (Adams and Seinfeld, 2002; Adams and Seinfeld, 2003; Spracklen et al., 2005a, b; Pierce and Adams, 2006; Spracklen et al., 2008; Makkonen et al., 2009; Pierce and Adams, 2009; Wang and Penner, 2009; Yu and Luo, 2009). The exact value of f_{sulfate} at each grid cell is expected to depend on OH concentration (e.g., Wilson, 1981) and model resolution. Primary sulfate emitted in the power plant plume is generally distributed into two lognormal modes with geometric number-based mean diameters (d_g) of 10 nm (nucleation mode) and 70 nm (accumulation or condensation mode) with standard deviations (σ) of 1.6 and 2.0, respectively (Whitby, 1978; Whitby et al., 1978). All the above mentioned global aerosol studies except Yu and Luo (2009) assume that 15% of primary sulfate mass is emitted in the nucleation mode while the rest in the accumulation mode. However, as pointed in Yu and Luo (2009), the limited observations presented in Whitby et al. (1978) suggested that the fraction of primary sulfate in the nucleation mode (f_{nucl}) is 5% (called Whitby78 value thereafter) while the remaining (95%) of sulfate is in the accumulation mode (i.e., $f_{\text{accu}} = 95\%$). The impacts of the two different assumptions of primary sulfate particle number size distribution (or f_{nucl} values) on simulated global aerosol properties have not yet been assessed.

Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Sensitivity of global
CCN to primary
sulfate emissions**

G. Luo and F. Yu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



For any given f_{sulfate} and f_{nucl} values, the contribution of primary sulfate emission to the global CCN abundance depends strongly on the particle nucleation schemes used in the model to predict particle nucleation rates in the ambient troposphere. Previous global aerosol simulations, with new particle formation calculated based on binary homogeneous nucleation (BHN), showed that a few percent of anthropogenic sulfur emitted as primary sulfate can increase CCN number concentrations in polluted areas by up to ~ 200 – 500% (Adams and Seinfeld, 2003; Spracklen et al., 2005b). Pierce and Adams (2009) reported that global mean CCN number concentrations increased 17% in the whole troposphere and 40% in the boundary layer, when primary aerosol emissions (sulfate, BC, OC and so on) were increased by a factor of 3. These changes drop to 3% and 22%, respectively, when the nucleation rates were calculated based on the ternary homogeneous nucleation (THN) model. Wang and Penner (2009) estimated the impacts of primary sulfate emission on CCN abundance using two different nucleation schemes: one based on BHN for the whole troposphere and the other using the empirical nucleation mechanism in the boundary layer and the BHN mechanism above boundary layer (BHN_EMP). Wang and Penner (2009) showed that, when f_{sulfate} increases from 0% to 2%, CCN concentration in the lower boundary layer increases by 53% based on BHN while it increases by 23% based on BHN_EMP parameterization. It is clear from these previous investigations that the impact of primary sulfate emission on the global CCN abundance based on different nucleation schemes can differ by a factor of up to ~ 6 .

It is well known that uncertainty in CCN concentration could lead to large uncertainty in aerosol indirect radiative forcing. Based on the relationship between cloud albedo (A) and CCN concentration (N) given in Platnick and Twomey (1994) ($\Delta A/A = (1-A)/3 \times \Delta N/N$), a 10% of increase in CCN concentrations can lead to $\sim 2\%$ increase in average cloud albedo (assuming global average A of 0.42, Han et al., 2001). Since clouds on average reflect about 50 W m^{-2} of incoming solar radiation back to space (Hartmann, 1993), a 2% increase in the average cloud albedo could lead to a radiative forcing of $\sim 1 \text{ W m}^{-2}$. While these estimations

are crude, they do indicate the importance in reducing the uncertainty in global CCN predictions.

The main objective of this study is to carry out a comprehensive investigation of how primary sulfate emission may influence CCN abundance in the global atmosphere, using a global size-resolved (sectional) aerosol microphysics model (Yu and Luo, 2009). Different from previous assessments on the impact of primary sulfate emission parameterization, we use the ion-mediated nucleation (IMN) scheme (Yu, 2010a) to calculate new particle formation in the troposphere and explicitly consider the condensation of low volatile secondary organic gas (SOG) on nucleated particles based on the recently developed extended secondary organic aerosol formation mechanism (Yu, 2010b). The effects of f_{nucl} values on simulated CCN concentrations, which have not been investigated in previous studies, are also investigated in this study. The physics behind the impacts of primary sulfate emission is analyzed in detail by comparing condensation sinks, nucleation rates, and particle size distributions simulated with different primary sulfate emission assumptions.

2 Model description and overview of simulations

The GEOS-Chem 3-D chemical transport model, with an advanced particle microphysics (APM) model incorporated (Yu and Luo, 2009), is employed for this study. Large-scale atmospheric transport is specified from the assimilated meteorological observations from the Goddard Earth Observing System (GEOS) of the NASA Global Modeling Assimilation Office (GMAO). The GEOS-Chem model includes a detailed simulation of tropospheric ozone-NO_x-hydrocarbon chemistry as well as of aerosols and their precursors (Park et al., 2004). Aerosol and gas-phase simulations are coupled through sulfate and nitrate formation, heterogeneous chemistry (Evans and Jacob, 2005), aerosol effects on photolysis rates (Martin et al., 2003), and secondary organic aerosol (SOA) formation (Liao et al., 2007). The ISORROPIA II thermodynamic equilibrium model (Fountoukis and Nenes, 2007) is used to calculate

Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



partitioning of total ammonia and nitric acid between the gas and aerosol phases. A detailed description of the GEOS-Chem model (including the treatment of various emission sources, chemistry and aerosol schemes) can be found in the model web-page (<http://acmg.seas.harvard.edu/geos/index.html>).

5 The present version of the APM module in GEOS-Chem employs 40 bins for sulfate to cover the dry diameter size range of 0.0012 μm to 12 μm , and 20 bins for sea salt to cover the dry diameter size range of 0.012 μm to 12 μm (Yu and Luo, 2009). The first bin of sulfate particles corresponds to the sizes of freshly nucleated particles in the atmosphere which generally have a dry diameter of $\sim 0.0012 \mu\text{m}$ (Yu and Turco, 2008),
10 therefore it does not need to scale the nucleated particles to the first bin particles. The sulfate aerosols are internally mixed with nitrate, ammonium and organics in our model, and the contributions of nitrate, ammonium, and SOAs to sulfate particle growth are considered through equilibrium uptake (Yu and Luo, 2009). An extended secondary organic aerosol formation mechanism developed by Yu (2010b) has been employed in
15 this study to transfer medium-volatile SOG to semi-volatile SOG and semi-volatile SOG to low-volatile SOG associated with oxidation aging (e.g., Jimenez et al., 2009). This treatment significantly increases the growth rates of nucleated particles associated with the condensation of low-volatile SOG (in addition to H_2SO_4), which is consistent with many field measurements and important for properly accounting for the contribution of
20 nucleated particles to CCN (Yu, 2010b). When we calculate the CCN activation, we use the dry size (sulfate + uptaken equilibrium nitrate, ammonium, and SOA) to reflect the contribution of multiple components (Yu and Luo, 2009). The model also considers the scavenging of secondary particles by primary particles such as BC, OC, dust, and sea salt. In the current model, the condensation of sulfuric acid, in-cloud SO_2 oxidation and the coagulation scavenging of secondary sulfate particles by primary particles are simulated by using four separate tracers (BC_sulfate, OC_sulfate, Dust_sulfate, and SeaSalt_sulfate) to keep track of the bulk sulfate mass associated with BC, OC, dust, and sea salt, respectively. When some of BC, OC, dust, and sea salt are removed
25 from the atmosphere via dry and wet deposition, the corresponding portions of sulfate

Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

associated with these particles are also removed. A more detailed description of the APM module, which will be added to the standard version of GEOS-Chem in the near future after further improvements, can be found in Yu and Luo (2009).

In the simulations presented here, we use a horizontal resolution of $4^\circ \times 5^\circ$ and 47 vertical layers up to 0.01 hpa (GEOS-5 meteorological fields). The global sulfur emission is based on the Emissions Database for Global Atmospheric Research (EDGAR) inventory v3.2 (Olivier et al., 2001), while the emissions in the United States, Canada, Mexico, Europe and East Asia are replaced by the regional emission inventories including the Environmental Protection Agency's National Emission Inventory 2005, the Criteria Air Contaminants (CAC) Emissions Inventory, the Big Bend Regional Aerosol and Visibility Observational (BRAVO) Study Emissions Inventory (Kuhns et al., 2003), the European Monitoring and Evaluation Programme (EMEP) Emissions Inventory, and the Transport and Chemical Evolution over the Pacific (TRACE-P) Emissions Inventory (Streets et al., 2003). The model also includes other primary particle emissions such as black carbon (BC), primary organic carbon (POC), dust, and sea salt. A detailed description of other key gaseous species and parameterizations of various primary particle emissions can be found in Yu and Luo (2009).

Four simulations, which are summarized in Table 1, have been run to quantify the impacts of primary sulfate emissions on nucleation, particle number size distribution, and CCN number concentration. In our study, following the parameterizations given in Whitby (1978) and Whitby et al. (1978), we assume that f_{nucl} of primary sulfate mass is emitted in the nucleation mode ($d_g = 10$ nm, $\sigma = 1.6$) while the rest in the accumulation mode ($d_g = 10$ nm, $\sigma = 1.6$) which has been treated as condensation on BC and primary OC particles (Yu and Luo, 2009). The FS0 simulation assumes $f_{\text{sulfate}} = 0$ (i.e., no primary sulfate emission). The FS2.5FN5 simulation, which assumes $f_{\text{sulfate}} = 2.5\%$ with $f_{\text{nucl}} = 5\%$ and $f_{\text{accu}} = 95\%$, is used as the baseline simulation. The FS2.5FN15 simulation corresponds to the case for $f_{\text{sulfate}} = 2.5\%$ with $f_{\text{nucl}} = 15\%$ and $f_{\text{accu}} = 85\%$. The comparison of the FS2.5FN5 and FS2.5FN15 simulations allows us to see how the f_{nucl} value may affect the simulations of CCN number concentrations. The FS5FN5

Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

simulation, which is the same as the FS2.5FN5 simulation except that f_{sulfate} is increased from 2.5% to 5%, is designed to provide further insight into how f_{sulfate} value may impact the simulated particle properties and CCN number concentrations. The net effects of f_{sulfate} and f_{nucl} on the primary sulfate emission can be easily interpreted by comparing FS2.5FN5 to FS0, FS2.5FN15 and FS5FN5, respectively. The spin-up time of our simulations is two months (November–December 2004). All annually averaged results shown in this paper are for year 2005.

3 Results and discussion

3.1 Impact of primary sulfate emissions on nucleation rates

In addition to directly injected particles into the grid boxes, primary sulfate emissions indirectly impact the particle number budget via influencing sulfur dioxide concentration (noting that $1-f_{\text{sulfate}}$ of anthropogenic sulfur emitted as SO_2), condensation sink (CS), sulfuric acid gas concentration ($[\text{H}_2\text{SO}_4]$), and hence new particle formation rates. Figure 1 shows how different primary sulfate emission assumptions affect CS in the atmosphere. With the exclusion of primary sulfate emission, CS near source regions is decreased by up to ~ 10 – 20% (Fig. 1b), while the zonally averaged relative changes at the Northern Hemisphere mid-latitude regions (30°N – 60°N) is decreased by about 5 – 10% in the low boundary layer (Fig. 1f). In northwestern Canada, CS is increased by ~ 0 – 5% due to the increased concentration of freshly nucleated particles. It can be clearly seen from Fig. 1c and g (the FS2.5FN15 case) that CS is very sensitive to the f_{nucl} values. The increased primary sulfate mass in the nucleation mode significantly increases the CS and thus decreases $[\text{H}_2\text{SO}_4]$ (not shown). Figure 1d indicate that CS is increased by 5 – 10% over source regions when f_{sulfate} is increased from 2.5% to 5%. It is clear that the impact of f_{sulfate} increased from 2.5% to 5% on CS is much smaller than that of f_{nucl} changed from 5% to 15%.

Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Compared to the baseline case (FS2.5FN5), nucleation rate is increased in the FS0 case and decreased in the FS2.5FN15 and FS5FN5 cases (Fig. 2) as a response to the changes of CS and $[\text{H}_2\text{SO}_4]$. Figure 2a shows the high values of low boundary layer nucleation rate are mainly located over the Northern Hemisphere mid-latitude regions (30°N – 60°N). Nucleation is sensitive to the changes of $[\text{H}_2\text{SO}_4]$. The decrease of $[\text{H}_2\text{SO}_4]$, caused by the enhanced CS associated with the increased primary sulfate particles and reduced fraction of anthropogenic sulfur emitted as SO_2 gas, decreases the intensity of nucleation in the atmosphere. In and near the anthropogenic sulfur source regions, nucleation rates increase by up to ~ 10 – 30% when f_{sulfate} changes from 2.5% to 0% (Fig. 2b and f), and decrease by ~ 10 – 30% when f_{nucl} changes from 5% to 15% (at fixed $f_{\text{sulfate}} = 2.5\%$, Fig. 2c and g). The FS5FN5 case has less impact on nucleation rates compared to the FS2.5FN15 case. It should be noted that the lower nucleation rates over tropical lower troposphere (annual mean $J < \sim 0.001\text{ cm}^3\text{ s}^{-1}$, Fig. 2a and e) lead to high percentage changes in J values in the regions but such changes are insignificant to particle number budget there and thus have been blocked out in Fig. 2b–d and f–h to highlight the changes in major source regions.

The results given in Fig. 2 clearly show a compensation effect of nucleation to primary sulfate emission. Reduced primary sulfate emission (lower f_{sulfate} and/or f_{nucl}) reduces the surface area of primary particles directly injected into the grid boxes resulting in an increase in new particle formation rates near the source regions. The increased nucleation rates compensate for some of the primary sulfate number emission reductions. Nevertheless, as we will show in the next several sections, the increased concentrations of freshly nucleated particles are less than the decreased concentrations of primary sulfate particles over the source regions. In other words, enhanced nucleation cannot fully replace the role of sub-grid scale sulfur oxidant and nucleation processes within the plume. Therefore, proper parameterizations of these sub-grid processes are still needed.

Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

3.2 Impact of primary sulfate emissions on particle number size distributions

To analyze how different primary sulfate emission assumptions and associated nucleation rate changes may impact the global aerosol properties, we plot in Fig. 3 the simulated annual mean particle number size distributions in the boundary over land (b) and over ocean (c) as well as those averaged in the global boundary layer (Fig. 3a). It can be clearly seen that the impacts of primary sulfate emissions on aerosol number size distributions are substantial in the boundary layer over land (Fig. 3b) but are small over the ocean (Fig. 3c). In addition to the direct influence on the concentrations of primary sulfate particles emitted into certain grid boxes, primary sulfate parameterizations also affect the total particle size distributions by influencing condensation sink, H₂SO₄ vapor concentration, and particle nucleation and growth rate. Figure 3a shows that boundary layer aerosol number size distribution has three modes. The first mode appears at the dry diameter of ~2.5 nm. Particles in this mode are formed via nucleation of sulfuric acid gas and water vapor. We call this the freshly nucleated mode. The second and third modes appear at the dry diameters of ~20 nm and ~120 nm, which are known as Aitken and accumulation modes, respectively. The FS0 simulation (black line) indicates that freshly nucleated particles can grow to the size of Aitken mode particles. The number concentrations of freshly nucleated mode particles decrease while those of Aitken mode particles and accumulation mode particles increase with the increase of f_{sulfate} and f_{nuc1} . Based on the FS2.5FN5 (red line), FS2.5FN15 (orange line) and FS5FN5 (blue line) simulations, we find that the change of f_{nuc1} from 5% to 15% has a more significant impact on boundary layer aerosol number size distributions than that of f_{sulfate} within the range of 2.5–5%. As shown in Fig. 3b and c, the impacts of primary sulfate emissions on boundary layer aerosol number size distribution over land and ocean are quite different. The difference in the boundary layer aerosol number size distribution among all four cases is negligible over the oceans but is substantial over land. In the boundary layer over land, the number concentrations of freshly nucleated mode, nucleation mode and accumulation mode particles in the FS2.5FN15 simulation

Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

are ~ 0.8 , ~ 1.9 and ~ 1.2 of those in the FS0 simulation (Fig. 3b). The changes of primary sulfate emissions have negligible impact on aerosol number size distribution averaged over the whole troposphere (not shown).

3.3 Impact of primary sulfate emissions on number concentrations of particles larger than 10 nm (CN10) and comparisons with measurements

Figure 4 shows the horizontal and vertical spatial distributions of annual mean CN10 (Fig. 4a, e) and its responses to the changes of primary sulfate emissions (Fig. 4b–d, f–h). Direct injection of nucleation mode particles with median diameter of 10 nm into the grid boxes, which is used to account for sub-grid scale nucleation, has a significant impact on the CN10 budget. As expected, the spatial patterns in the changes of CN10 associated with different primary sulfate emission assumptions are similar to that of CS (Fig. 1). Compared to the baseline case (FS2.5FN5), CN10 around the source regions decreases by ~ 10 – 30% when the primary sulfate emission is excluded (i.e., case FS0) (Figs. 4b and f) and increases by ~ 10 – 30% when f_{sulfate} is increased from 2.5% to 5% (Fig. 4d and h). Again, the increase of f_{nuc} value by a factor of 3 (from 5% to 15%) has significant impact on CN10 near source regions (20–30% change with peak value reaching $\sim 50\%$). The exclusion of primary sulfate emission reduces the number concentration of primary emitted sulfate particles, increases the nucleation rate, and then increases the number concentration of freshly nucleated particles. These new particles grow to larger size and have significant contribution to the particle number concentration over clean remote regions. Near the anthropogenic sulfur source regions, the CN10 increased by nucleation is less than that of particles decreased by the exclusion of primary sulfate emission. As a result, CN10 is increased over remote regions and decreased over polluted regions when primary sulfate emission is excluded. The FS2.5FN15 and FS5FN5 simulations indicate that both increases of f_{sulfate} and f_{nuc} can significantly increase the CN10 over source regions.

To assess the importance of proper primary particle emission parameterizations, we compare the predicted CN10 values with those observed at a number of surface-based

Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



stations which have long term continuous measurements. Figure 5 shows the CN10 values observed at the 21 boundary layer sites listed in Table 2 and the corresponding simulated values for the FS0, FS2.5FN5, FS2.5FN15 and FS5FN5 cases. As expected, primary sulfate emissions have significant impact on the CN10 values at the polluted sites but have relatively small influence on CN10 at remote sites. The statistic comparisons of the observations and the four sensitivity study cases at the 21 sites are also presented in Table 2. The FS2.5FN5 simulation is the baseline case. The coefficient of determination (R^2) is simply the square of the sample correlation coefficient between the observations and their predicted values. The statistic comparisons indicate that the coefficient of determination is within 0.1 for all the four studied cases but the difference in the mean absolute error and standard deviation of error is much bigger. While the mean absolute error and standard deviation of error of the FS0 case are very close to the values of FS2.5FN5 case, those for the FS2.5FN15 and FS5FN5 simulations are much larger. For example, the mean absolute error and standard deviation of error of the FS2.5FN15 case are 1285.9 and 1856.9, respectively, which are about two times higher than those of the FS2.5FN5 case. As shown in Fig. 5b, the FS2.5FN15 and FS5FN5 cases over predict the CN10 at most polluted sites. The impacts of primary emission of sulfate on aerosol nucleation, particle number concentration and CCN number concentration mainly depend on the total number and surface area of primary sulfate emitted in nucleation model which is associated both with f_{sulfate} and f_{nucl} . The total number and surface area of primary sulfate emitted in nucleation model when f_{nucl} is changed from 5% to 15% are a factor of $3/2 = 1.5$ larger than those when f_{sulfate} is changed from 2.5% to 5%. A comparison of (FS2.5FN15-FS2.5FN5)/FS2.5FN5 (Fig. 4c) with (FS5FN5-FS2.5FN5)/FS2.5FN5 (Fig. 4d) shows that the changes of CN10 at emission source regions in the FS2.5FN15 case are generally more than 2–3 times larger than those in the FS5FN5 case. This extra enhancement (above the factor of 1.5) is caused by the nonlinear contributions associated with aerosol nucleation and other microphysical processes. Our simulations indicate that the increase of the f_{nucl} from Whitby78 value of 5% to widely used value of 15% has

Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

more distinct impact on CN10 than that of the increase of f_{sulfate} from 2.5% to 5%. It is clear from Table 2 that, statistically, the increase of f_{sulfate} from 0% to 2.5% (with $f_{\text{nucl}} = 5\%$) does not improve the agreement of predicted CN10 at the 21 sites with those of observations but further increase of either f_{sulfate} from 2.5% to 5% (with $f_{\text{nucl}} = 5\%$) or f_{nucl} from 5% to 15% (with $f_{\text{sulfate}} = 2.5\%$) substantially deteriorates the agreement.

3.4 Impact of primary sulfate emissions on CCN number concentrations

The horizontal and vertical spatial distributions of CCN number concentration at 0.2% supersaturation (CCN0.2) and the changes of CCN0.2 associated with primary sulfate emissions are shown in Fig. 6. Our simulations show that CCN0.2 in the boundary layer decreases by $\sim 6\text{--}18\%$ over Europe, South Asia and East Asia when primary sulfate emission is excluded (Fig. 6b), while the maximum decrease of the zonally-averaged CCN0.2 appears at $\sim 30^\circ \text{N}$ in the boundary layer (Fig. 6f). It is interesting to note that the CCN0.2 over several small regions is increased by $\sim 0\text{--}3\%$ with the exclusion of primary sulfate emission, in large contrast with other locations within the boundary layer. Our analysis indicates that this is a result of complex interactions or competitions among nucleation, growth, transport, and scavenging. To assess the impact of f_{nucl} , we compare the CCN0.2 simulated by the FS2.5FN15 simulation with that simulated by the FS2.5FN5 simulation. As shown in Fig. 6c, the increase of f_{nucl} from 5% to 15% increases CCN0.2 by $\sim 6\text{--}18\%$ over source regions and $\sim 0\text{--}3\%$ over remote regions. The FS5FN5 simulation shows that the increase of f_{sulfate} from 2.5% to 5% generally increases CCN0.2 (up to 3–12% over source regions over Asia and Sothern America). The impact of f_{sulfate} increased from 2.5% to 5% on CCN0.2 (Fig. 6d, h) is much less than that of f_{nucl} increased from 5% to 15% (Fig. 6c, g), especially over high latitude regions in the Northern Hemisphere.

Table 3 shows that the change of f_{sulfate} from 2.5% to 5% has relatively small impact on the change of global CCN0.2 ($\sim 3\%$). Compared to the FS0 case, FS2.5FN5 and FS5FN5 simulations indicate only 8–11% increase of CCN0.2 in the boundary layer and 3–5% increase in the whole troposphere. The changes of f_{nucl} from 5% to 15%

Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



can double the increase in the changes of the global CCN0.2 both in the boundary layer and in the whole troposphere. The strongest impact of primary sulfate emissions on global CCN0.2 appears in the FS2.5FN15 case, with the mean change of global CCN0.2 as high as 16% in the boundary layer and 8% in the whole troposphere. It is clear that f_{nucl} value presents more significant impact on global CCN0.2 than that of f_{sulfate} , highlighting the importance in reducing the uncertainty associated with f_{nucl} in order to reduce the uncertainties in the predicted global CCN0.2 concentrations.

Compared to the results from Wang and Penner (2009) ($f_{\text{sulfate}} = 2\%$ with $f_{\text{nucl}} = 15\%$), the change of the CCN0.2 in the boundary layer of our baseline simulation (FS2.5FN5) is a factor of 3–7 smaller than those of previous works. The differences mainly arise from the different f_{nucl} assumed and the nucleation scheme employed, although other factors (such as parameterizations of other primary particles, particle growth and scavenging processes, etc.) could also contribute to the differences. When we use the widely used value of $f_{\text{nucl}} = 15\%$ instead of the Whitby78 value of 5%, the change of global CCN0.2 in boundary layer is increased from 8% to 16%, but still lower than the changes predicted by Wang and Penner (2009). The differences among these simulations are associated with the different treatments of aerosol microphysics, especially the nucleation process.

4 Summary and discussions

The parameterization of sub-grid sulfur oxidation and sulfate particle formation in anthropogenic SO₂ plumes, which has been named “primary sulfate” emissions in the literature, represents one of the major uncertainties in global cloud condensation nuclei (CCN) predictions. Both the fraction of sulfur mass emitted as primary sulfate particles (f_{sulfate}) and the fraction of primary sulfate mass distributed into the nucleation mode particles (f_{nucl}) have large uncertainties. Here, we use a global size-resolved aerosol microphysics model to explore how the different parameterizations of primary sulfate particle emission affect nucleation, particle size distributions, and CCN number

Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



concentrations in the global troposphere. Different from previous studies, we use the ion-mediated nucleation scheme to calculate new particle formation in the troposphere and explicitly consider the condensation of low volatile secondary organic gas on nucleated particles. The effects of f_{nucl} values on simulated CCN concentrations, which have not been investigated in previous studies, are investigated in the present study.

Our simulations clearly show a compensation effect of nucleation to primary sulfate emission. The inclusion of primary sulfate emission increases the surface area of pre-existing particles, reduces $[\text{H}_2\text{SO}_4]$, and thus decreases nucleation rates near the source regions. When primary sulfate emission is reduced, the number concentrations of sulfate particles directly injected into the grid boxes decrease yet new particle formation rates near the source regions increase. This increase of nucleation rates compensates for some of the primary sulfate number emission reductions. However, enhanced nucleation cannot fully replace the role of sub-grid plume scale sulfur oxidant and nucleation processes, highlighting the necessity for proper parameterizations of these sub-grid processes in global aerosol models.

Our sensitivity studies indicate that the change of f_{nucl} from the Whitby78 value of 5% to the widely used value of 15% has a more significant impact on nucleation rates, particle size distributions, and CCN concentrations than that of f_{sulfate} within the range of 2.5–5%. The predicted annual mean CN10 values based on different primary particle emission parameterizations have been compared to those observed at 21 surface-based stations. Our statistical analysis shows that the increase of f_{sulfate} from 0% to 2.5% (with $f_{\text{nucl}} = 5\%$) does not improve the agreement between predicted and observed CN10 at the 21 sites but further increase of either f_{sulfate} from 2.5% to 5% (with $f_{\text{nucl}} = 5\%$) or f_{nucl} from 5% to 15% (with $f_{\text{sulfate}} = 2.5\%$) substantially deteriorates the agreement.

When $f_{\text{sulfate}} = 2.5\%–5\%$ and $f_{\text{nucl}} = 5\%$ are assumed, our simulations indicate that the global CCN0.2 increases by 8–11% in the boundary layer and 3–5% in the whole troposphere (compared to the case without primary sulfate emission, i.e., $f_{\text{sulfate}} = 0$). Our predicted global CCN0.2 changes associated with primary sulfate emission are

Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a factor of 3–7 smaller than other recent works (Pierce and Adams, 2009; Wang and Penner, 2009). The differences mainly arise from the different f_{nucl} values assumed and the nucleation schemes employed. When we use the widely used value of $f_{\text{nucl}} = 15\%$ instead of the Whitby78 value of 5%, our predicted change of global CCN0.2 doubles but is still lower than the values found in previous studies.

The impact of primary sulfate emission on global CCN abundance derived in the present investigation, while much smaller than those from previous studies, is still significant especially in the boundary layer near source regions. Due to high sensitivity of aerosol indirect radiative forcing to CCN concentrations, it is necessary to further improve the representation of sulfur oxidation and sulfate particle formation in sub-grid scale SO_2 plumes in regional and global models. All the existing global studies (including the present study) assume that both f_{sulfate} and f_{nucl} have no diurnal, seasonal, and spatial variations. However, both f_{sulfate} and f_{nucl} in the real atmosphere depend on OH concentrations and other parameters which have significant diurnal, seasonal, and spatial variations (e.g., Wilson, 1981). In order to reduce the uncertainties in the predicted global CCN concentrations and aerosol radiative forcing, further research (including observation) is needed to determine the dependence of f_{sulfate} and f_{nucl} on key parameters and properly represent such a dependency within climate models. It should be noted that there may exist large uncertainty in the f_{nucl} value derived by Whitby et al. (1978) because of instrument limitation. More measurements of particle size distributions and compositions in fresh power plant plumes using the modern instruments are needed to reduce the uncertainty in primary sulfate emission parameterizations. In addition to the primary sulfate emissions, many other processes (such as primary emissions of other species, particle nucleation and growth rates, particle scavenging rates, etc.) will also affect simulated global CCN concentrations and the uncertainties in these processes may affect the conclusion of this study. It is an ongoing task to improve the representation of these processes in global aerosol models.

Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Acknowledgements. This study is supported by NASA under grant NNX08AK48G and NSF under grant 0942106. The GEOS-Chem model is managed by the Atmospheric Chemistry Modeling Group at Harvard University with support from the NASA Atmospheric Chemistry Modeling and Analysis Program.

References

- Adams, P. J. and J. H. Seinfeld, Predicting global aerosol size distributions in general circulation models, *J. Geophys. Res.*, 107(D19), 4370, doi:10.1029/2001JD001010, 2002.
- Adams, P. J., and J. H. Seinfeld, Disproportionate impact of particulate emissions on global cloud condensation nuclei concentrations, *Geophys. Res. Lett.*, 30(5), 1239, doi:10.1029/2002GL016303, 2003.
- Albrecht, B. A.: Aerosols, Cloud Microphysics, and Fractional Cloudiness, *Science*, 245(4923), 1227–1230, 1989.
- Dal Maso, M., Hyvarinen, A., Komppula, M., et al.: Annual and interannual variation in boreal forest aerosol particle number and volume concentration and their connection to particle formation, *Tellus*, 60B, 495–508, 2008.
- Evans, M. J. and Jacob, D. J.: Impact of new laboratory studies of N₂O₅ hydrolysis on global model budgets of tropospheric nitrogen oxides, ozone, and OH, *Geophys. Res. Lett.*, 32, L09813, doi:10.1029/2005GL022469, 2005.
- Fountoukis, C. and Nenes, A.: ISORROPIA II: a computationally efficient thermodynamic equilibrium model for $K^+ - Ca^{2+} - Mg^{2+} - NH_4^+ - Na^+ - SO_4^{2-} - NO_3^- - Cl^- - H_2O$ aerosols, *Atmos. Chem. Phys.*, 7, 4639–4659, doi:10.5194/acp-7-4639-2007, 2007.
- Han, Q., Rossow, W. B., Chou, J., and Welch, R. M.: Global survey of the relationships of cloud albedo and liquid water path with droplet size using ISCCP, *J. Climate*, 11, 1516–1528, 1998.
- Hartmann, D. L.: Radiative effects of clouds on Earth's climate, in *Aerosol-Cloud-Climate Interactions*, edited by: Bobbs, P. V., Academic Press, 1993.
- IPCC, *Climate Change 2007: The Physical Scientific Basis*, Rep., 996 pp., Cambridge Univ. Press, New York, USA, 2007.
- Jacobson, M. Z., Turco, R. P., Jensen, E. J., et al.: Modeling coagulation among particles of different composition and size, *Atmos. Environ.*, 28(7), 1327–1338, 1994.

Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Sensitivity of global
CCN to primary
sulfate emissions**

G. Luo and F. Yu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., et al.: Evolution of Organic Aerosols in the Atmosphere, *Science*, 326, 1525–1529, 2009.
- Kivekäs, N., Sun, J., Zhan, M., Kerminen, V.-M., Hyvärinen, A., Komppula, M., Viisanen, Y., Hong, N., Zhang, Y., Kulmala, M., Zhang, X.-C., Deli-Geer, and Lihavainen, H.: Long term particle size distribution measurements at Mount Waliguan, a high-altitude site in inland China, *Atmos. Chem. Phys.*, 9, 5461–5474, doi:10.5194/acp-9-5461-2009, 2009.
- Komppula, M., Lihavainen, H., Lihavainen, A.-P. Hyvärinen, V.-M., et al.: Physical properties of aerosol particles at a Himalayan background site in India, *J. Geophys. Res.*, 114, D12202, doi:10.1029/2008jd011007, 2009.
- Kuhns, H., Green, M., and Etyemezian, V.: Big Bend Regional Aerosol and Visibility Observational (BRAVO) Study Emissions Inventory Rep., Desert Research Institute, Las Vegas, 2003.
- Laakso, L., Laakso, H., Aalto, P. P., Keronen, P., Petäjä, T., Nieminen, T., Pohja, T., Siivola, E., Kulmala, M., Kgabi, N., Molefe, M., Mabaso, D., Phalatse, D., Pienaar, K., and Kerminen, V.-M.: Basic characteristics of atmospheric particles, trace gases and meteorology in a relatively clean Southern African Savannah environment, *Atmos. Chem. Phys.*, 8, 4823–4839, doi:10.5194/acp-8-4823-2008, 2008.
- Liao, H., Henze, D. K., Seinfeld, J. H., et al.: Biogenic secondary organic aerosol over the United States: comparison of climatological simulations with observations, *J. Geophys. Res.*, 112, D06201, doi:10.1029/2006jd007813, 2007.
- Makkonen, R., Asmi, A., Korhonen, H., Kokkola, H., Järvenoja, S., Räisänen, P., Lehtinen, K. E. J., Laaksonen, A., Kerminen, V.-M., Jrvinen, H., Lohmann, U., Bennartz, R., Feichter, J., and Kulmala, M.: Sensitivity of aerosol concentrations and cloud properties to nucleation and secondary organic distribution in ECHAM5-HAM global circulation model, *Atmos. Chem. Phys.*, 9, 1747–1766, doi:10.5194/acp-9-1747-2009, 2009.
- Martin, R. V., Jacob, D. J., Yantosca, R. M., et al.: Global and regional decreases in tropospheric oxidants from photochemical effects of aerosols, *J. Geophys. Res.*, 108, 4097, doi:10.1029/2002jd002622, 2003.
- Olivier, J. G. J., Berdowski, J. J. M., Peters, J. A. H. W., et al.: Applications of EDGAR. Including a description of EDGAR 3.0: reference database with trend data for 1970–1995, Rep. RIVM report no. 773301 001/NOP report no. 410200 051, 142 pp., RIVM, Bilthoven, 2001.
- Park, R. J., Jacob, D. J., Field, B. D., et al.: Natural and transboundary pollution influences on sulfate-nitrate-ammonium aerosols in the United States: Implications for policy, *J. Geophys.*

Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Res., 109, D15204, doi:10.1029/2003jd004473, 2004.

Pierce, J. R. and P. J. Adams: Global evaluation of CCN formation by direct emission of sea salt and growth of ultrafine sea salt, *J. Geophys. Res.*, 111, D06203, doi:10.1029/2005jd006186, 2006.

5 Pierce, J. R. and Adams, P. J.: Uncertainty in global CCN concentrations from uncertain aerosol nucleation and primary emission rates, *Atmos. Chem. Phys.*, 9, 1339–1356, doi:10.5194/acp-9-1339-2009, 2009.

Platnick, S. and Twomey, S.: Determining the susceptibility of cloud albedo to changes in droplet concentration with the advanced very high resolution radiometer, *J. Appl. Meteorol.*, 10 33, 334–347, 1994.

Spracklen, D. V., Pringle, K. J., Carslaw, K. S., Chipperfield, M. P., and Mann, G. W.: A global off-line model of size-resolved aerosol microphysics: I. Model development and prediction of aerosol properties, *Atmos. Chem. Phys.*, 5, 2227–2252, doi:10.5194/acp-5-2227-2005, 2005a.

15 Spracklen, D. V., Pringle, K. J., Carslaw, K. S., Chipperfield, M. P., and Mann, G. W.: A global off-line model of size-resolved aerosol microphysics: II. Identification of key uncertainties, *Atmos. Chem. Phys.*, 5, 3233–3250, doi:10.5194/acp-5-3233-2005, 2005b.

Spracklen, D. V., Carslaw, K. S., Kulmala, M., et al.: Contribution of particle formation to global cloud condensation nuclei concentrations, *Geophys. Res. Lett.*, 5(12), 3233–3250, 2005b.

20 Streets, D. G., Bond, T. C., Carmichael, G. R., et al.: An inventory of gaseous and primary aerosol emissions in Asia in the year 2000, *J. Geophys. Res.*, 108, L06808, doi:10.1029/2002jd003093, 2003.

Suni, T., Kulmala, M., Hirsikko, A., Bergman, T., Laakso, L., Aalto, P. P., Leuning, R., Cleugh, H., Zegelin, S., Hughes, D., van Gorsel, E., Kitchen, M., Vana, M., Hörrak, U., Mirme, S., 25 Mirme, A., Sevanto, S., Twining, J., and Tardos, C.: Formation and characteristics of ions and charged aerosol particles in a native Australian Eucalypt forest, *Atmos. Chem. Phys.*, 8, 129–139, doi:10.5194/acp-8-129-2008, 2008.

Toon, O. B., Turco, R. P., Westphal, D., et al.: A Multidimensional Model for Aerosols: Description of Computational Analogs, *J. Atmos. Sci.*, 45(15), 2123–2144, 1988.

30 Trivitayanurak, W., Adams, P. J., Spracklen, D. V., and Carslaw, K. S.: Tropospheric aerosol microphysics simulation with assimilated meteorology: model description and intermodel comparison, *Atmos. Chem. Phys.*, 8, 3149–3168, doi:10.5194/acp-8-3149-2008, 2008.

Turco, R. P., Hamill, P., Toon, O. B., et al.: A One-Dimensional Model Describing Aerosol For-

**Sensitivity of global
CCN to primary
sulfate emissions**

G. Luo and F. Yu

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

mation and Evolution in the Stratosphere: I. Physical Processes and Mathematical Analogs, *J. Atmos. Sci.*, 36(4), 699–717, 1979.

Twomey, S.: The Influence of Pollution on the Shortwave Albedo of Clouds, *J. Atmos. Sci.*, 34(7), 1149–1152, 1977.

5 Wang, M. and Penner, J. E.: Aerosol indirect forcing in a global model with particle nucleation, *Atmos. Chem. Phys.*, 9, 239–260, doi:10.5194/acp-9-239-2009, 2009.

Whitby, K. T.: The physical characteristics of sulfur aerosols, *Atmos. Environ.*, 12(1–3), 135–159, 1978.

10 Whitby, K. T., Cantrell, B. K., and Kittelson, D. B.: Nuclei formation rates in a coal-fired power plant plume, *Atmos. Environ.*, 12(1–3), 313–321, 1978.

Wilson, W. E.: Sulfate formation in point source plumes: a review of recent studies, *Atmos. Environ.*, 15, 2573–2581, 1981.

Yu, F.: Ion-mediated nucleation in the atmosphere: Key controlling parameters, implications, and look-up table, *Atmos. Environ.*, 115, D03206, doi:10.1029/2009JD012630, 2010a.

15 Yu, F.: An extended secondary organic aerosol formation model: effect of oxidation aging and implications, *Atmos. Chem. Phys. Discuss.*, 10, 19811–19844, doi:10.5194/acpd-10-19811-2010, 2010b.

20 Yu, F. and Turco, R.: Case studies of particle formation events observed in boreal forests: implications for nucleation mechanisms, *Atmos. Chem. Phys.*, 8, 6085–6102, doi:10.5194/acp-8-6085-2008, 2008.

Yu, F., Wang, Z., Luo, G., and Turco, R.: Ion-mediated nucleation as an important global source of tropospheric aerosols, *Atmos. Chem. Phys.*, 8, 2537–2554, doi:10.5194/acp-8-2537-2008, 2008.

25 Yu, F. and Luo, G.: Simulation of particle size distribution with a global aerosol model: contribution of nucleation to aerosol and CCN number concentrations, *Atmos. Chem. Phys.*, 9, 7691–7710, doi:10.5194/acp-9-7691-2009, 2009.

Yu, F., Luo, G., Bates, T., et al.: Spatial distributions of particle number concentrations in the global troposphere: Simulations, observations, and implications for nucleation mechanisms, *J. Geophys. Res.*, 115, D17205, doi:10.1029/2009JD013473, 2010.

30 Ziemba, L. D., Griffin, R. J., and Talbot, R. W.: Observations of elevated particle number concentration events at a rural site in New England, *J. Geophys. Res.*, 111, D23S34, doi:10.1029/2006JD007607, 2006.

Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu

Table 1. Overview of the four simulations presented in this work.

	FS0	FS2.5FN5	FS2.5FN15	FS5FN5
f_{sulfate}	0	2.5%	2.5%	5%
f_{nucl}	0	5%	15%	5%
f_{accu}	0	95%	85%	95%

f_{sulfate} is the fraction of anthropogenic sulfur mass emitted as primary sulfate particles, which is used to represent sub-grid SO_2 oxidation and sulfate particle formation in anthropogenic sulfur plumes; f_{nucl} is the fraction of primary sulfate mass in the nucleation mode; f_{accu} is the fraction of primary sulfate mass distributed into the accumulation mode.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Mean number concentrations of condensation nuclei larger than 10 nm (CN10) measured in 21 surface stations around the globe. The simulated values for the FS0, FS2.5FN5, FS2.5FN15, and FS5FN5 sensitivity study cases are from the corresponding locations in GEOS-Chem.

Location or Station Name		Obs.	FS0	FS2.5FN5	FS2.5FN15	FS5FN5
A	Point Barrow, AK, USA	231 ^a	293	317	377	341
B	Pallas-Sammaltunturi, Finland	802 ^a	713	877	1202	999
C	Värriö, Finland	823 ^b	944	1215	1731	1407
D	Hyytiälä, Finland	2016 ^b	1972	2485	3334	2794
E	Uto, Baltic Sea	2921 ^b	1972	2485	3334	2794
F	Aspvreten, Sweden	2567 ^b	2047	2386	2991	2591
G	Mace Head, Ireland	1907 ^a	1015	1165	1488	1287
H	Melpitz, Germany	4664 ^c	4135	5714	8335	6749
I	Sable Island, NS, Canada	869 ^d	1141	1174	1253	1191
J	Thompson Farm, NH, USA	7039 ^f	4299	4437	4749	4504
K	Trinidad Head, California, USA	918 ^a	867	1129	1657	1367
L	Bondville, IL, USA	5038 ^a	3542	6440	11074	8220
M	Southern Great Plains, OK, USA	5064 ^a	1688	2963	4936	3747
N	Mount Waliguan, China	2030 ^j	915	1096	1385	1208
O	Mukteshwar, India	2730 ^j	2421	3789	5564	4616
P	American Samoa, USA	270 ^a	207	206	210	208
Q	Botsalano game reserve, S. Africa	2340 ^g	1230	2334	4035	3099
R	Bago State forest, Australia	1800 ^h	3509	4102	5740	4738
S	Cape Grim, Australia	1203 ^a	558	618	769	677
T	Neumayer, Antarctica	324 ^a	310	309	306	307
U	South Pole, Antarctica	156 ^a	261	260	258	259
Statistic Coefficient						
	Mean Absolute Error	0	772.0	720.0	1285.9	925.9
	Standard Deviation of Error	0	1070.7	1077.0	1856.9	1331.7
	Coefficient of Determination	1	0.697	0.692	0.595	0.644

Sources of CN10 data: ^a (World Data Centre for Aerosols, <http://wdca.jrc.ec.europa.eu>); ^b (Dal Maso et al., 2008); ^c (CREATE Aerosol Database at NILU, <http://tarantula.nilu.no/projects/ccc/create/index.htm>); ^d (NOAA ESRL/GMD Aerosol Database, <http://www.cmdl.noaa.gov/aero/>); ^f (Ziamba et al., 2006); ^g (Laakso et al., 2008); ^h (Sun et al., 2008); ⁱ (Kivekäs et al., 2009); ^j (Komppula et al., 2009).

Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu

Table 3. A comparison of simulated impacts of different assumptions of primary sulfate emissions on global annual mean CCN_{0.2}. The percentage given in the table is the change relative to the case without any primary sulfate emission (i.e., FS = 0).

Cases	Nucleation theories	Global Boundary Layer	Whole Troposphere
FS2.5FN5, this study	IMN	8%	3%
FS2.5FN15, this study	IMN	16%	8%
FS5FN5, this study	IMN	11%	5%
FS2FN15, Wang and Penner, 2009	BHN	53%	
FS2FN15, Wang and Penner, 2009	BHN.EMP	23%	

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Sensitivity of global
CCN to primary
sulfate emissions

G. Luo and F. Yu




Fig. 1. (a) Horizontal distribution of annual mean condensational sink (CS) in the boundary layer (0–1 km) and (e) zonally-averaged vertical profile of annual mean CS for the baseline case simulation (i.e., case FS2.5FN5 in Table 1). Panels (b–d) and (f–h) give the corresponding horizontal and vertical percentage changes of CS induced by different primary sulfate emission parameterizations. FS0, FS2.5FN5, FS2.5FN15 and FS5FN5 are the cases defined in Table 1.

Sensitivity of global
CCN to primary
sulfate emissions

G. Luo and F. Yu




Fig. 2. Same as Fig. 1, except for the nucleation rate (J).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Sensitivity of global
CCN to primary
sulfate emissions

G. Luo and F. Yu




Fig. 3. Simulated boundary layer aerosol particles (BLAP) number size distributions over **(a)** global, **(b)** land, and **(c)** ocean. Black, red, yellow and blue curves represent the FS0, FS2.5FN5, FS2.5FN15, and FS5FN5 cases, respectively.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 4. Same as Fig. 1, except for the number concentration of condensation nuclei large than 10 nm (CN10).

Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu




Fig. 5. (a) Locations of aerosol number concentration measurements used for simulation evaluation. Letters correspond to those listed in Table 2. (b) Comparison of observed and simulated annual mean number concentrations of CN10 at the 21 low boundary layer sites. The solid line shows a 1:1 ratio and the dashed lines show ratios of 1:2 and 2:1.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sensitivity of global CCN to primary sulfate emissions

G. Luo and F. Yu




Fig. 6. Same as Fig. 1, except for the concentrations of cloud condensation nuclei at water supersaturation ratio of 0.2% (CCN0.2).

Title Page

Abstract	Introduction
Conclusions	References
Tables	Figures

⏪
⏩

◀
▶

Back
Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

