

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Dry deposition of reactive nitrogen to European ecosystems: a comparison of inferential models across the NitroEurope network

C. R. Flechard¹, E. Nemitz², R. I. Smith², D. Fowler², A. T. Vermeulen³, A. Bleeker³, J. W. Erisman³, D. Simpson^{4,5}, L. Zhang⁶, Y. S. Tang², and M. A. Sutton²

¹INRA, Agrocampus Ouest, UMR1069 Sol Agro et hydrosystème Spatialisation, 65, Rue de Saint-Brieuc, 35042 Rennes, France

²Center for Ecology and Hydrology (CEH) Edinburgh, Penicuik, UK

³ECN, Netherlands Energy Research Foundation, Petten, The Netherlands

⁴EMEP MSC-W, Norwegian Meteorological Institute, Norway

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



⁵Dept. Earth and Space Sciences, Chalmers University of Technology, Gothenburg, Sweden

⁶Environment Canada, Toronto, Canada

Received: 28 October 2010 – Accepted: 18 November 2010 – Published: 1 December 2010

Correspondence to: C. R. Flechard (chris.flechard@rennes.inra.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Inferential models have long been used to determine pollutant dry deposition to ecosystems from measurements of air concentrations and as part of national and regional atmospheric chemistry and transport models, and yet models still suffer very large uncertainties. An inferential network of 55 sites throughout Europe for atmospheric reactive nitrogen (N_r) was established in 2007, providing ambient concentrations of gaseous NH_3 , NO_2 , HNO_3 and HONO and aerosol NH_4^+ and NO_3^- as part of the NitroEurope Integrated Project.

Network results providing modelled inorganic N_r dry deposition to the 55 monitoring sites are presented, using four existing dry deposition routines, revealing inter-model differences and providing ensemble average deposition estimates. Dry deposition is generally largest over forests in regions with large ambient NH_3 concentrations, exceeding $30\text{--}40\text{ kg N ha}^{-1}\text{ yr}^{-1}$ over parts of The Netherlands and Belgium, while some remote forests in Scandinavia receive less than $2\text{ kg N ha}^{-1}\text{ yr}^{-1}$. Turbulent N_r deposition to short vegetation ecosystems is generally smaller than to forests due to reduced turbulent exchange, but also because NH_3 inputs to fertilised, agricultural systems is limited by the presence of a substantial NH_3 source in the vegetation, leading to periods of emission as well as deposition.

Differences between models reach a factor 2–3 and are often greater than differences between monitoring sites. For soluble N_r gases such as NH_3 and HNO_3 , non-stomatal pathways are responsible for most of the annual uptake over many surfaces, especially the non-agricultural land uses, but parameterisations of the sink strength vary considerably among models. For aerosol NH_4^+ and NO_3^- , discrepancies between theoretical models and field flux measurements lead to much uncertainty in dry deposition rates for fine particles ($0.1\text{--}0.5\text{ }\mu\text{m}$). The validation of inferential models at the ecosystem scale is best achieved by comparison with direct long-term micrometeorological N_r flux measurements, but too few such datasets are available, especially for HNO_3 and aerosol NH_4^+ and NO_3^- .

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

The environmental effects of excess atmospheric reactive nitrogen (N_r) deposition to terrestrial ecosystems include soil acidification, the eutrophication of water bodies, nutrient imbalances, the leaching of base cation and nitrate, loss of biodiversity, direct toxicity to plants, increased N_2O emissions, and the inhibition of soil CH_4 oxidation (Galloway et al., 2003; Erisman et al., 2007). Elevated N_r deposition rates are the result of increased ambient concentrations and emissions by intensive farming (mostly reduced N_r) and by traffic and industry (mostly oxidised N_r). A role of N_r deposition as a strong driver of carbon sequestration by temperate and boreal forests has been suggested (Magnani et al., 2007) but the magnitude of the effect (kg C sequestered/kg N deposited) has been contested (de Vries et al., 2008; Sutton et al., 2008). Dry and wet deposition control the atmospheric life times and mean transport distances of N_r species downwind from point and diffuse sources and therefore affect pollutant transport across borders. This is evaluated at the European scale within the framework of the 1979 Convention on Long Range Transboundary Air Pollution (CLRTAP) (UNECE, 1999, www.unece.org/env/lrtap/) and the associated European Monitoring and Evaluation Programme (EMEP, www.emep.int), using gas and particle concentration monitoring networks to validate atmospheric model simulations (e.g. Fagerli and Aas, 2008; Simpson et al., 2006a). In North America, the Canadian Air and Precipitation Monitoring Network (CAPMoN; <http://www.ec.gc.ca/rs-mn/>) and the US Clean Air Status and Trends Network (CASTNet; <http://www.epa.gov/castnet>) have also been monitoring air concentrations for more than three decades.

The dry deposition of N_r , present in air in various inorganic species such as gaseous NH_3 , HNO_3 , $HONO$, NO , NO_2 and aerosol NH_4^+ and NO_3^- , as well as in a range of organic molecules in both phases (e.g. gaseous peroxyacetyl nitrate (PAN) and other organic nitrates, amines), typically contributes between one third and two thirds of total atmospheric N deposition (Erisman et al., 1996; Simpson et al., 2006a; Zimmermann et al., 2006, 2009). The partitioning between dry, wet and occult (i.e. cloud water)

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

deposition depends on atmospheric gas and aerosol N_r concentrations, weather patterns as well as land use/vegetation characteristics such as surface roughness, canopy leaf surface area and vegetation wetness. Unlike wet deposition, which is widely monitored in regional networks of wet-only or bulk precipitation collectors, measurements of dry (turbulent) N_r exchange fluxes have largely remained experimental and limited to selected research sites and to measurement campaigns of typically a few days to a few months, due to technical complexity and to the large equipment and operational costs involved. N_r concentration detectors that are reliable, sturdy, interference-free, fast and precise have proved elusive so far, at least as far as long-term micrometeorological flux measurements are concerned. Additional issues concerning inlet design, sampling losses and air column chemical reactions for highly reactive and soluble N_r species further indicate that large-scale dry deposition monitoring networks remain as yet impracticable.

Inferential modelling has been used extensively as an operational tool to obviate the absence of measured dry deposition data at regional scales (Baumgardner et al., 2002; Sickles and Shadwick, 2007; Erisman et al., 2005; Zhang et al., 2009). The method was originally developed to assess ecosystem damage in areas subjected to acid (sulphur) deposition and to compute regional pollutant mass balances (e.g. Wesely and Hicks, 1977; Garland, 1977).

Dry deposition, or bi-directional surface/atmosphere exchange, may be inferred from the knowledge of (measured) atmospheric gaseous or particulate pollutant concentration above vegetation (or any roughness element at the Earth's surface), using various assumptions regarding transfer rates through the air and the surface. A number of increasingly complex inferential schemes have been implemented in atmospheric transport chemical models (Meyers et al., 1998; Wesely and Hicks, 2000; Wu et al., 2003; Zhang et al., 2003), or are being proposed for implementation (Wu et al., 2009, Zhang et al., 2010, Massad et al., 2010 in the case of NH_3), and these can also be used to interpret micrometeorological field flux measurements. These models have been parameterised on the basis of measured field flux data, but specific exchange processes

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

and pathways are still poorly understood and their parameterisations remain crude and largely empirical. Also, model development has taken place in different countries with different pollution climates, so that parameterisations derived from field data may not be universally valid. Thus, much variation in dry deposition estimates may be expected between models, hinting that uncertainties remain rather large.

In 2006 the EU-sponsored NitroEurope Integrated Project (NEU for short) established a continent-wide network of 55 sites to monitor monthly ambient inorganic N_r concentrations over a large range of ecosystems and to estimate dry deposition fluxes using inferential techniques (Sutton et al., 2007; Tang et al., 2009), with the final aim to interpret CO_2 and greenhouse gas exchange across the network in relation to atmospheric N_r inputs. The primary objective of this paper is to provide an ensemble average estimate of N_r deposition for monitoring sites across the network, based on measured concentration data from the first two years of the project (2007–2008), and obtained by running four existing dry deposition schemes at the ecosystem scale. The wide range of vegetation types, meteorological conditions and pollution climates described by all monitoring sites provide a comprehensive dataset, with which to explore the differences in model responses to input data. The four routines are described and compared with a view to point out the similarities and the major differences in the approaches adopted by each model, to identify priority areas for potential improvement, and to assess current uncertainties in dry deposition estimates from inferential networks.

2 Materials and methods

2.1 Dry deposition models

The four dry deposition routines implemented in this study, which are currently used as modules within chemical transport models (CTMs) at national or continental scales in Europe and N. America, include the UK CBED scheme (Smith et al., 2000), the Dutch

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



IDEM model (Bleeker et al., 2004; Erisman et al., 1994), the dry deposition module of the Environment Canada model (Zhang et al., 2001, 2003), termed “CDRY” here, and the surface exchange scheme of the EMEP model used under the CLRTAP (Simpson et al., 2003; see also Tuovinen et al., 2009, and refs therein). It should be noted that here we use the deposition module of EMEP version rv3.1 as documented in Simpson et al. (2003). The latest code (rv3.7, Simpson et al., 2010) carries a considerably different formulation for aerosol deposition, but is still undergoing testing. To distinguish these schemes we refer to the rv3.1 version as EMEP-03.

For the CTM versions of the models, in which the dry deposition schemes are embedded, the spatial patterns of dispersion, transport, chemistry and wet and dry deposition, as well as the whole regional mass balance of pollutants, are computed using input meteorological data from numerical weather prediction (NWP) models, prescribed emissions and land-use data. In the present application, however, the dry deposition routines are decoupled from any regional framework; they are driven instead at each individual site of the NEU network by local (field-scale) measurements of atmospheric concentrations, turbulence and meteorology. Thus, deposition estimates that are provided in this paper for any of the 4 models refer by default to “local” or ecosystem-scale runs of the dry deposition routines, rather than to the grid square average (e.g. 50 × 50 km) that could be provided by the CTM version (unless otherwise specified). Consequently, this analysis only assesses the parameterisations of the dry deposition models, and not the ability of their respective CTM frameworks to predict meteorology, concentrations or the built-in representations of vegetation characteristics.

2.1.1 Trace gases

The surface-vegetation-atmosphere transfer (SVAT) models use broadly similar resistance frameworks for pollutant trace gas exchange. The atmospheric aerodynamic resistance, noted $R_a(z_{\text{ref}}, d + z_0)$ or $R_a(z_{\text{ref}})$ for short, characterises the efficiency of turbulent transfer from a reference height z_{ref} in the surface layer down to $d + z_0$, d being the displacement height and z_0 being the momentum roughness length; the quasi-

laminar sublayer resistance (R_b) accounts for the transfer across a viscous, pseudo-laminar sub layer in the immediate vicinity of the vegetation or soil surface; and the surface or canopy resistance (R_c) characterises the surface affinity for pollutant uptake (Baldocchi et al., 1987; Monteith and Unsworth, 1990; Seinfeld and Pandis, 2006).

5 Mathematical expressions for R_a and R_b are well documented; the method of calculation is very similar in the models, and the reader is referred to the literature for the various formulations. The main differences between dry deposition models reside in the parameterisations for R_c . Differences in R_a and R_b do arise between models due to e.g. the use of marginally different atmospheric stability corrections, different assumptions regarding the viscous sublayer, and above all due to the model default value for z_0 , which controls the magnitude of friction velocity (u_*). The CBED model does not actually compute stability corrections for R_a , based on the postulate that neutral conditions largely prevail over the windy British Isles (Smith et al., 2000). For the sake of model comparability, however, they are included here in the base runs of the CBED
10 module and computed in an identical fashion to the EMEP-03 scheme. Alternative runs of the CBED model, in which the stability corrections were not implemented, are compared with the base runs in Fig. A2 of the Supplement published online, showing that stability corrections have little impact on annually averaged modelled fluxes.

In its simplest form the dry deposition flux F_χ is given as the product of concentration at the reference height $\chi(z_{\text{ref}})$ by the deposition velocity at the same level $V_d(z_{\text{ref}})$:

$$20 \quad F_\chi = -\chi(z_{\text{ref}}) \cdot V_d(z_{\text{ref}}) \quad (1)$$

with, by convention, negative fluxes denoting deposition, and V_d the inverse sum of resistances in series:

$$V_d(z_{\text{ref}}) = [R_a(z_{\text{ref}}, d + z_0) + R_b + R_c]^{-1} \quad (2)$$

25 The canopy resistance for vegetated surfaces results from a network of sub-resistances within the canopy (Seinfeld and Pandis, 2006), with foliar stomatal (R_s), mesophyll (R_m), and non-stomatal (R_{ns}) or cuticular (R_{cut}) or water film (R_w) or external (R_{ext})

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



resistances, as well as non foliage terms, e.g. the soil or ground surface resistance (R_{gr}). Most models (EMEP-03, IDEM, CDRY) also include an in-canopy aerodynamic resistance (R_{ac}), acting between the assumed big-leaf and ground surface, while the CBED approach is strictly single-layered. The main sub-resistances of R_c are briefly presented here; for details the reader is referred to the original publications. Note that all resistances are expressed in $s\ m^{-1}$ by default throughout this paper.

Gaseous transfer through stomata

Stomatal resistances to gaseous transfer are typically derived in the different models using a light-response function of the generic type (Jarvis, 1976):

$$R_s = R_{s,\min} \left[1 + \frac{b'}{I_p} \right] / (f_e f_w f_T f_s) \quad (3)$$

Here I_p is light intensity taken either as the photosynthetically active radiation (PAR) or global radiation (S_t) as its proxy; b' is an empirical constant, $R_{s,\min}$ is a minimum value of the stomatal resistance to water vapour, with b' and $R_{s,\min}$ taking characteristic values for each vegetation type or land use; the correction factors f_e , f_w and f_T account for the effects of increasing vapour pressure deficit (vpd), plant water stress and temperature, respectively (Jarvis, 1976). The last factor f_s is a scaling factor to account for the difference in molecular diffusivity between water vapour and the trace gas considered. For the EMEP-03 model a further factor for phenology is also included (Emberson et al., 2001; Simpson et al., 2003).

Note that R_s in Eq. (3) is expressed on a unit leaf area (projected) basis, or equivalent to a unity leaf area index (LAI). All models except IDEM split PAR into its direct and diffuse fractions and compute the sunlit and shaded components of LAI, such that total (or bulk) stomatal resistance is calculated from sunlit and shaded resistances weighted by their respective LAI fractions (Baldocchi et al., 1987). Thus in CBED, CDRY and EMEP-03, the bulk stomatal conductance $G_s (= R_s^{-1})$ does not increase linearly with

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



total LAI but tends to saturate for larger LAI levels. By contrast, IDEM uses by default a simplified version, in which LAI is not split into sunlit and shaded fractions, but where G_s is proportional to total LAI. The R_s routine by Wesely et al. (1989), which only requires global radiation and surface temperature as input, may be used as an option in IDEM when land use and vegetation characteristics are not well known.

Non-stomatal resistances

Although non-stomatal pathways, either on leaf cuticles or other non-foliar surfaces (stems, bark, ground, etc), provide an important, and often dominant, sink for atmospheric gases on an annual basis (Fowler et al., 2001, 2009; Flechard et al., 1998), there are as yet no consensual, generic and fully mechanistic parameterisations for non-stomatal resistances, which are variously termed R_{ns} , R_{ext} , R_w , R_{cut} , R_{gr} in different models. This is partly due to the much greater technical and methodological difficulties, and larger uncertainties, involved in measuring trace N_r gas (e.g. NH_3 , HNO_3 , HONO, PAN) fluxes, let alone non-stomatal resistances, and also due to the resulting relative scarcity of reliable field observations, as compared with water vapour fluxes and R_s . Also, in addition to the many environmental factors that have been shown or surmised to be involved in the control of non-stomatal resistances (e.g. wetness, temperature, vegetation type, pollution climate, soil pH, leaf surface chemistry), it appears that hysteresis or “memory” effects control the rate of charge or discharge of the surface N_r pool, especially in the case of NH_3 (Sutton et al., 1998; Flechard et al., 1999; Neirynek and Ceulemans, 2008; Burkhardt et al., 2009; Wichink Kruit et al., 2010), challenging the applicability of a (static) resistance approach.

For NH_3 , the four models use widely different empirical schemes for non-stomatal resistances, reflecting the spread in mean values and functional relationships found in the literature. This is consistent with the different ecosystems and pollution climates in which the original NH_3 flux measurements were made (Nemitz et al., 2001; Massad et al., 2010). CBED actually uses a constant R_c of 20 s m^{-1} for forests and moorland, while for grasslands and crops the following R_w function is implemented (Smith et al.,

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2000):

$$R_w(\text{CBED}) = 10^{\log_{10}(T_s+2)} \cdot \exp\left(\frac{100-\text{RH}}{7}\right) \quad (4)$$

with T_s surface temperature ($^{\circ}\text{C}$) and RH surface relative humidity (in %). In frozen conditions R_w takes a constant value of either 1000 s m^{-1} ($T_s < -5^{\circ}\text{C}$) or 200 s m^{-1} ($-5^{\circ}\text{C} < T_s < 0^{\circ}\text{C}$). The EMEP-03 model uses the same basic formula (EMEP-03's F_1 factor is the same as R_w (CBED)), but then modulates R_w by a correction factor F_2 such that (Simpson et al., 2003):

$$F_2 = 0.0455 \times 10^{(-1.1099 \times a_{\text{SN}} + 1.6769)} \quad (5)$$

$$R_{\text{ns}}(\text{EMEP} - 03) = \min[200, \max[2, R_w(\text{CBED}) \cdot F_2]] \quad (6)$$

where a_{SN} is the ratio of atmospheric SO_2 to NH_3 mixing ratios. F_2 was quantified following the synthesis by Nemitz et al. (2001), who showed that R_{ext} observations across 8 UK and Dutch sites declined exponentially with a_{SN} , thus supporting the co-deposition hypothesis (Erisman and Wyers, 1993) that surface NH_3 uptake was most efficient (i.e. R_{ext} was smallest) at sites with a relative abundance of atmospheric SO_2 .

The R_{ext} parameterisation for NH_3 in IDEM also uses a functional dependence on RH (Eq. 7), although this is often supplanted by default values in given circumstances related to land use, season, snow cover, surface wetness, and surface acidity as quantified by the proxy a_{SN} (Erisman et al., 1994; Bleeker et al., 2004). Default R_{ext} values range from typically $10\text{--}20 \text{ s m}^{-1}$ in forests, moorland, crops and ungrazed pasture in wet conditions, to $200\text{--}1000 \text{ s m}^{-1}$ in fertilised systems in dry summer night-time (Bleeker et al., 2004).

$$R_{\text{ext}}(\text{IDEM}) = 19257 \cdot \exp(-0.094 \cdot \text{RH}) + 5 \quad (7)$$

In CDRY, explicit and specific parameterisations of R_{cut} exist only for SO_2 and O_3 as functions of leaf wetness (dry vs. wet; dew vs. rain), relative humidity, leaf area index and friction velocity. Values of R_{cut} for other gases are calculated as multipliers of

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$R_{\text{cut}}(\text{SO}_2)$ or $R_{\text{cut}}(\text{O}_3)$ or a combination of both. For NH_3 , R_{cut} is taken to be identical to that for SO_2 , in both the wet (Eq. 8a) and dry (Eq. 8b) cases:

$$R_{\text{cut},w}(\text{CDRY}) = \frac{R_{\text{cut},w0}}{u_* \cdot \text{LAI}^{0.5}} \quad (8a)$$

$$R_{\text{cut},d}(\text{CDRY}) = \frac{R_{\text{cut},d0}}{\exp^{0.03 \cdot \text{RH}} u_* \cdot \text{LAI}^{0.25}} \quad (8b)$$

5 with $R_{\text{cut},w0}$ and $R_{\text{cut},d0}$ being land-use specific reference values (Zhang et al., 2003).

For HNO_3 , the scarcity of field flux measurements to date means that there are few data from which to derive parameterisations, and two models use near-zero R_c values in most cases (CBED, EMEP-03). By contrast IDEM implements a substantial R_c of 10 s m^{-1} by default and of 50 s m^{-1} for frozen or snow-covered surfaces, while CDRY models R_{cut} (HNO_3) on the basis of the reference values for SO_2 and O_3 (Zhang et al., 2003), resulting in R_c values that are an order of magnitude smaller than those for SO_2 . For HONO, there are even less data available, and it is only treated by CDRY using an R_w value a factor 5 larger than that for HNO_3 .

15 Nitrogen dioxide exchange is assumed by all models to be exclusively downward (deposition only), and mostly (CDRY, EMEP-03, IDEM) or entirely (CBED) controlled by stomatal opening. In the EMEP-03 model, however, NO_2 dry deposition is switched off whenever the ambient concentration falls below 4 ppb. This reflects the pseudo compensation point behaviour of NO_2 exchange, due to NO emissions from the soil and conversion within plant canopies to NO_2 through reaction with O_3 , that leads to net (NO_x) emissions in the field at small ambient concentrations (Simpson et al., 2003).

2.1.2 NH_3 compensation point modelling

20 One exception to the deposition-only (R_c) paradigm prevalent in surface/atmosphere N_r exchange modelling is the bi-directional canopy compensation point approach for

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NH₃ (Sutton et al., 1998), implemented in the CBED model for crops and grass land use classes (LUC) (Smith et al., 2000). Here, a non-zero canopy-equivalent potential, termed the canopy compensation point χ_c , determines the direction and sign of the flux when compared with the atmospheric concentration $\chi(z_{\text{ref}})$, such that:

$$F_{\chi} = -\frac{\chi(z_{\text{ref}}) - \chi_c}{R_a(z_{\text{ref}}) + R_b} \quad (9)$$

The canopy compensation point is a function of, and quantifies the net bulk effect of, all source and sink terms within the canopy, but it is also a weak function of the atmospheric concentration $\chi(z_{\text{ref}})$ itself (Nemitz et al., 2000a). In CBED, a basic version is implemented, where the *stomatal* compensation point (χ_s) provides the only potential NH₃ source, the dissolved NH₃ and NH₄⁺ pool in the apoplast of sub-stomatal cavities (Farquhar et al., 1980; Schjøerring et al., 1998; Massad et al., 2008) being mediated by the stomatal resistance R_s , while R_w characterises the sink strength of non-stomatal foliar surfaces. Other mechanistic models (e.g. Nemitz et al., 2000a, 2000b, 2001; Personne et al., 2009) consider additional NH₃ sources in e.g. seed pods of oilseed rape and in the leaf litter and soil under winter wheat and grassland. Such approaches have not been implemented in CTMs to date, partly because this would require detailed (and generally unavailable) knowledge of sub-grid variations in NH₃ concentrations, vegetation/crop type and fertilisation practices.

The stomatal compensation point in CBED is calculated following Eq. (10) assuming an apoplastic pH of 6.8 and intercellular NH₄⁺ concentration of 600 μmol l⁻¹, i.e. an apoplastic Γ_s ratio ($=[\text{NH}_4^+]/[\text{H}^+]$) of 3785:

$$\chi_s = \frac{K_a(T_s)}{K_h(T_s)} \Gamma_s \quad (10)$$

with $K_a(T_s)$ the dissociation constant of NH₃ in water (Bates and Pinching, 1950) and $K_h(T_s)$ the Henry coefficient for NH₃ (Dasgupta and Dong, 1986). The canopy

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



compensation point itself is given as (Sutton et al., 1998):

$$\chi_c = \frac{\frac{\chi(z_{\text{ref}})}{[R_a(z_{\text{ref}}) + R_b]} + \frac{\chi_s}{R_s}}{[R_a(z_{\text{ref}}) + R_b]^{-1} + R_s^{-1} + R_w^{-1}} \quad (11)$$

2.1.3 Aerosol deposition

Aerosol dry deposition fluxes are computed as the product of air particle concentration by deposition velocity (Eq. 1). Parameterisations for aerosol V_d range from the strongly mechanistic to the fully empirical, depending on the model and the ion species considered. The 2003 version of the unified EMEP model (EMEP-03), the CDRY scheme and to some extent the IDEM model, are originally based on Slinn's approach (Slinn, 1982), but have distinctly different features. In EMEP-03, V_d is calculated as (Simpson et al., 2003):

$$V_d(z_{\text{ref}}) = \frac{1}{R_a(z_{\text{ref}}) + R_b + [R_a(z_{\text{ref}}) \cdot R_b \cdot V_g]} + V_g \quad (12)$$

where V_g is the gravitational settling (or sedimentation) velocity (Seinfeld and Pandis, 2006), calculated as a function of particle diameter (D_p), and R_b is calculated from explicit formulations from the literature that are particle size- and vegetation/land use-dependent.

By contrast, CDRY does not explicitly compute R_b but uses an overall surface resistance (R_{surf}) concept such that (Zhang et al., 2001):

$$V_d(z_{\text{ref}}) = \frac{1}{R_a(z_{\text{ref}}) + R_{\text{surf}}} + V_g \quad (13)$$

$$R_{\text{surf}} = \frac{1}{\varepsilon_0 u_* R_1 (E_B + E_{\text{IN}} + E_{\text{IM}})} \quad (14)$$

where ε_0 is an empirical constant and R_1 the fraction of particles that stick to the surface. Parameters used to calculate aerosol collection efficiencies E_B (Brownian diffusion), E_{IN} (interception) and E_{IM} (impaction) are land-use and season-dependent.

In IDEM, the deposition velocity for particulate NH_4^+ and NO_3^- is calculated according to Wesely et al. (1985) for short vegetation and other areas with a momentum roughness length smaller than 0.5 m. For forests and other areas with $z_0 > 0.5$ m, the scheme by Ruijgrok et al. (1997) is used, such that:

$$V_d(z_{\text{ref}}) = \frac{1}{R_a(z_{\text{ref}}) + V_{\text{ds}}^{-1}} + V_g \quad (15)$$

$$V_{\text{ds}} = E \frac{u_*^2}{U_{hc}} \quad (16)$$

with V_{ds} the surface deposition velocity, E the overall collection efficiency and U_{hc} the wind speed at canopy height (h_c). It can readily be seen that V_{ds} is equivalent to R_{surf}^{-1} of CDRY (Eq. 13), but Ruijgrok et al. (1997) derived simplified relationships for the overall collection efficiency E and V_{ds} for the chemical species NH_4^+ , SO_4^{2-} , NO_3^- and Na^+ and other base cations under various conditions. For $\text{RH} < 80\%$ E is of the form:

$$E = \alpha u_*^\beta \quad (17)$$

where the empirical constants α and β are chemical species- and surface wetness-dependent. For relative humidity above 80% they introduce a dependence on relative humidity to account for the observed increased V_{ds} with growing particle diameter (D_p). In IDEM, the calculation scheme for the settling velocity V_g (implemented for large particles only) is similarly simplified. Note that gravitational settling is included conceptually in Eqs. (13), (14) and (16), although it is negligible for the fine aerosol fraction (aerodynamic diameter $< 1 \mu\text{m}$), where most of NH_4^+ and NO_3^- mass is likely found, and only becomes relevant for coarse particles.

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The CBED model currently calculates NH_4^+ , NO_3^- and SO_4^{2-} aerosol deposition velocities using a simple, empirically-derived scheme, whereby V_d is the product of u_* times a tabulated land use- and chemical species-specific constant (α). The parameter α is of the order of 0.005 for grassland and semi-natural vegetation, of 0.01 for arable land, and of 0.02–0.03 for forests (for u_* and V_d expressed in the same unit e.g. m s^{-1}); also, $\alpha(\text{NO}_3^-)$ is 49%, 36% and 60% larger than $\alpha(\text{NH}_4^+)$ for grassland/semi-natural, arable land and forests, respectively (Smith and Nemitz, CEH Edinburgh, unpublished). These α values were derived by weighting measured curves of $V_d(D_p)/u_*$ over different ecosystems (Gallagher et al., 1997; Nemitz et al., 2002; Joutsenoja, 1992) with typical size-distributions of nitrate and ammonium.

2.2 NitroEurope inferential network sites

Reactive nitrogen dry deposition was estimated by field-scale inferential modelling at the 55 monitoring sites of the NitroEurope network (Sutton et al., 2007; Tang et al., 2009) where all necessary input data, including N_r atmospheric concentrations, meteorological and/or micrometeorological data, were available for the two years 2007–2008, or at least one full year. The network included 29 forest (F) stations; 9 semi-natural short vegetation ecosystems (SN) e.g. semi-arid steppe, alpine or upland grasslands, moorlands and fens; 8 fertilised, productive grasslands (G); and 9 cropland (C) sites (Table 1). All NEU inferential sites, with the exception of DE-Hoe, FI-Lom, NL-Spe and UA-Pet, were also CO_2 flux monitoring stations of the EU-funded CarboEurope Integrated Project (<http://www.carboeurope.org/>), which aimed at an assessment of the European terrestrial carbon balance (Dolman et al., 2008). Sites locations and vegetation characteristics are summarised in Table 1, and details and photographs may be obtained from the CarboEurope-IP database (<http://gaia.agraria.unitus.it/database/carboeuropeip/>), or from the list of selected references provided in Table A1 in the online Supplement to this article. The study sites were distributed across Europe from Ireland to Russia and from Finland to Portugal,

with mean annual temperatures ranging from -0.1°C (FI-Lom) to 17.8°C (ES-ES1), and mean annual rainfall ranging from 464 mm (UA-Pet) to 1450 mm (IE-Dri). Sites elevations range from -2 m a.m.s.l. (NL-Hor) to 1765 m a.m.s.l. (ES-VDA). Measured maximum canopy heights (h_c) and LAI are on average $20.2\text{ m}/4.9\text{ m}^2\text{ m}^{-2}$ for forests, $0.8\text{ m}/3.2\text{ m}^2\text{ m}^{-2}$ for semi-natural vegetation, $0.4\text{ m}/5.5\text{ m}^2\text{ m}^{-2}$ for grasslands and $1.8\text{ m}/7.0\text{ m}^2\text{ m}^{-2}$ for crops.

2.3 Input data and model implementation

2.3.1 Ecosystem and micrometeorological data

For a detailed description of the management of input data and model implementation at the ecosystem scale for all NEU monitoring sites, the reader is referred to the online Supplement. Briefly, the model base runs used measured values of h_c as inputs, whereas for LAI inputs the model default values were used preferentially, due to the uncertainties in measured estimates of LAI. A comparison of model default values of LAI and h_c with actual measurements is shown in Fig. 1c and d.

For u_* and sensible heat flux (H), actual measurements from EC datasets at each site were used whenever possible, and data were otherwise gap-filled from standard meteorological data (cf. Sect. A3 in Supplement). Measurements of canopy wetness were available at very few sites, and thus a dynamic surface wetness energy balance model was coupled to the modelling framework for most sites; a comparison with actual measurements is shown in Sect. A5 of Supplement.

Alternative model runs were computed to investigate the sensitivity of annual fluxes to input values of h_c and LAI and to surface temperature and relative humidity, as detailed in Sects. A2 and A4 of the Supplement, with the characteristics of the base and sensitivity runs being summarised in Table A2 therein.

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.3.2 Atmospheric N_r concentration data

Pollutant monitoring by denuder and filter sampling

Ambient N_r concentrations of gaseous NH₃, HNO₃ and HONO and aerosol NH₄⁺ and NO₃⁻ were monitored monthly at the 55 sites of the inferential network from early 2007 onwards using DELTA systems (DENuder for Long-Term Atmospheric sampling, described in detail in Sutton et al., 2001 and Tang et al., 2009) (Table 2). Briefly, the DELTA sampling “train” consists of two coated borosilicate glass denuder tubes in series for scrubbing acidic trace gases (HNO₃, SO₂, HCl, HONO), followed by two denuders for NH₃ and finally by a filter-pack assembly with a first impregnated filter to capture aerosol phase anions (NO₃⁻, SO₄²⁻, Cl⁻) as well as base cations (Na⁺, Mg²⁺, Ca²⁺), and a second filter to collect the evolved particulate NH₄⁺. Air is sampled at a rate of 0.3–0.4 l min⁻¹ and directly into the first denuder with no inlet line to avoid sampling losses. Denuders for acid gases and filters for aerosol anions and base cations are coated/impregnated with potassium carbonate/glycerol, while for gaseous NH₃ and aerosol NH₄⁺ citric acid or phosphorous acid is used. The empirically determined effective size cut-off for aerosol sampling is of the order of 4.5 μm (Nemitz, unpublished data).

The DELTA sampling trains were prepared and assembled in seven coordinator laboratories (CEAM, Spain; CEH, United Kingdom; FAL/vTI, Germany; INRA, France; MHSC, Croatia; NILU, Norway; and SHMU, Slovakia), sent out to the inferential sites for monthly field exposure, then sent back to the laboratories for denuder/filter extraction and analysis. The DELTA systems thus provided monthly mean ambient N_r concentrations for each site of the network; this paper deals, unless otherwise stated, with the data collected during the first two years (2007–2008) of the whole monitoring period (2007–2010).

To ensure comparability of data provided by the different laboratories, DELTA inter-comparison campaigns were carried out at yearly intervals at selected sites as part of a

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

defined QA/QC programme, whereby seven sample trains (one provided by each laboratory) were exposed side by side for a month and then extracted and analysed by each laboratory (Tang et al., 2009). In addition to this full intercomparison exercise, in which the whole sample train management (preparation, coating, impregnation, assembly, dispatching, exposure, field handling, extraction, analysis) was tested, each laboratory also regularly received synthetic solutions for “blind” analysis from three chemical intercomparison centres: CEH, Scotland; EMEP/NILU, Norway; and the Global Atmospheric Watch program (GAW) of the WMO. The results of the first DELTA intercomparisons were presented in Tang et al. (2009); an in-depth analysis of the full concentration dataset will be published in a companion paper (Tang et al., 2011).

In addition to the monthly denuder and filter N_r concentration data provided by DELTA systems, ambient NO_2 concentrations were monitored by chemiluminescence on an hourly or half-hourly basis at a number of sites (BE-Bra, FI-Hyy, IT-Ren, NL-Spe, FI-Lom, HU-Bug, UK-AMo, CH-Oe1, UK-EBu, FR-Gri, IT-Cas). Although NO_2 concentrations were not measured at all sites, the available data are useful to assess the likely magnitude of ecosystem NO_2 uptake relative to total N_r dry deposition and the variability between model predictions for NO_2 deposition. For the remaining sites, mean modelled NO_2 concentrations from the EMEP 50 km \times 50 km model output for the year 2004 were used.

Aerosol size distribution

The extraction of DELTA filters yielded total aerosol concentrations, as the fractions of fine vs. coarse aerosols could not be determined for each of NH_4^+ , NO_3^- or other chemical species. For the two aerosol V_d schemes (CBED, IDEM) that do not explicitly model aerosol size-dependent deposition velocities, but instead calculate a species-specific mean V_d across the aerosol size range, this was not an issue. However, in both the EMEP-03 and CDRY models, aerosol V_d is a function of particle diameter D_p . In EMEP-03 two deposition velocities are calculated, one for each of fine ($D_p = 0.3 \mu m$) and coarse ($D_p = 4 \mu m$) aerosols, independent of the chemical species considered. In

29309

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

CDRY, species-specific values of the geometric mean mass diameter (DG) and geometric standard deviation (GSD) are attributed to both fine and coarse aerosol modes, and two log-normal particle size distributions are generated on the basis of DG and GSD, one for each mode. In both models, therefore, the fine and coarse fractions of the total aerosol loading measured on the DELTA filters need to be estimated, so that modelled V_d is applied to the concentration in the appropriate size range. In the CTM versions of EMEP-03 and CDRY, fine and coarse fractions are calculated dynamically within the regional chemical model, but in the present local-scale application such data are not available. By default, and in a first approximation, fine aerosol was assumed to account for 94% of total NH_4^+ , 81% of total NO_3^- and 90% of total SO_4^{2-} following Ruijgrok et al. (1997), realising that in reality this ratio will be site specific, especially for NO_3^- , which has a larger contribution from coarse NaNO_3 at coastal sites.

Corrections for within-canopy concentration data

At most sites of the NEU network, air sampling by DELTA systems provided concentrations at least 1 m above the canopy. However, at 10 forest sites (BE-Vie, DE-Hai, DE-Tha, ES-ES1, ES-LMa, FI-Sod, IT-Ren, PT-Mi1, SE-Nor, SE-Sk2), the DELTA system was actually set up in a clearing or in the trunk space, typically 1.5 to 2 m above the forest floor. This was for practical reasons, mostly to facilitate the safe exchange of sampling trains in challenging winter conditions or windy weather. The inferential method requires atmospheric concentrations and turbulence intensity *above* the canopy to predict rates of dry deposition to the forest, and thus the validity of clearing or below-crown concentrations as proxies of above-canopy concentrations can be questioned and needs to be examined (Zhang et al., 2009; Tuovinen et al., 2009). There are very few published within-canopy (vertical) NH_3 and HNO_3 concentration profiles in the literature for forests. Within-canopy profile data for NH_3 have been obtained mostly in grasslands (Nemitz et al., 2009) and crops such as oilseed rape (Nemitz et al., 2000b) and maize (Bash et al., 2010). These data showed consistently larger concentrations near the ground and below canopy, compared with above the canopy, indicative of NH_3

correlations on daily time scales between concentration and V_d could lead to significant systematic bias in the modelled fluxes at some sites, but this was not investigated here.

For cases when all input data were available throughout the 2-year measurement period, the monthly and annual fluxes can simply be obtained by adding up all modelled half-hourly fluxes. In practise, however, there were at most sites periods of a few hours to a few days or weeks during which at least one key variable (such as windspeed, temperature or relative humidity) was missing, e.g. due to instrument malfunction, breakdown, power cuts or theft/vandalism, such that mechanistic gap-filling for fluxes was precluded. A simple upscaling procedure based on the arithmetic mean of all modelled fluxes multiplied by the total number of 30-min time intervals in the year potentially leads to a statistical bias. Thus, the approach adopted here consists of computing for each month the arithmetic mean diurnal cycle from all modelled half-hourly flux data, then scaling up to the whole month, and adding up 12 monthly fluxes for the annual total.

At intensively managed grassland and cropland sites of the NEU network, fertilisation occurred once to several times a year, in which net NH_3 emissions typically ensued over one or several weeks, and where elevated ambient NH_3 concentrations occurred as a result (e.g. Flechard et al., 2010). Here the modelled (inferential) NH_3 flux data from the fertilisation months were not included in the annual deposition total, the reason being twofold; first, inferential models are primarily deposition models and are not suited to situations with large NH_3 emissions e.g. from applied fertiliser, but to background conditions (Flechard et al., 2010); the special case of fertiliser- or manure-induced NH_3 losses requires a different kind of modelling approach (e.g. Genermont and Cellier, 1997) and is not considered here. Second, applying an inferential model to months when fertilisation occurred would result in a large deposition flux (due to the elevated NH_3 concentration) when net emission actually occurred, thus over-estimating annual deposition.

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Results and discussion

3.1 Model evaluation for key exchange variables

3.1.1 Gap-filling of friction velocity data

Measured values of u_* from EC datasets were used preferentially for flux modelling whenever possible; the prediction of u_* based on an assumed value of z_0 for vegetation and on meteorological conditions (Sect. A3, Supplement) was used only when measured turbulence data were missing. This represented on average 21% of the time across the network, although u_* data capture was close to 100% at some sites and less than 60% at others for the period 2007–2008.

For the gap-filling of u_* , the model base runs used measured values of h_c to calculate z_0 , while inferential models within the framework of CTMs would normally predict u_* from their own default h_c values. The discrepancies in modelled u_* are shown in Fig. 1, with the different default values of h_c leading to different u_* estimates between models across the sites (Fig. 1a). The actual use of measured h_c naturally suppressed these differences between models (Fig. 1b), with residual inter-model discrepancies being due to slightly different stability correction functions in the four models. Not surprisingly, the use of measured h_c (as opposed to model defaults) also considerably improved the agreement between modelled and measured u_* , and reduced the scatter in the relationship (Fig. 1b), even if there was a marked tendency to overestimate u_* over forests at the higher end of the scale. The three forest sites whose mean measured u_* was around 0.65 m s^{-1} (DE-Hai, DE-Wet, DK-Sor), and whose mean modelled u_* were 0.76 , 0.83 and 0.91 m s^{-1} , respectively, were 33 m tall beech, 22 m tall spruce and 31 m tall beech forests, respectively. The other forest site whose mean u_* (0.51 m s^{-1}) was largely overestimated (0.75 m s^{-1}) was NL-Spe, a mixed species 32 m tall stand, dominated in the near field by Douglas fir. These four forests have comparatively large maximum leaf area indices, in the range $5\text{--}11 \text{ m}^2 \text{ m}^{-2}$ (Table 1), which may reduce frictional retardation of wind. Further, the underlying model assumption that

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



z_0 increases linearly with h_c (with z_0 being normally calculated as one tenth of h_c in CBED, EMEP-03 and IDEM) is probably not valid, depending on canopy structure and leaf morphology, and z_0 values of 3 m for the aforesaid 30 m tall forests are therefore unrealistic. Another explanation is that most anemometers over forest are operated within the roughness sublayer, where wind speed is larger than would be predicted on the basis of the logarithmic wind profile, thus leading to larger modelled u_* values. This may be different for CTMs where the reference height is higher (e.g. 50 m).

Note that in this study, “modelled” u_* means a value derived from the measured wind speeds and stability functions; values of u_* in the regional application of these models depend on the NWP model and sub-grid treatment, and might be quite different. While the CTMs aim to capture h_c , u_* and other features relevant for dry deposition over representative landscapes, the comparison shown in Fig. 1a is only fully meaningful to the extent that the limited number of NEU sites may be considered as statistically representative of their land-use class.

3.1.2 Stomatal conductance

Stomatal conductance ($G_s = 1/R_s$) is controlled by leaf surface area and by PAR, as well as temperature, soil moisture and ambient relative humidity, and therefore strong seasonal cycles are expected in European conditions. The four models do show some temporal correlation with respect to G_s as shown in Fig. 2. Over forests, the mean daytime G_s was modelled to be generally largest in summer, with values of typically 5 to 10 mm s⁻¹. There were clear discrepancies between models in summer for forests, with G_s in CBED and IDEM typically a factor of two larger than in CDRY and EMEP-03 for coniferous forests, but the agreement was much better for deciduous forests (e.g. DE-Hai, DK-Sor, FR-Fon, FR-Hes). During the other seasons, the IDEM model stands out over the coniferous sites, with mean daytime G_s values of typically 10 mm s⁻¹, almost regardless of the season except in the more northerly regions, while the other three models are rather consistent and show reduced values compared with summer. At selected mediterranean or Southern European coniferous sites, where summer heat

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



stress and drought reduce stomatal exchange in summer, G_s values predicted by IDEM are actually marginally larger in winter than in summer (e.g. ES-ES1, FR-LBr, IT-SRo).

Over short vegetation, the seasonal picture is much more pronounced than in the NEU forest network, in which evergreen forests were dominant. Strong seasonal cycles in LAI in SN, G and C land uses, as well as in solar radiation, drive the annual variations in G_s , with logically annual maxima in summer (Fig. 2). The IDEM model predicts much larger (a factor 2 to 4) summer daytime stomatal conductances, typically 15–30 mm s⁻¹, than the other models with typically 5–10 mm s⁻¹. By contrast the EMEP-03 model yields the smallest summer G_s values, particularly in crops in spring and summer, owing in part to a rather short predicted growing season, typically 100 days, outside of which the soil is assumed to be bare (LAI = 0, G_s = 0). The four models are otherwise roughly consistent during the rest of the year, with residual stomatal exchange in spring and autumn and a near zero G_s in winter.

3.1.3 Trace gas and aerosol deposition velocities

Deposition velocities were calculated for the height of the DELTA system inlets in order to infer exchange fluxes directly from DELTA concentrations (Eq. 1), but since sampling- and canopy- heights varied between sites, and for comparability's sake, we present in this section mean V_d data evaluated at a standard height of 3 m above $d + z_0$ for all F, SN, G and C ecosystems (Fig. 3). With the exception of NO₂, for the N_r species considered here, V_d was substantially larger over forests than over short vegetation, regardless of the model, due to the reduced aerodynamic resistance R_a for rougher forest surfaces (over the same vertical path of 3 m). For NO₂ this had no noticeable effect on V_d , as R_c made up the bulk of the total resistance to dry deposition, with uptake being largely limited to the stomatal pathway in the four models.

For HNO₃ over short vegetation, the mean V_d was of the order of 10–12 mm s⁻¹ and very similar between models since the non-stomatal resistance was generally considered to be small, though not necessarily negligible (CDRY, IDEM), and V_d could be approximated to $1/(R_a + R_b)$ as the sum of atmospheric resistances was much larger

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



than R_c . The spread in mean V_d values for each vegetation type (F, SN, G, C), as shown by the range of mean site V_d values from the 5-th to the 95-th percentile in Fig. 3, thus reflected the range of mean windspeeds measured at the different sites, so that mean V_d could exceed 15 mm s^{-1} at the windier sites. Over forests, by contrast, mean V_d for HNO_3 was typically $25\text{--}35 \text{ mm s}^{-1}$, with the mean R_a at 3 m above $d + z_0$ being of the order of a few s m^{-1} . Here, the differences between models in R_c for HNO_3 (Sect. 2.1.1) became significant, so that the mean V_d across sites in CDRY and IDEM was substantially smaller ($\sim 25 \text{ mm s}^{-1}$) than in CBED and EMEP-03 ($> 35 \text{ mm s}^{-1}$).

The most significant absolute inter-model differences, however, were found for NH_3 for all vegetation types, and for aerosol NH_4^+ and NO_3^- over forests (Fig. 3). For unfertilised vegetation (F, SN), V_d for NH_3 was a factor 2–3 larger in CBED and IDEM than in CDRY and EMEP-03. The CDRY scheme systematically predicted the smallest V_d of the four models for NH_3 due to a generally much larger non-stomatal resistance, which was taken to be equal to that for SO_2 . There was a relatively small spread of mean site V_d values in CBED for F and SN, compared with EMEP-03 and IDEM, as the CBED scheme used a constant R_c of 20 s m^{-1} for unfertilised vegetation, while in the other models variations in R_c were controlled by RH, T and sometimes by the ratio of SO_2 to NH_3 ambient concentrations. Remarkably, however, the mean V_d for NH_3 across sites was almost identical in CBED and IDEM for F and SN. There are very few long-term micrometeorological NH_3 flux datasets over (European) forests, from which comprehensive and robust parameterisations may be derived, with the bulk of NH_3 flux measurements stemming from mainly coniferous stands in the high N environment of The Netherlands (Wyers and Erisman, 1998), Belgium (Neiryck et al., 2005, 2007; Neiryck and Ceulemans, 2008) and Denmark (Andersen et al., 1999) (see also Zhang et al., 2010 and Massad, 2010, for reviews), and this is clearly reflected in the wide range of deposition velocities provided by the different models.

Over fertilised systems (G, C), no V_d is provided for NH_3 in CBED since this uses a compensation point approach, but for the other three models the same hierarchy in V_d estimates is found as for F and SN, with IDEM providing the largest values, about

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

10 mm s⁻¹) and CDRY the smallest, around 5 mm s⁻¹ (Fig. 3).

Aerosol N_r deposition velocities were predicted to be very small for short vegetation, typically 2–3 mm s⁻¹, with little variation between models. All models consistently showed slightly larger V_d for NO₃⁻ than for NH₄⁺, reflecting the larger fraction of NO₃⁻ found in the coarse aerosol mode compared with NH₄⁺. By contrast to short vegetation, V_d estimates over forests varied widely different between the four routines, with theoretical (Slinn-type) models (CDRY, EMEP-03) providing similar estimates of the order of 2–5 mm s⁻¹, and the more empirical, measurement-based or simplified models (CBED, IDEM) yielding much larger estimates (typically 10–25 mm s⁻¹). Publications from the last 10 years have also demonstrated that, over forests, deposition velocities for particles in the size range 0.1–1 μm, which contain most of the atmospheric N_r, are much larger than would be expected on the basis of theory, with values of typically 10 mm s⁻¹ (Gallagher et al., 1997) or even 50–100 mm s⁻¹ (Wolff et al., 2009). Gallagher et al. (2002) further showed from a compilation of published V_d data for small aerosols (0.1–0.2 μm) that V_d was strongly dependent on the roughness of vegetation and that measured V_d was typically a factor 10 larger than Slinn-type models, not only for forests but across the range of z₀ values from the various datasets over heathland, grassland and arable land. However, it should also be noted that many of the larger deposition velocities (e.g. Gallagher et al., 1997) have been measured over Speulder forest (NL-Spe in Table 1), which is a Douglas fir forests with a projected LAI >10; this canopy is far denser than the typical Scots pine or Norway spruce canopies (LAI ~3–5), and hence large V_d would be expected (Petroff et al., 2008a, b). Further, apparent emission fluxes are common in flux datasets, and there are significant difficulties in interpreting how far such data are real or represent artifacts (Pryor et al., 2007, 2008a,b). Emerging evidence from chemically resolved particle flux measurements suggests that the volatilisation of NH₄NO₃ during deposition may increase effective deposition rates of these compounds and that effective deposition rates for NO₃⁻ may therefore be significantly larger than for thermodynamically stable SO₄²⁻ (Fowler et al., 2009). Such large model/measurement discrepancies, as well as the large differences between models

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(Fig. 3), hint at much uncertainty regarding aerosol V_d , especially to forests, where the large roughness may potentially mean much larger aerosol dry deposition than assumed heretofore.

3.2 Dry deposition of N_r to European ecosystems

5 Modelled annual dry deposition fluxes of atmospheric N_r are summarised in Table 3 and Fig. 4. We approximate N_r dry deposition to the sum of the dominant inorganic species, i.e. gas NH_3 , HNO_3 and NO_2 and aerosol NH_4^+ and NO_3^- fluxes, as no data were available for organic N_r . As expected from the model inter-comparison for V_d (Fig. 3), the annual N_r dry deposition estimates are very model dependent, with variations
10 between the largest and smallest estimates at any given site reaching typically a factor 2 to 3 (Fig. 4). There was nonetheless a strong correlation across the sites between models, which was logically driven by the measured atmospheric concentrations and meteorology.

Note that the results discussed hereafter were obtained from model base runs as
15 outlined earlier (Sect. 2.3.1), and detailed in Sections A2-A5 and Table A2 of the Supplement. Alternative runs shown therein (Fig. A2) demonstrate that the choice of measured or model default LAI and h_c as inputs to the models has a significant impact on annual fluxes, generally of the order of ± 10 to 20% of the base run flux, but sometimes reaching $\pm 50\%$. Likewise, the use of temperature and relative humidity data
20 estimated at canopy level ($d + z_0'$) where exchange processes take place, rather than data in the ambient air above the canopy (base run), has a very large impact on NH_3 emissions by stomata of grass and crops in the CBED model (Fig. A2 of Supplement). A full sensitivity analysis of the models is beyond the scope of this paper, but these results show that models have different sensitivities to input data and that the various
25 land use classes respond differently.

Over F and SN ecosystems, the largest N_r dry deposition estimates were consistently given by CBED and IDEM, which were largely in agreement, while the EMEP-03 and CDRY fluxes were typically a factor of 2 smaller. The largest annual N_r dry

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



deposition to forests was derived for The Netherlands (NL-Loo, NL-Spe) and Belgium (BE-Bra), while remote boreal forests (FI-Hyy, FI-Sod, SE-Nor) received the smallest inputs. Similar differences occurred in SN ecosystems, with less than $1 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ at FI-Lom compared with about $15\text{--}25 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ at NL-Hor. Dry deposition of N_r to short semi-natural vegetation was dominated by NH_3 , except in CDRY, contributing typically 50–75% of total dry deposition inputs, depending on the model (Fig. 5). Despite similar concentrations overall (Table 2), the relative contribution of NH_3 was less over F than over SN, typically only 30–40%, either because aerosol deposition rates were larger, especially in CBED and IDEM (Fig. 3), or because HNO_3 fluxes were large, being of the same order as NH_3 over forests in the CDRY and EMEP-03 models (Fig. 5).

Although the deposition velocity of NO_2 was small compared with that of NH_3 and HNO_3 (Fig. 3), the comparatively large ambient NO_2 concentrations at a few sites (BE-Bra, FI-Hyy, IT-Ren, NL-Spe, CH-Oe1, FR-Gri) resulted in NO_2 contributing a large – and sometimes dominant – fraction of total N_r dry deposition at some sites (Fig. 5), especially with CDRY. In a scoping study of 14 short-term inferential campaigns over 8 CAPMoN sites in Eastern and Central Canada, Zhang et al. (2009) estimated that the combined dry deposition of NO_2 , PAN and other NO_y species contributed between 4% and 18% of total (dry + wet) N_r deposition. Most sites of the NEU network, however, were located in remote or rural landscapes, and although NO_2 concentrations were not measured everywhere, it may be assumed that NO_2 generally contributed less than 10–15% of dry N_r deposition, as observed at e.g. IT-Col, FI-Lom, UK-AMo, HU-Bug (Fig. 5). The estimated NO_2 contribution was especially small, and often even nought, with the EMEP-03 routine due to the implementation of the 4 ppb threshold (Sect. 2.1.1). HONO was generally not detectable except at roadside (e.g. CH-Oe1) and suburban sites (FR-Gri, FR-Fon), but concentrations were very small and may partly have resulted from a sampling artefact, and HONO deposition is neglected here, also given that inferential modelling of HONO is very uncertain due to the possibility of heterogeneous production at surfaces.

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

10–15 mm s⁻¹ over coniferous forest in daytime, equivalent to a canopy resistance of the order of 100 s m⁻¹ (Turnipseed et al., 2006; Wolfe et al., 2009), suggesting that PAN deposition to forests may be much faster than predicted by current algorithms (e.g. Zhang et al., 2009). With typical PAN concentrations of 0.1–1 ppb, Turnipseed et al. (2006) calculated that PAN contributed about 20% of daytime NO_y (NO + NO₂ + HNO₃ + NO₃⁻ + PAN) dry deposition at their forest site. However, considering the strong control of PAN deposition by stomatal opening and uptake (Doskey et al., 2004), and the consequently reduced V_d at night and in winter, the contribution of PAN and other atmospheric organic nitrates to total N_r inputs must be minor on the annual time scale.

3.3 Comparison with flux monitoring datasets

The surface/atmosphere exchange of reactive nitrogen has been investigated and measured at numerous sites in Europe and elsewhere, yet this has been done most often campaign-wise, with measurements lasting typically a few days to a few weeks. The data thus obtained are invaluable for understanding exchange processes and developing parameterisations for atmospheric models, but they typically cover only a limited range of meteorological conditions, atmospheric concentrations and vegetation development stages. The validation of inferential models at the ecosystem scale benefits much from comparisons with long-term flux measurement datasets, as the wide range of environmental conditions covered is useful for highlighting deficiencies in process understanding and for comparing scaled-up, annual estimates with actual, measured dry deposition. Such long-term flux datasets are rare in the case of NH₃ and NO_x, and almost non-existent for HNO₃ and aerosol NH₄⁺ and NO₃⁻.

Many forest sites of the NEU network have been monitoring wet-only or bulk deposition and throughfall as part of national or international initiatives (e.g. the ICP-Forests programme of the CLRTAP; <http://www.icp-forests.org/>), which, by difference between above- and below-canopy fluxes, may provide estimates of dry deposition,

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



though uncertainties are large due to canopy interactions (Lovett and Lindberg, 1993; Zimmermann et al., 2006; Neiryndck et al., 2007; Simpson et al., 2006b). Only two forest sites (BE-Bra, NL-Spe) within the NEU network have actually monitored annual NH_3 dry deposition in the past using the flux-gradient technique (Fig. 6). The measurements by Neiryndck et al. (2007) at BE-Bra suggested an annual deposition input of nearly $-20 \text{ kg N ha}^{-1} \text{ yr}^{-1}$, which is larger than the output of any of the four models in the present study (Table 3), whose ensemble average is only of the order of $-10 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ (Fig. 6). Only part of the difference may be explained by the larger mean NH_3 concentration ($3.0 \mu\text{g m}^{-3}$) at the time of the flux measurements in 1999–2001 (Neiryndck et al., 2007) than in the NEU DELTA dataset ($2.3 \mu\text{g m}^{-3}$) in 2007–2008. A clear indication that especially CDRY and EMEP-03 both largely under-estimated NH_x ($\text{NH}_3 + \text{NH}_4^+$) dry deposition at BE-Bra, with annual fluxes of the order of -6 to $-8 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ (Table 3), is provided by a comparison with throughfall data. Measured wet deposition of NH_x was $7 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ at BE-Bra, which together with dry deposition from CDRY or EMEP-03, would total around $15 \text{ kg NH}_x\text{-N ha}^{-1} \text{ yr}^{-1}$, while the measured throughfall was actually $18 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ over the same time period (J. Neiryndck, personal communication, 2010).

The comparison is more favourable at NL-Spe, where the measured total deposition in 1994–1995 of $-17.9 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ (Erismann et al., 1996) is well in the range of the four model estimates in the NEU dataset and close to the ensemble mean ($-16.9 \text{ kg N ha}^{-1} \text{ yr}^{-1}$) (Fig. 6), with the difference in mean concentrations between the two periods being consistent with the model/measurement difference. A striking element in the comparison of BE-Bra with NL-Spe is the roughly equal measured annual NH_3 dry deposition at the two sites (-19.6 vs. $-17.9 \text{ kg N ha}^{-1} \text{ yr}^{-1}$) while the mean concentration was about 50% larger at NL-Spe, pointing to a much smaller R_c at BE-Bra, since the annual mean u_* was identical (0.51 m s^{-1}) at the two sites. The much smaller mean NH_3/SO_2 molar ratio at BE-Bra (2.9) than at NL-Spe (11.1) has been held responsible for the difference in measured R_c for NH_3 (Neiryndck et al., 2005), but the effect of leaf surface chemistry on deposition rates is not adequately reflected in most

dry deposition models. Flux measurements at BE-Bra during 1999–2001 showed a reduced R_c for NH_3 and larger R_c for SO_2 during winter when the NH_3/SO_2 molar ratio was below 1; in summer this ratio was larger than 3 and R_c for SO_2 was correspondingly smaller, while R_c for NH_3 was increased (J. Neyrinck, personal communication, 2010). Because in Europe the total acid concentration is not necessarily dominated by SO_2 , the molar ratio of NH_3 to the sum of the main atmospheric strong acids (SO_2 , HNO_3 , HCl) is actually a better proxy for linking surface resistance to the pollution climate (Flechard et al., 1999); this ratio was almost a factor of 3 smaller at BE-Bra (1.6) than at NL-Spe (4.5), with BE-Bra being the second most acidic site of the NEU network, after CZ-BK1.

At the only NEU semi-natural site with a long-term NH_3 flux dataset (UK-AMo) (Flechard, 1998), measured annual dry deposition in 1995 ($-2.5 \text{ kg N ha}^{-1} \text{ yr}^{-1}$) is compatible with the range of model estimates in NEU for the 2007–2008 reference period and within 10% of the models ensemble mean ($-2.2 \text{ kg N ha}^{-1} \text{ yr}^{-1}$). For agricultural systems in the NEU network, comparisons can only be made at the managed grasslands CH-Oe1 and UK-EBu. For these fertilised, cut and/or grazed systems, a comparison of measurements with inferential models is only meaningful in conditions of background NH_3 exchange, i.e. discarding measured NH_3 emission fluxes that follow the application of manure, slurry or mineral fertilisers, as these processes are not currently considered nor implemented in inferential routines. At CH-Oe1, the overall net measured NH_3 budget was $+17 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ and driven by a gross annual NH_3 emission by applied cattle slurry of $+20 \text{ kg N ha}^{-1} \text{ yr}^{-1}$, but during most of the year background exchange amounted to a deposition of $-3 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ (Flechard et al., 2010), which is in the range of model predictions within NEU of -3.2 to $+0.1 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ (Table 3). Equally, at UK-EBu, the overall annual measured NH_3 flux was a net emission of $+1.9 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ but, discarding the gross NH_3 emissions of $+4.2 \text{ kg N ha}^{-1} \text{ yr}^{-1}$ mostly due to mineral fertiliser and urea applications (Milford et al., 2004), one may calculate a background annual dry deposition of $-2.3 \text{ kg N ha}^{-1} \text{ yr}^{-1}$, also within the range of the four model estimates based on NEU 2007–2008 data (-2.6 to

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

$-0.2 \text{ kg N ha}^{-1} \text{ yr}^{-1}$, Table 3).

3.4 Reducing uncertainties in N_r dry deposition

The uncertainty of modelled N_r dry deposition at the regional scale results from the combined uncertainties in concentrations of N_r species and in their respective deposition (or exchange) velocities. Establishing a monitoring network for NH_3 , HNO_3 , HONO and aerosol NH_4^+ and NO_3^- concentrations at the continental scale in Europe has been a significant step forward, even if the basic setup did not include NO_x and other N_r species except at a few more intensive measurement stations. Continent-scale networks of a similar size, e.g. EMEP (EMEP, 2009; Torseth et al., 2001), CASTNet (Sickles and Shadwick, 2007; Baumgardner et al., 2002) and CAPMoN (Zhang et al., 2009), have long placed the emphasis on acidifying gases (SO_2 , HNO_3 , NO_x) deposition and aerosol-phase N_r (NH_4^+ and NO_3^-), but have not included the gas/particle partitioning of NH_x . This has been measured at selected sites as part of research projects (Erisman et al., 1996; Zimmermann et al., 2006; Neiryneck et al., 2007) and has been used to evaluate the output of regional atmospheric models at selected sites (Zhang et al., 2009), but data on speciated NH_3 and NH_4^+ concentrations at regional scales have been sparse and irregular outside of a few national initiatives (Bleeker et al., 2009). High time-resolution measurements with aerosol mass spectrometer measurements are also becoming available (Laj et al., 2009), although one limitation is that to date only (ultra-) fine particles can be captured: coarse nitrate is not typically measured at the same sites. The monitoring data gathered as part of NEU allow a large-scale investigation of the relative contributions of NH_3 and NH_4^+ as well as HNO_3 and NO_3^- to total dry deposition, despite the large uncertainties and discrepancies associated with inferential models, and they also provide important ground validation data for CTMs.

The differences in deposition velocities between models (Fig. 3) results from both the natural variability in surface resistances found in existing N_r flux datasets, leading to different parameterisations, and from the rarity and complexities of flux datasets. The

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



physical, biological and chemical exchange mechanisms involved are too complex to model explicitly and completely from first principles, so that parameterisations tend to be empirical but dependent on few datasets, without the confidence that the statistics of large or robust numbers afford. The recent efforts by Zhang et al. (2010) and Massad et al. (2010) to bring together the existing NH_3 flux and compensation point datasets into coherent and comprehensive exchange schemes for the main ecosystem types point in the right direction. Significant gaps in knowledge remain, especially with respect to surface chemistry, canopy cycling, soil/litter/vegetation interactions, management practices for agricultural systems, which will not be bridged without a more extensive coverage of NH_3 fluxes. Within the NitroEurope IP, intensive N_r flux measurements to improve process understanding at a few core sites of the network have been complemented at other sites by low-cost methods for N_r concentrations (DELTA) and also for fluxes (COTAG, or COnditional Time-Averaged Gradient; Famulari et al., 2010); this could serve as a blueprint for a future European N_r monitoring and modelling strategy.

4 Conclusions

Inferential modelling with four dry deposition routines was applied to estimate annual N_r fluxes at the ecosystem scale across the NitroEurope inferential network. Differences between models were reviewed in terms of canopy characteristics for the main land use types, of derived friction velocity, of stomatal conductance, and of deposition velocities and exchange rates for five dominant inorganic N_r chemical species in the atmosphere (NH_3 , HNO_3 , NO_2 , and aerosol NH_4^+ and NO_3^-). Differences in stomatal conductances between models are large, but this is only decisive for NO_2 , which is assumed to be mainly deposited through stomata. However, these models are also routinely used for other pollutant gases such as SO_2 and O_3 , for which the stomatal share in the total deposition is also large (see Fowler et al., 2009, and references therein). For water-soluble gases such as NH_3 and HNO_3 , parameterisations of non-stomatal resistances are the main sources of inter-model discrepancies in deposition velocities, which can

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



reach a factor of 3 between models for NH_3 . For aerosol N_r deposition to forests, empirical and measurement-oriented parameterisations predict deposition rates that are a factor 5–10 larger than theoretical models. As a result, both the total modelled N_r fluxes and the shares of individual N_r species in the overall N_r dry deposition are extremely model-dependent. The few NH_3 flux datasets available for comparison within this study were within the range of models and broadly comparable with the ensemble average, but model validation generally suffers from a serious lack of long-term N_r flux monitoring data over different vegetation types.

Inferential modelling was originally based on the concept of uni-directional exchange (deposition from the atmosphere), and has traditionally viewed vegetation elements and soil more or less as physical receptors with a given surface roughness, chemical sink strength and aerosol capture efficiency, with little regard to underlying biological and biochemical processes. The discipline is currently undergoing a paradigm shift, recognising the need to increasingly couple ecosystem modelling, including soil/litter/vegetation cycling, as well as crop/grass management and fertilisation, to surface/atmosphere bi-directional exchange frameworks, especially with respect to NH_3 and NO_x . Here, compensation points need to be made dependent on the N status of the ecosystem, whether fertilised or unfertilised, characterising emission potentials that interact with advected air masses. Major developments are also needed to better deal with in-canopy air chemistry and phase partitioning that affect the net exchange of NH_3 and HNO_3 versus NH_4NO_3 aerosol. Similarly, the roles of O_3 deposition and emission of biogenic volatile organic compounds on net NO_x fluxes in ecosystems need to be better understood. Although not considered in this study, uncertainties in wet deposition estimates add to the total uncertainty in the N_r deposition predicted by CTMs.

Supplementary material related to this article is available online at:
[http://www.atmos-chem-phys-discuss.net/10/29291/2010/
acpd-10-29291-2010-supplement.pdf](http://www.atmos-chem-phys-discuss.net/10/29291/2010/acpd-10-29291-2010-supplement.pdf).

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Acknowledgements. The authors gratefully acknowledge financial support by the European Commission (NitroEurope-IP, project 017841). Many thanks are due to all environmental chemists and scientists involved in running the DELTA network, not least N. van Dijk, I. Simons (CEH), U. Dämmgen, J. Conrad (FAL/vTI), V. Djuricic, S. Vidic (MHSC), M. Mitosinkova (SHMU), M.J. Sanz, P. Sanz, J.V. Chorda (CEAM), M. Ferm (IVL), Y. Fauvel (INRA), T.H. Ungerud and the late J. E. Hanssen (NILU). The assistance of A. Freibauer and D. Papale in establishing this successful partnership with CarboEurope-IP and facilitating access to micrometeorological data was much appreciated, and we also thank all principal investigators, managers and operators of CarboEurope-IP sites for their individual contributions, including R. Ceulemans, J. Neiryneck, M. Aubinet, C. Moureaux, N. Buchmann, W. Eugster, S. Etzold, C. Ammann, A. Neftel, C. Spirig, M. Jocher, M.V. Marek, R. Czerny, W. Kutsch, E.D. Schultze, O. Kolle, C. Bernhofer, C. Rebmann, A. Don, C. Don, S. Müller, K. Butterbach-Bahl, N. Brüggemann, L. Lotte Soerensen, K. Pilegaard, A. Ibrom, K. Larsen, P. Sørensen, H. Soegaard, A. Carrara, M.T. Sebastià, T. Vesala, T. Laurila, A. Lohila, E. Dufrière, J.Y. Pontailier, P. Cellier, B. Loubet, S. Masson, A. Granier, B. Longdoz, P. Gross, D. Loustau, C. Lambrot, J.F. Soussana, K. Klumpp, O. Darsonville, S. Rambal, J.M. Ourcival, Z. Tuba, L. Horvath, M. Jones, M. Saunders, F. Albanito, B. Roth, G. Kiely, P. Leahy, N. Foley, R. Valentini, G. Matteucci, P. Stefani, D. Gianelle, A. Cescatti, V. Magliulo, T. Bertolini, G. Seufert, G. Manca, A. Meijide Orive, S. Minerbi, L. Montagnani, W. Baumgartner, C. Valtingoier, E. Moors, W.C.M. van den Bulk, A. Hensen, A. Frumau, A. J. Dolman, D. Hendriks, J. Olejnik, B.H. Chojnick, G.B.A. Pita, J.S. Pereira, R. Lobo do Vale, J. Banza, A. Varlagin, A. Lindroth, M. Mölder, P. Vestin, J. Moncrieff, R. Clement, W. Medinets and S. Medinets. The meteorological data of the state forest De Inslag (Brasschaat, Belgium) have been processed from data kindly provided by the Research Institute for Nature and Forest (INBO, Belgium).

References

- Andersen, H. V., Hovmand, M., Hummelshøj, P., and Jensen, N. O.: Measurements of ammonia concentrations, fluxes and dry deposition velocities to a spruce forest 1991–1995, *Atmos. Environ.*, 33, 1367–1383, 1999.
- Baldocchi, D. D., Hicks, B. B., and Camara, P.: A canopy stomatal resistance model for gaseous deposition to vegetated surfaces, *Atmos. Environ.*, 21, 91–101, 1987.

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Bash, J. O., Walker, J. T., Katul, G. G., Jones, M. R., Nemitz, E., and Robarge, W. P.: Estimation of In-Canopy Ammonia Sources and Sinks in a Fertilized *Zea mays* Field, *Environ. Sci. Technol.*, 44, 1683–1689, 2010.

Bates, R. G. and Pinching, G. D.: Dissociation constant of aqueous ammonia at 0 to 50 °C from E.m.f. studies of the ammonium salt of a weak acid, *Am. Chem. J.*, 72, 1393–1396, 1950.

Baumgardner Jr., R. E., Lavery, T. F., Rogers, C. M., and Isil, S. S.: Estimates of the Atmospheric Deposition of Sulfur and Nitrogen Species: Clean Air Status and Trends Network, 1990–2000, *Environ. Sci. Technol.*, 36, 2614–2629, 2002.

Bleeker, A., Reinds, G. J., Vermeulen, A. T., de Vries, W. and Erisman, J. W.: Critical loads and resent deposition thresholds of nitrogen and acidity and their exceedances at the level II and level I monitoring plots in Europe, ECN report ECN-C–04-117, Petten, The Netherlands, December 2004.

Bleeker, A., Sutton, M. A., Acherman, B., Alebic-Juretic, A., Aneja, V. P., Ellermann, T., Erisman, J. W., Fowler, D., Fagerli, H., Gauger, T., Harlen, K. S., Hole, L. R., Horvath, L., Mitosinkova, M., Smith, R. I., Tang, Y. S. and van Pul, A.: Linking Ammonia Emission Trends to Measured Concentrations and Deposition of Reduced Nitrogen at different Scales, in: *Atmospheric Ammonia, Detecting emission changes and environmental impacts*, edited by: Sutton, M. A., Reis, S., and Baker, S. M. H., Springer, 123–180, 2009.

Burkhardt, J., Flechard, C. R., Gressens, F., Mattsson, M. E., Jongejan, P. A. C., Erisman, J. W., Weidinger, T., Meszaros, R., Nemitz, E., and Sutton, M. A.: Modelling the dynamic chemical interactions of atmospheric ammonia and other trace gases with measured leaf surface wetness in a managed grassland canopy, *Biogeosciences* 6, 67–84, 2009

Dasgupta, P. K. and Dong, S.: Solubility of ammonia in liquid water and generation of trace levels of standard gaseous ammonia, *Atmos. Environ.*, 20, 565–570, 1986.

de Vries, W., Solberg, S., Dobbertin, M., Sterba, H., Laubhahn, D., Reinds, G. J., Nabuurs, G. J., Gundersen, P. and Sutton, M.A.: Ecologically implausible carbon response, *Nature*, 451, E1–E3, doi:10.1038/nature06579, 2008.

Dolman, A. J., Valentini, R., and Freibauer, A. (Eds.): *The Continental-Scale Greenhouse Gas Balance of Europe*, Springer Ecological Series 203, 390 pp., ISBN:978-0-387-76568-6, Springer, New York, USA, 2008.

Doskey, P. V., Rao Kotamarthi, V., Fukui, Y., Cook, D. R., Breitbeil, F. W. and Wesely, M. L.: Air-surface exchange of peroxyacetyl nitrate at a grassland site, *J. Geophys. Res.*, 109, D10310, doi:10.1029/2004JD004533, 2004.

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Emberson, L., Ashmore, M., Simpson, D., Tuovinen, J.-P., and Cambridge, H.: Modelling and mapping ozone deposition in Europe, *Water Air Soil Poll.*, 130, 577–582, 2001.
- EMEP (European Monitoring and Evaluation Programme): Transboundary Acidification, Eutrophication and Ground Level Ozone in Europe in 2007, EMEP Report 1/2009, http://www.emep.int/publ/reports/2009/status_report_1_2009.pdf, Norwegian Meteorological Institute, 2009.
- Erismann, J. W. and Wyers, G. P.: Continuous measurements of surface exchange of SO₂ and NH₃; implications for their possible interaction in the deposition process, *Atmos. Environ.*, 27A, 1937–1949, 1993.
- Erismann, J. W., van Pul, A., and Wyers, P.: Parametrization of surface resistance for the quantification of atmospheric deposition of acidifying pollutants and ozone, *Atmos. Environ.*, 28, 2595–2607, 1994.
- Erismann, J. W., Mennen, M. G., Fowler, D., Flechard, C. R., Spindler, G., Grüner, A., Duyzer, J. H., Ruigrok, W., and Wyers, G. P.: Towards development of a deposition monitoring network for air pollution in Europe, Report n 722108015, RIVM, The Netherlands, <http://rivm.openrepository.com/rivm/bitstream/10029/10432/1/722108015.pdf>, 1996.
- Erismann, J. W., Vermeulen, A., Hensen, A., Flechard, C., Dammggen, U., Fowler, D., Sutton, M., Grunhage, L., and Tuovinen, J.P.: Monitoring and modelling of biosphere/atmosphere exchange of gases and aerosols in Europe, *Environ. Pollut.*, 133, 403–413, 2005.
- Erismann, J. W., Bleeker, A., Galloway, J., and Sutton, M. A.: Reduced nitrogen in ecology and the environment, *Environ. Pollut.*, 150, 140–149, 2007.
- Fagerli, H. and Aas, W.: Trends of nitrogen in air and precipitation: Model results and observations at EMEP sites in Europe, 1980–2003, *Environ. Pollut.*, 154, 448–461, 2008.
- Famulari, D., Fowler, D., Nemitz, E., Hargreaves, K.J., Storeton-West, R. L., Rutherford, G., Tang, Y. S., Sutton, M. A. and Weston, K. J.: Development of a low-cost system for measuring conditional time-averaged gradients of SO₂ and NH₃, *Environ. Monit. Assess.*, 161, 11–27, doi:10.1007/s10661-008-0723-6, 2010.
- Farquhar, G. D., Firth, P. M., Wetselaar, R., and Weir, B.: On the gaseous exchange of ammonia between leaves and the environment: determination of the ammonia compensation point, *Plant. Physiol.*, 66, 710–714, 1980.
- Flechard, C. R.: Turbulent exchange of ammonia above vegetation, Nottingham University, 231 pp., 1998.
- Flechard, C. R. and Fowler, D.: Atmospheric ammonia at a moorland site. II: Long term sur-

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



face/atmosphere micrometeorological flux measurements, Q. J. Roy. Meteor. Soc., 124, 759–791, 1998.

Flechard, C. R., Fowler, D., Sutton, M. A., and Cape, J. N.: A dynamic chemical model of bi-directional ammonia exchange between semi-natural vegetation and the atmosphere, Q. J. Roy. Meteor. Soc., 125, 2611–2641, 1999.

Flechard, C. R., Spirig, C., Neftel, A., and Ammann, C.: The annual ammonia budget of fertilised cut grassland – Part 2: Seasonal variations and compensation point modelling, Biogeosciences, 7, 537–556, 2010, <http://www.biogeosciences.net/7/537/2010/>.

Fowler, D., Flechard, C., Cape, J. N., Storeton-West, R. L., and Coyle, M.: Measurements of ozone deposition to vegetation quantifying the flux, the stomatal and non-stomatal components, Water Air Soil. Poll., 130, 63–74, 2001.

Fowler, D., Pilegaard, K., Sutton, M.A., Ambus, P., Raivonen, M., Duyzer, J., Simpson, D. and 50 others: Atmospheric composition change: Ecosystems–Atmosphere interactions, Atmos. Environ., 43, 5193–5267, 2009.

Gallagher, M. W., Beswick, K. M., Duyzer, J., Westrate, H., Choularton, T. W., and Hummelshøj, P.: Measurements of aerosol fluxes to Speulder forest using a micrometeorological technique, Atmos. Environ., 31, 359–373, 1997.

Gallagher, M. W., Nemitz, E., Dorsey, J. R., Fowler, D., Sutton, M. A., Flynn, M. and Duyzer, J. H.: Measurements and parameterisations of small aerosol deposition velocities to grassland, arable crops, and forests: Influence of surface roughness length on deposition, J. Geophys. Res., 107(D12), doi:10.1029/2001JD000817, 2002.

Galloway, J. N., Aber, J. D., Erisman, J. W., Seitzinger, S. P., Howarth, R. W., Cowling, E. B., and Cosby, B. J.: The Nitrogen Cascade, BioScience, 53, 4, 341–356, 2003.

Garland, J. A.: The dry deposition of sulphur dioxide to land and water surfaces, P. R. Soc. London, A354, 245–268, 1977.

Genermont, S. and Cellier, P.: A mechanistic model for estimating ammonia volatilization from slurry applied to bare soil, Agr. Forest Meteorol., 88, 145–167, 1997.

González Benítez, J. M., Cape, J. N. and Heal, M. R.: Gaseous and particulate water-soluble organic and inorganic nitrogen in rural air in southern Scotland, Atmos. Environ., 44, 1506–1514, 2010.

Hicks, B. B.: Dry deposition to forests – On the use of data from clearings, Agr. Forest Meteorol., 136, 214–221, 2006.

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Jarvis, P. G.: The interpretation of the variations in leaf water potential and stomatal conductance found in canopies in the field, *Philos. T. Roy. Soc.*, B273, 593–610, 1976.
- Joutsenoja, T.: Measurements of aerosol deposition to a cereal crop, in: *Measurements and Modelling of Gases and Aerosols to Complex Terrain*, NERC Report GR3/7259, edited by: Choularton, T. W., *Nat. Environ. Res. Counc.*, UK, 1992.
- Laj, P., Klausen, J., Bilde, M., Pla-Duelmer, C., Pappalardo, G., Clerbaux, C., Baltensperger, U., Hjorth, J., Simpson, D., Reimann, S., Coheur, P.-F., Richter, A., De Mazière, M., Rudich, Y., McFiggans, G., Torseth, K., Wiedensohler, A., Morin, S., Schulz, M., Allan, J.D., Attié, J.-L., Barnes, I., Birmili, W., Cammas, J. P., Dommen, J., Dorn, H.-P., Fowler, D., Fuzzi, S., Glasius, M., Granier, C., Hermann, M., Isaksen, I.S.A., Kinne, S., Koren, I., Madonna, F., Maione, M., Massling, A., Moehler, O., Mona, L., Monks, P.S., Müller, D., Müller, T., Orphal, J., Peuch, V.-H., Stratmann, F., Tanré, D., Tyndall, G., Abo Riziq, A., Van Roozendaal, M., Villani, P., Wehner, B., Wex, H., and Zardini, A. A.: Measuring atmospheric composition change, *Atmos. Environ.*, 43, 5351–5414, 2009.
- Lovett, G. M. and Lindberg, S. E.: Atmospheric deposition and canopy interactions of nitrogen in forests, *Can. J. Forest Res.*, 23, 1603–1616, 1993.
- Magnani, F., Mencuccini, M., Borghetti, M., Berbigier, P., Berninger, F., Delzon, S., Grelle, A., Hari, P., Jarvis, P.G., Kolari, P., Kowalski, A. S., Lankreijer, H., Law, B. E., Lindroth, A., Loustau, D., Manca, G., Moncrieff, J. B., Rayment, M., Tedeschi, V., Valentini, R., and Grace, J.: The human footprint in the carbon cycle of temperate and boreal forests, *Nature*, 447, 848–851, 2007.
- Massad, R. S., Loubet, B., Tuzet, A., and Cellier, P.: Relationship between ammonia stomatal compensation point and nitrogen metabolism in arable crops: Current status of knowledge and potential modelling approaches, *Environ. Pollut.*, 154, 390–403, 2008.
- Massad, R.-S., Nemitz, E., and Sutton, M. A.: Review and parameterisation of bi-directional ammonia exchange between vegetation and the atmosphere, *Atmos. Chem. Phys. Discuss.*, 10, 10335–10398, doi:10.5194/acpd-10-10335-2010, 2010.
- Meyers, T. P., Finkelstein, P. L., Clarke, J., Ellestad, T. G. and Sims, P.: A multilayer model for inferring dry deposition using standard meteorological measurements, *J. Geophys. Res.*, 103, 22, 645–661, 1998.
- Milford, C.: Dynamics of atmospheric ammonia exchange with intensively-managed grassland, PhD Thesis, University of Edinburgh, 230 pp., 2004.
- Monteith, J. L. and Unsworth, M. H.: *Principles of Environmental Physics*, 2nd edn., Edward

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Arnold, London, 291 pp., 1990.

Nemitz, E., Sutton, M. A., Schjoerring, J. K., Husted, S., and Wyers, G. P.: Resistance modelling of ammonia exchange over oilseed rape, *Agr. Forest Meteorol.*, 105, 405–425, 2000a.

Nemitz, E., Sutton, M. A., Gut, A., San Jose, R., Husted, S., and Schjørring, J. K.: Sources and sinks of ammonia within an oilseed rape canopy, *Agr. Forest Meteorol.*, 105, 385–404, 2000b.

Nemitz, E., Milford, C., and Sutton, M. A.: A two-layer canopy compensation point model for describing bi-directional biosphere-atmosphere exchange of ammonia, *Q. J. Roy. Meteor. Soc.*, 127, 815–833, 2001.

Nemitz, E., Gallagher, M. W., Duyzer, J. H. and Fowler, D.: Micrometeorological measurements of particle deposition velocities to moorland vegetation, *Q. J. Roy. Meteor. Soc.*, 128A, 2281–2300, 2002.

Nemitz, E., Loubet, B., Lehmann, B. E., Cellier, P., Neftel, A., Jones, S. K., Hensen, A., Ihly, B., Tarakanov, S. V. and Sutton, M. A.: Turbulence characteristics in grassland canopies and implications for tracer transport, *Biogeosciences*, 6, 1519–1537, 2009, <http://www.biogeosciences.net/6/1519/2009/>.

Neirynek, J. and Ceulemans, R.: Bidirectional ammonia exchange above a mixed coniferous forest, *Environ. Pollut.*, 154, 424–438, 2008.

Neirynek, J., Kowalski, A. S., Carrara, A., and Ceulemans, R.: Driving forces for ammonia fluxes over mixedforest subjected to high deposition loads, *Atmos. Environ.*, 39, 5013–5024, 2005.

Neirynek, J., Kowalski, A. S., Carrara, A., Genouw, G., Berghmans, P., and Ceulemans, R.: Fluxes of oxidised and reduced nitrogen above a mixed coniferous forest exposed to various nitrogen emission sources, *Environ. Pollut.*, 149, 31–43, 2007.

Personne, E., Loubet, B., Herrmann, B., Mattsson, M., Schjoerring, J. K., Nemitz, E., Sutton, M. A., and Cellier, P.: SURFATM-NH₃ : a model combining the surface energy balance and bi-directional exchanges of ammonia applied at the field scale, *Biogeosciences*, 6, 1371–1388, 2009, <http://www.biogeosciences.net/6/1371/2009/>.

Petroff, A., Mailliat, A., Amielh, M., and Anselmet, F.: Aerosol dry deposition on vegetative canopies. Part I: Review of present knowledge, *Atmos. Environ.*, 42, 3625–3653, 2008a.

Petroff, A., Mailliat, A., Amielh, M., and Anselmet, F.: Aerosol dry deposition on vegetative canopies. Part II: A new modelling approach and applications, *Atmos. Environ.*, 42, 3654–

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

3683, 2008b.

Pryor, S. C., Larsen, S. E., Sørensen, L. L., Barthelmie, R. J., Grönholm, T. and co-authors: Particle fluxes over forests: analyses of flux methods and functional dependencies. *J. Geophys. Res. Atmos.*, 112, D07205, doi:10.1029/2006JD008066, 2007.

5 Pryor, S. C., Gallagher, M., Sievering, H., Larsen, S. E., Barthelmie, R. J., Birsan, F., Nemitz, E., Rinne, J., Kulmala, M., Groenholm, T., Taipale, R., and Vesala, T.: A review of measurement and modelling results of particle atmosphere-surface exchange, *Tellus*, 60, 42–75, 2008a.

Pryor, S. C., Larsen, S. E., Sorensen, L. L., and Barthelmie, R. J.: Particle fluxes above forests: Observations, methodological considerations and method comparisons, *Environ. Poll.*, 152, 667–678, 2008b.

10 Ruijgrok, W., Tieben, H., and Eisinga, P.: The dry deposition of particles to a forest canopy: a comparison of model and experimental results, *Atmos. Environ.*, 31, 399–415, 1997.

Schjøerring, J. K., Husted, S., and Mattsson, M.: Physiological parameters controlling plant-atmosphere ammonia exchange, *Atmos. Environ.*, 32, 491–498, 1998.

15 Seinfeld, J. H. and Pandis, S. N.: Atmospheric chemistry and physics. From Air Pollution to climate Change, 2nd edn., John Wiley and Sons Inc., 2006.

Sickles, J. E. and Shadwick, D. S.: Seasonal and regional air quality and atmospheric deposition in the eastern United States, *J. Geophys. Res.*, 112, D17302, doi:10.1029/2006JD008356, 2007.

20 Simpson, D., Fagerli, H., Jonson, J. E., Tsyro, S., Wind, P., and Tuovinen, J.-P.: Transboundary Acidification, Eutrophication and Ground Level Ozone in Europe. Part I: Unified EMEP Model Description, EMEP Status Report 2003, ISSN 0806–4520, Det Meteorologisk Institutt, Oslo, 2003.

25 Simpson, D., Butterbach-Bahl, K., Fagerli, H., Kesik, M., Skiba, U. and Tang, S.: Deposition and Emissions of Reactive Nitrogen over European Forests: A Modelling Study, *Atmos. Environ.*, 40, 5712–5726, 2006a.

Simpson, D., Fagerli, H., Hellsten, S., Knulst, J., and Westling, O.: Comparison of modelled and monitored deposition fluxes of sulphur and nitrogen to ICP-forest sites in Europe, *Biogeosciences*, 3, 337–355, 2006b,

30 <http://www.biogeosciences.net/3/337/2006/>.

Slinn, W. G. N.: Predictions for particle deposition to vegetative canopies, *Atmos. Environ.*, 16, 1785–1794, 1982.

Smith, R. I., Fowler, D., Sutton, M. A., Flechard, C., and Coyle, M.: Regional estimation of pol-

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



lutant gas deposition in the UK: model description, sensitivity analyses and outputs, *Atmos. Environ.*, 34, 3757–3777, 2000.

Sutton, M. A., Fowler, D., Moncrieff, J. B., and Storeton-West, R. L.: The exchange of atmospheric ammonia with vegetated surfaces. II. Fertilized vegetation, *Q. J. Roy. Meteor. Soc.*, 119, 1047–1070, 1993.

Sutton, M. A., Burkhardt, J.K., Guerin, D., Nemitz, E., and Fowler, D.: Development of resistance models to describe measurements of bi-directional ammonia surface atmosphere exchange, *Atmos. Environ.*, 32(3), 473–480, 1998.

Sutton, M. A., Tang, Y. S., Miners, B., and Fowler, D.: A new diffusion denuder system for long-term, regional monitoring of atmospheric ammonia and ammonium, *Water Air Soil Poll.: Focus*, 1, 145–156, 2001.

Sutton, M. A., Nemitz, E., Erisman, J. W., Beier, C., Butterbach-Bahl, K., Cellier, P., de Vries, W., Cotrufo, F., Skiba, U., Di Marco, C., Jones, S., Laville, P., Soussana, J. F., Loubet, B., Twigg, M., Famulari, D., Whitehead, J., Gallagher, M. W., Neftel, A., Flechard, C. R., Herrmann, B., Calanca, P. L., Schjoerring, J. K., Daemmgen, U., Horvath, L., Tang, Y. S., Emmett, B. A., Tietema, A., Peñuelas, J., Kesik, M., Brüeggemann, N., Pilegaard, K., Vesala, T., Campbell, C. L., Olesen, J. E., Dragosits, U., Theobald, M. R., Levy, P., Mobbs, D. C., Milne, R., Viovy, N., Vuichard, N., Smith, J. U., Smith, P., Bergamaschi, P., Fowler, D., and Reis, S.: Challenges in quantifying biosphere-atmosphere exchange of nitrogen species, *Environ. Pollut.*, 150, 125–139, 2007.

Sutton, M. A., Simpson, D., Levy, P. E., Smith, R. I., Reis, S., van Oijen, M., and de Vries, W.: Uncertainties in the relationship between atmospheric nitrogen deposition and forest carbon sequestration, *Glob. Change Biol.*, 14, 2057–2063, 2008.

Tang, Y. S., Simmons, I., van Dijk, N., Di Marco, C., Nemitz, E., Dämmgen, U., Gilke, K., Djuricic, V., Vidic, S., Gliha, Z., Borovecki, D., Mitosinkova, M., Hanssen, J. E., Uggerud, T.H., Sanz, M.J., Sanz, P., Chorda, J.V., Flechard, C.R., Fauvel, Y., Ferm, M., Perrino, C., Sutton, M. A.: European scale application of atmospheric reactive nitrogen measurements in a low-cost approach to infer dry deposition fluxes, *Agr. Ecosyst. Environ.*, 133, 183–195, 2009.

Tang, Y. S., Dämmgen, U., Conrad, J., Djuricic, V., Vidic, S., Flechard, C. R., Mitosinkova, M., Sanz, M. J., Uggerud, T. H., Borovecki, D., Chorda, J. V., Fauvel, Y., Gilke, K., Gliha, Z., Sanz, P., Simmons, I., van Dijk, N., Ferm, M., Nemitz, E., and Sutton, M. A.: Temporal and spatial variability in reactive inorganic trace gas and aerosol concentrations across Europe,

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

in preparation, *Atmos. Chem. Phys. Discuss.*, 2011.

Torseth, K., Aas, W., and Solberg, S.: Trends in airborne sulphur and nitrogen compounds in Norway during 1985–1996 in relation to air mass origin, *Water Air Soil Poll.*, 130, 1493–1498, 2001.

5 Tuovinen, J.-P., Emberson, L., and Simpson, D.: Modelling ozone fluxes to forests for risk assessment: status and prospects, *Ann. For. Sci.*, 66, 401.1–401.14, 2009.

Turnipseed, A. A., Huey, L. G., Nemitz, E., Stickel, R., Higgs, J., Tanner, D. J., Slusher, D. L., Sparks, J. P., Flocke, F., and Guenther, A.: Eddy covariance fluxes of peroxyacetyl nitrates (PANs) and NO_y to a coniferous forest, *J. Geophys. Res.*, 111, D09304, doi:10.1029/2005JD006631, 2006.

10 UNECE: Protocol to the 1979 Convention on long-range transboundary air pollution to abate acidification, eutrophication and ground-level ozone. United Nations Economic Commission for Europe, Geneva, http://www.unece.org/env/lrtap/multi_h1.htm, 1999.

Wesely, M. L.: Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical models, *Atmos. Environ.*, 23, 1293–1304, 1989.

15 Wesely, M. L. and Hicks, B.B.: Some factors that affect the deposition rates of sulfur dioxide and similar gases on vegetation, *J. Air Poll. Cont. Assoc.*, 27, 1110–1116, 1977.

Wesely, M. L. and Hicks, B. B.: A review of the current status of knowledge on dry deposition, *Atmos. Environ.*, 34, 2261–2282, 2000.

20 Wesely, M. L., Cook, D. R., Hart, R. L., and Speer, R. E.: Measurements and parameterization of particle sulfur deposition over grass, *J. Geophys. Res.*, 90, 2131–2143, 1985.

Wichink Kruit, R. J., van Pul, W. A. J., Sauter, F. J., van den Broek, M., Nemitz, E., Sutton, M. A., Krol, M., and Holtslag, A. A. M.: Modelling the surface-atmosphere exchange of ammonia, *Atmos. Environ.*, 44(7), 945–957, 2010.

25 Wolfe, G. M., Thornton, J. A., Yataavelli, R. L. N., McKay, M., Goldstein, A. H., LaFranchi, B., Min, K.-E., and Cohen, R. C.: Eddy covariance fluxes of acyl peroxy nitrates (PAN, PPN and MPAN) above a Ponderosa pine forest, *Atmos. Chem. Phys.*, 9, 615–634, doi:10.5194/acp-9-615-2009, 2009.

30 Wolff, V., Trebs, I., Foken, T., and Meixner, F. X.: Exchange of reactive nitrogen compounds: concentrations and fluxes of total ammonium and total nitrate above a spruce canopy, *Biogeosciences*, 7, 1729–1744, 2010, <http://www.biogeosciences.net/7/1729/2010/>.

Wu, Y., Brashers, B., Finkelstein, P. L., and Pleim, J. E.: A multiplayer biochem-

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

ical dry deposition model, I. Model formulation, *J. Geophys. Res.*, 108(D1), 4013, doi:10.1029/2002JD00293, 2003.

5 Wu, Y., Walker, J., Schwede, D., Peters-Lidard, C., Dennis, R., and Robarge, W.: A new model of bi-directional ammonia exchange between the atmosphere and biosphere: Ammonia stomatal compensation point, *Agr. Forest Meteorol.*, 149, 263–280, 2009.

Wyers, G. P. and Erisman, J. W.: Ammonia exchange over coniferous forest, *Atmos. Environ.*, 32, 441–451, 1998.

Zhang, L., Gong, S., Padro, J., and Barrie, L.: A size-segregated particle dry deposition scheme for an atmospheric aerosol module, *Atmos. Environ.*, 35, 549–560, 2001.

10 Zhang, L., Brook, J. R., and Vet, R.: A revised parameterization for gaseous dry deposition in air-quality models, *Atmos. Chem. Phys.*, 3, 2067–2082, doi:10.5194/acp-3-2067-2003, 2003.

Zhang, L., Vet, R., O'Brien, J.M., Mihele, C., Liang, Z., and Wiebe, A.: Dry deposition of individual nitrogen species at eight Canadian rural sites, *J. Geophys. Res.*, 114, D02301, doi:10.1029/2008JD010640, 2009.

15 Zhang, L., Wright, L. P., and Asman, W. A. H.: Bi-directional air-surface exchange of atmospheric ammonia – A review of measurements and a development of a big-leaf model for applications in regional-scale air-quality models, *J. Geophys. Res. Atmos.*, doi:10.1029/2009JD013589, 115, D20310, doi:10.1029/2009JD013589, 2010.

20 Zimmermann, F., Plessow, K., Queck, R., Bernhofer, C., and Matschullat, J.: Atmospheric N- and S-fluxes to a spruce forest-Comparison of inferential modelling and the throughfall method, *Atmos. Environ.*, 40, 4782–4796, doi:10.1016/j.atmosenv.2006.03.056, 2006.

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. NitroEurope inferential network monitoring sites^a.

Site Code	Site Name	Land use/Dominant vegetation	LU ^b	Lat. ° N	Long. ° E	Altitude m a.m.s.l.	Temp. ° C	Rainfall mm	h ^c m	LAI ^d m ² m ⁻²
BE-Bra	Brasschaat	Scots pine, pedunculate oak	F	51.31	4.52	16	11.2	770	22	2
BE-Vie	Vielsalm	Eur. beech, coast douglas fir	F	50.31	6.00	450	8.4	1000	30	5
CH-Lae	Laegeren	Ash, sycamore, beech, spruce	F	47.48	8.37	689	7.6	1100	30	6
CZ-BK1	Bily Kriz	Norway spruce	F	49.50	18.54	908	8.3	1200	13	9
DE-Hai	Hainich	Eur. beech, maple, ash	F	51.08	10.45	430	8.7	775	33	7
DE-Hoe	Höglwald	Norway spruce	F	48.30	11.10	540	7.8	870	35	6
DE-Tha	Tharandt	Norway spruce, scots pine	F	50.96	13.57	380	9.2	820	27	8
DE-Wet	Wetzstein	Norway spruce	F	50.45	11.46	785	6.7	950	22	8
DK-Sor	Soroe	Eur. beech	F	55.49	11.65	40	9.0	730	31	5
ES-ES1	El Saler	Aleppo pine, stone pine, macchia	F	39.35	-0.32	5	17.8	551	10	3
ES-LMa	Las Majadas	Open holm oak, shrubs	F	39.94	-5.77	258	15.8	528	8	1
FI-Hyy	Hyytiälä	Scots pine	F	61.85	24.30	181	4.8	709	14	7
FI-Sod	Sodankylä	Scots pine	F	67.36	26.64	180	0.7	499	13	1
FR-Fon	Fontainbleau	Oak	F	48.48	2.78	92	11.3	690	28	5
FR-Hes	Hesse	Eur. beech	F	48.67	7.07	300	10.3	975	16	5
FR-LBr	Le Bray	Maritime pine	F	44.72	-0.77	61	12.9	972	22	3
FR-Pue	Puechabon	Holm oak	F	43.74	3.60	270	13.7	872	6	3
IT-Col	Collelongo	Eur. beech	F	41.85	13.59	1560	7.5	1140	22	7
IT-Ren	Renon	Norway spruce, stone pine	F	46.59	11.43	1730	4.9	1010	29	5
IT-Ro2	Roccarespampani	Turkey oak, downy oak	F	42.39	11.92	224	15.1	876	17	4
IT-SRo	San Rossore	Maritime pine, holm oak	F	43.73	10.28	4	15.2	920	18	4
NL-Loo	Loobos	Scots Pine	F	52.17	5.74	25	10.4	786	17	2
NL-Spe	Speulderbos	Douglas fir, Jap. larch, Eur. Beech	F	52.25	5.69	52	9.7	966	32	11
PT-Esp	Espirra	Eucalyptus coppice	F	38.64	-8.60	95	16.2	709	20	5
PT-Mi1	Mitra II (Evora)	Cork oak	F	38.54	-8.00	264	15.4	665	7	3
RU-Fyo	Fyodorovskoye	Norway spruce	F	56.46	32.92	265	5.3	711	21	3
SE-Nor	Norunda	Norway spruce, scots pine	F	60.08	17.47	45	7.0	527	25	5
SE-Sk2	Skyttorp	Scots pine, Norway spruce	F	60.13	17.84	55	5.5	527	14	3
UK-Gri	Griffin	Sitka Spruce	F	56.62	-3.80	340	7.8	1200	9	8
DE-Meh	Mehrstedt	Afforested grassland	SN	51.28	10.66	293	8.5	547	0.5	2.9
ES-VDA	Vall d'Alinyà	Upland grassland	SN	42.15	1.45	1765	7.1	1064	0.1	1.4
FI-Lom	Lompolojännkkä	Sedge fen	SN	68.21	24.35	269	-0.1	500	0.4	1.0
HU-Bug	Bugac	Semi-arid grassland	SN	46.69	19.60	111	10.8	500	0.5	4.7
IT-Amp	Amplero	Grassland	SN	41.90	13.61	884	9.6	1365	0.4	2.5
IT-MBo	Monte Bondone	Upland grassland	SN	46.03	11.08	1550	5.5	1189	0.3	2.5
NL-Hor	Horstermeer	Natural fen (peat)	SN	52.03	5.07	-2	10.8	800	2.5	6.9
PL-wet	POLWET	Wetland (reeds, carex, sphagnum)	SN	52.76	16.31	54	8.9	550	2.1	4.9
UK-AMo	Auchencorth Moss	Blanket bog	SN	55.79	-3.24	270	7.6	798	0.6	2.1

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. Continued.

Site Code	Site Name	Land use/Dominant vegetation	LU ^b	Lat. ° N	Long. ° E	Altitude m a.m.s.l.	Temp. ° C	Rainfall mm	h^c m	LAI ^d m ² m ⁻²
CH-Oe1	Oensingen	Cut Grassland	G	47.29	7.73	450	9.4	1200	0.6	6.6
DE-Gri	Grillenburg	Cut Grassland	G	50.95	13.51	375	9.0	861	0.7	6.0
DK-Lva	Rimi	Cut Grassland	G	55.70	12.12	8	9.2	600	0.5	3.5
FR-Lq2	Laqueuille	Grazed Grassland	G	45.64	2.74	1040	7.5	1100	0.2	2.4
IE-Ca2	Carlow	Grazed Grassland	G	52.85	-6.90	56	9.4	804	0.2	5.7
IE-Dri	Dripsey	Grazed Grassland	G	51.99	-8.75	187	9.6	1450	0.5	4.0
NL-Ca1	Cabauw	Grazed Grassland	G	51.97	4.93	-1	11.1	786	0.2	9.9
UK-EBu	Easter Bush	Grazed Grassland	G	55.87	-3.21	190	9.0	870	0.2	5.5
BE-Lon	Lonzee	Crop rotation	C	50.55	4.74	165	9.1	772	0.9	6
DE-Geb	Gebesee	Crop rotation	C	51.10	10.91	162	10.1	492	1.0	5.5
DE-Kli	Klingenberg	Crop rotation	C	50.89	13.52	478	8.1	850	2.2	5.0
DK-Ris	Risbyholm	Crop rotation	C	55.53	12.10	10	9.0	575	1.0	4.6
FR-Gri	Grignon	Crop rotation	C	48.84	1.95	125	11.1	600	2.4	6.2
IT-BCi	Borgo Cioffi	Crop rotation	C	40.52	14.96	20	16.4	490	3.0	7.3
IT-Cas	Castellaro	Maize/Rice rotation	C	45.06	8.67	89	13.2	984	2.8	4.9
UA-Pet	Petrodolinskoye	Crop Rotation	C	42.50	30.30	66	10.1	464	0.6	4.2
UK-ESa	East Saltoun	Crop rotation	C	55.90	-2.84	97	8.5	700	na	na

^aSee Table A1 in the online Supplement for literature references for each site.

^bEcosystem type: F: forest; SN: semi-natural short vegetation; G: grassland (G); C: cropland.

^cCanopy height: mean tree height for F; annual maximum value for SN, G and C.

^dLeaf area index: annual maximum; the measurement type (single-sided, projected, total) is not specified.

"na": not available.

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Table 2. Summary of ambient N_r concentrations across the NEU inferential network (unit: $\mu\text{g N m}^{-3}$). Data for NH_3 , HNO_3 , NH_4^+ and NO_3^- are arithmetic means, minima and maxima of 24 monthly values over the 2007–2008 period. Data for NO_2 are calculated from hourly concentration measurements for some sites (see text), or from modelled EMEP 50×50 km data.

Site Code	NH_3			HNO_3			NO_2			NH_4^+		NO_3^-			
	Mean	Min	Max	Mean	Min	Max	Mean	Min	Max	Mean	Max	Mean	Max		
BE-Bra	2.28	0.03	9.43	0.46	0.10	1.28	8.98	3.96	16.69	1.34	0.04	3.89	0.87	0.01	5.21
BE-Vie	0.37	0.09	1.51	0.13	0.01	0.31	3.38	na	na	0.66	0.12	1.82	0.53	0.03	3.06
CH-Lae	1.14	0.37	2.55	0.36	0.26	0.64	2.49	na	na	0.95	0.43	2.12	0.60	0.18	1.58
CZ-BK1	0.51	0.12	0.95	0.40	0.21	0.78	2.75	na	na	0.89	0.12	1.38	0.40	0.22	0.76
DE-Hai	0.57	0.06	1.64	0.22	0.11	0.52	2.65	na	na	0.94	0.35	1.86	0.44	0.17	0.92
DE-Hoe	1.91	0.60	3.31	0.34	0.13	0.77	2.85	na	na	1.02	0.39	2.57	0.50	0.20	0.99
DE-Tha	0.62	0.11	1.37	0.28	0.17	0.60	2.82	na	na	0.87	0.56	1.35	0.40	0.14	0.84
DE-Wet	0.43	0.10	1.01	0.26	0.16	0.42	2.51	na	na	0.80	0.43	1.46	0.43	0.19	0.83
DK-Sor	1.32	0.37	4.74	0.22	0.06	0.78	2.47	na	na	0.72	0.16	2.21	0.77	0.01	2.94
ES-ES1	1.56	0.80	2.57	0.32	0.10	0.45	1.88	na	na	0.90	0.34	1.94	0.99	0.52	1.95
ES-LMa	1.03	0.52	2.08	0.23	0.10	0.50	0.50	na	na	0.46	0.15	1.64	0.38	0.18	0.84
FI-Hyy	0.10	0.02	0.27	0.09	0.00	0.16	2.72	0.91	8.83	0.19	0.04	0.52	0.07	0.01	0.30
FI-Sod	0.13	0.00	0.61	0.04	0.01	0.12	0.21	na	na	0.12	0.00	0.55	0.02	0.00	0.06
FR-Fon	0.90	0.27	2.95	0.41	0.24	0.80	2.12	na	na	0.96	0.38	2.20	0.68	0.32	1.59
FR-Hes	0.89	0.26	2.42	0.35	0.21	0.61	1.99	na	na	0.80	0.37	1.54	0.48	0.21	0.94
FR-LBr	1.16	0.46	5.17	0.28	0.14	0.45	1.01	na	na	0.58	0.24	1.40	0.45	0.26	0.88
FR-Pue	0.43	0.12	0.82	0.23	0.11	0.52	0.95	na	na	0.46	0.19	1.19	0.30	0.14	0.60
IT-Col	0.42	0.12	0.98	0.13	0.05	0.31	1.11	na	na	0.47	0.16	0.83	0.25	0.06	0.48
IT-Ren	0.26	0.05	0.50	0.09	0.03	0.21	1.10	0.30	2.18	0.52	0.03	1.29	0.26	0.02	0.62
IT-Ro2	1.83	0.77	7.51	0.24	0.13	0.34	0.86	na	na	0.86	0.51	1.53	0.51	0.30	0.78
IT-SRo	0.84	0.30	5.71	0.31	0.11	0.51	1.12	na	na	0.90	0.38	1.93	0.62	0.31	1.04
NL-Loo	3.44	0.99	6.67	0.27	0.08	0.51	7.41	na	na	1.60	0.70	5.26	0.79	0.26	1.42
NL-Spe	3.91	1.58	6.74	0.36	0.24	0.52	5.10	2.56	9.74	1.32	0.63	2.21	0.91	0.16	1.62
PT-Esp	1.86	0.86	4.40	0.39	0.15	0.82	2.63	na	na	0.84	0.45	1.73	0.51	0.04	0.93
PT-Mi1	0.94	0.26	2.49	0.25	0.06	0.96	0.89	na	na	0.69	0.24	2.10	0.38	0.20	0.88
RU-Fyo	0.28	0.05	0.51	0.14	0.07	0.29	0.50	na	na	0.45	0.18	0.79	0.15	0.06	0.31
SE-Nor	0.22	0.02	0.68	0.05	0.01	0.17	0.66	na	na	0.25	0.03	1.02	0.10	0.01	0.31
SE-Sk2	0.16	0.02	0.95	0.06	0.02	0.12	0.63	na	na	0.21	0.01	0.64	0.10	0.01	0.45
UK-Gri	0.27	0.04	1.47	0.12	0.02	0.47	0.48	na	na	0.39	0.02	1.76	0.29	0.03	1.49
Mean (F)	1.03	0.32	2.83	0.24	0.11	0.51	2.15	1.25	6.92	0.73	0.26	1.75	0.45	0.15	1.19
DE-Meh	1.48	0.21	4.08	0.29	0.18	0.48	2.67	na	na	1.12	0.03	1.66	0.55	0.20	0.92
ES-VDA	0.90	0.07	5.28	0.12	0.04	0.49	0.83	na	na	0.70	0.09	3.42	0.27	0.02	0.75
FI-Lom	0.09	0.01	0.31	0.03	0.00	0.26	0.19	0.00	0.48	0.21	0.00	0.65	0.02	0.00	0.07
HU-Bug	2.27	0.71	5.16	0.30	0.12	0.48	2.61	1.53	4.65	1.25	0.63	2.40	0.46	0.15	1.03
IT-Amp	0.56	0.19	1.20	0.14	0.07	0.36	1.11	na	na	0.48	0.25	1.05	0.22	0.10	0.40
IT-MBo	0.74	0.14	1.84	0.22	0.12	0.38	1.77	na	na	0.74	0.06	2.18	0.47	0.02	1.13
NL-Hor	2.49	0.77	5.28	0.33	0.12	0.52	9.45	na	na	1.37	0.54	2.97	0.94	0.43	1.85
PL-wet	0.95	0.24	2.39	0.25	0.02	0.41	1.45	na	na	1.09	0.42	2.85	0.46	0.12	1.13
UK-AMo	0.63	0.30	1.22	0.09	0.03	0.23	1.45	0.71	2.46	0.38	0.09	0.97	0.23	0.05	0.56
Mean (SN)	1.12	0.29	2.97	0.20	0.08	0.40	2.39	0.75	2.53	0.82	0.24	2.02	0.40	0.12	0.87

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Table 2. Continued.

Site Code	NH ₃			HNO ₃			NO ₂			NH ₄ ⁺		NO ₃ ⁻			
	Mean	Min	Max	Mean	Min	Max	Mean	Min	Max	Mean	Min	Max	Mean	Min	Max
CH-Oe1	2.68	0.71	6.51	0.41	0.20	0.71	10.89	5.53	19.01	1.15	0.50	2.05	0.66	0.34	1.25
DE-Gri	0.70	0.12	1.28	0.36	0.17	1.22	2.82	na	na	0.89	0.49	2.94	0.47	0.17	1.89
DK-Lva	1.26	0.27	3.71	0.20	0.02	0.35	2.47	na	na	0.56	0.22	1.37	0.79	0.05	3.08
FR-Lq2	1.11	0.37	1.81	0.14	0.06	0.48	0.65	na	na	0.44	0.19	1.36	0.25	0.11	0.70
IE-Ca2	1.56	0.81	3.04	0.10	0.04	0.22	0.75	na	na	0.59	0.10	1.87	0.33	0.11	1.08
IE-Dri	2.03	0.72	4.94	0.07	0.01	0.17	0.45	na	na	0.53	0.05	2.24	0.29	0.05	0.93
NL-Ca1	5.93	3.10	10.79	0.41	0.25	0.98	9.45	na	na	1.66	0.35	4.95	1.10	0.09	2.16
UK-EBu	1.08	0.32	2.17	0.12	0.04	0.24	0.85	0.20	1.96	0.38	0.08	0.87	0.26	0.05	0.59
Mean (G)	2.04	0.80	4.28	0.23	0.10	0.55	3.54	2.87	10.48	0.78	0.25	2.21	0.52	0.12	1.46
BE-Lon	3.93	1.00	14.46	0.29	0.05	0.47	4.31	na	na	1.08	0.04	2.58	0.73	0.09	2.41
DE-Geb	4.14	0.50	13.41	0.25	0.15	0.33	2.65	na	na	1.41	0.05	6.73	0.56	0.18	1.18
DE-Kli	1.32	0.24	2.49	0.31	0.14	0.49	2.82	na	na	1.05	0.61	2.56	0.53	0.18	1.94
DK-Ris	4.32	0.15	14.26	0.14	0.02	0.27	2.47	na	na	0.58	0.01	1.66	0.44	0.07	0.90
FR-Gri	3.16	0.92	10.24	0.45	0.18	0.98	4.99	1.95	11.01	0.94	0.26	2.56	0.76	0.30	2.01
IT-BCi	7.18	2.58	21.63	0.38	0.22	0.82	1.26	na	na	3.12	0.37	14.81	0.73	0.33	1.23
IT-Cas	3.42	1.30	5.91	0.44	0.22	0.86	1.12	0.54	1.61	2.38	0.32	4.81	1.43	0.35	3.05
UA-Pet	2.50	0.62	5.35	0.36	0.18	0.68	1.00	na	na	1.44	0.34	9.52	0.48	0.21	0.76
UK-ESa	2.92	0.80	13.57	0.12	0.06	0.20	2.39	na	na	0.71	0.15	3.18	0.24	0.10	0.41
Mean (C)	3.65	0.90	11.26	0.30	0.13	0.57	2.56	1.25	6.31	1.41	0.24	5.38	0.66	0.20	1.54

“na”: not available

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 3. Summary of modelled annual dry deposition fluxes to the sites of the NEU inferential network (unit: $\text{kg N ha}^{-1} \text{ yr}^{-1}$), averaged over the two years 2007–2008. A minus “-” sign denotes net deposition; positive numbers for NH_3 in CBED indicate a net emission.

Site	CBED					CDRY					EMEP-03				IDEM					
	NH_3	HNO_3	NO_2	NH_4^+	NO_3^-	NH_3	HNO_3	NO_2	NH_4^+	NO_3^-	NH_3	HNO_3	NO_2	NH_4^+	NO_3^-	NH_3	HNO_3	NO_2	NH_4^+	NO_3^-
BE-Bra	-16.7	-6.6	-2.5	-4.2	-4.3	-5.3	-3.6	-8.0	-1.0	-0.7	-7.4	-6.5	-2.7	-1.0	-1.5	-12.6	-4.3	-7.9	-6.6	-5.3
BE-Vie	-3.7	-2.4	-1.3	-2.2	-3.0	-1.5	-1.4	-4.6	-0.5	-0.4	-2.1	-2.2	-0.4	-0.4	-0.3	-3.2	-1.6	-1.3	-3.9	-4.3
CH-Lae	-7.9	-4.3	-0.9	-2.6	-2.6	-3.1	-2.4	-2.9	-0.6	-0.4	-3.5	-3.7	-0.1	-0.5	-0.6	-6.3	-2.9	-0.9	-6.1	-4.8
CZ-BK1	-3.4	-4.2	-1.0	-2.2	-1.6	-1.3	-2.8	-2.9	-0.5	-0.2	-2.6	-3.2	-0.3	-0.6	-0.6	-3.2	-3.0	-2.4	-3.9	-2.1
DE-Hai	-4.8	-4.6	-1.0	-4.3	-3.2	-1.4	-2.3	-2.4	-0.9	-0.4	-2.4	-4.2	-0.1	-0.5	-0.4	-4.4	-3.0	-1.1	-5.7	-3.4
DE-Hoe	-14.8	-4.4	-1.3	-3.3	-2.6	-5.4	-2.6	-3.2	-0.8	-0.4	-5.5	-4.3	-0.3	-1.0	-1.2	-9.6	-3.0	-3.2	-4.5	-2.5
DE-Tha	-5.6	-4.3	-1.2	-2.9	-2.2	-2.2	-2.7	-3.1	-0.7	-0.4	-3.2	-4.0	-0.2	-0.7	-0.8	-4.5	-3.0	-2.8	-3.6	-2.0
DE-Wet	-3.7	-4.5	-0.9	-3.2	-2.8	-1.7	-2.7	-4.2	-0.8	-0.5	-2.5	-3.6	-0.1	-0.9	-1.3	-3.9	-2.8	-2.4	-5.3	-3.7
DK-Sor	-9.8	-3.5	-1.0	-3.1	-5.3	-3.3	-1.8	-2.3	-0.7	-0.6	-4.5	-3.3	0.0	-0.3	-0.4	-7.1	-2.2	-1.0	-3.4	-4.7
ES-ES1	-12.6	-4.5	-1.3	-2.5	-4.3	-4.3	-3.0	-1.8	-0.6	-0.6	-4.9	-4.5	0.0	-0.7	-1.8	-8.0	-3.3	-1.8	-3.1	-3.8
ES-LMa	-6.2	-1.9	-0.2	-0.8	-1.1	-1.1	-0.6	-0.2	-0.2	-0.1	-2.2	-1.9	0.0	-0.2	-0.4	-2.5	-1.4	-0.3	-0.7	-0.6
FI-Hyy	-0.8	-1.1	-0.4	-0.6	-0.3	-0.3	-0.7	-2.8	-0.1	-0.1	-0.4	-0.8	-1.0	-0.1	-0.1	-0.7	-0.7	-1.6	-1.1	-0.4
FI-Sod	-1.0	-0.5	-0.1	-0.3	-0.1	-0.4	-0.4	-0.2	-0.1	0.0	-0.5	-0.2	0.0	-0.1	0.0	-1.0	-0.3	-0.2	-0.5	-0.1
FR-Fon	-6.0	-4.3	-1.0	-2.8	-3.2	-1.7	-2.2	-1.6	-0.6	-0.4	-2.7	-4.2	0.0	-0.3	-0.5	-4.4	-3.1	-1.0	-3.8	-3.2
FR-Hes	-5.8	-3.7	-0.8	-2.2	-2.1	-1.6	-1.8	-1.3	-0.5	-0.3	-2.4	-3.5	0.0	-0.2	-0.3	-4.4	-2.6	-1.0	-3.2	-2.3
FR-LBr	-7.9	-3.0	-0.5	-1.4	-1.8	-3.2	-1.9	-1.2	-0.3	-0.3	-3.1	-3.0	0.0	-0.4	-0.7	-5.9	-2.1	-1.0	-2.7	-2.6
FR-Pue	-2.7	-2.3	-0.5	-1.1	-1.2	-0.6	-0.9	-0.6	-0.3	-0.1	-0.9	-2.3	0.0	-0.3	-0.5	-1.8	-1.6	-0.6	-1.5	-1.1
IT-Col	-2.8	-1.4	-0.5	-1.2	-1.0	-0.8	-0.6	-0.7	-0.3	-0.1	-1.1	-1.2	0.0	-0.4	-0.6	-2.0	-1.0	-0.5	-2.9	-1.9
IT-Ren	-2.2	-1.2	-0.3	-1.3	-1.0	-0.8	-0.7	-0.9	-0.3	-0.2	-1.0	-0.9	0.0	-0.5	-0.5	-1.6	-0.8	-1.0	-2.4	-1.4
IT-Ro2	-12.3	-2.5	-0.5	-2.0	-1.8	-2.9	-1.0	-0.5	-0.5	-0.2	-4.4	-2.5	0.0	-0.7	-0.9	-6.8	-1.7	-0.7	-3.7	-2.4
IT-SRo	-5.6	-3.2	-0.6	-2.1	-2.2	-1.8	-1.9	-1.0	-0.5	-0.3	-2.2	-3.2	0.0	-0.6	-1.0	-4.0	-2.2	-1.0	-3.8	-3.0
NL-Loo	-25.7	-3.6	-3.1	-5.1	-4.1	-11.5	-2.4	-9.4	-1.2	-0.7	-11.1	-3.5	-2.3	-1.2	-1.5	-20.6	-2.4	-7.6	-8.0	-4.6
NL-Spe	-28.4	-4.6	-1.5	-4.4	-5.0	-9.9	-2.6	-5.2	-1.0	-0.6	-10.0	-4.5	-1.7	-1.0	-1.7	-19.2	-3.1	-4.3	-5.7	-4.7
PT-Esp	-6.9	-1.9	-1.2	-1.3	-1.3	-1.2	-0.5	-1.0	-0.3	-0.2	-2.3	-1.9	-0.1	-0.3	-0.4	-2.8	-1.5	-1.8	-1.0	-0.7
PT-Mi1	-5.2	-2.0	-0.4	-1.3	-1.2	-1.5	-0.8	-0.5	-0.3	-0.2	-2.1	-2.2	0.0	-0.4	-0.5	-3.0	-1.6	-0.6	-2.0	-1.2
RU-Fyo	-2.2	-2.0	-0.2	-1.4	-0.7	-0.8	-1.2	-0.6	-0.3	-0.1	-1.2	-1.3	0.0	-0.3	-0.3	-2.3	-1.2	-0.4	-2.7	-1.0
SE-Nor	-2.1	-0.9	-0.3	-0.8	-0.5	-0.9	-0.6	-0.8	-0.2	-0.1	-0.9	-0.7	0.0	-0.2	-0.2	-1.5	-0.6	-0.6	-1.2	-0.6
SE-Sk2	-1.5	-1.1	-0.2	-0.7	-0.5	-0.5	-0.7	-0.7	-0.2	-0.1	-0.7	-0.9	0.0	-0.1	-0.2	-1.0	-0.8	-0.6	-1.2	-0.7
UK-Gri	-2.1	-1.6	-0.2	-1.0	-1.3	-0.7	-0.9	-0.4	-0.2	-0.2	-1.0	-1.5	0.0	-0.2	-0.5	-2.2	-0.9	-0.4	-2.1	-2.1
DE-Meh	-5.6	-1.3	-0.8	-0.4	-0.3	-2.6	-1.2	-2.5	-0.5	-0.2	-2.4	-1.3	-0.1	-0.4	-0.5	-5.6	-1.1	-2.1	-0.6	-0.3
ES-VDA	-1.9	-0.3	-0.2	-0.2	-0.1	-0.7	-0.3	-0.5	-0.2	-0.1	-0.9	-0.3	0.0	-0.2	-0.2	-2.0	-0.3	-0.5	-0.4	-0.2
FI-Lom	-0.3	-0.1	0.0	-0.1	0.0	-0.1	-0.1	0.0	-0.1	0.0	-0.1	-0.1	0.0	0.0	0.0	-0.1	-0.1	0.0	-0.1	0.0
HU-Bug	-5.8	-0.9	-0.5	-0.3	-0.2	-2.1	-0.7	-1.1	-0.4	-0.1	-2.6	-0.9	-0.1	-0.3	-0.2	-5.7	-0.8	-1.3	-0.7	-0.2
IT-Amp	-1.1	-0.3	-0.4	-0.1	0.0	-0.4	-0.2	-0.4	-0.1	0.0	-0.5	-0.3	0.0	-0.2	-0.1	-1.1	-0.3	-0.8	-0.5	-0.2
IT-MBo	-2.1	-0.6	-0.4	-0.1	-0.1	-0.7	-0.5	-0.7	-0.2	-0.1	-1.1	-0.6	0.0	-0.3	-0.3	-2.4	-0.5	-1.1	-0.6	-0.4
NL-Hor	-13.9	-2.6	-3.7	-0.7	-0.7	-6.0	-2.2	-9.7	-0.8	-0.5	-8.0	-2.6	-1.7	-0.6	-1.0	-16.9	-2.0	-10.1	-1.2	-0.9
PL-wet	-4.5	-1.2	-0.4	-0.4	-0.2	-1.4	-0.8	-0.9	-0.5	-0.2	-2.1	-1.1	0.0	-0.4	-0.3	-4.0	-1.0	-1.2	-0.9	-0.4
UK-AMo	-2.9	-0.6	-0.5	-0.2	-0.2	-1.2	-0.5	-1.1	-0.2	-0.1	-1.5	-0.5	0.0	-0.1	-0.2	-3.2	-0.4	-1.1	-0.2	-0.1

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Table 3. Continued.

Site	CBED				CDRY				EMEP-03				IDEM							
	NH ₃	HNO ₃	NO ₂	NH ₄ ⁺	NO ₃ ⁻	NH ₃	HNO ₃	NO ₂	NH ₄ ⁺	NO ₃ ⁻	NH ₃	HNO ₃	NO ₂	NH ₄ ⁺	NO ₃ ⁻	NH ₃	HNO ₃	NO ₂	NH ₄ ⁺	NO ₃ ⁻
CH-Oe1	0.1	-1.0	-3.8	-0.2	-0.2	-1.4	-0.9	-5.9	-0.3	-0.2	-1.9	-1.0	-2.7	-0.2	-0.3	-3.2	-0.9	-6.6	-0.8	-0.5
DE-Gri	5.6	-1.1	-1.1	-0.2	-0.2	-0.9	-1.0	-2.0	-0.2	-0.1	-1.3	-1.1	-0.2	-0.2	-0.2	-1.8	-1.0	-1.8	-0.4	-0.2
DK-Lva	1.9	-0.9	-1.0	-0.2	-0.5	-2.1	-0.8	-2.2	-0.2	-0.3	-2.1	-0.9	0.0	-0.2	-0.4	-4.5	-0.8	-2.0	-0.3	-0.2
FR-Lq2	-0.2	-0.7	-0.2	-0.2	-0.2	-1.3	-0.6	-0.6	-0.2	-0.1	-1.5	-0.6	0.0	-0.1	-0.2	-2.9	-0.6	-0.4	-0.2	-0.1
IE-Ca2	1.3	-0.4	-0.3	-0.2	-0.1	-1.8	-0.3	-0.6	-0.2	-0.1	-1.8	-0.4	0.0	-0.1	-0.2	-4.3	-0.3	-0.5	-0.3	-0.2
IE-Dri	-2.2	-0.4	-0.2	-0.2	-0.2	-3.5	-0.4	-0.5	-0.2	-0.1	-3.8	-0.4	0.0	-0.2	-0.3	-9.3	-0.4	-0.4	-0.3	-0.1
NL-Ca1	-5.8	-1.6	-4.0	-0.6	-0.6	-9.5	-1.5	-8.3	-0.6	-0.4	-9.8	-1.6	-2.2	-0.4	-0.6	-19.1	-1.4	-7.0	-0.7	-0.5
UK-EBu	-0.2	-0.5	-0.3	-0.1	-0.1	-1.2	-0.5	-0.7	-0.2	-0.1	-1.1	-0.5	0.0	-0.1	-0.2	-2.6	-0.4	-0.6	-0.2	-0.1
BE-Lon	0.5	-1.1	-1.4	-0.7	-0.6	-3.0	-0.9	-3.6	-0.4	-0.3	-3.1	-1.1	-0.4	-0.2	-0.1	-6.1	-0.9	-3.0	-0.7	-0.4
DE-Geb	1.5	-0.9	-0.9	-0.8	-0.4	-2.5	-0.7	-1.9	-0.5	-0.2	-2.2	-0.9	-0.1	-0.2	-0.1	-5.5	-0.7	-1.8	-0.8	-0.3
DE-Kli	1.9	-1.2	-0.9	-0.8	-0.5	-1.7	-1.0	-2.4	-0.4	-0.2	-2.0	-1.2	-0.1	-0.2	-0.1	-3.6	-1.0	-1.9	-0.6	-0.3
DK-Ris	-1.9	-0.3	-0.7	-0.3	-0.3	-3.7	-0.3	-1.6	-0.2	-0.1	-4.1	-0.3	0.0	-0.2	-0.1	-6.7	-0.3	-1.5	-0.7	-0.3
FR-Gri	1.0	-1.5	-0.8	-0.6	-0.6	-2.9	-1.2	-2.8	-0.3	-0.3	-2.9	-1.5	-0.6	-0.1	-0.2	-6.2	-1.3	-2.0	-0.4	-0.4
IT-BCi	1.6	-1.8	-0.5	-2.3	-0.7	-6.2	-1.3	-1.1	-1.3	-0.3	-5.7	-1.8	0.0	-0.5	-0.2	-16.3	-1.5	-1.1	-2.0	-0.4
IT-Cas	4.1	-0.8	-0.6	-0.5	-0.4	-1.9	-0.6	-0.6	-0.3	-0.3	-2.1	-0.8	0.0	-0.5	-0.4	-3.9	-0.7	-1.0	-2.9	-1.8
UA-Pet	-0.2	-0.8	-0.2	-0.7	-0.3	-1.4	-0.6	-0.6	-0.4	-0.1	-1.9	-0.7	0.0	-0.4	-0.1	-2.9	-0.7	-0.5	-1.4	-0.4
UK-ESa	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na	na

“na”: not available.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.




Fig. 1. Comparison of measured u_* (long-term means at each observation site) with inferential model estimates, using as input either model default values of h_c (A) or measured h_c at each site (B). Panels (C) and (D): comparison of mean observations and model default values of h_c and LAI for the different land use types (F: forests; SN: semi-natural; G: grasslands; C: croplands). Note that the CDRY model uses tabulated ecosystem-specific values of z_0 , and does not require h_c as a predictor of z_0 ; thus, for comparability's sake, the h_c values presented for CDRY in C were actually calculated by multiplying model z_0 by 10, since the other three models all use $z_0 = h_c/10$.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.




Fig. 2. Comparison of mean modelled daytime bulk stomatal conductances from the four inferential schemes.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.




Fig. 3. Overview of mean modelled deposition velocities (evaluated at $d + z_0 + 3$ m for all sites) for dominant atmospheric N_r species. Data are medians and 5-th and 95-th percentiles across the sites of mean V_d values at each site. Note the different scale on the vertical axis for NO_2 .

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.




Fig. 4. Modelled annual N_r dry deposition to NEU monitoring sites. Data are calculated as the sum of NH_3 , HNO_3 , aerosol NH_4^+ and NO_3^- fluxes from DELTA measurements, plus NO_2 dry deposition from modelled (EMEP 50×50 km) or measured NO_2 concentrations.




Fig. 5. Relative contributions of N_r species to total inorganic N dry deposition. For G and C data in CBED (top panel), negative percentages for NH_3 denote net NH_3 emissions, which are expressed relative to the sum of dry deposition fluxes for the other four N_r species.

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 6. Comparison of modelled annual NH₃ exchange from NEU network DELTA data with measured estimates from historical long-term micrometeorological flux datasets. For five monitoring sites, the ensemble average of CBED, CDRY, EMEP-03 and IDEM is shown with error bars showing the range (min, max) of model estimates. (1): Neiryneck et al., 2007; (2): Erisman et al. (1996); (3): Flechard (1998); (4) and (5): data from Flechard et al. (2010), showing (4) the annual NH₃ flux for background conditions (outside fertilisation events) and (5) the net emission flux from the whole dataset; (6) and (7): data from Milford (2004), with (6) the annual dry deposition, calculated from the net overall flux (7) minus the gross annual emission of 4.2 kg N ha⁻¹ yr⁻¹ due to grassland management activities (fertilisation, cuts). The secondary axis shows the mean concentrations during the NEU reference period (2007–2008) as well as during the flux monitoring periods.

Dry deposition of reactive nitrogen to European ecosystems

C. R. Flechard et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion