

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

**Water vapor budget  
associated to  
overshoots in the  
tropical stratosphere**

X. M. Liu et al.

# Water vapor budget associated to overshoots in the tropical stratosphere: mesoscale modelling study of 4–5 August 2006 during SCOUT-AMMA

X. M. Liu<sup>1</sup>, E. D. Rivière<sup>1</sup>, V. Marécal<sup>2</sup>, G. Durry<sup>1</sup>, A. Hamdouni<sup>1</sup>, J. Arteta<sup>2,\*</sup>, and S. Khaykin<sup>3</sup>

<sup>1</sup>Groupe de Spectrométrie moléculaire et Atmosphérique (GSMA), Université de Reims Champagne-Ardenne (URCA) and CNRS, UMR6089, Reims, France

<sup>2</sup>Laboratoire de Physique et Chimie de l'Environnement et de l'Espace (LPC2E), CNRS and Université d'Orléans, France

<sup>3</sup>Central Aerological Observatory of Roshydromet 3, Pervomayskaya str. Dolgoprudny, Moscow region 141700, Russian Federation, Russia

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

\* now at: Centre National de Recherche Météorologique (CNRM), Météo-France and CNRS, Toulouse, France

Received: 9 December 2009 – Accepted: 21 December 2009 – Published: 11 February 2010

Correspondence to: X. M. Liu (xiaoman.liu@univ-reims.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

ACPD

10, 3975–4025, 2010

**Water vapor budget associated to overshoots in the tropical stratosphere**

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

The aim of this paper is to study the impacts of overshooting convection at a local scale on the water distribution in the tropical UTLS. Overshooting convection is likely to be one of the key processes controlling the entry of water vapour amount in the stratosphere by injecting ice crystals above the tropopause which later sublime and hydrate the lower stratosphere. For this purpose, we quantify the individual impact of two overshooting cases in Africa observed during SCOUT-AMMA: the case of 4 August 2006 over Southern Chad which is likely to have influenced the water vapour measurements by micro-SDLA and FLASH-B from Niamey on 5 August, and the case of a mesoscale convective system over Aïr on 5 August 2006. We make use of high resolution (down to 1 km horizontally) three nested grid simulations with the three-dimensional regional atmospheric model BRAMS (Brazilian Regional Atmospheric Modelling System). In both cases, BRAMS succeeds in simulating the main features of the convective activity, as well as overshooting convection, though the exact position and time of the overshoots indicated by MSG brightness temperature difference is not fully reproduced (typically  $1^\circ$  in latitude compared with the overshoots indicated by brightness temperature difference from satellite observations for both cases, and several hours shift the Aïr case on 5 August 2006). Total water budgets associated with these two events show a significant injection of ice particles above the tropopause with maximum values of about  $3.7 \text{ ton s}^{-1}$  for the Chad case (4 August) and  $1.4 \text{ ton s}^{-1}$  for the Aïr case (5 August), and a total cross tropopause transport of about  $3300 \text{ ton h}^{-1}$  for the Chad case and  $2400 \text{ ton h}^{-1}$  for the Aïr case in the third domain of simulation. The order of magnitude of these modelled fluxes is lower but comparable with similar studies in other tropical areas based on models. These two estimations exhibit significant differences and highlight variability among the cases of the impact of overshooting convection in hydrating the lower stratosphere. We show that the regional enhancement of water above the tropopause is between 0.21 to 0.67 ppmv between 380 K and 400 K, in the range of other model estimations. Finally we emphasize that as long as the model resolution

ACPD

10, 3975–4025, 2010

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

is high, the hydrated area in the LS by overshooting convection can be advected relatively far away from the overshoot initial location, with locally mixing ratios of more than 3 ppmv higher than the background level, which is compatible with the balloon borne measurements performed above Niamey in the same air mass, 30 h after the overshoot. However, in the model, when exiting the highest resolution grid, the hydrating signal is lost rapidly.

## 1 Introduction

Water vapour is a key component of the stratosphere, both for climate and chemistry. As the most important greenhouse gas in the atmosphere, the amount of water vapour in the stratosphere can significantly affect the earth's climate. It also affects the ozone layer chemistry since water vapour is one of the main sources of OH hydroxyl radicals in the stratosphere and favours ozone depletion in winter by polar stratospheric cloud formation. A 1% increase of water vapour per year in the stratosphere was observed (Oltmans et al., 2000; Rosenlof et al., 2001) during the second half of the last century, which is believed to be partially due to water vapour transport across the tropical tropopause. The understanding and the prediction of water vapour distribution in the tropical upper troposphere (UT) and lower stratosphere (LS) is currently a key issue since this region is likely to control the entry of water vapour in the stratosphere. At local scale, one important process controlling the water amount in the LS is overshooting convection, injecting directly ice crystals above the tropopause which later sublime and hydrate the LS (Pommereau and Held, 2007).

The most accepted picture until recently is due to the extremely low water vapour mixing ratios reported by the few measurements available at the high altitude cold point tropopause over the Western Pacific and is also supported by trajectory model calculations. This is a dehydration process of air masses by freezing followed by sedimentation during the very slow ascent (6 months between 15 and 20 km) of tropospheric air due to positive radiative heating (Holton and Gettleman, 2001; Randel et al., 2001;

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**Water vapor budget associated to overshoots in the tropical stratosphere**X. M. Liu et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Gettleman et al., 2002; Fueglistaler et al., 2004, 2005) in the Tropical Tropopause Layer (hereafter TTL, Fueglistaler et al., 2009). This process is called the cold trap hypothesis. In contrast to this hypothesis, recent total water and water vapour measurements from aircraft and balloons show the presence of thin layers of enhanced water vapour or sometimes ice crystals up to several kilometres above the tropopause (Pommereau and Held, 2007; Nielsen et al., 2007; Chaboureaud et al., 2007; Corti et al., 2008; Khaykin et al., 2009). They are attributed to convective overshoots and lead to the hydration of the lower stratosphere. As also seen by the TRMM Precipitation Radar and Lightning Imaging Sensor, these injections occur mainly over land. (Liu and Zipser, 2005, Zipser, 2006). Although such events can be simulated by Cloud Resolving Models (Chaboureaud et al., 2007; Grosvenor et al. 2007; Chemel et al., 2009) they have a too small horizontal size (10–20 km) and are of too short duration (typically of one hour or less) for being captured by global Numerical Weather Prediction (NWP) models in which their impact cannot be taken into account explicitly. The existence of such events is now widely accepted but what is still unknown is their importance at the seasonal scale and at global scale, which depends of their frequency in a given convective area. Also unknown is their relative quantitative impact with respect to large-scale mechanisms such as the cold trap. Aiming at a deeper investigation of the mechanism of hydration of the lower stratosphere by convective overshoots, a series of simultaneous water vapour, particle and ozone measurements has been carried out within a SCOUT-AMMA (Stratosphere-Climate Links With Emphasis On The UTLS – African Monsoon Multidisciplinary Analysis) campaign in August 2006 from Niamey, Niger (13.6° N) in West Africa (Cairo et al., 2009). This is a location of frequent overshoots during the monsoon season according to Liu and Zipser (2005).

Although during SCOUT-AMMA the set of water measurements by several types of instruments onboard different platforms provides an unprecedented documentation of the UTLS tropical water vapour distribution, they are not sufficient to provide a full picture of the relative impact of the different processes affecting this distribution in the tropics. However, the modelling approach, validated by this set of measurements, can be

used to complement these observations and to quantify the impact of these processes. In particular three-dimensional limited-area meteorological models (mesoscale models) with fine resolution, down to cloud resolving simulations can solve explicitly deep convection, and are able to represent in a consistent manner dynamical and microphysical processes responsible for the water amount in the LS. Quantifying precisely the impact of single overshooting deep convection events on the LS hydration is a key step before those results can be extrapolated at a wider scale and before parameterizations of overshoots can be developed for global models. The relative impact of overshoots with respect to the freezing/drying mechanism in the TTL on the water budget could be then estimated. Recent studies (Schiller et al., 2009; James et al., 2008; Fueglistaler et al., 2005) conclude that the slow ascent leading to freezing/drying in the cold trap is predominant, trajectory calculations roughly explaining the water distribution above the tropical tropopause. Nonetheless Schiller et al. (2009) also conclude that the impact of overshooting convection is non negligible at local scale up to 420 K. Another important argument showing that the cold trap cannot explain alone the amount of water entering the tropical stratosphere concerns the trends of the tropical tropopause temperature: a decrease by  $0.5 \text{ K decade}^{-1}$  reported in Siedel et al. (2001) should lead to an enhancement of the dehydration at the tropopause, and thus to a decrease of the water amount in the stratosphere. This is not consistent with the observed trends of water vapour (Oltmans et al., 2000), suggesting that other mechanisms must play a significant role.

Only few estimations of the impact of single overshooting events are available in the literature and correspond to different areas of the globe – Maritime continent for Chemel et al. (2009), Brazil for Chaboureaud et al. (2007) and Grosvenor et al. (2007) – and different models, leading sometimes to different impact at local scale. The present work focuses on Africa, an area known for the relatively high frequency of overshoots according to Liu and Zipser (2005), and where no estimation of LS water injection by overshoot has been provided by CRMs up to now. Thus the aim of this study is to quantify the amount of water injected into the stratosphere by two different overshooting events in Africa: the first one is composed of several cloud clusters, the second one is

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

a well organised MCS. We also aim at complementing the seldom estimations already available in the literature which are necessary before parameterization of overshoots in global models can be developed.

The present work is based on the use of the three-dimensional regional and cloud resolving model BRAMS (Brazilian Regional Atmospheric Modelling System) (Freitas et al., 2009). It will be used to examine how a single overshooting deep convective system affects the water content of the TTL at a local scale in Africa. Here we study two overshooting cases observed in Niger and Chad during SCOUT-AMMA (Cairo et al., 2009), a European funded campaign part of the SCOUT-O3 programme, synchronous with the international AMMA project (Redelsperger et al., 2006): the case of 4 August 2006 over Southern Chad that likely influenced the water vapour balloon-borne measurements of 5 August from Niamey (Khaykin et al., 2009, hereafter K2009), and the case of 5 to 6 August 2006 of a mesoscale convective system (MCS) generated over central Niger which later propagated toward Niamey and Burkina Faso.

The paper is organized as follows. Section 2 gives an overview of the convective events. Section 3 describes the numerical tool used in this study as well as the experimental data. Section 4 discusses the ability of the model to simulate the observed cases. Section 5 is dedicated to the calculation of water vapour transport to the lower stratosphere by overshooting convection and its impact at local/regional scale in the LS. The conclusions are given in Sect. 6.

## 2 Overview of the observation of the convective event

Two cases of overshooting convection are presented in this paper. Both of them occurred in a relatively active period. The first one took place over Southern Chad on 4 August 2006 and is referred hereafter as the “Chad case”. The second case occurred in Air (central Niger) on 5 August 2006. In the following, it will be referred as the “Air case”.

### Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 2.1 Description of the Chad case

Balloon borne water vapor measurements of 5 August 2006 from Niamey military airport (13°29' N, 02°19' E) by micro-SDLA onboard the “water vapor flight no 1” launched at 18:40 UT and by FLASH-B launched at 18:52 UT are shown in Fig. 1. Micro-SDLA is a tunable diode laser spectrometer devoted to the in situ measurement of H<sub>2</sub>O, CH<sub>4</sub> and CO<sub>2</sub> by infrared absorption spectroscopy (Durry et al., 2004). Three near-infrared telecommunication-type InGaAs laser diodes are connected by means of optical fibres to an open multi-path optical cell providing an absorption path-length of 28 m. The laser beam is absorbed by ambient gas molecules as it is bouncing back and forth between the cell mirrors. CH<sub>4</sub> is monitored at 1.65 μm, CO<sub>2</sub> at 1.60 μm and H<sub>2</sub>O at 1.39 μm using the differential detection technique. The payload also includes pressure and temperature sensors. The accuracy of the H<sub>2</sub>O measurements within 160 ms is 5%. Micro-SDLA was already flown successfully in the tropics during the HIBISCUS campaign (Marécal et al., 2007). FLASH is a Lyman- $\alpha$  hygrometer. The instrument as well as its measurements performed on 5 August 2006, 12 min later than micro-SDLA is described in K2009. Although the fine structures of the micro-SDLA profile shown in Fig. 1 are qualitatively consistent with FLASH observations on the same day, micro-SDLA shows a still unexplained dry bias. We suspect this artifact could be due to a mis-adjustment of the optical multipass cell so that a wrong value for absorption path length was used in the concentration retrieval process for the water vapor channel.

The study of this case was motivated by a layer of enhanced water vapour around 17–17.5 km altitude, seen by both instruments in Fig. 1. The enhancement of water vapor with respect to the background value in this layer is about 2 ppbv for both instruments, and the corresponding mixing ratios are 8 ppmv for FLASH and 7 ppmv for micro-SDLA. This layer is located significantly above the tropopause (~16.5 km). Using backward trajectories based on ECMWF operational analysis winds, K2009 have related this enhancement to overshooting convection from deep convection over Southern Chad on 4 August 2006, 14:30 UT.

### Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Water vapor budget  
associated to  
overshoots in the  
tropical stratosphere**

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Figure 2 shows a series of Meteosat Second Generation (MSG) IR images in the 10° E to 30° E and 0° N to 20° N domain describing the time evolution of the convective activity in the Sudan/Chad area for this “Chad case”. The convective activity begins at 10:30 UT on the Sudan-Chad border. It develops further in a cluster of several cloud systems, South-East of Chad at 12:00 UT (Fig. 2b). At ~12:14 UT, the lidar instrument CALIOP onboard the Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observation (CALIPSO) satellite scanned the convective cloud cluster (not shown). The CALIPSO measurements highlight that the cloud top of the cloud cluster is typically at 16.5 km in the latitude range 10° N to 13° N and that some parts of this cloud top are classified as “stratospheric features” in the version 2.0.1 of the feature mask products. Although the class “stratospheric features” is not always unambiguous (see CALIPSO Quality Statement Lidar level 2 vertical feature mask at [http://eosweb.larc.nasa.gov/PRODOCS/calipso/Quality\\_Summaries/CALIOP\\_L2VFMPProducts\\_2.01.html](http://eosweb.larc.nasa.gov/PRODOCS/calipso/Quality_Summaries/CALIOP_L2VFMPProducts_2.01.html)), CALIOP measurements confirm that at about 12:12 UT the cloud top is very close to the tropopause. At 14:30 UT (Fig. 2c), the convective area is composed of an East-West band of clouds (from 22° E to 16° E; ~11° N). North-East of this band, there is another cloud system area that joins the first one while it is decaying. East of the Chad/Sudan border, cluster of several convective clouds are growing. This is the time when K2009 identify overshooting convection from the 11° N East-West band of cloud from brightness temperatures analysis. They identified from a trajectory analysis that the LS air coming from the overshoot moved toward the Niamey area and was sampled by the FLASH-B instrument above Niamey (Fig. 1). At 17:00 UT (Fig. 2d), the 11° N East-West band spreads South and North while convective activity develops on the Eastern to North-Western part of it. At 20:00 UT (Fig. 2e) the convective activity is mainly around 17° E, 12.5° N, coming both from the North-East and from the former 11° N band that join together.

To identify the location and time of the overshoots associated with the convective activity in the Southern Chad area we use a detection method of overshoots from Meteosat Second Generation (MSG) observations. This method (Schmetz et al., 1997) is

---

**Water vapor budget associated to overshoots in the tropical stratosphere**X. M. Liu et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

based on the brightness temperature difference (hereafter BTM) between the 6.2  $\mu\text{m}$  and the 10.8  $\mu\text{m}$  channel. The first channel is sensitive to the water vapour emission in the warmer temperature of the lower stratosphere with respect to the adiabatically cooled overshooting turret temperature. The BTM threshold for the detection of overshoot is set to 3 K as in Chaboureau et al. (2007) and K2009. In a study of deep convection over the Indian Ocean, Roca et al. (2002) use a larger BTM threshold of 5 K. However, it has been checked for a specific overshoot observational case of SCOUT-AMMA in the afternoon of 23 August 2006, that the MSG BTM was of about 3.2 K while the MIT radar at Niamey sampled a convective cell reaching 18.3 km well above the tropopause ( $\sim 16.5$  km). This proves that the threshold of BTM=3 K for the detection of overshoot is realistic in this region of Africa. Figure 3 shows the MSG pixels with BTMs higher than 3 K for the 12:15 UT to 19:15 UT with a 15 min timestep period. It shows that the overshooting convection activity mainly occurs in the 11° N East-West band of clouds, and moves with time westward. 14:30 UT is the time when K2009 identify 23 overshooting pixels in this area, to be of the same air mass as the one sample by FLASH 30 h later above Niamey. From 15:45 UT, overshooting pixels appear in the North of this band (typically 12° N, 20.5° E). From 17:00 UT, an area of overshoot is given by MSG observations in West Sudan (12° N, 23° E) and corresponds to a large red area in Fig. 2d. From 19:00 UT, another area of overshooting pixels appears around 12.3° N, 17.8° E. It correspond to the high convective activity shown in Fig. 2e coming both from the spreading of the former East-West cloud band and the Northern cloud cluster.

To summarize, the period after 12:00 UT on 4 August 2006 in Southern Chad is a period of severe deep convection including overshoots in several subareas: the 11° N East-West band of clouds at 14:30 UT, the area in Sudan close to the Chad border at 17:00 UT, and later the North-West area at 20:00 UT. Satellite images (Fig. 2) show that the different cloud bands interact with each other to create a large cloud cluster.

## 2.2 Description the Air case

A mesoscale convective system was observed East/North-East of Niamey at the time of the balloon-borne measurements on 5 August 2006. Figure 4 shows the time evolution of this “Air case” MCS as seen by MSG. Convection triggers in Air mountain area (centre Niger) at about 13:30 UT (Fig. 4a) in the 8.5° E, 17° N to 19° N area. Another active cloud area is observed close to the Niger/Nigeria/Chad border. The Air system grows and extends to 15° N and 7° E on 5 August (Fig. 4b) becoming an organized Mesoscale Convective System (MCS) at 18:00 UT. At the same time, the Niger/Nigeria cloud system grows and propagates westward. The Air MCS grows and moves westward until 18:00 UT (Fig. 4c) when it reaches its top activity. At the same time the Niger/Nigeria/Chad cloud cluster has decayed and split into two different parts: the first one stays at the Chad Niger border while the second, at the Niger/Nigerian border (10° E), continue to propagate westward. It later joins the Air MCS while the latter propagates south-westward towards Niamey (Fig. 4d). The total system reaches Niamey in the morning of 6 August and will later move to Burkina Faso while decaying.

Using the same MSG brightness temperature difference technique as for the “Chad case”, we have checked the occurrence of overshooting activity of the Air MCS. Figure 5 shows the overshooting pixels in central Niger for the period from 14:45 UT to 17:30 UT. It shows that the Air area has experienced overshooting convection during this period. The overshooting area moves westward (slightly northward) with time, which is compatible with the westward propagation of the MCS during its maximum of activity. At 15:30 UT, 16:15 and 16:30 UT the number of pixels with  $BTD > 3\text{K}$  is 13, 14, and 16, respectively. These numbers are smaller than for the Chad case: at 14:30 UT, K2009 have reported a total of 23 pixels with  $BTD > 3\text{K}$  (see Table 2 of K2009 for details about the Chad Case BTDs). Table 1 summarizes the time evolution of the overshoot size in pixels. It shows that the most intense overshooting activity is between 15:30 UT and 16:30 UT with a maximum BTD of 4.8K which is close to the overshoot criteria proposed by Roca et al. (2002).

### Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

In the following we study both the Chad case and the Air case with a mesoscale model. It will give an opportunity to compare the impact of different types of system on the lower stratosphere hydration by overshooting convection. The next section describes the modelling tools and model setup used in this study.

### 3 Modelling tools and simulation setup

The Brazilian Regional Atmospheric Modelling System (BRAMS) model is a mesoscale model tailored to the tropics. It is the Brazilian version of the RAMS model (Cotton et al., 2003) of the University of Colorado/ATMET. BRAMS (Freitas et al., 2009) was developed at CPTEC (<http://brams.cptec.inpe.br/>). It is designed to simulate atmospheric circulations at different scales from large regional simulations down to large eddy simulations. It includes a full set of parameterizations for surface processes, radiative scheme, dynamics, sub grid scale convection and microphysics. BRAMS differs from RAMS in improvements concerning cumulus convection parameterization, soil moisture initialization and surface scheme. BRAMS/RAMS offer the possibility to use nested grids, each grid communicating in a two way process with its mother or son grid. BRAMS was successfully used to simulate water vapour distribution in the tropical UTLS in a deep convective environment over Brazil (Marécal et al., 2007).

Two simulations were run using the BRAMS model for each of the case studies described in Sect. 2. For both simulations, the radiative scheme of Harrington (1997) was used for short and long wavelengths. The sub grid scale parameterization of shallow and deep convection of Grell and Devenyi (2002) was chosen. The setup for microphysics makes use of the two moment scheme (Meyers et al., 1997). Seven types of hydrometeors are taken into account in the model: liquid cloud droplets and rain for liquid particles, and pristine, snow, hail, graupel and aggregates for ice particles. In this scheme, both hydrometeor concentrations and mass mixing ratios are prognostic variables of the model, assuming that each hydrometeor size distributions follows gamma function. In the simulation the shape parameter of the gamma function is assumed to

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



be 2 for each category of hydrometeors as successfully used with BRAMS in Pénide et al. (2010) for AMMA case simulations. Although considered as a bulk scheme, this relatively detailed scheme is particularly suitable for the quantification of water transport across the tropopause by overshooting convection, in which ice particles are supposed to play an important role (Pommereau and Held, 2007; Grosvenor et al., 2007; Chemel et al., 2009): as a matter of fact, the number of hydrometeors is higher than for other CRMs already used for studying the role of overshooting convection on the LS hydration. The Meso-NH model used in Chaboureau et al. (2007) and the ARW model used in Chemel et al. (2009) has 5 types of hydrometeors (including 3 classes of ice particles). Furthermore double moment schemes used in BRAMS and ARW are said to be more precise than single moment schemes as in Meso-NH. These differences among models are likely to induce differences in the simulated amount of water injected by overshooting convection.

Global ECMWF (European Centre for Medium-range Weather Forecasts) 6-hourly analyses were used in this study, both for the initialization and the nudging at the lateral boundaries of the larger BRAMS domain (Grid 1). The set of analyses used are the special AMMA reanalyses done at ECMWF (Augusti-Panareda et al., 2009), which take into account all the water vapour sonde measurements of the AMMA campaign in West Africa. Sensitivity tests have been performed using operational analyses instead of AMMA reanalyses and showed a significant improvement of the BRAMS results using the reanalyses in term of system formation, severity, lifetime and propagation.

### 3.1 Specific setting for the Chad case

The 4 August 2006 convective system simulation includes three nested grids as shown in Fig. 6a. The coarse grid (Grid 1) covers a domain from  $\sim 8.20^\circ$  E to  $25.94^\circ$  E in longitude and from  $2.39^\circ$  N to  $17.78^\circ$  N in latitude including the Atlantic Ocean in the South-West corner. Its horizontal grid spacing is 20 km. A finer resolution second grid (Grid 2), including the area where most of the deep convection was observed, extends from  $15.18^\circ$  E to  $24.28^\circ$  E in longitude and from  $8.88^\circ$  N to  $14.68^\circ$  N in latitude with

4 km grid spacing. The finest grid has 1 km grid spacing and covers a 214 km×110 km domain from 19.98° E to 21.93° E in longitude and from 11.15° N to 12.19° N in latitude focused in the area of the East-West cloud band in its mature phase in order to properly take into account the overshoot area highlighted by K2009. The topography is also reported in Fig. 6, indicating within Grid 3 a region with hills higher than 800 m, and within Grid 2 an area in Sudan with topography higher than 1300 m. For all these grids, the vertical coordinate is terrain-following height coordinate with 68 levels from the surface to 30 km altitude with 300 m spacing in the UTLS. In order to damp the gravity waves at the top of the domain, an absorbing layer of 5 km thickness was prescribed at the top boundary. The time steps used are 30 s, 10 s and 2 s for Grid 1, Grid 2 and Grid 3, respectively. The simulation was initialized at 18:00 UT on 3 August 2006 and ended on 5 August at 12:00 UT. Model outputs were saved every hour except during the most intense period of convection in the model from 4 August, 12:00 UT to 20:00 UT when model outputs were saved every five minutes. The choice of the model configuration was a trade-off between high resolution and computing cost.

### 3.2 Specific setup for the Air case

The Air system was also simulated with triply nested grids (Fig. 6b). Grid 1 covers a domain from 7.942° W to 22.05° E in longitude and from 2.02° S to 26.10° N in latitude with 20 km grid spacing. This includes a large part of West Africa. A significant part of Grid 1 includes the Atlantic Ocean in order to better account for the monsoon flux. The domain of Grid 2 is from 5.80° E to 12.14° E in longitude and from 13.16° N to 20.36° N in latitude. This grid, with a relatively fine horizontal resolution (4 km grid spacing) covers the Air Mountains with height higher than 1600 m in the central Niger where the MCS under the scope of our study developed, as shown by the topography in Fig. 6b. Grid 3 has 1 km grid spacing and covers a 106 km by 118 km domain from 7.84° E to 8.83° E in longitude and from 15.36° S to 16.45° N in latitude, is located at the South of the Air Mountains. The location of Grid 3 is chosen in the area where the modeled convection is the most severe from a previous simulation with two grids only.

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The vertical coordinate and the time steps are the same as in the Chad simulation. The simulation was initialized at 00:00 UT on 5 August 2006 and was run for 24 h saving model outputs every hour except during the period 18:00 UT to 00:00 UT when model outputs were saved every 5 min to follow the detailed evolution of the overshoots.

## 4 Validation of the simulations

In this section, we evaluate the results of the two simulations against observations. We focus on the ability of the model to reproduce the observed deep convective systems/clusters at the regional scale in which the overshoots are embedded and also the observed overshoots locally. We first compare BRAMS outputs with TRMM (Tropical Rainfall Measuring Mission) estimates of surface rainrates. The TRMM dataset used was produced by the 3B42 algorithm (Huffman et al., 2007, <http://trmm.gsfc.nasa.gov>). It is 3-hourly and  $0.25^\circ \times 0.25^\circ$  resolution. The comparison is done using the Grid 1 results which include the mesoscale convective systems affecting the overshoot area. Moreover Grid 1 resolution is close to TRMM products (20 km for Grid 1 and  $0.25^\circ$  for TRMM). Secondly, we compare the model results to the MSG overshoot observations. This is done using the simulation fields from the finest resolution grid (Grid 3 with 1 km horizontal resolution). Overshooting convection can only be simulated when convective dynamical fields are explicitly resolved in the model, i.e. using a resolution  $\leq 1$  km, as shown in previous studies (Grosvenor et al., 2007; Chaboureau et al., 2007). Note that the observed overshoots are far from the AMMA/SCOUT-AMMA campaign zone. Therefore it was not possible to make comparisons with balloon borne or aircraft measurements close to the overshoots.

### 4.1 Chad case

Figure 7 shows comparisons of surface rainrates (expressed in  $\text{mm h}^{-1}$ ) between TRMM (Fig. 7a, c, e) and BRAMS model outputs (Fig. 7b, d, f) for 15:00 UT (between

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

13:30 UT and 16:30 UT), 18:00 UT (between 16:30 UT and 19:30 UT), and for 21:00 UT (between 19:30 UT and 22:30 UT). Thus rainrates are accumulated over three hour periods. The focus being on the observed overshoots, we have displayed the model and observed surface rainrates over the sub-domain of Grid 1 including the convective systems that are likely to affect the overshoots. Note that this sub-domain is larger than Grid 2 area and includes the whole Grid 2 domain. During the period of interest, several convective systems develop and interact while advected as illustrated by the time evolution of the many features appearing in the TRMM surface precipitation fields (Fig. 8a, c, f) and already discussed on MSG observations (see Sect. 2). The features of interest have been indentified in Fig. 8a, c, e by grey and green color contours.

The first one is highlighted by the green ellipse corresponding to an area of precipitation in the North-East of the domain close to the Sudan-Chad border. This rainband hardly moves between 13:30 UT to 22:30 UT. Only the intensity of the individual convective cells varies with time. BRAMS fairly well simulates the position of this rain area (Fig. 7b). However, BRAMS overestimates the intensity of this rain area with maxima over  $10 \text{ mm h}^{-1}$  while the maxima from TRMM are usually between 5 and  $10 \text{ mm h}^{-1}$ .

The second important precipitation feature is highlighted by a grey domain. At 15:00 UT both BRAMS and TRMM show a East-West band of precipitation at  $\sim 11\text{--}11.5^\circ \text{ N}$  with the same intensity and distribution. This area corresponds to the zone where the K2009 Chad overshoot has been highlighted and corresponding to the purple to blue overshooting pixel zone in Fig. 3. South of this band, BRAMS simulates a local maximum of the same intensity as the TRMM estimation. BRAMS only fails in the North of the domain where an isolated low precipitating cell seen by TRMM is simulated by BRAMS much further West in a larger area. Later at 18:00 UT (Fig. 7c) the TRMM products show that the precipitation zone spread North and South (Fig. 7c). This is also well captured by BRAMS (Fig. 7d), both in location and intensity. The Northern cell in the West of the domain given in TRMM products intensifies, as well as the cell located more West in the BRAMS simulation which is at the West edge of the Grid 2 domain. At 21:00 UT the area of precipitation in the centre of the domain at  $11^\circ \text{ N}$ ,

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

19° E for TRMM and 11.5° N, 20° E for BRAMS is decaying. Concurrently, further West, both for TRMM and BRAMS, there is an intensification of the precipitation from North to South. A very intense area in the North-West is seen by TRMM (Fig. 7e) and corresponds to the orange overshooting pixels in Fig. 3. A detailed analysis of the MSG infrared images shows that this area is initially formed from an interaction between the Northern precipitating cells and the 11° N precipitating band. In the BRAMS simulation (Fig. 7f), intensification of the precipitation in the West from North to South is also computed, though the location is too far West.

To validate the overshooting activity in BRAMS with respect to the MSG BTDO observations, Fig. 8 shows the Grid 3 points for which there is a significant ice amount (mass mixing ratio above  $0.05 \text{ g kg}^{-1}$ ) above the tropopause (380 K level) during the time of overshoots in BRAMS. Dependence with time is indicated by the color scale, chosen the same as in Fig. 3. Note that the horizontal resolution of the model in Grid 3 (1 km) is higher than the MSG resolution ( $\sim 3.3 \text{ km}$ ). It shows that the overshooting activity begins at  $\sim 13:15 \text{ UT}$  and ends at  $14:45 \text{ UT}$ . Except for a few overshooting points at  $11.6^\circ \text{ N}$  from  $\sim 13:15 \text{ UT}$ , most of them are along the  $12^\circ \text{ N}$  latitude and move with time from East to West (from  $13:15$  to  $14:15 \text{ UT}$ ). This tendency both in time and location is well in agreement with the BTDO overshooting pixels shown in Fig. 3, though the observed overshooting pixels are located along the  $11.1^\circ \text{ N}$  latitude. This good result must be balanced by the fact that ice which is calculated to be produced in overshoots at  $14:00 \text{ UT}$  is advected westward in the LS before it sublimates totally. Thus not the totality of the grid points with ice mixing ratio higher than  $0.05 \text{ g kg}^{-1}$  should be considered as overshooted grid points but as grid points with ice of overshooting origin. On the other hands, it is possible that the BTDO signature is also sensitive to advection of sublimating ice injected by overshoot when the cloud system below remains very active. Due to the limited size of the Grid 3 domain to save computing resources, the overall overshooting activity observed by MSG in all the southern part of Chad cannot fully be reproduced by BRAMS. We chose to restrict our goal to reproduce the overshoots occurring in the time range and the location pointed out by K2009, shown to

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

influence the FLASH-B/ $\mu$ -SDLA measurements.

Figure 9 shows a vertical cross section of total water along the 12° N latitude in the BRAMS Grid 3 domain at 14:15 UT. It highlights that the simulated cloud system overshoots the tropopause up to 17.8 km, above the 400 K level. Note that the position and the time of this overshooting plume is in very good agreement with the overshoot highlighted by K2009.

To summarize, BRAMS succeeds in reproducing the main features of the observed precipitation, except in the West part of the domain from 18:00 UT where BRAMS compute precipitation which is seen by the satellite much further East. The model provides good results in the center of the domain for the East-West band of precipitation where the K2009 overshoots are highlighted. The overshooting activity in the time range from 12:00 UT to 15:00 UT is in good agreement with the observations.

## 4.2 Air case

As for the Chad case we validate the simulation of the Air MCS of 5 August 2006 by comparing model outputs to TRMM rainrates and to BTD from MSG. Figure 10 shows comparisons of surface rainrates (expressed in  $\text{mm h}^{-1}$ ) between TRMM (Fig. 10a, c, e) at 15:00 UT, 18:00 UT and 21:00 UT, respectively, and BRAMS model outputs (Fig. 10b, d, f) for 14:00–17:00 UT, 17:00–20:00 UT, and 20:00–23:00 UT respectively, thus 30 min time-shifted with respect to the TRMM products. BRAMS outputs are expressed in the Grid 1 resolution in the whole geographical domain to be comparable to the TRMM products resolution. Panel 10a shows a large area of precipitation in central Niger with a maximum at  $\sim 17^\circ \text{N}$ ,  $8.5^\circ \text{N}$ . Another area of precipitation is sampled in the South-East of the domain close to the Niger/Nigeria border. The corresponding BRAMS simulation (Fig. 10b) shows a smaller area of precipitation in central Niger because the convective system in BRAMS develops later than in the observations. However the maximum of precipitation is located at the same place with a similar intensity. It is worth noticing that the convection triggers in a region of mountains as shown in Fig. 6. Sensitivity tests on orography parameters used in the simulation show

### Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



that orography play an important role in the convection triggering. The area of precipitation close to the Nigeria border is also reproduced by BRAMS, although it is located more West than the observations.

Around 18:00 UT the observed and modelled rainrates both indicate an intensification of the MCS North to South (close to 8° E and between 16° N and 18° N), and a westward propagation. The core of high rainrates is comparable in both cases. The area of precipitation in the South-East slightly propagates westward with a core of high rainrates similar in both cases but more West in BRAMS, as at 15:00 UT. Around 21:00 UT both panels 10e (TRMM) and 10f (BRAMS) show a south-westward propagation of the MCS but slower in the model. This is due to the fact that while propagating westward, the BRAMS MCS gets out of Grid 2 and enters in the Grid 1 domain where convection is determined by a subgrid scale parameterisation, coarsening the key parameters which determine the MCS lifetime and propagation speed. The Niger/Nigeria border area of precipitation propagates westward in TRMM and BRAMS. Due to the south-westward propagation of the MCS, this area converges to the MCS and later forms a unique cloud system, both shown in satellite observations and modelling. The TRMM outputs at 21:00 UT show that the activity of the MCS decays as highlighted by a maximum rainrate lower than 10 mm h<sup>-1</sup> in the core of the system while BRAMS still computes a very active core. This is partially due to the fact that BRAMS outputs are 30 min later than the TRMM outputs and that the observed MCS, after a decaying phase, re-intensify later from 23:00 UT. This time is taken into account into BRAMS in the evaluation of the rainrates but not taken into account in the TRMM outputs of the corresponding period which ends at 22:30 UT. This also could be due to the fact that the modeled MCS is late with respect to the observed one, and thus is still active at that time.

Figure 11 illustrates the overshooting activity as seen by the BRAMS simulations within Grid 3. It shows latitude/longitude grid points having ice mass mixing ratio higher than 0.1 g kg<sup>-1</sup> above the 380 K level in the time range between 18:30 UT and 21:15 UT. The threshold of 0.1 g kg<sup>-1</sup> is twice the one chosen for the Chad to make Fig. 11 more

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

readable. Figure 11 uses the same color timescale as in Fig. 5 but shifted in time by 3 h and 15 min with respect to the BTD overshooting pixels observations. This is done to account for the time shift of the model MCS compared to observations. Figure 11b shows that the modeled MCS injects ice above the tropopause during a time range similar to the overshooting observations, but later than the observed MCS. The longitude range of the modeled overshoot is also comparable with the observed one in the first 75 min (Fig. 11a). Contrarily to the Chad case, the displacement of the overshooting signal in BRAMS is not only due to the horizontal advection of ice injected by the overshoot, since there are several stratospheric penetrations (first at 8.35° E, and later at 8.18° E) in our simulation. After this time, BRAMS still computes ice injection above the tropopause but the geographical distribution of overshoots differs from the observations. Overshoots are computed in the South-West part of Grid while the observations only show a displacement of the overshooting pixels with time from East to West-North-West. Nonetheless the overshooting time evolution is very well reproduced by BRAMS in the first 75 min of the modelled overshooting period (see results in Fig. 11a) with the same direction of displacement, highlighting that during this period, the overshoots in BRAMS are realistic even if late with respect to the observations. Contrarily to Chad case, the displacement of the overshooting signal with time is not only due to the horizontal advection of lower stratospheric ice particle injected by overshoots since for the Air case, there are several stratospheric penetration at 19:10 UT and at 19:45 UT. Examination of the cloud top during the overshooting period shows that overshoots can reach 18.5 km (Fig. 12).

To summarize, the key parameters of the 5 August 2006 Air MCS (location, propagation direction, intensity of precipitation, overshooting activity) are generally well reproduced by the BRAMS simulation. Nevertheless the modeled MCS develops later than the observations and the overshooting activity is overestimated at the end of the simulation. The end of the simulation will not be taken into account in the calculation of the water fluxes across to tropopause presented in the next section.

---

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## 5 Overshooting convection and cross-tropopause water vapour transport

In Sect. 4 we have shown that the simulations performed for the two overshoot case studies are in good agreement with observations. This allows us to calculate water budget across the tropopause. These two overshooting case studies are interesting to compare since they have different characteristics (organization, formation process, duration, size, intensity).

The aim of this study is to study the impact of African overshooting convection on the water amount in the lower stratosphere, to quantify the amount of water injected into the stratosphere by overshooting convection and to complement the seldom estimations already available in the literature. The two cases described above give an opportunity in the same study to compare the impact of two different overshoots by their lifetime and area. This might be helpful to later determine a size/duration dependence of the amount of ice injected by such extreme events.

Since the tropopause height is locally disturbed by deep convection, especially during convective events when waves generated by convection can impact the tropopause height, calculating the water flux across the tropopause needs to account for this variation. This tropopause height variation with the convective activity is highlighted in Figs. 9 and 12. Therefore the isentropic vertical coordinate should be used to locate the tropopause. Here we use the 380 K level (cf. Holton et al., 1995).

To compute the water flux across isentropic levels which are potentially moving, the vertical velocity of the  $\theta$ -levels needs to be determined, so that the relative vertical wind speed with respect to the  $\theta$ -levels displacement can be used. To compute the vertical speed of the  $\theta$ -level, the variation of height with time of a  $\theta$ -level is calculated. For this, a time-step of 5 min is used, justifying why the time frequency of BRAMS outputs are of 5 min during the overshooting period.

The total water fluxes across the  $\theta$ -level are computed within the third grid of simulation through grid mesh with positive relative wind speed. The same technique has been applied by Chaboureaud et al. (2007), but these authors have used a time-step for

### Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

model outputs of 10 min. instead of 5 min. in our case. All variables needed for the flux calculation are interpolated in a new grid using potential temperature as vertical coordinate.

## 5.1 Chad case

5 In order to estimate the impact of overshooting convection on the water composition at mesoscale, Fig. 13 shows a Grid 3 scatter plot of UTLS total water at different times of the simulation: initial state, overshooting period, just after the overshooting period, and after convection. At 14:15 UT (panel b), the range of total water values is widely spread above the 380 K level, with some values over 100 ppmv. This is due to the very  
10 low altitude of the 380 K at that time in the vicinity of overshooting plume as shown in Fig. 9. Values as high as 50 ppmv have already been observed by the FISH instrument onboard the Geophysika M55 aircraft at 400 K above the Hector cloud in the maritime continent (Corti et al., 2008; Chemel et al., 2009) on 30 November 2005. During AMMA in Africa, lower values are reported but reach sometimes 10 ppmv in the lower  
15 stratosphere (Schiller et al., 2009). In Fig. 12b purple crosses highlight grid points in the overshooting plume area (latitude from 11.95° N to 12.06° N and longitude from 20.9° E to 21.1° E). They show that most of the high water mixing ratios in the stratosphere are due to this overshoot, enhancing the water amount of 2 to 3 ppmv above the 400 K level.

20 Important to notice is that the mean profile at 14:15 UT is higher at any altitude than the initial profile, showing a mean hydration at all level. The same conclusion is reached when comparing the pre-overshoot mean profile at 12:00 UT on 4 August and the mean profile at 14:15 UT. After the overshooting activity (panel c) stratospheric water values are less scattered than during the overshoot. But there is still a local enhancement of 2 ppmv with respect to the mean profile between 380 and 390 K. A  
25 striking feature of panel b and c is the scatter of the very low water mixing ratios in the TTL below the tropopause, showing that deep convection can also lead to dehydration in the upper troposphere. This tendency is visible in the mean profile at 15:55 UT where

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



it is lower than the initial state in the TTL from 360 K to 370 K. It is not seen at 14:15 UT since deep convection leads to a local large hydration that influences the mean profile. When convection disappears from Grid 3, there is almost no variation of the water amount in the stratosphere, even if the mean profile remains significantly higher than the initial one. Above the tropopause, the mean profile is typically 0.21 ppmv at 400 K to 0.67 ppmv at 380 K higher than the pre-overshoot mean profile at 12:00 UT between 380 K and 400 K. This is higher than the estimation of the 0.12 ppmv hydration of the Hector cloud by the ARW model (Chemel et al., 2009) in the maritime continent and comparable with the 0.26 ppmv of the idealized case of Grosvenor et al. (2007) over Brazil but significantly lower than the 5.6 ppmv of the Hector cloud simulation by running the Unified Model (Chemel et al., 2009). Let's note however that the above-mentioned estimations are from other areas and continents.

Water fluxes of the Chad case are calculated for the most active period in Grid 3 from 12:00 UT to 16:00 UT on 4 August 2006. Figure 14 presents the times series of upward total water flux through isentropic surface from 375 K to 400 K. For all levels except 400 K several peaks are found at 13:50 UT, 14:00 UT and 14:15 UT. They correspond well to the time of overshooting plumes in Grid 3, as shown in Fig. 9 for 14:15 UT. The overshooting signal is not obvious at the 400 K level. This can be explained by the fact that the overshooting plumes shown in Fig. 9 cross the 400 K level in a more limited area and that the ice mixing ratios are lower, in contrast to lower  $\theta$ -levels. Focusing on the tropopause level (380 K), the maximum instantaneous flux is of  $3.77 \text{ ton s}^{-1}$ , which is 3 times less than for the 370 K (not shown) and 1.66 times less than for the 375 K level. The order of magnitude of the 380 K flux of a few  $\text{ton s}^{-1}$  is compatible with that reported in Chaboureau et al. (2007). However our maximum estimation is lower. In Chaboureau et al. (2007) the 380 K instantaneous flux is  $8 \text{ ton s}^{-1}$  for water vapour alone and thus is expected to be even higher for total water. This might be due to the use in our study of a double moment microphysical scheme and a budget calculation timestep of 5 minutes instead of a single moment scheme and a 10 min timestep for Chaboureau et al. (2007). Marécal et al. (2009) have shown in a convective cell

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

simulation that the amount of ice of the cell core was higher for a single moment microphysical scheme than for a similar simulation using a double moment microphysical scheme. At the same time, note the instantaneous flux of more than  $1 \text{ ton s}^{-1}$  across the 395 K level. After 14:15 UT, the instantaneous fluxes decrease rapidly to tend to the same value at levels higher than 375 K when overshoot activity has disappeared in Grid 3, to tend later to zero.

Integrated on the 4 h period encompassing the overshooting event and on the surface area of Grid 3, the mass of total water crossing the  $\theta$ -levels are given in Table 3. The corresponding numbers per unit of time are also given. For the 380 K level which corresponds to the tropopause, a total mass of water of  $\sim 13 \text{ kton}$  has crossed this level during the 4 h period. This corresponds to a mean value of  $\sim 3300 \text{ ton h}^{-1}$  or  $0.91 \text{ ton s}^{-1}$ . This computation does not take into account potential removal of ice crystal by sedimentation. However, the total water relative humidity with respect to ice computed by BRAMS highlights generally unsaturated conditions ( $< 80\%$ ) in the LS except a local maximum of 110% at 14:20 UT at 17 km. These conditions provide sublimation of ice crystal before they can be removed from the lower stratosphere by sedimentation. Nevertheless, we cannot rule out that this mechanism leads to a significant loss of water in the stratosphere as it was suggested by Corti et al. (2008): the budget computed here should be seen as an upper limit.

Figure 15 shows the time evolution of the total water at the 390 K isentropic levels in the Grid 3 domain from the time of the most severe overshoot (14:15 UT). It shows that the hydrated signal is advected with the wind for one hour in Grid 3. While travelling westward, the signal spreads North and South and the maximum of total water decreases due to its horizontal diffusion, and possibly its vertical diffusion and sedimentation of larger ice particles. Figure 15 also shows that the ratio of ice to the total water decreases with time while the hydrated bulge travels westward. It varies at the maximum from  $\sim 98\%$  at 14:15 UT to  $\sim 15\%$  at 15:10 UT, showing that despite of slightly subsaturated conditions with respect to ice, ice particles remains in the lower stratosphere one hour after the overshoot. However, before going out of Grid 3, the

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

hydrated spot is still 3 ppmv higher than the background water content at the maximum (10.2 ppmv at 15:20 UT), 1h and 5 min after the overshoot, ~100 km away from it. Note that the maximum of total water decreases with time from the time of the overshoot to the time when the hydrated spot goes out of the Grid 3 domain. The decrease is fast immediately after the overshoot, and it is relatively slow long after the overshoot. The maximum value decreases from 10.8 ppmv to 10.2 ppmv between 15:10 UT and 15:20 UT. This indicates that a local enhancement of stratospheric water due to overshoot can travel relatively far from the overshoot. This is compatible with the K2009 hypothesis which explains that the local enhancement of water vapour above Niamey is due to the overshooting convection of the Chad system. However, once in Grid 2 (4 km horizontal resolution), the hydrated signal is lost rapidly. The maximum value at 15:55 UT is <2 ppmv higher than the background level, due to a larger horizontal and vertical mixing in the model due in the coarser resolution. Thus a high resolution is mandatory to properly transport the hydrated maximum far from the overshoot.

## 5.2 Air case and comparison with the Chad case

Similar calculations were carried out for the case of 5 August. The period of 18:00 UT to 19:50 UT has been used to compute water fluxes across different isentropic levels in the Grid 3 domain. We limit our calculation to this period in order to keep only the period when there is a good agreement between the MSG overshooting activity and the BRAMS overshooting activity (see Sect. 4.2). The results are given in Fig. 16 and flux values are summarized in Table 3. A two peak structure is shown in Fig. 16, highlighting two different short periods of intense overshooting convection. The first peak, at 19:10 UT is visible for all  $\theta$ -levels showing that overshooting convection directly injects ice particles above the 400 K, as shown in Fig. 12. At 380 K, this peak reaches a maximum of  $1.1 \text{ ton s}^{-1}$ , which is less than the peak described for the Chad case. The second peak, at the end of the calculation period, reaches a higher value of  $1.4 \text{ ton s}^{-1}$  that is still lower than for the Chad case. The difference is likely due to the smaller surface area of overshoots for the Air case. Note for the second peak that

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the overshooting signal is not seen at the 390 K level and above. This illustrates that this overshoot penetrates those level on smaller surface area that for the Chad case. The integrated fluxes during the whole period given in Table 3 highlight lower values per unit of time than for the Chad case, though the Air case is shorter. For  $\theta$ -levels in the stratosphere ( $\geq 380$  K), the Air case budget is typically 0.75 times the Chad budget. This shows the variability of the impact of overshooting convection on the amount of water injected in the stratosphere among the cases.

## 6 Conclusions

The aim of this study was to quantify the impact of two different overshooting events on the water budget in tropical tropopause region during the African monsoon 2006. One of these events was directly upstream of a pair of balloon flights from Niamey measuring water vapour in the UTLS in the frame of the SCOUT-AMMA campaign. During these flights a layer of enhanced water vapour of more than 2 ppmv was measured. Three nested grids regional down to cloud resolving simulations were conducted with the BRAMS model in order to achieve this goal. A double moment bulk microphysical scheme is used in our simulation. This study confirms previous estimation from model simulations that overshooting convection injects a large amount of water via ice crystals which later sublime rapidly in a sub saturated environment, while sedimenting. However, our estimation is lower than Chaboureau et al. (2007) who studied an overshoot over Brazil. This could be due to the use of a double moment scheme in BRAMS in contrast with the single moment scheme of meso-NH: single moment schemes are known to overestimate precipitation and the concentration of ice crystals with respect to double moment schemes which are more precise. For the case of 4 August 2006 over Southern Chad, a total amount of 13 kton is injected into the stratosphere leading locally to an enhancement of total water of 3 ppmv at 400 K and to mixing ratios higher than 100 ppmv at the tropopause. At a regional scale, significantly after the overshoot, the mean water amount in the Grid 3 domain above the tropopause is still enhanced

### Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**Water vapor budget associated to overshoots in the tropical stratosphere**

---

X. M. Liu et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

by 0.21 ppmv at 400 K to 0.67 ppmv at 380 K. This result is between the ARW model result of 0.12 ppmv (Chemel et al., 2009) or the Grosvenor et al. (2007) model estimation (0.26 ppmv) and the UM model (5.91 ppmv) result reported in Chemel et al. (2009) from a simulation of the Hector cloud in the West Pacific. This highlights the variability among mesoscale models to simulate the hydration of the lower stratosphere by deep convection, together with the variability of the impact of overshoots depending on the continent considered. But whatever this variability is, all these modeling studies conclude to a potential of overshooting convection to hydrate the LS.

After being advected for more than one hour and  $\sim 100$  km away from the overshoot, the hydrated area at the 390 K level is still 3 ppmv higher than the background water amount, showing that the impact of the overshoot can be seen far away from its location. Thus, it is likely that the water vapour enhancement of more than 2 ppmv sampled by FLASH-B and micro-SDLA on 5 August above Niamey is due to overshooting convection over South Chad 30 h earlier, as proposed by Khaykin et al. (2009).

Another more organised overshooting system was studied over Air and the associated water budget across the tropopause was computed. It shows a total mass of water injected in the stratosphere 3 times lower than for the Chad system. The difference is less but still significant when the result is expressed per unit of time:  $0.66 \text{ ton s}^{-1}$  for the Air case instead of  $0.91 \text{ ton s}^{-1}$  for the Chad case. This shows the variability of the impact of overshooting convection on the hydration potential of the stratosphere, which was not highlighted in previous studies.

The relative impact of overshooting convection with respect to the cold trap dehydration is not assessed yet. The impact of such mesoscale hydration at a larger space and timescale needs to be studied first, in order to reconcile the small scale and the larger scale  $\text{H}_2\text{O}$  transport pathways which are still highly debated. We propose as a first step to better estimate the variability of the impact of overshooting convection among a larger set of different cases. We also propose to check the impact of key parameters in the model such as the microphysical scheme or the model resolutions to put to the fore the reasons of the differences among models on the LS hydration. Concurrently, it

would be interesting in the future to compare water budgets of the same case but with different mesoscale models. From this, typical impact of overshoots could be obtained and be included in larger models to upscale the impact of overshooting convection on the whole tropical lower stratosphere.

5 *Acknowledgements.* This work was supported by the European integrated project SCOUT-O3 (GOCE-CT-2004-505390) and by the program LEFE/INSU in France (projects UTLS-tropicale and Tropopause 2009). The computer resources were provided by CRIHAN (Centre de Ressource Informatique de Haute Normandie) project no. 2008006 .This work was also granted  
10 access to the HPC resources of CINES under the allocation 20080125036 and 2009015036 made by GENCI (Grand Equipement National de Calcul Intensif). The TRMM data were provided by GSFC/DAAC, NASA. The CALIPSO/CALIOP image was distributed by Atmospheric Science Data Center, at NASA Langley Research Center. Meteosat Second Generation data were provided by the AMMA database and the Climserv database at Institut Pierre Simon Laplace (IPSL). One of the authors, Xiaoman Liu is funded by Région Champagne-Ardenne.  
15 G. Pénide and V. Giraud at Laboratoire de Météorologie Physique (LaMP) are acknowledged for their advices in the BRAMS setting. Finally, we are grateful to the micro-SDLA technical team of DT-INSU (N. Amarouche, F. Blouzon and J. Deléglise) for their investment during the SCOUT-AMMA campaign in Niamey.



20 The publication of this article is financed by CNRS-INSU.

## References

Agusti-Panareda, A., Beljaars, A., Cardinali, C., Genkova, I, and Thorncroft, C.: Impact of assimilating AMMA soundings on ECMWF analyses and forecasts, Weather Forecast., in

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



review, 2009.

Agusti-Panareda, A., Vasiljevic, D., Beljaars, A., Bock, O., Guichard, F., Nuret, M., Garcia Mendez, A., Anderson, B. E., Bechtold, P., Fink, A., Hersbach, H., Lafore, J.-P., Ngamini, J.-B., Parker, D. J., Redelsperger, J.-L., and Tompkins, A. M.: Radiosonde humidity bias correction over the West African region for the special AMMA reanalysis at ECMWF, Q. J. Roy. Meteorol. Soc., 135, 595–617, 2009.

Cairo, F., Pommereau, J. P., Law, K. S., Schlager, H., Garnier, A., Fierli, F., Ern, M., Streibel, M., Arabas, S., Borrmann, S., Berthelot, J. J., Blom, C., Christensen, T., D'Amato, F., Di Donfrancesco, G., Deshler, T., Diedhiou, A., Durry, G., Engelsen, O., Goutail, F., Harris, N. R. P., Kerstel, E. R. T., Khaykin, S., Konopka, P., Kylling, A., Larsen, N., Lebel, T., Liu, X., MacKenzie, A. R., Nielsen, J., Oulanowski, A., Parker, D. J., Pelon, J., Polcher, J., Pyle, J. A., Ravegnani, F., Rivière, E. D., Robinson, A. D., Röckmann, T., Schiller, C., Simões, F., Stefanutti, L., Stroh, F., Some, L., Siegmund, P., Sitnikov, N., Vernier, J. P., Volk, C. M., Voigt, C., von Hobe, M., Viciani, S., and Yushkov, V.: An overview of the SCOUT-AMMA stratospheric aircraft, balloons and sondes campaign in West Africa, August 2006: rationale, roadmap and highlights, Atmos. Chem. Phys. Discuss., 9, 19713–19781, 2009, <http://www.atmos-chem-phys-discuss.net/9/19713/2009/>.

Chaboureau, J.-P., Cammas, J.-P., Duron, J., Mascart, P. J., Sitnikov, N. M., and Voessing, H.-J.: A numerical study of tropical cross-tropopause transport by convective overshoots, Atmos. Chem. Phys., 7, 1731–1740, 2007, <http://www.atmos-chem-phys.net/7/1731/2007/>.

Chemel, Ch., Russo, M., Pyle, J. A., Sokhi, R. S., and Schiller, C.: Quantifying the imprint of a severe Hector thunderstorm during ACTIVE/SCOUT-O3 onto the water content in the upper troposphere/lower stratosphere, Mon. Weather Rev., 103, 2493–2514, doi:10.1175/2008MWR2666.1, 2009.

Corti, T., Luo, B. P., de Reus, M., Brunner, D., Cairo, F., Mahoney, M. J., Martucci, G., Matthey, R., Mitev, V., dos Santos, F. H., Schiller, C., Shur, G., Sitnikov, N. M., Spelten, N., Vossing, H. J., Borrmann, S., and Peter, T.: Unprecedented evidence for overshooting convection hydrating the tropical stratosphere, Geophys. Res. Lett., 35, L10810, doi:10.1029/2008GL033641, 2008.

Cotton, W. R., Pielke Sr., R. A., Walko, R. L., Liston, G. E., Tremback, C. J., Jiang, H., McAnelly, R. L., Harrington, J.-Y., Nicholls, M. E., Carrio, G. G., and McFadden, J. P.: RAMS 2001: Current status and future directions, Meteorol. Atmos. Phys., 82, 5–29, doi:10.1007/s00703-

**Water vapor budget associated to overshoots in the tropical stratosphere**

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

001-0584-9, 2003.

Durry, G., Amarouche, N., Zéninari, V., Parvitte, B., Lebarbu, T., and Ovarlez, J.: In situ sensing of the middle atmosphere with balloonborne near-infrared laser diodes, *Spectrochim. Acta*, 60(4), 3371–3379, 2004.

5 Freitas, S. R., Longo, K. M., Silva Dias, M. A. F., Chatfield, R., Silva Dias, P., Artaxo, P., Andreae, M. O., Grell, G., Rodrigues, L. F., Fazenda, A., and Panetta, J.: The Coupled Aerosol and Tracer Transport model to the Brazilian developments on the Regional Atmospheric Modeling System (CATT-BRAMS) – Part 1: Model description and evaluation, *Atmos. Chem. Phys.*, 9, 2843–2861, 2009,

10 <http://www.atmos-chem-phys.net/9/2843/2009/>.

Fueglistaler, S., Wernli, H., and Peter, T.: Tropical troposphere-to-stratosphere transport inferred from trajectory calculations, *J. Geophys. Res.*, 109, D03108, doi:10.1029/2003JD004069, 2004.

15 Fueglistaler, S., Bonazzola, M., Haynes, P. H., and Peter, T.: Stratospheric water vapor predicted from Lagrangian temperature history of air entering the stratosphere in the tropics, *J. Geophys. Res.*, 110, D08107, doi:10.1029/2004JD005516, 2005.

Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Folkins, I., Fu, Q., and Mote, P. W.: The tropical tropopause layer, *Rev. Geophys.*, 47, RG1004, doi:10.1029/2008RG0002672009, 2009.

20 Gettleman, A., Salby, M. L., and Sassi, F.: The distribution and influence of convection in the tropical tropopause region, *J. Geophys. Res.*, 107, 4080, doi:10.1029/2001JD001048, 2002.

Grell, G. A. and Devenyi, D.: A generalized approach to parameterizing convection combining ensemble and data assimilation techniques, *Geophys. Res. Lett.*, 29, 1693, doi:10.1029/2002GL015311, 2002.

25 Grosvenor, D. P., Choularton, T. W., Coe, H., and Held, G.: A study of the effect of overshooting deep convection on the water content of the TTL and lower stratosphere from Cloud Resolving Model simulations, *Atmos. Chem. Phys.*, 7, 4977–5002, 2007, <http://www.atmos-chem-phys.net/7/4977/2007/>.

Harrington, J. Y.: The effects of radiative and microphysical processes on simulated warm and transition season Arctic stratus, PhD Diss., Atmospheric Science Paper No 637, Colorado State University, Department of Atmospheric Science, Fort Collins, CO 80523, 289 pp., 1997.

30 Holton, J., Haynes, P., McIntyre, M., Douglass, A., Rood, R., and Pfister, L.: Stratosphere-Troposphere Exchange, *Rev. Geophys.*, 33(4), 403–439, 1995.

---

**Water vapor budget associated to overshoots in the tropical stratosphere**

X. M. Liu et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Water vapor budget associated to overshoots in the tropical stratosphere**

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Holton, J. R. and Gettleman, A.: Horizontal transport and the dehydration of the stratosphere, *Geophys. Res. Lett.*, 28, 2799–2802, 2001.
- Huffman, G. J., Adler, R. F., Bolvin, D. T., Gu, G., Nelkin, E. J., Bowman, K. P., Hong, Y., Stocker, E. F., and Wolff, D. B.: The TRMM multi-satellite precipitation analysis: quasi-global, multi-year, combined-sensor precipitation estimates at fine scale, *J. Hydrometeor.*, 8(1), 38–55, 2007.
- James, R., Bonazzola, M., Legras, B., Surbled, K., and Fueglistaler, S.: A Lagrangian analysis of a the Asian Monsoon Water Vapor Maximum at 110 hPa, *Geophys. Res. Lett.*, 35, L20810, doi:10.1029/2008GL035441, 2008.
- Khaykin, S., Pommereau, J.-P., Korshunov, L., Yushkov, V., Nielsen, J., Larsen, N., Christensen, T., Garnier, A., Lukyanov, A., and Williams, E.: Hydration of the lower stratosphere by ice crystal geysers over land convective systems, *Atmos. Chem. Phys.*, 9, 2275–2287, 2009, <http://www.atmos-chem-phys.net/9/2275/2009/>.
- Liu, C. and Zipser, E. J.: Global distribution of convection penetrating the tropical tropopause, *J. Geophys. Res.*, 110(D23), D23104, doi:10.1029/2005JD006063, 2005.
- Marécal, V., Durry, G., Longo, K., Freitas, S., Rivière, E. D., and Pirre, M.: Mesoscale modelling of water vapour in the tropical UTLS: two case studies from the HIBISCUS campaign, *Atmos. Chem. Phys.*, 7, 1471–1489, 2007, <http://www.atmos-chem-phys.net/7/1471/2007/>.
- Marécal, V., Pirre, M., Rivière, E. D., Pouvesle, N., Crowley, J. N., Freitas, S. R., and Longo, K. M.: Modelling the reversible uptake of chemical species in the gas phase by ice particles formed in a convective cloud, *Atmos. Chem. Phys. Discuss.*, 9, 24361–24410, 2009, <http://www.atmos-chem-phys-discuss.net/9/24361/2009/>.
- Meyers, M. P., Walko, R. L., Harrington, J. Y., and Cotton, W. R.: New RAMS cloud microphysics parameterization. Part II: The two-moment scheme, *Atmos. Res.*, 45, 3–39, 1997.
- Nielsen, J. K., Larsen, N., Cairo, F., Di Donfrancesco, G., Rosen, J. M., Durry, G., Held, G., and Pommereau, J. P.: Solid particles in the tropical lowest stratosphere, *Atmos. Chem. Phys.*, 7, 685–695, 2007, <http://www.atmos-chem-phys.net/7/685/2007/>.
- Oltmans, S. J., Vömel, H., Hofmann, D. J., Rosenlof, K. H., and Kley, D.: The increase in stratospheric water vapor from balloonborne frostpoint hygrometer measurements at Washington D.C., and Boulder, Colorado, *Geophys. Res. Lett.*, 27, 3453–3456, 2000.
- Pénide, G., Giraud, V., Bouniol, D., Dubuisson, P., Duroure, C., Protat, A., and Cautenet, S.:

**Water vapor budget associated to overshoots in the tropical stratosphere**

X. M. Liu et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Numerical Simulation of the 7 to 9 September 2006 AMMA mesoscale convective system: Evaluation of the Dynamics and Cloud Microphysics using Synthetic observations, Q. J. Roy. Meteorol. Soc., in press, 2010.

Pommereau, J.-P. and Held, G.: Is there a stratospheric fountain?, Atmos. Chem. Phys. Discuss., 7, 8933–8950, 2007,  
5 <http://www.atmos-chem-phys-discuss.net/7/8933/2007/>.

Randel, W. J., Wu, F., Gettelman, A., Russell, J. M., Zawodny, J. M., and Oltmans, S. J.: Seasonal variation of water vapor in the lower stratosphere observed in Halogen Occultation Experiment data, J. Geophys. Res., 106, 14313–14326, 2001.

10 Redelsperger, J.-L., Thorncroft, C. D., Diedhiou, A., Lebel, T., Parker, D. J., and Polcher, J.: African monsoon multidisciplinary analysis: An international research project and field campaign, B. Am. Meteor. Soc., 87, 1739–1746, 2006.

Roca, R., Viollier, M., Picon, L., and Desbois, M.: A multi satellite analysis of deep convection and its moist environment over the Indian Ocean during the winter monsoon, J. Geophys. Res., 107, 8012, doi:10.1029/2000JD000040, 2002.

15 Rosenlof, K., Oltmans, S., Kley, D., Russell III, J. M., et al.: Stratospheric water vapor increases over the past half-century, Geophys. Res. Lett., 28(7), 1195–1198, 2001.

Schiller, C., Grooß, J.-U., Konopka, P., Plöger, F., Silva dos Santos, F. H., and Spelten, N.: Hydration and dehydration at the tropical tropopause, Atmos. Chem. Phys., 9, 9647–9660,  
20 <http://www.atmos-chem-phys.net/9/9647/2009/>.

Schmetz, J., Tjemkes, S. A., Gube, M., and van de Berg, L.: Monitoring deep convection and convective overshooting with METEOSAT, Adv. Space Res., 19, 433–441, 1997.

Siedel, D. J., Ross, R. J., Angell, J. K., and Read, G. C.: Climatological characteristics of the tropical tropopause as revealed by radiosondes, J. Geophys. Res., 106, 7857–7878, 2001.

25 Stohl, A., and Seibert, P.: Accuracy of trajectories as determined from the conservation of meteorological tracers, Q. J. Roy. Meteor. Soc., 124, 1465–1484, 1998.

Zipser, E. J., Cecil, D. J., Liu, C., Nesbitt, S. W., and Yorlhy, D. P.: Where are the most intense thunderstorms on Earth?, B. Am. Meteor. Soc., 87, 1058–1072, 2006.

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

**Table 1.** Time evolution of the overshoot size in MSG pixels for the Air case, from 14:45 UT to 17:30 UT. At each time, the maximum BTD is also reported.

Local time of overshoot (UT)	Size of overshoot (number of pixels)	Maximum BTD value of the overshoot
5 August 14:45	3	3.49
5 August 15:00	3	3.06
5 August 15:15	3	3.27
5 August 15:30	13	4.8
5 August 15:45	5	3.87
5 August 16:00	8	3.52
5 August 16:15	14	3.87
5 August 16:30	16	3.89
5 August 16:45	11	3.87
5 August 17:00	6	3.59
5 August 17:15	3	3.79
5 August 17:30	2	3.92

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

**Table 2.** Model setting used for the Chad (4 August 2006) and Air (5 August 2006) case simulations. Size, resolution and starting time for each grid are reported.

Cases	Grid setting			
	Grid	Horizontal resolution (km)	Grid points (NX, NY, NZ)	Model start (UT)
Chad	G1	20×20	93, 83, 68	3 August, 18:00
	G2	4×4	242, 157, 68	
	G3	1×1	214, 110, 68	
Air	G1	20×20	152, 154, 68	5 August, 00:00
	G2	4×4	162, 197, 68	
	G3	1×1	106, 118, 68	

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.

**Table 3.** Water budget associated to the overshooting convection for the Chad and the Air cases across different isentropic surfaces. The integrated flux is the total mass of water crossing the  $\theta$  levels upward during the integration period encompassing the overshoot (4 h for the Chad case and 1 h 50' for the Air case). Also shown are the same results divided by the period of integration.

cases	Isentropic levels (K)	370 K	375 K	380 K	385 K	390 K	395K	400 K
Chad	Integrated flux (ton)	29 558	18 171	13 107	10 581	8963	7726	6841
	Per unit time (ton h <sup>-1</sup> )	7389	4542	3276	2645	2240	1931	1710
	Per unit time (ton s <sup>-1</sup> )	2.1	1.3	0.9	0.7	0.6	0.5	0.5
Air	Integrated flux (ton)	12 172	6313	4332	3465	2966	2677	2471
	Per unit time (ton h <sup>-1</sup> )	6639	3443	2363	1890	1618	1460	1348
	Per unit time (ton s <sup>-1</sup> )	1.8	0.9	0.7	0.5	0.4	0.4	0.4

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion


**Water vapor budget associated to overshoots in the tropical stratosphere**

X. M. Liu et al.



**Fig. 1.** UTLS water vapor profiles from Balloon-borne micro-SDLA (red line) and FLASH-B (green) measurements from Niamey at 18:40 UT and 18:52 UT (launch time) on 5 August 2006, respectively.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



**Fig. 2.** Meteosat Second Generation Infrared images in the 10° E to 30° E longitude range and in the 0° N to 20° N latitude range on 4 August 2006 at **(a)** 10:30 UT **(b)** 12:00 UT **(c)** 14:30 UT **(d)** 17:00 UT **(e)** 20:00 UT.

**Water vapor budget associated to overshoots in the tropical stratosphere**

X. M. Liu et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Water vapor budget  
associated to  
overshoots in the  
tropical stratosphere**

X. M. Liu et al.




**Fig. 3.** MSG pixels with brightness temperature difference (BTD) higher than 3 K between the  $6.2\ \mu\text{m}$  channel and the  $10.8\ \mu\text{m}$  channel in Southern Chad from 12:30 UT to 19:30 UT on 4 August 2006. The time resolution of MSG data is 15 min.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Water vapor budget  
associated to  
overshoots in the  
tropical stratosphere**

X. M. Liu et al.




**Fig. 4.** MSG Infrared images centered on Niger on 5 August 2006. **(a)** 13:30 UT **(b)** 16:00 UT **(c)** 18:00 UT **(d)** 23:30 UT.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Water vapor budget  
associated to  
overshoots in the  
tropical stratosphere**

X. M. Liu et al.




**Fig. 5.** Same as Fig. 3 but for the Air case from 14:45 UT to 17:30 UT on 5 August 2006.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Water vapor budget associated to overshoots in the tropical stratosphere**

X. M. Liu et al.




**Fig. 6.** Topography of the 3 nested grids used for BRAMS simulation (iso-lines for Grid 1, and shaded contour for Grids 2 and 3. **(a)** Chad case. **(b)** Air case.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.



**Fig. 7.** Comparison between TRMM (a, c, e) and BRAMS (b, d, f) surface rainrate accumulated on 3h periods for the Chad case. Panel (a) and (b) : from 13:30 UT to 16:30 UT on 4 August 2006. Panel (c) and (d) from 16:30 to 19:30 UT. Panel (e) and (f) from 19:30 UT to 21:30 UT.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Water vapor budget associated to overshoots in the tropical stratosphere**

X. M. Liu et al.




**Fig. 8.** Chad case BRAMS simulation. Grid 3 time evolution of the grid points having an ice mixing ratio higher than  $0.05 \text{ g kg}^{-1}$  above the tropopause (380 K). The same color scale as in Fig. 3 is chosen.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Water vapor budget  
associated to  
overshoots in the  
tropical stratosphere**

X. M. Liu et al.




**Fig. 9.** Longitude/Altitude cross-section of condensed water mixing ratio ( $\text{g kg}^{-1}$ ) around the overshoot location with iso-theta levels inside the Grid 3 at  $= 12^\circ \text{N}$  latitude and 14:15 UT on 4 August 2006.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Water vapor budget  
associated to  
overshoots in the  
tropical stratosphere

X. M. Liu et al.




**Fig. 10.** Comparison between TRMM (a, c, e) and BRAMS (b, d, f) surface rainrate accumulated on 3 h periods for the Aircase. Panel (a) from 13:30 UT to 16:30 UT on 4 August 2006. (b) From 14:00 to 17:00 UT. Panel (c) from 16:30 to 19:30 UT, and (d) from 17:00 to 20:00. Panel (e) from 19:30 UT to 21:30 UT and (f) from 20:00 to 23:00 UT.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Water vapor budget  
associated to  
overshoots in the  
tropical stratosphere**

X. M. Liu et al.




**Fig. 11.** Same as Fig. 8 but for the Air case. The same color scale as in Fig. 5 is chosen. **(a)** Zoom on the Grid 3 from 18:30 to 19:50 UT. **(b)** From 18:30 UT to 21:15 UT.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.



**Fig. 12.** Grid 3 vertical cross-section of condensed water ( $\text{g kg}^{-1}$ ) for the Air case BRAMS simulation. Isentropic levels are also shown. **(a)** Longitude/Altitude vertical cross-section at  $16.15^\circ$  E latitude, at 19:10 UT. **(b)** Longitude/Altitude vertical cross-section at  $15.9^\circ$  E latitude, at 19:45 UT.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.



**Fig. 13.** Scattergram of total  $\text{H}_2\text{O}$ /potential temperature within the Grid 3 domain of BRAMS at different times of the simulation. **(a)** Initial state at 18:00 UT on 3 August. **(b)** 14:15 UT on 4 August at the maximum of the overshooting activity. **(c)** 15:55 UT after the overshooting activity. **(d)** 21:00 UT after the convective activity. The black solid line is the mean mixing ratio within the grid. The blue dashed line is the initial mean mixing ratio.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Water vapor budget associated to overshoots in the tropical stratosphere**

X. M. Liu et al.




**Fig. 14.** Upward total water flux across different isentropic levels within the BRAMS Grid 3 domain between 12:00 UT and 15:55 UT for the Chad case.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Water vapor budget associated to overshoots in the tropical stratosphere

X. M. Liu et al.



**Fig. 15.** Total water at the 390K isentropic level in the BRAMS Grid 3 domain at 14:15 UT (a), 14:30 UT (b), 14:45 UT (c) and 15:10 UT (d) for the Chad case simulation. Also shown are contours of ratio (%) of ice with respect to total water (black solid lines): (a) and (b) 10% and 50%, (c) 10% and 20%, and (d) 10%. The maximum total water mixing ratio as well as the maximum ratio (%) of ice with respect to total water are reported in each panel. The same color scale applies to each panel.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Water vapor budget associated to overshoots in the tropical stratosphere**

X. M. Liu et al.



**Fig. 16.** Same as Fig. 14 but for the Air case, computed between 18:00 UT and 19:50 UT on 5 August 2006.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)