

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.

Atmospheric deposition of mercury and major ions to the Pensacola Bay (Florida) watershed: spatial, seasonal, and inter-annual variability

J. M. Caffrey¹, W. M. Landing², S. D. Nolek³, K. Gosnell², S. S. Bagui⁴, and S. C. Bagui⁵

¹Center for Environmental Diagnostics and Bioremediation University of West Florida, 11000 University Parkway Pensacola, FL 32514, USA

²Department of Oceanography Florida State University 1015 West Call Street Tallahassee, FL 32306-4320, USA

³PetroAlgae 1901 S. Harbor City Blvd., Suite 300 Melbourne, FL 32901, USA

⁴Department of Computer Science University of West Florida 11000 University Parkway Pensacola, FL 32514, USA

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



⁵Department of Mathematics and Statistics University of West Florida 11000 University Parkway Pensacola, FL 32514, USA

Received: 24 November 2009 – Accepted: 8 February 2010 – Published: 15 February 2010

Correspondence to: J. M. Caffrey (jcaffrey@uwf.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Abstract

ACPD

10, 4593–4616, 2010

Atmospheric deposition was measured at three sites in the Pensacola Bay watershed, Florida, between November 2004 and December 2007. Mercury deposition in the Pensacola Bay watershed was similar to that from nearby Mercury Deposition Network sites along the Northern Gulf of Mexico coast. Mercury deposition during the summer months is higher than other months due to higher concentrations in rainfall throughout the region. Deposition of constituents like H^+ , sulfate, nitrate, ammonium, chloride and sodium, were much higher in Pensacola Bay than at National Atmospheric Deposition Program (NADP) sites. Chloride and sodium fluxes are higher because Pensacola Bay sites are closer to the Gulf of Mexico which is a source of sea salt aerosols. Acid rain constituents, H^+ , sulfate, nitrate and ammonium are most likely higher at Pensacola Bay sites because these sites are much closer to emission sources of these constituents than NADP sites, particularly two Florida NADP sites, FL14 and FL23, which are located in rural counties far from major industrial activities.

15 1 Introduction

Atmospheric concentrations of many compounds including nitrogen, mercury, copper, zinc, sulfur, and polycyclic aromatic hydrocarbons have increased since the industrial revolution due to combustion of fossil fuels, industrial and manufacturing processes and agricultural operations. Mercury emissions to the atmosphere are primarily associated with coal burning power plants, industrial processes such as chlor-alkali production, and waste incinerators (Pirrone et al., 1996; Pacyna et al., 2005). Mercury in the atmosphere occurs in 3 different forms: elemental gaseous mercury (Hg^0), reactive gaseous mercury (RGM or Hg^{2+}) which is the oxidized form, and mercury associated with particles (HgP). Reactive gaseous mercury is usually deposited near the source of the emission (Dvorch et al., 1998; Lin and Pehkonen, 1999), while elemental gaseous mercury has a residence time in the atmosphere of 1–2 yr, thus deposition may be

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



worldwide (Lin and Pehkonen, 1999). Reactive gaseous mercury can be adsorbed directly onto plant and soil surfaces, as well as associated with soot, fog droplets or rain drops (Lin and Pehkonen, 1999).

Deposition of atmospheric constituents to land and water surfaces occurs at a wide range of scales, from less than a kilometer to hundreds of kilometers from the emission sources. Thus, airsheds encompass huge areas, usually 10 to 100 times greater than watershed area (Valigura, 2001). Airshed boundaries are amorphous, dynamic and dependent on regional weather patterns. Atmospheric deposition represents a significant source of nutrients and other contaminants in many watersheds (Valigura, 2001). Between 50–90% of the mercury load to US waters originates from atmospheric deposition (<http://nadp.sws.uiuc.edu/>). Deposition of these compounds onto land and water surfaces has a negative impact on plant and animal communities, ecosystem function and human health. Accumulation of toxic contaminants in animal tissues can lead to reproductive failures. Toxic compounds such as mercury, selenium and organic chemicals accumulate in food chains increasing human health risks when fish or waterfowl are consumed. Currently, 42 states have mercury consumption advisories for freshwater fish while the Southeast and Gulf coasts have coastal fish advisories (<http://nadp.sws.uiuc.edu/>).

The growth of monitoring networks (e.g. National Atmospheric Deposition Program (NADP)) since 1978 has provided an increasingly detailed picture on atmospheric deposition at the national and regional level in the United States. The National Atmospheric Deposition Program, which includes the Mercury Deposition Network (MDN), is a national network of collaborating state and federal agencies that collect data to examine long-term trends in atmospheric deposition at regional scales. Regional and national trends provide critical information necessary for determining policies to improve air and water quality at the national level (Lynch and Kerchner, 2004; Lindberg et al., 2007). However, in order to improve water quality at the watershed level, the importance of local sources in addition to regional sources needs to be determined. Urban areas frequently have higher mercury deposition than rural areas (Mason et al., 1997;

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Keeler et al., 2006; Van Arsdale et al., 2007), particularly areas downwind of coal-fired utilities (White et al., 2009). In addition, sampling individual rain events provides a more detailed assessment of rain composition that can be used for source tracking

Our primary objective was to measure mercury, trace metals and major ions in rainfall over annual cycles. We evaluated the temporal and spatial patterns in atmospheric deposition by comparing the results among 3 sites in the Pensacola Bay watershed and comparing them with data collected at MDN and NADP sites along the Central Gulf of Mexico coast. Our primary research question was what is the spatial distribution of mercury, trace metal and nutrient deposition in the Pensacola Bay watershed? Trace metal data are presented in Landing et al. (2009).

2 Study area

This research was conducted as part of the Partnership for Environmental Research and Community Health in order to understand the contribution of mercury and trace metals from atmospheric sources to the Pensacola Bay region. Air quality in Pensacola Bay is affected by emissions from industrial sources, automobiles and other vehicles as well as other stationary area sources such as dry cleaners, gas stations, agriculture, and construction sites (FDEP 1998). Industrial sources include a coal fired electric generating plant, a paper mill, chemical and other manufacturing companies (EPA website: <http://www.epa.gov/air/data/reports.html>). NO_x and SO₂ emissions are dominated by power generation, although automobile and other vehicles are also a large source of NO_x. The major source of mercury emissions in 2005 was from Plant Crist, the coal-fired electric generating plant, which was estimated to release approximately 87 kg. Other mercury emission sources in the Pensacola Bay watershed include coal-fired industrial boilers at the International Paper Plant (0.9 kg), medical waste incineration (0.0009 kg) and landfills (0.018 kg) (T. Rogers, personal communication, 2009). Nearby Mobile and Escambia Counties in Alabama have significant sources of mercury emissions, approximately 350

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



and 400 kg respectively in 2002 (<http://www.epa.gov/air/data/repsco.html?co~01053%2001097~Escambia%20Co%2C%20Mobile%20Co%2C%20Alabama>).

3 Methods

Rain collectors were installed at 3 sites between November and December 2004 (Fig. 1) for event based collection of rain samples. Rain samples were collected within 24 h of 0.5 cm rain events using Aerochemetrics or Loda rain collection devices. Buckets were modified to hold 3 bottles with Teflon collars as in the Florida Atmospheric Mercury Study (Landing et al., 1995). Duplicate Teflon bottles for mercury and trace metal analysis were deployed along with a single polyethylene bottle for pH and major ions analysis. Approximately 10% of the deployments had a single Teflon bottle for mercury and trace metal samples and duplicate polyethylene bottles for pH and major ions. Trace metal results are reported in Landing et al. (2009). Sampling in 2005 was suspended for 2 weeks by Hurricane Dennis because collectors were removed from the sites to keep them from being damaged or destroyed.

15 3.1 Mercury analysis

Rain samples (in FEP Teflon bottles) were acidified after collection to 0.045 M HCl plus 0.048 M HNO₃, then placed in a low wattage UV digestion box for at least 48 h to completely solubilize the collected Hg. The digested samples were analyzed by CVAFS using EPA method 1631 using a Tekran 2600 Mercury Analyzer at Florida State University.

20 3.2 Major ions and pH

The pH of the samples was measured in the laboratory immediately following collection. The major ions, nitrate (NO₃⁻), ammonium (NH₄⁺), phosphate (PO₄³⁻), sulfate

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀
Back

▶
Close

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



(SO_4^{2-}), sodium (Na^+), chloride (Cl^-), magnesium (Mg^{2+}), calcium (Ca^{2+}), and potassium (K^+) were measured in polyethylene sample bottles. Major ion analyses were conducted using a Dionex DX500 ion chromatograph using EPA method 300.0 for anions and Dionex protocols (Application Note 141, http://www.dionex.com/en-us/webdocs/4211-AN141_V15.pdf) for cations at the University of West Florida. Phosphate, magnesium, calcium and potassium were often below detection limits (0.03 mg/L PO_4^{3-} , 0.02 mg/L Mg^{2+} , 0.04 mg/L Ca^{2+} , 0.02 mg/L K^+), so values are not reported here.

3.3 Calculations and statistics

All deposition fluxes are calculated by multiplying the measured concentrations (grams per cubic meter) by the rainfall amounts (meters per event), and are expressed as mass

10 per square meter (per event). We tested for differences between sites, seasons and years using ANOVA. The seasons are winter (December, January, February), spring (March, April, May), summer (June, July, August) and fall (September, October, November). We used the event based data to compare our three sites. A correlation analysis

15 of mercury and major ion fluxes was performed on the data. In addition, we compared our sites with Mercury Deposition Network (MDN) and National Atmospheric Deposition Program (NADP) sites in Northern Gulf of Mexico region (Fig. 1, NADP 2009) using ANOVA to test for differences among sites. For this comparison, we summed our data over the same weekly intervals as the NADP and MDN sites. The level of significance

20 equal or less than 0.05 was used for these analyses.

4 Results

Over the course of this study from 19 November 2004 through 2 January 2008, we collected 565 rain samples from 3 sites. These samples represent about 225 separate rain events in the Pensacola Bay watershed. The amount of precipitation was not significantly different among the 3 sites, nor were there any significant inter-annual dif-

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



ferences. However, seasonal differences were significant with the higher volume rainfall events in the spring and fall and the lower volume rainfall during winter. Approximately 33% of the total rainfall occurs in the summer, 28% in the fall, 23% in the spring and only 16% in the winter.

5 4.1 Mercury concentrations in rainfall

Concentrations of mercury and most major ions, particularly sulfate and ammonium, were generally higher in low volume samples (Fig. 2). pH values were generally more variable in low volume samples, with many samples less than 4 or greater than 5.5. With larger storms, mercury and major ion concentrations were diluted. This 10 washout effect was most obvious for mercury. Mercury deposition showed no site differences between the three Pensacola Bay watershed sites (Fig. 3). Mercury fluxes from individual events ranged between 13 and 2050 ng/m²/event. There were significant seasonal differences. The highest mercury deposition occurred in the summer (403±19.9 ng/m²/event) compared to other seasons (fall 286±18 ng/m²/event, winter 15 245±15 ng/m²/event, spring 333±24 ng/m²/event). Interannual differences were significant with the highest average deposition in 2007 (400.7 ng/m²/event) and the lowest in 2006 (286 ng/m²/event).

4.2 Precipitation chemistry

For each of the major ion fluxes (H⁺, sulfate, nitrate, ammonium, chloride and sodium), 20 we found no significant differences among the 3 Pensacola Bay sites. There were significant seasonal differences in H⁺, nitrate and ammonium deposition, but not sulfate, chloride or sodium deposition. The highest H⁺ flux occurred in summer and the lowest during winter and fall (Fig. 3). Nitrate fluxes were also significantly lower during the winter (16.5 mg NO₃/m²/event) and fall (26.0 mg NO₃/m²/event) compared to spring 25 (56.4 mg NO₃/m²/event) and summer (45.1 mg NO₃/m²/event) (Fig. 4). The seasonal pattern in ammonium was the same as nitrate fluxes with higher fluxes in the spring

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

(9.2 mg NH₄/m²/event) and summer (5.2 mg NH₄/m²/event) and lower fluxes during the fall (1.9 mg NH₄/m²/event) and winter (2.5 mg NH₄/m²/event) (Fig. 4).

The average sulfate flux was 40.43 mg SO₄/m²/event and values from individual rain events ranged from 0.5 to 438 mg SO₄/m²/event. (Fig. 3). The average nitrate flux was 35.6 and values ranged from 0.5 to 285 mg NO₃/m²/event (Fig. 4). Nitrate fluxes in 2007 (19.5 mg NO₃/m²/event) were significantly lower than in 2005 (45.5 mg NO₃/m²/event) or 2006 (43.2 mg NO₃/m²/event). Ammonium fluxes were about ten percent of nitrate fluxes (Fig. 4). Average ammonium flux was 4.35 and values ranged from 0.04 to 46.3 mg NH₄/m²/event.

Chloride fluxes had a larger range than the other major element fluxes, with a few fluxes exceeding 1000 mg/m²/event, particularly at Ellyson (Fig. 5). This is likely due to the accumulation of sea salt aerosols in rainwater. The average chloride flux was 58.76 and values ranged from 0.6 to 2070 mg/m²/event. There were no significant differences in chloride fluxes among the different sites, years or seasons. Sodium fluxes also had a large range and generally tracked the chloride concentrations (Fig. 5). As observed with the chloride fluxes, sodium fluxes had no significant differences among the different sites, years or seasons. Average sodium fluxes were 24.5 and values ranged from 0.2 to 938 mg/m²/event.

4.3 Comparisons with MDN and NADP sites

Mercury fluxes along the Northern Gulf of Mexico were similar among MDN and Pensacola Bay sites (Table 1). There were no interannual differences in mercury fluxes, although seasonal fluxes were significantly different from one another. As we observed with the Pensacola Bay data alone, summer fluxes were significantly higher than fluxes during the other months. Summer mercury fluxes represented an average of 44% of the total annual mercury flux at these sites and could represent as much as 60% for some sites in some years.

Sites in Pensacola Bay area were often higher than NADP sites for H⁺, sulfate,

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



nitrate, ammonium, chloride, and sodium fluxes (Table 1). H^+ fluxes at Ellyson and Pace were significantly higher than at all other NADP sites, while fluxes at Molino were significantly higher than FL14 and LA30. Sulfate fluxes at Ellyson were significantly higher than all NADP sites, Pace was significantly higher than NADP sites in Florida and Louisiana, while Molino was significantly higher than the NADP sites in Florida. 5 Nitrate fluxes at all Pensacola Bay sites were significantly greater than NADP sites. Ammonium flux at Ellyson was greater than FL14 and FL23. Ammonium flux at the NADP site LA30 was also greater than those at FL23. Chloride fluxes at Ellyson were greater than those at Molino and all NADP sites, while Pace was greater than the 10 NADP sites in Florida, Louisiana and at AL02. Sodium fluxes were significantly higher at Ellsyon than at all NADP sites. Pace also had significantly higher sodium fluxes than at all NADP sites except for AL24.

There were significant seasonal differences in sulfate, nitrate, ammonium and chloride fluxes with sulfate, nitrate and ammonium having higher fluxes in the spring and 15 summer than the other seasons. Chloride fluxes were highest in the fall. Inter-annual differences were significant for sulfate and nitrate fluxes with higher fluxes in 2005 and 2006 compared to 2007. H^+ and ammonium flux declined over the three years of observations.

5 Discussion

20 These results have shown that there were no significant differences in mercury deposition among our three sampling sites or between our sites and MDN sites in the Northern Gulf of Mexico between 2005 and 2007. The mercury deposition at the Ellyson site, which is about 4.8 km away and generally downwind of the Crist coal fired power plant (Fig. 1), was not significantly higher than at the Molino site which is furthest away 25 from local mercury emission sources (24 km from Crist power plant and 14 km from International Paper). Our results contrast with studies in Michigan and Maryland which found higher mercury deposition in urban areas compared to rural areas (Mason et al.,

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



2000; Lawson and Mason, 2001; Landis et al., 2002; Keeler and Dvonch, 2005) and those finding higher concentrations of mercury in rainwater close to coal-fired utilities (White et al., 2009). We did observe seasonal differences, with the highest deposition rate during the summer at Pensacola Bay and MDN sites. Summer deposition at our sites and the nearby MDN sites represents about 43% of the total annual mercury flux, similar to finds in Maryland (Mason et al., 2000). This contrasts with the FAMS study which found that summer deposition in south Florida was between 80–90% of total mercury flux (Guentzel et al., 1995). Higher summer mercury deposition is not driven by higher rainfall amounts, but by higher concentrations in the rain since the highest rainfall amounts are usually in the spring and fall. Several mechanisms leading to enhanced mercury concentrations in rain during the summer include greater emissions associated with higher electricity use, scavenging of reactive gaseous mercury from the free troposphere by tall convective thunderstorms, and the concentration of reactive gaseous mercury by the sea breeze effect, where the diurnal alternation of onshore and offshore winds can lead to a buildup of pollutants in the air mass. In the Pensacola Bay area, SO_2 levels increased as the sea breeze move urban air masses offshore and then back onshore (Chang et al., 2007). Keeler and Dvonch (2005) also observed strong seasonal patterns in atmospheric mercury deposition in Michigan with higher deposition in the summer due to higher concentrations of mercury in rain droplets compared to ice crystals that form in winter. As with seasonal differences, inter-annual variability was not strictly driven by rainfall amount since the year with the highest rainfall, 2005, had an intermediate mercury deposition rate.

The deposition of major ions (sulfate, chloride, nitrate, sodium, H^+ , or ammonium) was similar among the Pensacola Bay sites and often significantly higher than NADP sites. The NADP sites in Florida (FL14 and FL23) often had the lowest values of all the sites examined. This is consistent with other studies that have shown higher deposition in urban settings (Elliott et al., 2007; Lohse et al., 2008). In particular, stationary sources are a significant source of NO_x which results in high nitrate fluxes in wet deposition (Elliott et al., 2007). The Florida NADP sites (FL14 and FL23) are

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

much further from urban centers and major pollution sources than either the Alabama NADP or our Pensacola Bay sites. Pensacola Bay sites were not chosen by NADP criteria because we were interested in capturing deposition from local sources. The Molino site is farthest away from pollution sources in Pensacola, about 14 km from International Paper and 20 km away from the urban center, yet it still is high for sulfate and nitrate, which are often associated with fossil fuel combustion.

5 Sea salt aerosols are be important in sequestering reactive gaseous and particulate mercury in the Gulf Coast (Engle et al., 2008). In this study, mercury deposition was significantly correlated with sodium flux, however, the correlation coefficient was quite 10 low ($r=0.22$, $p=0.02$). Sea salt aerosols have a significant impact on chemistry of rain water in the Pensacola Bay area. Sodium and chloride, elements associated with sea water, were highly correlated ($r=0.73$, $p <0.001$). About 21% of the rainfall events had Cl:Na ratios near the seawater Cl:Na ratio of 1.79 (Fig. 6). However, chloride fluxes were enriched relative to the seawater ratio in about 63% of the rain events, suggesting 15 that there is another source of chloride in rainwater. The distance of the sampling site to the Gulf of Mexico determines how significant an effect the sea salt aerosols have on rain water composition. Sites that are further from the Gulf have lower average annual sodium fluxes than those near to it (Fig. 7). There was a similar pattern for chloride fluxes (data not shown). Fifty km was the distance where the line becomes asymptotic 20 and the sea salt effect is lost.

Element ratios are a useful tool for examining potential sources in rain water. Our samples were highly enriched in some constituents relative to sodium. The sulfate to sodium ratio was 4 to 100 times greater than seawater ratio (Fig. 6). Only 2% of the rain events had sulfate:sodium ratios near the seawater ratio. The highest ratios generally 25 occurred in the summer, particularly in 2005 (Fig. 6). Several of the major ions associated with acid rain constituents were significantly correlated with one another. Sulfate was strongly correlated with H^+ and chloride, with correlation coefficients of 0.62 and 0.72, respectively ($p <0.001$). The relationships between sulfate and nitrate or ammonium were also significant, but only had correlation coefficients of 0.24 and 0.32,

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

respectively. The ratio of nitrate to non-sea salt sulfate in our samples was about 0.5, which is similar to that found during the FAMS study (Landing et al., 1995). Mercury was strongly correlated with H⁺ and sulfate concentrations, with correlation coefficients of 0.6 and 0.73, respectively ($p < 0.001$). This is similar to the findings from Michigan and the Ohio valley which also observed strong associations between sulfate, pH and mercury deposition (Landis et al., 2002; Keeler et al., 2006). These relationships are consistent with mercury sources from coal combustion.

The three years of data from this study have shown consistent seasonal trends in mercury and major ion deposition. Because sampling is ongoing at these sites, future papers will be able to address the factors controlling interannual variability and long term trends, particularly the effect of installing scrubbers at Plant Crist.

6 Conclusions

In contrast to studies from other regions, mercury deposition at urban sites in the Pensacola Bay watershed was similar to mercury deposition at Mercury Deposition Network sites along the Gulf Coast. However, deposition of acid rain constituents such as H⁺, sulfate, nitrate, chloride and ammonium was much higher at Pensacola Bay sites than rural National Atmospheric Deposition Network sites along the Gulf Coast. Summer deposition of mercury represented approximately 40% of the annual deposition, but only 33% of the annual rainfall.

Acknowledgements. This study was a component of the “Assessment of Environmental Pollution and Community Health in Northwest Florida”, supported by a U.S. EPA Cooperative Agreement award X-9745502 to the University of West Florida (Project Director: K. Ranga Rao). We thank Fran Aftanas, Nathaniel Davila, Autumn Dunn, Elizabeth Gaige, Brad Kuykendall, Tanner Martin, Melissa Overton, and Pam Vaughan for their assistance in field collections. We thank Jan Macauley, Jeremy Bosso and the Wetland Research Laboratory for laboratory analyses. The Florida Department of Environmental Protection, Pace Water Systems, and Carol Hatcher provided access to sampling sites. We thank Jessie Brown for her assistance with the database development.

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Chang, M. E., Baumann, K., Bostrom, A., and Russell, A.: PERCH Air Quality Study: an assessment of particulate matter, ozone, and air toxics in Escambia and Santa Rosa Counties, Final Report, Partnership for Community Health and Environmental Research, 5 <http://cure.eas.gatech.edu/~chang/perch/>, 2007.

Dvonch, J. T., Graney, J. R., Marsik, F. J., Keeler, G. J., and Stevens, R. K.: An investigation of source-receptor relationships for mercury in south Florida using event precipitation data, *The Science of the Total Environment*, 213, 95–108, 1998.

Elliott, E. M., Kendall, C., Wankel, S. D., Burns, D. A., Boyer, E. W., Harlin, K., Bain D. J., 10 and Butler, T. J.: Nitrogen isotopes as indicators of NO_x source contributions to atmospheric nitrate deposition across the Midwestern and northeastern United States, *Environ. Sci. Technol.*, 41, 7661–7667, 2007.

Engle, M. A., Tate, M. T., Krabbenhoft, D. P., Kolker, A., Olsen, M. L., Edgerton, E. S., DeWild, 15 J. F., and McPherson, A. K.: Characterization and cycling of atmospheric mercury along the central US Gulf Coast, *Appl. Geochem.*, 23, 419–437, 2008.

Guentzel, J. L., Landing, W. M., Gill, G. A., and Pollman, C. D.: Atmospheric deposition of mercury in Florida: the FAMS project (1992–1994), *Water, Air, Soil Pollut.* 80, 393–402, 1995.

Keeler, G. J. and Dvonch, J. T.: Chapter 25, Atmospheric mercury: a decade of 20 observations in the Great lakes, in: *Dynamics of mercury pollution on regional and global scales: atmospheric processes and human exposures around the world*, edited by: Pirrone, N. and Mahaffey, K. R., Springer, 611–636, 2005.

Keeler, G. J., Landis, M. S., Norris, G. A., Christianson, E. M., and Dvonch, J. T.: Sources of mercury wet deposition in Eastern Ohio, USA, *Environ. Sci. Technol.*, 40, 5874–5881, 2006.

Landing, W. M., Perry Jr., J. J., Guentzel, J. L., Gill, G. A., and Pollman, C. D.: Relationships 25 between atmospheric deposition of trace elements, major ions and mercury in Florida: The FAMS project (1992–1993), *Water, Air, Soil Pollut.*, 80, 343–352, 1995.

Landing, W. M., Caffrey, J. M., Nolek, S. D., and Gosnell, K.: Atmospheric wet deposition of mercury and other trace elements in Pensacola, Florida, *Atmos. Chem. Phys. Discuss.*, 9, 30 27649–27674, 2009, <http://www.atmos-chem-phys-discuss.net/9/27649/2009/>.

Landis, M. S., Vette, A. F., and Keeler, G. J.: Atmospheric mercury in the Lake Michigan Basin:

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Influence of the Chicago/Gary Urban Area, *Environ. Sci. Technol.*, 36, 4508–4517, 2002.

Lawson, N. M. and Mason, R. P.: Concentration of mercury, methylmercury, cadmium, lead, arsenic and selenium in the rain and stream water of two contrasting watersheds in western Maryland, *Water Res.*, 35, 4039–4052, 2001.

5 Lin, C.-J. and Pehkonen, S. O.: The chemistry of atmospheric mercury: a review, *Atmos. Environ.*, 33, 2067–2079, 1999.

Lindberg, S., Bullock, R., Ebinghaus, R., Engstrom, D., Feng, X., Fitzgerald, W., Pirrone, N., Prestbo, E., and Seigneur, C.: A synthesis of progress and uncertainties in attributing the sources of mercury in deposition, *Ambio*, 36, 19–32, 2007

10 Lohse, K. A., Hope, D., Sponsell, R., Allen, J. O., and Grimm, N. B.: Atmospheric deposition of carbon and nutrients across an arid metropolitan area, *Sci. Tot. Environ.*, 402, 95–105, 2008.

Lynch, J. A. and Kerchner, M.: The National Atmospheric Deposition Program: 25 years of monitoring in support of science and policy: an ammonia workshop: the state of science and future needs, *Environ. Pollut.*, 135, 343–346, 2005.

15 Mason, R. P., Lawson, N. M., and Sullivan K. A.: The concentration, speciation and sources of mercury in Chesapeake Bay precipitation, *Atmos. Environ.*, 31, 3541–3550, 1997.

Mason, R. P., Lawson, N. M., and Sheu, G. R.: Annual and seasonal trends in mercury deposition in Maryland, *Atmos. Environ.*, 34, 1691–1701, 2000.

20 National Atmospheric Deposition Program (NRSP-3)/National Trends Network, NADP Program Office, Illinois State Water Survey, 2204 Griffith Drive, Champaign, IL 61820, 2009.

Pacyna, J. M., Munthe, J., Larjava, K., and Pacyna, E. G.: Chapter 3. Mercury emissions from anthropogenic sources: estimates and measurements for Europe, in: *Dynamics of mercury pollution on regional and global scales: atmospheric processes and human exposures around the world*, edited by: Pirrone, N. and Mahaffey, K. R., Springer, 51–64, 2005

25 Pirrone, N., Keeler, G. J., and Nriagu, J. O.: Regional differences in worldwide emissions of mercury to the atmosphere, *Atmos. Environ.*, 30, 2981–2987, 1996.

Valigura, R. A.: An introduction to the first assessment of nitrogen loads to US estuaries with an atmospheric perspective, in: *Nitrogen loading in coastal water bodies: an atmospheric perspective*, edited by: Valigura, R. A., Alexander, R. B., Castro, M. S., Meyers, T. P., Paerl, H. W., Stacey, P. E., and Turner, R. E., AGU. Washington, D.C, 1–10, 2001.

30 VanArsdale, A., Weiss, J., Keeler, G., Miller, E., BOulet, G., Brulotte, R., and Poissant, L.: Patterns of mercury deposition and concentration in Northeastern North America (1996–

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



2002), Ecotoxicology, 14, 37–52, 2007.

White, E. M., Keeler, G. J., and Landis, M. S.: Spatial variability of mercury wet deposition in Eastern Ohio: Summertime meteorological case study analysis of local source influences, Environ. Sci. Technol., 43, 4946–4953, 2009.

ACPD

10, 4593–4616, 2010

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.

Table 1. Analysis of variance in mercury and major ion fluxes for Pensacola Bay sites, National Atmospheric Deposition Program (NADP) sites and Mercury Deposition Network (MDN) sites along the central Gulf of Mexico. P-values reported, NS is non significant. Seasons are spring (Sp), summer (Su), fall (Fa) and winter (W).

Constituent	Site	Season	Year
Mercury	NS	$p < 0.001$ (Su>Sp, Fa & W)	NS
H^+	$p < 0.001$	$p < 0.0001$ (Su>Sp, Fa & W)	$p = 0.028$ (2005>2006 & 2007)
Sulfate	$p < 0.001$	$p < 0.0001$ (Sp & Su>Fa & W)	$p < 0.0001$ (2005>2006>2007)
Nitrate	$p < 0.001$	$p < 0.001$ (Sp & Su>Fa & W)	$p < 0.0001$ (2005 & 2006>2007)
Chloride	$p < 0.001$	$p = 0.03$ (Fa>Sp & Su & W)	NS
Sodium	$p < 0.001$	NS	NS
Ammonium	$p = 0.03$	$p < 0.001$ (Sp>Su>Fa & W)	$p < 0.001$ (2005>2006>2007)

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)





Fig. 1. Map of Pensacola Bay showing sampling locations: Ellyson, Molino and Pace. Mercury emission sources: Crist Plant, International Paper, Sacred Heart Hospital medical waste incinerator. Inset map shows location of Mercury Deposition Network (MDN) and National Atmospheric Deposition Program (NADP) sampling sites along the central Gulf Coast.

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[|◀](#)

[▶|](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.




Fig. 2. Mercury (Hg ng/L), pH, sulfate, and nitrate concentrations versus the volume of rain (mL) collected in sample bottles at Pensacola Bay sites.

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.




Fig. 3. Mercury (Hg) deposition ($\text{ng}/\text{m}^2/\text{event}$) (top panel), H^+ flux ($\text{mole}/\text{m}^2/\text{event}$) (middle panel), and sulfate (SO_4^{2-}) deposition (bottom panel) ($\text{mg } \text{SO}_4^{2-}/\text{m}^2/\text{event}$) at Ellyson (brown), Molino (orange) and Pace (green) between November 2004 and December 2007 for individual rain events (left side). Box plot (right side) for each site, showing median, 25th and 75th percentiles (box boundary), 10th and 90th percentiles (whiskers), and outliers (circles).

Printer-friendly Version

Interactive Discussion

[Title Page](#) [Abstract](#) [Introduction](#)
[Conclusions](#) [References](#)
[Tables](#) [Figures](#)
[◀](#) [▶](#)
[◀](#) [▶](#)
[Back](#) [Close](#)
[Full Screen / Esc](#)

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.




Fig. 4. Nitrate (NO_3^-) deposition ($\text{mg NO}_3^-/\text{m}^2/\text{event}$) (top panel) and ammonium (NH_4^+) deposition ($\text{mg NH}_4^+/\text{m}^2/\text{event}$) (bottom panel) as in Fig. 3.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.





Fig. 5. Chloride (Cl^-) deposition ($\text{mg}/\text{m}^2/\text{event}$) (top panel) and sodium (Na^+) deposition ($\text{mg}/\text{m}^2/\text{event}$) (bottom panel) as in Fig. 3.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.




Fig. 6. Plot of the ratio of chloride (Cl) and sodium (Na) fluxes (top panel) and ratio of sulfate (SO_4) and sodium (Na) fluxes (bottom panel) at Pensacola Bay sites between November 2004 and December 2007. The seawater mass ratios (1.79 Cl:Na and 0.25 SO_4 :Na) are plotted as dashed lines.




Fig. 7. Plot of average sodium (Na^+) deposition ($\text{mg/m}^2/\text{y}$) versus distance from Gulf (km) for Pensacola Bay and NADP sites in the Gulf of Mexico.

Atmospheric deposition in the Pensacola Bay watershed

J. M. Caffrey et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)