

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Chemistry of hydrogen oxide radicals (HO_x) in the Arctic troposphere in spring

J. Mao¹, D. J. Jacob^{1,2}, M. J. Evans³, J. R. Olson⁴, X. Ren⁵, W. H. Brune⁶, J. M. S. Clair⁷, J. D. Crounse⁷, K. M. Spencer⁷, M. R. Beaver⁷, P. O. Wennberg^{8,9}, M. J. Cubison¹⁰, J. L. Jimenez¹⁰, A. Fried¹¹, P. Weibring¹¹, J. G. Walega¹¹, S. R. Hall¹², A. J. Weinheimer¹², R. C. Cohen¹³, G. Chen⁴, J. H. Crawford⁴, L. Jaeglé¹⁴, J. A. Fisher², R. M. Yantosca¹, P. Le Sager^{1,*}, and C. Carouge¹

¹School of Engineering and Applied Sciences, Harvard University, Cambridge, MA, USA

²Department of Earth and Planetary Sciences, Harvard University, Cambridge, MA, USA

³School of Earth and the Environment, University of Leeds, Leeds, LS2 9JT, UK

⁴Science Directorate, NASA Langley Research Center, Hampton, VA, USA

⁵Rosenstiel School of Marine and Atmospheric Science, University of Miami, Miami, FL, USA

⁶Department of Meteorology, Pennsylvania State University, University Park, PA, USA

⁷Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, CA, USA

⁸Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA, USA

ACPD

10, 6955–6994, 2010

**Chemistry of
hydrogen oxide
radicals in the Arctic
troposphere in spring**

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

⁹Division of Engineering and Applied Science, California Institute of Technology, Pasadena, CA, USA

¹⁰Department of Chemistry and Biochemistry and Cooperative Institute for Research in the Environmental Sciences (CIRES), University of Colorado at Boulder, Boulder, CO, USA

¹¹Earth Observing Laboratory, National Center for Atmospheric Research, Boulder, CO, USA

¹²Atmospheric Chemistry Division, National Center for Atmospheric Research, Boulder, CO, USA

¹³Department of Chemistry and Department of Earth and Planetary Science, University of California Berkeley, Berkeley, CA, USA

¹⁴Department of Atmospheric Sciences, University of Washington, Seattle, Washington, USA

*now at: KNMI, Chemistry and Climate Division, De Bilt, The Netherlands

Received: 3 March 2010 – Accepted: 4 March 2010 – Published: 11 March 2010

Correspondence to: J. Mao (mao@fas.harvard.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

**Chemistry of
hydrogen oxide
radicals in the Arctic
troposphere in spring**

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

We use observations from the April 2008 NASA ARCTAS aircraft campaign to the North American Arctic, interpreted with a global 3-D chemical transport model (GEOS-Chem), to better understand the sources and cycling of hydrogen oxide radicals ($\text{HO}_x \equiv \text{H} + \text{OH} + \text{peroxy radicals}$) and their reservoirs ($\text{HO}_y \equiv \text{HO}_x + \text{peroxides}$) in the springtime Arctic atmosphere. We find that a standard gas-phase chemical mechanism overestimates the observed HO_2 and H_2O_2 concentrations. Computation of HO_x and HO_y gas-phase chemical budgets on the basis of the aircraft observations also indicates a large missing sink for both. We hypothesize that this could reflect HO_2 uptake by aerosols, favored by low temperatures and relatively high aerosol loadings, through a mechanism that does not produce H_2O_2 . Such a mechanism could involve HO_2 aqueous-phase reaction with sulfate (58% of the ARCTAS submicron aerosol by mass) to produce peroxymonosulfate (HSO_5^-) that would eventually convert back to sulfate and return water. We implemented such an uptake of HO_2 by aerosol in the model using a standard reactive uptake coefficient parameterization with $\gamma(\text{HO}_2)$ values ranging from 0.02 at 275 K to 0.5 at 220 K. This successfully reproduces the concentrations and vertical distributions of the different HO_x species and HO_y reservoirs. HO_2 uptake by aerosol is then a major HO_x and HO_y sink, decreasing mean OH and HO_2 concentrations in the Arctic troposphere by 48% and 45% respectively. Circumpolar budget analysis in the model shows that transport of peroxides from northern mid-latitudes contributes 50% of the HO_y source above 6 km, and cloud chemistry and deposition of H_2O_2 account together for 40% of the HO_y sink below 3 km. Better rate and product data for HO_2 uptake by aerosol are needed to understand this role of aerosols in limiting the oxidizing power of the Arctic atmosphere.

ACPD

10, 6955–6994, 2010

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

1 Introduction

Radiative forcing by aerosol and tropospheric ozone pollution transported from mid-latitudes may be an important driver of recent Arctic warming (Quinn et al., 2008; Shindell et al., 2008). This pollution is strongest in spring (Scheuer et al., 2003) and is modulated by chemical reactions serving as sources or sinks of aerosols and ozone. The Arctic photochemical environment in spring is characterized by polar sunrise, low sun angles, intense cold, and underlying ice surface. Considerable attention has focused on halogen radical photochemistry under these conditions as a rapid sink for ozone (Simpson et al., 2007), but this appears to be important only in the shallow boundary layer where sea ice provides a halogen radical source (Wagner et al., 2001). Hydrogen oxide radicals ($\text{HO}_x \equiv \text{H} + \text{OH} + \text{peroxy radicals}$) have a more pervasive effect in the tropospheric column but the chemistry of these radicals in the Arctic spring has received little study. The OH radical is the principal atmospheric oxidant, affecting both aerosols and ozone in complex ways. Peroxy radical reactions with nitric oxide (NO) are the main chemical source of tropospheric ozone. We use here observations from the April 2008 NASA Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS) aircraft campaign (Jacob et al., 2009), interpreted with a global 3-D chemical transport model (GEOS-Chem CTM), to better understand the sources and cycling of HO_x radicals in the springtime Arctic atmosphere.

HO_x radicals originate from water vapor. The main pathway involves oxidation by the high-energy O (^1D) atom produced from photolysis of ozone:



The OH atoms cycle with peroxy radicals, driving various HO_x -catalyzed mechanisms for atmospheric oxidation and ozone formation and loss. Oxidation of methane and other volatile organic compounds (VOCs) yields formaldehyde (HCHO), which photolyzes to produce additional HO_x radicals and amplify the original source:

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Loss of HO_x eventually takes place by radical-radical reactions. These reactions can return water vapor in which case the loss is terminal, or they may produce reservoir species such as hydrogen peroxide (H_2O_2) and methyl hydrogen peroxide (CH_3OOH). The peroxides can photolyze to return HO_x radicals. Alternatively, they can be converted to water or deposited resulting in a terminal HO_x sink. It is useful to define an expanded hydrogen oxides family $\text{HO}_y \equiv \text{HO}_x + \text{reservoirs}$ (Jaeglé et al., 2001), where the reservoirs include mainly peroxides but also some other minor reservoir species such as nitrous acid (HONO). The HO_x budget can then be understood on the basis of the HO_y budget and the chemical cycling within the HO_y family, governed in part by reactions involving nitrogen oxide radicals ($\text{NO}_x \equiv \text{NO} + \text{NO}_2$) (Jaeglé et al., 2001). The lifetime of HO_y against conversion to water vapor is of the order of a few days, so that transport of HO_y reservoir species on convective and synoptic scales can modulate the supply of HO_x radicals (Jaeglé et al., 1997; Prather and Jacob, 1997; Müller and Brasseur, 1999).

Past studies of HO_x chemistry in Arctic spring have mainly been from surface sites. They have pointed out the importance of HO_x radical production from photochemically driven snow emissions of H_2O_2 (Hutterli et al., 2001; Jacobi et al., 2002, 2004), HCHO (Sumner and Shepson, 1999; Sumner et al., 2002), and HONO (Zhou et al., 2001). They have identified a large photochemical emission of NO_x from snow (Honrath et al., 1999; Ridley et al., 2000) that plays an important role in HO_x cycling (Yang et al., 2002; Chen et al., 2004). Another unique aspect of HO_x chemistry in the boundary layer is the interaction with halogen radicals. These interactions include HO_x production from $\text{Br} + \text{HCHO}$ (Evans et al., 2003), additional HO_y reservoirs such as HOBr (Bloss et al., 2005), and additional processes for cycling between HO_2 and OH (Simpson et al., 2007).

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The Arctic boundary layer is very shallow (~ 100 m) and capped by a strong thermal inversion (Kahl, 1990). The atmosphere above is more relevant for the impacts of HO_x chemistry on the Arctic troposphere. It had received little exploration prior to ARCTAS, due to the requirement of an aircraft with comprehensive chemical payload. The Tropospheric Ozone Production about the Spring Equinox (TOPSE) aircraft campaign conducted a series of flights in the North American Arctic from February to May of 2000 including measurements of total peroxy radicals (Cantrell et al., 2003a), HCHO (Fried et al., 2003), and peroxides (Snow et al., 2003) up to 8 km altitude. Photochemical model calculations constrained with these data found Reactions (R1) and (R3) to be the major HO_x sources (Wang et al., 2003). However, the model greatly overestimated the observed concentrations of peroxy radicals and H_2O_2 (Cantrell et al., 2003b; Wang et al., 2003) while underestimating HCHO (Fried et al., 2003).

Observations from the ARCTAS aircraft expand greatly on TOPSE in terms of both coverage and chemical payload. ARCTAS vertical profiles extend from the boundary layer to the stratosphere. The payload included measurements of HO_x radicals by two independent methods to resolve experimental uncertainty (Chen et al., 2010). It also included an extensive suite of HO_x precursors, reservoirs, and related species. As we will see, this ensemble of observations offers strong constraints and a new perspective on the factors controlling HO_x concentrations in the Arctic spring troposphere.

2 Data and model

2.1 The ARCTAS campaign

The ARCTAS spring campaign took place from 1–21 April 2008 (Jacob et al., 2009). It included a DC-8 aircraft with a detailed chemical and aerosol payload and a P-3 aircraft with a detailed aerosol and radiation payload. Both were based in Fairbanks, Alaska (65 N, 148 W). We focus our attention on the DC-8 chemical data but will also make reference to the P-3 aerosol data. The DC-8 conducted nine flights in the North

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



American Arctic totaling 73 flight hours. These included sorties out of Fairbanks as well as transit flights to and from Thule, Greenland (77 N, 69 W) and Iqaluit, Nunavut (64 N, 69 W). Almost all the data were collected north of 60 N. All flights included frequent vertical profiling from 100 m to 12 km altitude.

5 One of the major goals of ARCTAS was to better understand radical photochemistry in the Arctic. The DC-8 payload included measurements of HO_x radicals, NO_x radicals, H₂O₂, CH₃OOH, HNO₄, O₃, H₂O, VOCs, HCHO, aerosol composition, and photolysis frequencies (Jacob et al., 2009). OH and HO₂ concentrations were measured by two independent techniques, Laser Induced Fluorescence (LIF) and Chemical Ionization Mass Spectrometry (CIMS). There were instrumental difficulties with the CIMS HO₂ measurement but intercomparison for OH shows a median ratio (OH)_{CIMS}/(OH)_{LIF}=0.73 (*R*=0.51), which agrees within the stated accuracies (32% for LIF, 65% for CIMS) (Chen et al., 2010). We rely on the LIF measurements here as they covered 96% of the flight hours. We use 1-min average data with reported accuracies in parentheses: OH (32%), HO₂(32%), H₂O₂(±(40%+100 pptv)), CH₃OOH (±(40%+50 pptv)), ozone (3%), HCHO (12%±26 pptv), NO (10%±5 pptv), NO₂(5%±5 pptv), HNO₄(30%±15 pptv). We also make use of fine aerosol data including sulfate (34%) and ammonium (34%). The NO₂ LIF measurement includes methylperoxynitrate (CH₃OONO₂) decomposing in the inlet and this represents a significant positive artifact in the upper troposphere (Browne et al., 2010). We correct for it here by using local CH₃OONO₂/NO₂ ratios from the GEOS-Chem simulation. Mean ratios are 0.03 at 0–3 km, 0.17 at 3–6 km, and 1.4 above 6 km.

25 Several halogen-catalyzed ozone and mercury depletion events were observed during ARCTAS (Salawitch et al., 2010), but all were confined below 500 m. No obvious signature of either bromine or chlorine radical chemistry was found above 500 m from the measured soluble bromide (Liao et al., 2010), BrO (Neuman et al., 2010) or VOC indicators (Fried et al., 2010). We neglect the effect of halogen chemistry in our modeling of the ARCTAS HO_x data but will comment on its possible role in the boundary layer.

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2.2 GEOS-Chem model

GEOS-Chem is a global 3-D chemical transport model driven by assimilated meteorological observations from the Goddard Earth Observing System (GEOS-5) of the NASA Global Modeling and Assimilation Office (GMAO) (Bey et al., 2001). We apply here GEOS-Chem version 8-01-04 (<http://acmg.seas.harvard.edu/geos/index.html>) to simulation of the ARCTAS period (April 2008). The GEOS-5 meteorological data have 6-h temporal resolution (3-h for surface variables and mixing depths) with $0.5^\circ \times 0.667^\circ$ horizontal resolution and 72 vertical layers from the surface to 0.01 hPa. We regrid here the meteorological data to 2° latitude \times 2.5° longitude for input to GEOS-Chem. The model is initialized with a 1-year simulation from January 2007 to January 2008 with $4^\circ \times 5^\circ$ resolution, and from January 2008 on with $2^\circ \times 2.5^\circ$ resolution. Our analysis of the GEOS-Chem simulation focuses on HO_x chemistry. The same GEOS-Chem simulation is applied in companion papers to interpretation of ARCTAS observations for CO (Fisher et al., 2010a), SO_2 and sulfate (Fisher et al., 2010b), carbonaceous aerosols (Wang et al., 2010), methane (Pickett-Heaps et al., 2010), and mercury (Holmes et al., 2010).

We use the standard GEOS-Chem simulation of ozone- NO_x - HO_x -VOC-aerosol chemistry as described for example by Park et al. (2006). We updated the chemical mechanism with compiled data from the Jet Propulsion Laboratory (Sander et al., 2006) (hereinafter “JPL06”) and the International Union of Pure and Applied Chemistry (Atkinson et al., 2006) (hereinafter “IUPAC06”). We implemented the Fast-JX radiative transfer code for calculation of photolysis rate constants (Wild et al., 2000; Bian and Prather, 2002), including updates to absorption cross-sections and quantum yields from JPL06. Total ozone columns used as input to Fast-JX are from daily measurements by the Ozone Monitoring Instrument (OMI) satellite instrument with $1^\circ \times 1^\circ$ resolution (<ftp://toms.gsfc.nasa.gov/pub/omi/data/ozone/Y2008/>). The range of ozone columns during ARCTAS was 380~430 Dobson Units. Surface albedo used in Fast-JX is from the Total Ozone Mapping Spectrometer (TOMS) satellite monthly climatology

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

with $1^\circ \times 1.25^\circ$ resolution (Herman and Celarier, 1997).

A major topic of this paper will be the role of HO_2 uptake by aerosol. GEOS-Chem includes a general representation for first-order uptake of gases by aerosols parameterized by a reactive uptake coefficient γ and applied to the model aerosol fields (Martin et al., 2003). The standard model includes aerosol uptake of NO_2 , NO_3 , and N_2O_5 (Jacob, 2000; Evans and Jacob, 2005) and aqueous-phase reaction of H_2O_2 with SO_2 in cloud (Park et al., 2004). Earlier versions also included aerosol uptake of HO_2 (Martin et al., 2003a), but this was removed in v7-04-06 (and hence in the v8-01-04 version we used) on the basis of laboratory data indicating low γ values in the absence of transition metal catalysts (Thornton and Abbatt, 2005; Sauvage et al., 2007). More recent standard versions of GEOS-Chem (v8-02-01 and beyond), developed after this work was initiated, include HO_2 uptake following Thornton et al. (2008). As we will show below, the ARCTAS observations suggest an important role for aerosol uptake of HO_2 under the cold, low-light, and relatively aerosol-rich conditions of Arctic spring. The GEOS-Chem simulation of aerosols during ARCTAS is evaluated by Fisher et al. (2010b) and Wang et al. (2010). It provides in general a good match to observations.

Anthropogenic emissions in GEOS-Chem are as described in van Donkelaar et al. (2008). A prominent feature of ARCTAS flights was the influence of Siberian fire plumes (Warneke et al., 2009). Daily biomass burning emissions for 2008 with $1^\circ \times 1^\circ$ resolution are specified from the Fire Locating and Monitoring of Burning Emissions (FLAMBE) emission inventory (Reid et al., 2009) constrained by GOES and MODIS fire count data. Further details on model emissions are given by Fisher et al. (2010a).

The model wet deposition scheme is described by Liu et al. (2001). It includes wet scavenging in convective updrafts as well as grid-resolved first-order rainout and washout. Of particular interest here is the representation of peroxide and HCHO scavenging. For warm clouds ($T > 268\text{ K}$), H_2O_2 , CH_3OOH , and HCHO are scavenged by liquid water based on their Henry's law constants. For mixed clouds ($248 < T < 268\text{ K}$), precipitation is assumed to take place by riming of liquid cloud droplets with retention efficiencies $R_{\text{H}_2\text{O}_2}=0.05$, $R_{\text{CH}_3\text{OOH}}=0.02$, and $R_{\text{HCHO}}=0.02$ (Mari et al., 2000). In cold

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

clouds ($T < 248$ K), scavenging of H_2O_2 takes place by co-condensation on ice surfaces while scavenging of CH_3OOH and HCHO are considered negligible (Mari et al., 2000).

Dry deposition is calculated using a standard resistance-in-series scheme (Wesely, 1989) applied to the local surface. For snow and ice the deposition velocity of H_2O_2 is in the range $0.1\text{--}0.3\text{ cm s}^{-1}$, while dry deposition of CH_3OOH and HCHO are negligible. Snow emission could offset dry deposition for the above species (Frey et al., 2005; Frey et al., 2006) but is not taken into account here as its effect would be confined to the shallow boundary layer. For the same reason we find dry deposition to be unimportant from a tropospheric column perspective.

All comparisons between model and observations use model output sampled along the flight tracks and at the flight time with 15-min time resolution. We exclude all measurements in the stratosphere as diagnosed by an ozone/CO molar ratio greater than 1.25. This excludes 72% of the data above 10 km, 41% of the data at 8–10 km, and 9% of the data at 6–8 km. We also exclude all measurements at latitudes lower than 60N. In order to assess the effect of GEOS-Chem errors for species driving HO_x chemistry (such as ozone, CO, NO), we compare results to those of a gas-phase photochemical box model (Olson et al., 2004) constrained locally by the ARCTAS observations (Olson et al., 2010). Similar comparisons between GEOS-Chem and this box model have been conducted in previous aircraft campaigns (Olson et al., 2004; Hudman et al., 2007; Zhang et al., 2008).

Previous comparisons of the GEOS-Chem HO_x simulation to aircraft LIF measurements from the same Pennsylvania State University group have been reported for the NASA INTEX-A campaign over North America (summer 2004) and the NASA INTEX-B campaign over the North Pacific (spring 2006). Hudman et al. (2007) reported a model overestimate of 30~60% for both OH and HO_2 in INTEX-A, but subsequent recalibration of the measurements resolved the discrepancy (Ren et al., 2008). Zhang et al. (2008) found no model bias for HO_2 in INTEX-B but a 27% high bias for OH. The global mean tropospheric OH concentration in our ARCTAS simulation is $10.3 \times 10^5 \text{ molecules cm}^{-3}$, consistent with the multimodel annual mean of

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

$11.1 \pm 1.7 \times 10^5$ molecule cm^{-3} from the intercomparison by Shindell et al. (2006).

3 Median distributions

Figure 1 presents median vertical profiles of OH, HO₂, H₂O₂, CH₃OOH, NO, NO₂, HCHO and HNO₄ concentrations for 1 km vertical bins during ARCTAS. We compare here the observed profiles (black) to results from the standard GEOS-Chem simulation not including HO₂ uptake by aerosols (dashed green line). Also shown in Fig. 1 are model simulations including HO₂ uptake, which will be discussed in Sect. 4. Most data were collected under sunlit conditions, between 08:00 and 18:00 local time. Stratospheric data were excluded as described in Sect. 2. Mean observed temperature is 256 K at 0–3 km, 243 K at 3–6 km, and 226 K above 6 km, consistent with the model.

Observed concentrations of OH are relatively low (20 ppqv = 5×10^5 molecules cm^{-3} in surface air) reflecting the low water vapor, low solar elevation, and thick ozone columns characteristic of Arctic spring. The model reproduces the observed OH concentrations including the vertical gradient with no significant bias.

Observed concentrations of HO₂, H₂O₂, and CH₃OOH show little altitude dependence, consistent with TOPSE (Wang et al., 2003) but in contrast to measurements in the tropics and northern mid-latitudes that show decreases with altitude driven by water vapor (Cohan et al., 1999; O'Sullivan et al., 1999; Hudman et al., 2007; Snow et al., 2007; Zhang et al., 2008). The model reproduces this lack of vertical structure in the Arctic spring observations and attributes it to a strong vertical gradient of UV radiation (low solar angles, thick ozone columns) compensating for the water vapor gradient and the influx of peroxides from northern mid-latitude into upper troposphere.

The standard simulation overestimates HO₂ and H₂O₂ by up to a factor of 2. Olson et al. (2010) find a similar discrepancy for HO₂ in their box model results, and show that it cannot be resolved by adjusting observed concentrations of other species (such as NO or BrO) within their measurement uncertainties. A similar overestimate of HO₂ and H₂O₂ was previously found in TOPSE when comparing box model calculations to

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



observations (Cantrell et al., 2003b; Wang et al., 2003). We propose below that aerosol uptake of HO₂ leading to a terminal sink for HO_y could be particularly efficient in the Arctic and explain the discrepancy between model and observations.

Median observed NO_x concentrations increase from 13 pptv in the boundary layer to 21 pptv in the upper troposphere. The model is consistent, increasing from 16 pptv in the boundary layer to 33 pptv in the upper troposphere. April observations from TOPSE at 60–80 N showed a mean NO concentration of 6 pptv (Wang et al., 2003), consistent with the ARCTAS observations and with the model. A sensitivity model simulation with no fuel emissions shows a 40% mean decrease of NO_x along the ARCTAS flight tracks, while a sensitivity simulation with no biomass burning emissions shows only a 5% decrease. We conclude that a large fraction of the NO_x in ARCTAS was anthropogenic.

Peroxynitric acid (HNO₄) is an important reservoir for HO_x in the upper troposphere at northern mid-latitudes (Jaeglé et al., 2001). ARCTAS observations are below the detection limit in the lower troposphere due to thermal dissociation but increase to 30 pptv in the upper troposphere. The model reproduces the observed concentrations and vertical gradient within the measurement uncertainty.

Observed HCHO decreases with altitude, from 140 pptv near the surface to 25 pptv in the upper troposphere. TOPSE observations by a similar absorption spectrometer using a tunable lead-salt diode laser averaged 95 pptv below 0.2 km and 60 pptv at 6–8 km (Fried et al., 2003). A different laser source, based on tunable difference frequency generation (DFG), was employed in ARCTAS (Weibring et al., 2007; Richter et al., 2009). Model values decrease from 50 pptv near the surface to 30 pptv in the upper troposphere. The discrepancy below 3 km is outside the stated precision of the measurements (26 pptv). Olson et al. (2010) find a similar discrepancy in their box model simulation of the ARCTAS data (with the point-by-point median discrepancies range up to 75 pptv for altitudes between 0.5 and 3 km), and Fried et al. (2003) also reported a low model bias relative to the TOPSE data. Previous GEOS-Chem evaluation with observed HCHO vertical profiles at northern mid-latitudes using a tunable lead-salt diode laser by the same investigator show no such discrepancy (Palmer et al.,

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2003; Millet et al., 2006). Halogen radical chemistry and snow emissions of HCHO could provide additional HCHO that is not included in the model at perhaps altitudes as high as 0.5 km. However, this does not explain discrepancies between 0.5 and 3 km given the lifetime of HCHO (hours). Singh et al. (2000) suggested a heterogeneous conversion from CH₃OH to HCHO, but we find no correlation between these two species below 4 km ($R < 0.01$). Although these discrepancies are small, it is important that they are resolved. Fried et al. (2010) will further discuss this. As pointed out in Sect. 5, the source of HO_x and HO_y implied by the observed HCHO appears inconsistent with independent calculations of HO_x and HO_y sinks from the ARCTAS observations.

4 HO₂ uptake by aerosols

4.1 Parameterization of uptake

A prominent feature of the comparison between model and observations in Fig. 1 is the overestimate of HO₂. Such a discrepancy has been reported in a number of previous model studies and tentatively attributed to HO₂ uptake by aerosols (Cantrell et al., 1996a, 1996b; Plummer et al., 1996; Jaeglé et al., 2000; Kanaya et al., 2000; Sommariva et al., 2004; de Reus et al., 2005; Sommariva et al., 2006). This uptake has been reported in a number of laboratory studies but rates and mechanism are uncertain (Jacob, 2000). It could be particularly important in the Arctic because of the low temperature, relatively high aerosol, and slow photochemical cycling.

Figure 2 summarizes literature data of the $\gamma(\text{HO}_2)$ reactive uptake coefficient $\gamma(\text{HO}_2)$ for different surface types. $\gamma(\text{HO}_2)$ is defined as the fraction of HO₂ collisions with the aerosol surface resulting in reaction. Consistently high values ($\gamma(\text{HO}_2) > 0.2$) are observed for Cu-doped aqueous surfaces. Soluble copper is known to drive rapid catalytic conversion of $\gamma(\text{HO}_2)$ to H₂O₂ by redox chemistry (Jacob, 2000). Other values for aqueous surfaces at room temperature are also relatively high ($\gamma(\text{HO}_2)$ in the range 0.07–0.2) except for concentrated H₂SO₄ ($\gamma(\text{HO}_2) < 0.01$). However, $\gamma(\text{HO}_2)$ for con-

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

centrated H_2SO_4 increases rapidly with decreasing temperature and exceeds 0.2 at 240 K. Reactive uptake coefficients in general increase with decreasing temperature, reflecting negative temperature dependences of both the mass accommodation coefficient α on the surface and the solubility constant (Kolb et al., 1995). Solid surfaces do not take up HO_2 significantly. To our knowledge no data are available for $\gamma(\text{HO}_2)$ on organic surfaces.

A high reactive uptake probability ($\gamma(\text{HO}_2) > 0.1$) is needed for aerosol uptake to compete in the atmosphere with other chemical sinks for HO_x . This requires that the aerosol be aqueous. Relative humidity (RH) relative to liquid water was typically in the range 40–80% during ARCTAS. Aerosol measurements aboard the P-3 show that non-refractory submicron particles contributed more than 90% of total surface area (McNaughton et al., 2010). High-resolution Aerosol Mass Spectrometer (AMS; DeCarlo et al., 2006) measurements aboard the DC-8 show an average mass composition for non-refractory submicron particles of 58% sulfate, 32% organic aerosol, 6% ammonium, 3% nitrate, and 0.7% chloride (Cubison et al., 2008). An ammonium vs. sulfate molar plot for the AMS data (Fig. 3) shows dominance of the acidic NH_4HSO_4 form, although in some cases the aerosol was close to sulfuric acid while in other cases it was fully neutralized as $(\text{NH}_4)_2\text{SO}_4$. Because of metastability of the aqueous phase, both NH_4HSO_4 and $(\text{NH}_4)_2\text{SO}_4$ are expected to remain aqueous over the range of RH values experienced in ARCTAS (Onasch et al., 1999; Martin et al., 2003b; Colberg et al., 2004). Sulfuric acid is aqueous under all conditions. In addition, Parsons et al. (2004) found that the crystallization RH of ammonium sulfate aerosol decreases as the organic fraction increases. We assume therefore that the aerosol surface area in ARCTAS was mainly contributed by aqueous particles.

To investigate the role of HO_2 uptake by aerosol in our ARCTAS simulations, we included the $\gamma(\text{HO}_2)$ parameterization of Thornton et al. (2008) as implemented in the most recent standard versions of GEOS-Chem (v8-02-01 and beyond). This parameterization describes HO_2 uptake by aqueous aerosol as driven by acid-base dissociation followed by the $\text{HO}_2(\text{aq}) + \text{O}_2^-$ reaction at an assumed pH 5, producing H_2O_2 that

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

then volatilizes to the gas phase. It is clearly inconsistent with the prevailing acidic conditions observed in ARCTAS (Fig. 3), and is also theoretically incorrect since it describes HO₂ uptake as a second-order process while the $\gamma(\text{HO}_2)$ parameterization intrinsically describes a first-order process. Nevertheless, as shown in Fig. 2 (solid line), it yields values of $\gamma(\text{HO}_2)$, that are consistent with those measured in the laboratory for acidic surfaces, ganging from less than 0.05 near the surface to 0.4 in the upper troposphere. Its temperature dependence (mainly driven by the Henry's law constant for HO₂) is consistent with the laboratory data for concentrated H₂SO₄ (Fig. 2). We adopt the Thornton et al. (2008) scheme here to compute $\gamma(\text{HO}_2)$ for want of anything better and because it fits the overall laboratory data for acid aerosols.

The fate of HO₂ in aerosol phase is generally assumed to involve conversion to H₂O₂ followed by H₂O₂ volatilization (Jacob, 2000). However, this would exacerbate the overestimate of H₂O₂ in ARCTAS (Fig. 1). One possible solution would be protonation of H₂O₂ to HOOH₂⁺, a very strong oxidant (Oiestad et al., 2001), which would rapidly react and convert to H₂O. However, this requires normal acidity (pH<0) to be effective (Bach and Su, 1994). Such a mechanism could conceivably take place in concentrated H₂SO₄ aerosols, but not in the less acidic aerosol that prevailed under ARCTAS conditions (Fig. 3).

Another possible solution is that HO₂ uptake may not produce H₂O₂. Cooper and Abbatt (1996) proposed that HO₂ could react with HSO₄⁻, forming the peroxymonosulfate radical SO₅⁻:



The possible atmospheric chemistry of the SO₅⁻ radical is discussed by Jacob (1986), who proposed that the dominant sinks would be the exothermic reactions with O₂⁻, HCOO⁻, and HSO₃⁻, all producing the peroxide HSO₅⁻ (peroxymonosulfate). Peroxymonosulfate is the first dissociated form of Caro's acid (H₂SO₅, highly hygroscopic solid at room temperature, melting point 45 °C), which has first and second acid dissociation constants (pK_a) of 0.4 and 9.1 (Elias et al., 1994). As a peroxide, HSO₅⁻ is a member

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of the HO_y family and so its formation does not necessarily represent a terminal sink of HO_x. However, Jacob (1986) suggested that the subsequent fate of HSO₅⁻ in an acidic environment might follow Reactions (R7) and (R8), terminally losing HO_y by conversion to water:



- 5 Another possible mechanism for uptake of HO₂ by sulfate aerosols might involve the formation of an HO₂ complex. Miller and Francisco (2001) found from quantum chemical calculations that a stable HO₂-H₂SO₄ complex can be formed in the gas as well as in the aerosol phase. HSO₄⁻ has similar potential for bonding with HO₂ molecules (C. Miller, personal communication, 2009). The fate of these complexes is unknown.
- 10 They must not decompose to the original reactants if they are to represent an actual HO_x or HO_y sink. One possibility would be conversion to SO₅⁻ by Reaction (R6), with subsequent chemistry forming HSO₅⁻ and eventually returning water as described above.

4.2 Application to the ARCTAS simulation

- 15 Figure 1 shows the results of two simulations, one with HO₂ aerosol uptake producing H₂O₂ (γ(HO₂ → 0.5H₂O₂), dashed blue line), and the other with HO₂ aerosol uptake leading to a permanent HO_y sink as postulated above (γ(HO₂ → products), red line). The first simulation worsens the overestimate of H₂O₂, as aerosol uptake now competes with other HO_x sinks such as OH+HO₂ and CH₃O₂+HO₂ that do not produce
- 20 H₂O₂. The correction to HO₂ is also insufficient in the upper troposphere as H₂O₂ can be recycled to HO_x by photolysis. The second simulation provides a much better fit to the observations for both HO₂ and H₂O₂ (within their measurement uncertainties), and

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

also improves the fit for OH and CH₃OOH, while not significantly affecting the fits for the other species. OH concentrations decrease by up to 58% in the upper troposphere.

Figure 4 shows scatterplots of simulated vs. observed OH, HO₂, H₂O₂, and CH₃OOH concentrations for the model simulation with $\gamma(\text{HO}_2 \rightarrow \text{products})$ and the ensemble of tropospheric observations in ARCTAS. The slopes of the reduced-major-axis regression lines are within the measurement accuracy for all species. Correlation coefficients for OH ($R=0.61$) and HO₂ ($R=0.81$) are only slightly improved from the gas-phase-only simulation ($R=0.58$ and $R=0.78$ respectively). Correlations are weak for H₂O₂ ($R=0.45$) and CH₃OOH ($R=0.50$), which might reflect the narrow dynamic range. The HO₂ variability is mostly correlated with solar zenith angle in both the observations and the model ($R=-0.80$ in both cases), with additional significant correlations with water vapor ($R=0.30$ observed, $R=0.40$ model) and temperature ($R=0.30$ observed, $R=0.40$ model) above 4 km. The correlation of HO₂ concentrations with temperature offers some supporting evidence for a sink from aerosol uptake. The DC-8 did not include measurements of aerosol surface area that we could correlate to HO₂ concentrations, but in the model we find that temperature is a much stronger driver of $\gamma(\text{HO}_2)$ variability than aerosol surface area. No correlation is found between HO₂ and NO_x concentrations in either the model or the observations.

5 Budget of HO_x radicals in Arctic spring

We now proceed to quantify the budgets of HO_x and HO_y in Arctic spring as constrained by the ARCTAS observations and derived from the model. We use for that purpose the model including terminal loss of HO_y from reactive uptake of HO₂ by aerosols as described in Sect. 4.1. Even though the process is uncertain, it represents our best hypothesis for explaining the HO_x and peroxide observations in ARCTAS.

Figure 5 shows the median vertical profiles of HO_x sources and sinks computed from the observed ARCTAS concentrations, gas-phase reaction rate constants from JPL06 and IUPAC06, and $\gamma(\text{HO}_2 \rightarrow \text{products})$ from Thornton et al. (2008). The CH₃O₂ con-

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

centration is assumed to be 30% of that of HO_2 on the basis of the model $\text{HO}_2/\text{CH}_3\text{O}_2$ ratio. HO_2 aerosol uptake is computed using local model values for γ and aerosol surface area. $\text{O} (^1\text{D}) + \text{H}_2\text{O}$ and HCHO photolysis are the major HO_x sources below 4 km. Above that altitude the photolysis of H_2O_2 becomes dominant, a remarkable feature that has not been reported to our knowledge anywhere else in the troposphere. It reflects the low OH concentrations in Arctic spring and therefore the dominance of photolysis as a H_2O_2 sink rather than reaction with OH (Fig. 5).

The $\text{HO}_2 + \text{HO}_2$ reaction is the principal HO_x sink in the lower troposphere but HO_2 uptake by aerosol becomes dominant above 5 km. HO_x sinks from $\text{CH}_3\text{O}_2 + \text{HO}_2$ and $\text{OH} + \text{HO}_2$ are relatively small. The NO_x -based HO_x sinks including $\text{HO}_2 + \text{NO}_2$, $\text{OH} + \text{HNO}_4$, and $\text{OH} + \text{NO}_2$ are negligibly small because of the low NO_x concentrations, so that ozone production is NO_x -limited throughout the troposphere (Jaeglé et al., 2001). In contrast, previous aircraft campaigns at northern mid-latitudes (SONEX, INTEX-A) found that the NO_x -based reactions dominated the HO_x sink in the upper troposphere, implying NO_x -neutral or NO_x -saturated conditions for ozone production (Jaeglé et al., 2000; Ren et al., 2008). The observed NO_x concentrations in the upper troposphere in these campaigns (medians of 93 pptv in SONEX, 440 pptv in INTEX-A) are much higher than in ARCTAS (33 pptv), reflecting major sources at northern mid-latitudes from convective injection of pollution and lightning (Jaeglé et al., 1998; Allen et al., 2000; Hudman et al., 2007).

We see from Fig. 5 that HO_2 uptake by aerosols is crucial for balancing the HO_x sources with the HO_x sinks independently computed from observations, at least in the upper troposphere. There the main gas-phase sinks ($\text{HO}_2 + \text{HO}_2$, $\text{CH}_3\text{O}_2 + \text{HO}_2$, $\text{OH} + \text{HO}_2$) can balance only 20% of the HO_x source. With HO_2 uptake by aerosol included in the budget, the total HO_x sinks balance 50% of the HO_x sources in the lower troposphere and 70% in upper troposphere. The imbalance in the lower troposphere reflects the high observed HCHO concentrations (Fig. 1), for which we have no explanation. The aerosol sink is ineffective in the lower troposphere because of the relatively high temperatures (average $T = 258 \text{ K}$ at 0–2 km, resulting in $\gamma(\text{HO}_2) = 0.06$).

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Figure 5 also shows the median vertical profiles of instantaneous HO_y source and sink terms, again computed from the measurements. The main HO_y sources are O (¹D)+H₂O and HCHO photolysis. The main sinks are HO₂ uptake by aerosol and the gas-phase OH+CH₃OOH reaction. OH+H₂O₂ is relatively unimportant due to its much slower rate constant ($1.8 \times 10^{-12} \text{ cm}^3 \text{ molecules}^{-1} \text{ s}^{-1}$, no temperature dependence) compared to OH + CH₃OOH ($3.8 \times 10^{-12} \exp(200/T) \text{ cm}^3 \text{ molecules}^{-1} \text{ s}^{-1}$) as given by JPL06. Such a dominance of the gas-phase HO_y sink by the OH+CH₃OOH reaction has not been reported before to our knowledge, except in deep convective outflow where H₂O₂ has been scavenged but not CH₃OOH (Cohan et al., 1999). We attribute it to the low concentrations of NO_x in Arctic spring, suppressing the NO_x-based HO_x sinks (see discussion above) and promoting CH₃OOH formation.

Figure 5 shows an imbalance between HO_y chemical sources and sinks that reverses sign with altitude. The HO_y sink balances 30% of the source below 3 km, 92% at 3–6 km, and 170% above 6 km. The HO_y lifetime is 2–6 d, sufficiently long that additional terms may be relevant in the HO_y budget including long-range transport, wet and dry deposition, and aqueous-phase oxidation of SO₂ by H₂O₂ in clouds. To consider the effect of these terms, we conducted a HO_y budget analysis in the GEOS-Chem model averaged over the 60–90°N circumpolar Arctic cap. Results are shown in Fig. 6. The budget in the model is balanced by mass conservation; the excess of HO_y sources over HO_y sinks in the tropospheric column reflects accumulation of peroxides over the course of April.

We see from Fig. 6 that influx of peroxides from northern mid-latitudes in the model accounts for 50% of the total HO_y source above 6 km and 20% at 3–6 km. This explains the chemical imbalance in the HO_y budget constrained by the ARCTAS observations (Fig. 5). Considering the dominant role of H₂O₂ photolysis as a source of HO_x above 5 km (Fig. 5), this implies a significant contribution of northern mid-latitudes to the HO_x budget of the Arctic free troposphere. Below 3 km, we find in the model that cloud chemistry and deposition of H₂O₂ together account for 40% of the HO_y sink. This helps but is insufficient to correct the chemical imbalance in the HO_y budget constrained by

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the observations. As in the case of the HO_x budget, the residual imbalance reflects the high observed HCHO.

Figure 7 gives a summary diagram of the HO_x and HO_y cycling as represented by our model for Arctic spring. Primary sources include the O (¹D)+H₂O reaction within the region (70%) and transport of peroxides from northern mid-latitudes (30%). Photolysis of HCHO produced from oxidation of methane by OH is a major amplifying source of HO_y, of comparable magnitude to the primary source from O (¹D)+H₂O. HO₂ aerosol uptake accounts for 35% of the HO_y sink. Cycling within the HO_x family (between OH and peroxy radicals) is relatively efficient (chain length=3.4) given the low NO_x concentrations. This is because formation of peroxides to terminate the chain is slow as a result of the low HO_x concentrations.

6 Implications

The ARCTAS observations show a large missing sink of HO_x and HO_y in Arctic spring relative to current understanding. If our hypothesis that this reflects a fast terminal loss of HO₂ to aerosols is correct, then it implies a significant sensitivity of the oxidizing power of the Arctic atmosphere to aerosol perturbations. A measure of this effect is provided by the difference in Fig. 1 between our standard simulation (solid red line) and the gas-only simulation (dashed green line). In the absence of aerosols, OH and HO₂ concentrations would increase on average respectively by 48% and 45% in the tropospheric column, the largest effects being in the upper troposphere where uptake by aerosol is particularly efficient (low temperatures).

Laboratory data show a wide range for the reactive uptake coefficient $\gamma(\text{HO}_2)$, as summarized in Fig. 2. Values are generally high for aqueous surfaces and appear to have a strong temperature dependence. Both of these factors are important for HO₂ uptake in the Arctic and require further study. In particular, we have no information on HO₂ uptake by organic aerosol.

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The HO₂ chemistry in the aerosol phase is another critical issue to resolve. The only product study to our knowledge is that of Loukhovitskaya et al. (2009), who found H₂O₂ to be the main product for solid NaBr surfaces. It is conventionally assumed that uptake by aqueous aerosols would also produce H₂O₂ from the HO₂(aq)+O₂⁻ self-reaction (Thornton et al., 2008) or from catalytic cycles involving transition metal ions (Graedel et al., 1986). We find that an HO₂ uptake mechanism producing H₂O₂ would greatly overestimate the observed H₂O₂ concentrations in ARCTAS. In order to fit the observations, we propose an alternate mechanism involving reaction of HO₂ with acid sulfate to produce peroxymonosulfate (HSO₅⁻). This mechanism and its sensitivity to aerosol acidity need to be tested in the laboratory. HO₂ uptake by aerosols has opposite effects on H₂O₂ depending on whether H₂O₂ is produced or not as a result of uptake. Changes in aerosol types (biomass burning vs. fossil fuel) or aerosol acidity (sulfuric acid vs. ammonium) could thus have large effects on H₂O₂. This may be relevant to explaining the complex long-term trend of H₂O₂ observed in Greenland ice cores (Möller, 1999).

Acknowledgements. The authors would like to thank Scot T. Martin, Cameron S. McNaughton, Hongyu Liu, Charles E. Miller, Richard A. Ferrare, Karl D. Froyd and Daniel M. Murphy for helpful discussions. We also would like to thank Yuhang Wang for providing TOPSE dataset, Huisheng Bian for providing Fast-JX updates and Dirk Richter for contributing to HCHO measurement. J. Mao also thanks David M. Shelow and the NASA DC-8 crew for their generous help with making HO_x measurements on the aircraft. This work was supported by the NASA Tropospheric Chemistry Program.

References

- Allen, D., Pickering, K., Stenchikov, G., Thompson, A., and Kondo, Y.: A three-dimensional total odd nitrogen (NO_y) simulation during sonex using a stretched-grid chemical transport model, J. Geophys. Res.-Atmos., 105, 3851–3876, 2000.
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., Troe, J., and IUPAC Subcommittee: Evaluated kinetic and photochemical

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



data for atmospheric chemistry: Volume II –gas phase reactions of organic species, Atmos. Chem. Phys., 6, 3625–4055, 2006,
<http://www.atmos-chem-phys.net/6/3625/2006/>.

- Bach, R. D. and Su, M. D.: The transition-state for the hydroxylation of saturated-hydrocarbons with hydroperoxonium ion, J. Am. Chem. Soc., 116, 10103–10109, 1994.
- 5 Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. A., Field, B. D., Fiore, A. M., Li, Q. B., Liu, H. G. Y., Mickley, L. J., and Schultz, M. G.: Global modeling of tropospheric chemistry with assimilated meteorology: Model description and evaluation, J. Geophys. Res.-Atmos., 106, 23073–23095, 2001.
- 10 Bian, H. S. and Prather, M. J.: FAST-J2: Accurate simulation of stratospheric photolysis in global chemical models, J. Atmos. Chem., 41, 281–296, 2002.
- Bloss, W. J., Lee, J. D., Johnson, G. P., Sommariva, R., Heard, D. E., Saiz-Lopez, A., Plane, J. M. C., McFiggans, G., Coe, H., Flynn, M., Williams, P., Rickard, A. R., and Fleming, Z. L.: Impact of halogen monoxide chemistry upon boundary layer OH and HO₂ concentrations at a coastal site, Geophys. Res. Lett., 32(4), L06814, doi:10.1029/2004GL022084, 2005.
- 15 Browne E., Wooldrige, P., and Cohen, R.: Σ PNs, CH₃ONO₂ and NO_x Observations: Comparisons to TOPSE Σ PNs/O₃, in preparation, 2010.
- Cantrell, C. A., Shetter, R. E., Gilpin, T. M., and Calvert, J. G.: Peroxy radicals measured during mauna loa observatory photochemistry experiment 2: The data and first analysis, J. Geophys. Res.-Atmos., 101, 14643–14652, 1996a.
- 20 Cantrell, C. A., Shetter, R. E., Gilpin, T. M., Calvert, J. G., Eisele, F. L., and Tanner, D. J.: Peroxy radical concentrations measured and calculated from trace gas measurements in the Mauna Loa observatory photochemistry experiment 2, J. Geophys. Res.-Atmos., 101, 14653–14664, 1996b.
- 25 Cantrell, C. A., Edwards, G. D., Stephens, S., Mauldin, L., Kosciuch, E., Zondlo, M., and Eisele, F.: Peroxy radical observations using chemical ionization mass spectrometry during topse, J. Geophys. Res.-Atmos., 108(D6), 8371, doi:10.1029/2002jd002715, 2003a.
- Cantrell, C. A., Mauldin, L., Zondlo, M., Eisele, F., Kosciuch, E., Shetter, R., Lefer, B., Hall, S., Campos, T., Ridley, B., Walega, J., Fried, A., Wert, B., Flocke, F., Weinheimer, A., Hannigan, J., Coffey, M., Atlas, E., Stephens, S., Heikes, B., Snow, J., Blake, D., Blake, N., Katzenstein, A., Lopez, J., Browell, E. V., Dibb, J., Scheuer, E., Seid, G., and Talbot, R.: Steady state free radical budgets and ozone photochemistry during TOPSE, J. Geophys. Res.-Atmos., 108(D4), 8361, doi:10.1029/2002jd002198, 2003b.
- 30

**Chemistry of
hydrogen oxide
radicals in the Arctic
troposphere in spring**

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Chen, G., Davis, D., Crawford, J., Hutterli, L. M., Huey, L. G., Slusher, D., Mauldin, L., Eisele, F., Tanner, D., Dibb, J., Buhr, M., McConnell, J., Lefer, B., Shetter, R., Blake, D., Song, C. H., Lombardi, K., and Arnoldy, J.: A reassessment of HO_x south pole chemistry based on observations recorded during ISCAT 2000, *Atmos. Environ.*, 38, 5451–5461, doi:10.1016/j.atmosenv.2003.07.018, 2004.

Chen, G., Olson, J., and Crawford, J. H.: Intercomparison results during ARCTAS, in preparation, 2010.

Cohan, D. S., Schultz, M. G., Jacob, D. J., Heikes, B. G., and Blake, D. R.: Convective injection and photochemical decay of peroxides in the tropical upper troposphere: Methyl iodide as a tracer of marine convection, *J. Geophys. Res.-Atmos.*, 104, 5717–5724, 1999.

Colberg, C. A., Krieger, U. K., and Peter, T.: Morphological investigations of single levitated H₂SO₄/NH₃/H₂O aerosol particles during deliquescence/efflorescence experiments, *J. Phys. Chem. A*, 108, 2700–2709, doi:10.1021/jp037628r, 2004.

Cooper, P. L. and Abbatt, J. P. D.: Heterogeneous interactions of OH and HO₂ radicals with surfaces characteristic of atmospheric particulate matter, *J. Phys. Chem.*, 100, 2249–2254, 1996.

Cubison, M. J., Sueper, D., Dunlea, E. J., and Jimenez, J. L.: Submicron aerosol composition during the arctas campaign: Arctic haze, biomass burning, and california pollution, *EOS Transaction AGU*, 89, Fall Meet. Suppl. A11A-0081, 2008.

de Reus, M., Fischer, H., Sander, R., Gros, V., Kormann, R., Salisbury, G., Van Dingenen, R., Williams, J., Zllner, M., and Lelieveld, J.: Observations and model calculations of trace gas scavenging in a dense Saharan dust plume during MINATROC, *Atmos. Chem. Phys.*, 5, 1787–1803, 2005, <http://www.atmos-chem-phys.net/5/1787/2005/>.

DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T., Docherty, K. S., Worsnop, D. R., and Jimenez, J. L.: Field-deployable, high-resolution, time-of-flight aerosol mass spectrometer, *Anal. Chem.*, 78, 8281–8289, doi:10.1021/ac061249n, 2006.

Elias, H., Gotz, U., and Wannowius, K. J.: Kinetics and mechanism of the oxidation of sulfur(IV) by peroxomonosulfuric acid anion, *Atmos. Environ.*, 28, 439–448, 1994.

Evans, M. J., Jacob, D. J., Atlas, E., Cantrell, C. A., Eisele, F., Flocke, F., Fried, A., Mauldin, R. L., Ridley, B. A., Wert, B., Talbot, R., Blake, D., Heikes, B., Snow, J., Walega, J., Weinheimer, A. J., and Dibb, J.: Coupled evolution of BrO_x–ClO_x–HO_x–NO_x chemistry during bromine-

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

catalyzed ozone depletion events in the arctic boundary layer, J. Geophys. Res.-Atmos., 108(D4), 8368, doi:10.1029/2002jd002732, 2003.

Evans, M. J. and Jacob, D. J.: Impact of new laboratory studies of N₂O₅ hydrolysis on global model budgets of tropospheric nitrogen oxides, ozone, and OH, Geophys. Res. Lett., 32, L09813, doi:10.1029/2005gl022469, 2005.

Fisher, J. A., Jacob, D. J., Purdy, M. T., Kopacz, M., Le Sager, P., Carouge, C., Holmes, C. D., Yantosca, R. M., Batchelor, R. L., Strong, K., Diskin, G. S., Fuelberg, H. E., Holloway, J. S., Hyer, E. J., McMillan, W. W., Warner, J., Streets, D. G., Zhang, Q., Wang, Y., and Wu, S.: Source attribution and interannual variability of Arctic pollution in spring constrained by aircraft (ARCTAS, ARCPAC) and satellite (AIRS) observations of carbon monoxide, Atmos. Chem. Phys., 10, 977–996, 2010a, <http://www.atmos-chem-phys.net/10/977/2010/>.

Fisher J. A., Jacob, D. J., Dibb, J. E., Scheuer E., Jimenez, J. L., and Cubison, M. : Sources and acidity of sulfate-nitrate-ammonium aerosol in the Arctic in spring, in preparation, 2010b.

Frey, M. M., Stewart, R. W., McConnell, J. R., and Bales, R. C.: Atmospheric hydroperoxides in west antarctica: Links to stratospheric ozone and atmospheric oxidation capacity, J. Geophys. Res.-Atmos., 110, D23301, doi:10.1029/2005jd006110, 2005.

Frey, M. M., Bales, R. C., and McConnell, J. R.: Climate sensitivity of the century-scale hydrogen peroxide (H₂O₂) record preserved in 23 ice cores from west antarctica, J. Geophys. Res.-Atmos., 111, D21301, doi:10.1029/2005jd006816, 2006.

Fried, A., Wang, Y. H., Cantrell, C., Wert, B., Walega, J., Ridley, B., Atlas, E., Shetter, R., Lefer, B., Coffey, M. T., Hannigan, J., Blake, D., Blake, N., Meinardi, S., Talbot, B., Dibb, J., Scheuer, E., Wingenter, O., Snow, J., Heikes, B., and Ehhalt, D.: Tunable diode laser measurements of formaldehyde during the TOPSE 2000 study: Distributions, trends, and model comparisons, J. Geophys. Res.-Atmos., 108(D4), 8365, doi:10.1029/2002jd002208, 2003.

Fried A., Richter D., Weibring P., Apel, E. C., Gorham, K. A., Walega J., Blake, D. R., Blake, N. J., Orlando, J. J., Weinheimer, A., Knapp, D. J., Huey, L. G., and Meinardi, S.: Airborne Formaldehyde and VOC Measurements during Select Arctic Boundary Layer Runs in the 2008 ARCTAS Study and Estimates of Halogen Atom Mixing Ratios, in preparation, 2010.

Gershenson, Y. M., Grigorieva, V. M., Ivanov, A. V., and Remorov, R. G.: O₃ and OH sensitivity to heterogeneous sinks of HO_x and CH₃O₂ on aerosol particles, Faraday Discuss., 100, 83–100, 1995.

Graedel, T. E., Mandich, M. L., and Weschler, C. J.: Kinetic-model studies of atmospheric

**Chemistry of
hydrogen oxide
radicals in the Arctic
troposphere in spring**

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

droplet chemistry: 2. Homogeneous transition-metal chemistry in raindrops, J. Geophys. Res.-Atmos., 91, 5205–5221, 1986.

Hanson, D. R., Burkholder, J. B., Howard, C. J., and Ravishankara, A. R.: Measurement of OH and HO₂ radical uptake coefficients on water and sulfuric-acid surfaces, J. Phys. Chem., 96, 4979–4985, 1992.

Herman, J. R. and Celarier, E. A.: Earth surface reflectivity climatology at 340–380 nm from toms data, J. Geophys. Res.-Atmos., 102, 28003–28011, 1997.

Holmes, C. D., Jacob, D. J., Corbitt E. S., Soerensen, A. L., and Talbot, R.: Global atmospheric budget of mercury including oxidation of elemental mercury by bromine atoms, in preparation, 2010.

Honrath, R. E., Peterson, M. C., Guo, S., Dibb, J. E., Shepson, P. B., and Campbell, B.: Evidence of NO_x production within or upon ice particles in the greenland snowpack, Geophys. Res. Lett., 26, 695–698, 1999.

Hudman, R. C., Jacob, D. J., Turquety, S., Leibensperger, E. M., Murray, L. T., Wu, S., Gilliland, A. B., Avery, M., Bertram, T. H., Brune, W., Cohen, R. C., Dibb, J. E., Flocke, F. M., Fried, A., Holloway, J., Neuman, J. A., Orville, R., Perring, A., Ren, X., Sachse, G. W., Singh, H. B., Swanson, A., and Wooldridge, P. J.: Surface and lightning sources of nitrogen oxides over the united states: Magnitudes, chemical evolution, and outflow, J. Geophys. Res.-Atmos., 112, D12s05, doi:10.1029/2006JD007912, 2007.

Hutterli, M. A., McConnell, J. R., Stewart, R. W., Jacobi, H. W., and Bales, R. C.: Impact of temperature-driven cycling of hydrogen peroxide (H₂O₂) between air and snow on the planetary boundary layer, J. Geophys. Res.-Atmos., 106, 15395–15404, 2001.

Hutterli, M. A., McConnell, J. R., Chen, G., Bales, R. C., Davis, D. D., and Lenschow, D. H.: Formaldehyde and hydrogen peroxide in air, snow and interstitial air at south pole, Atmos. Environ., 38, 5439–5450, doi:10.1016/j.atmosenv.2004.06.003, 2004.

Jacob, D. J.: Chemistry of OH in remote clouds and its role in the production of formic-acid and peroxymonosulfate, J. Geophys. Res.-Atmos., 91, 9807–9826, 1986.

Jacob, D. J.: Heterogeneous chemistry and tropospheric ozone, Atmos. Environ., 34, 2131–2159, 2000.

Jacob, D. J., Crawford, J. H., Maring, H., Clarke, A. D., Dibb, J. E., Ferrare, R. A., Hostetler, C. A., Russell, P. B., Singh, H. B., Thompson, A. M., Shaw, G. E., McCauley, E., Pederson, J. R., and Fisher, J. A.: The ARCTAS aircraft mission: design and execution, Atmos. Chem. Phys. Discuss., 9, 17073–17123, 2009,

**Chemistry of
hydrogen oxide
radicals in the Arctic
troposphere in spring**

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



<http://www.atmos-chem-phys-discuss.net/9/17073/2009/>.

Jacobi, H. W., Frey, M. M., Hutterli, M. A., Bales, R. C., Schrems, O., Cullen, N. J., Steffen, K., and Koehler, C.: Measurements of hydrogen peroxide and formaldehyde exchange between the atmosphere and surface snow at summit, greenland, *Atmos. Environ.*, 36, 2619–2628, 2002.

Jaeglé, L., Jacob, D. J., Wennberg, P. O., Spivakovsky, C. M., Hanisco, T. F., Lanzendorf, E. J., Hints, E. J., Fahey, D. W., Keim, E. R., Proffitt, M. H., Atlas, E. L., Flocke, F., Schauffler, S., McElroy, C. T., Midwinter, C., Pfister, L., and Wilson, J. C.: Observed OH and HO₂ in the upper troposphere suggest a major source from convective injection of peroxides, *Geophys. Res. Lett.*, 24, 3181–3184, 1997.

Jaeglé, L., Jacob, D. J., Wang, Y., Weinheimer, A. J., Ridley, B. A., Campos, T. L., Sachse, G. W., and Hagen, D. E.: Sources and chemistry of NO_x in the upper troposphere over the united states, *Geophys. Res. Lett.*, 25, 1705–1708, 1998.

Jaeglé, L., Jacob, D. J., Brune, W. H., Faloon, I., Tan, D., Heikes, B. G., Kondo, Y., Sachse, G. W., Anderson, B., Gregory, G. L., Singh, H. B., Poeschel, R., Ferry, G., Blake, D. R., and Shetter, R. E.: Photochemistry of HO_x in the upper troposphere at northern midlatitudes, *J. Geophys. Res.-Atmos.*, 105, 3877–3892, 2000.

Jaeglé, L., Jacob, D. J., Brune, W. H., and Wennberg, P. O.: Chemistry of HO_x radicals in the upper troposphere, *Atmos. Environ.*, 35, 469–489, 2001.

Kahl, J. D.: Characteristics of the low-level temperature inversion along the alaskan Arctic coast, *Int. J. Climatol.*, 10, 537–548, 1990.

Kanaya, Y., Sadanaga, Y., Matsumoto, J., Sharma, U. K., Hirokawa, J., Kajii, Y., and Akimoto, H.: Daytime HO₂ concentrations at Oki Island, Japan, in summer 1998: Comparison between measurement and theory, *J. Geophys. Res.-Atmos.*, 105, 24205–24222, 2000.

Kolb, C. E., Worsnop, D. R., Zahniser, M. S., Davidovits, P., Keyser, L. F., Leu, M.-T., Molina, M. J., Hanson, D. R., and Ravishankara, A. R.: Laboratory studies of atmospheric heterogeneous chemistry, in: *Progress and problems in atmospheric chemistry*, edited by: Baker, J., *Advanced series in physical chemistry*, 3, 771–875, 1995.

Liao, J., Huey, L. G., Scheuer, E., and Dibb, J. E.: Characterization of soluble bromide measurements during ARCTAS, in preparation, 2010.

Liu, H. Y., Jacob, D. J., Bey, I., and Yantosca, R. M.: Constraints from Pb-210 and Be-7 on wet deposition and transport in a global three-dimensional chemical tracer model driven by assimilated meteorological fields, *J. Geophys. Res.-Atmos.*, 106, 12109–12128, 2001.

**Chemistry of
hydrogen oxide
radicals in the Arctic
troposphere in spring**

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Loukhovitskaya, E., Bedjanian, Y., Morozov, I., and Le Bras, G.: Laboratory study of the interaction of HO₂ radicals with the NaCl, NaBr, MgCl₂ center dot ⁶H(2)O and sea salt surfaces, PCCP Phys. Chem. Chem. Phys., 11, 7896–7905, doi:10.1039/b906300e, 2009.
- Mari, C., Jacob, D. J., and Bechtold, P.: Transport and scavenging of soluble gases in a deep convective cloud, J. Geophys. Res.-Atmos., 105, 22255–22267, 2000.
- Martin, R. V., Jacob, D. J., Yantosca, R. M., Chin, M., and Ginoux, P.: Global and regional decreases in tropospheric oxidants from photochemical effects of aerosols, J. Geophys. Res.-Atmos., 108(D3), 4097, doi:10.1029/2002jd002622, 2003a.
- Martin, S. T., Schlenker, J. C., Malinowski, A., Hung, H. M., and Rudich, Y.: Crystallization of atmospheric sulfate-nitrate-ammonium particles, Geophys. Res. Lett., 30(21), 2102, doi:10.1029/2003gl017930, 2003b.
- McNaughton, C. S., Clarke, A. D., Freitag, S., Kapustin, V. N., Kondo, Y., Moteki, N., Sahu, L. K., Takegawa, N., Schwarz, J. P., Spackman, J. R., Diskin, G. S., Sachse, G. W., Podolske, J. R., Holloway, J., Wisthaler, A., Mikoviny, T., De Gouw, J. A., Warneke, C., Jimenez, J. L., Cubison, M., Howell, S., Middlebrook, A. M., Bahreini, R., Anderson, B., Winstead, E. L., Thornhill, K. L., Lack, D. A., Cozic, J., and Brock, C. A.: Absorbing aerosol in the troposphere of the Western Arctic during the 2008 ARCTAS/ARCPAC airborne field campaigns, in preparation, 2010.
- Miller, C. E. and Francisco, J. S.: The formation of a surprisingly stable HO₂–H₂SO₄ complex, J. Am. Chem. Soc., 123, 10387–10388, 2001.
- Millet, D. B., Jacob, D. J., Turquety, S., Hudman, R. C., Wu, S. L., Fried, A., Walega, J., Heikes, B. G., Blake, D. R., Singh, H. B., Anderson, B. E., and Clarke, A. D.: Formaldehyde distribution over North America: Implications for satellite retrievals of formaldehyde columns and isoprene emission, J. Geophys. Res.-Atmos., 111, D24s02, doi:10.1029/2005jd006853, 2006.
- Möller, D.: Explanation for the recent dramatic increase of H₂O₂ concentrations found in Greenland ice cores, Atmos. Environ., 33, 2435–2437, 1999.
- Mozurkewich, M., McMurry, P. H., Gupta, A., and Calvert, J. G.: Mass accommodation coefficient for HO₂ radicals on aqueous particles, J. Geophys. Res.-Atmos., 92, 4163–4170, 1987.
- Müller, J. F. and Brasseur, G.: Sources of upper tropospheric HO_x: A three-dimensional study, J. Geophys. Res.-Atmos., 104, 1705–1715, 1999.
- Neuman, J. A., Nowak, J. B., Huey, L. G., Burkholder, J. B., Dibb, J. E., Holloway, J. S., Liao, J.,

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Peischl, J., Roberts, J. M., Ryerson, T. B., Scheuer, E., Stark, H., Stickel, R. E., Tanner, D. J., and Weinheimer, A.: Bromine measurements in ozone depleted air over the Arctic Ocean, *Atmos. Chem. Phys. Discuss.*, 10, 3827–3860, 2010, <http://www.atmos-chem-phys-discuss.net/10/3827/2010/>.

- 5 O'Sullivan, D. W., Heikes, B. G., Lee, M., Chang, W., Gregory, G. L., Blake, D. R., and Sachse, G. W.: Distribution of hydrogen peroxide and methylhydroperoxide over the pacific and south atlantic oceans, *J. Geophys. Res.-Atmos.*, 104, 5635–5646, 1999.
- Oiestad, A. M. L., Petersen, A. C., Bakken, V., Vedde, J., and Uggerud, E.: The oxidative power of protonated hydrogen peroxide, *Angew. Chem.-Int. Edit.*, 40, 1305–1309, 2001.
- 10 Olson, J. R., Crawford, J. H., Chen, G., Fried, A., Evans, M. J., Jordan, C. E., Sandholm, S. T., Davis, D. D., Anderson, B. E., Avery, M. A., Barrick, J. D., Blake, D. R., Brune, W. H., Eisele, F. L., Flocke, F., Harder, H., Jacob, D. J., Kondo, Y., Lefer, B. L., Martinez, M., Mauldin, R. L., Sachse, G. W., Shetter, R. E., Singh, H. B., Talbot, R. W., and Tan, D.: Testing fast photochemical theory during trace-p based on measurements of OH, HO₂, and CH₂O, *J. Geophys. Res.-Atmos.*, 109(16), D15s10, doi:10.1029/2003jd004278, 2004.
- 15 Olson, J. R., Crawford, J. H., Chen, G., Brune, W. H., Ren, X., Wennberg, P. O., Fried, A., and Mao, J.: Photochemical Analysis of ARCTAS Observations, in preparation, 2010.
- Onasch, T. B., Siefert, R. L., Brooks, S. D., Prenni, A. J., Murray, B., Wilson, M. A., and Tolbert, M. A.: Infrared spectroscopic study of the deliquescence and efflorescence of ammonium sulfate aerosol as a function of temperature, *J. Geophys. Res.-Atmos.*, 104, 21317–21326, 2004.
- 20 Palmer, P. I., Jacob, D. J., Fiore, A. M., Martin, R. V., Chance, K., and Kurosu, T. P.: Mapping isoprene emissions over north america using formaldehyde column observations from space, *J. Geophys. Res.-Atmos.*, 108(D6), 4180, doi:10.1029/2002jd002153, 2003.
- 25 Park, R. J., Jacob, D. J., Field, B. D., Yantosca, R. M., and Chin, M.: Natural and transboundary pollution influences on sulfate-nitrate-ammonium aerosols in the united states: Implications for policy, *J. Geophys. Res.-Atmos.*, 109, D15204, doi:10.1029/2003jd004473, 2004.
- Park, R. J., Jacob, D. J., Kumar, N., and Yantosca, R. M.: Regional visibility statistics in the united states: Natural and transboundary pollution influences, and implications for the regional haze rule, *Atmos. Environ.*, 40, 5405–5423, doi:10.1016/j.atmosenv.2006.04.059, 2006.
- 30 Parsons, M. T., Knopf, D. A., and Bertram, A. K.: Deliquescence and crystallization of ammonium sulfate particles internally mixed with water-soluble organic compounds, *J. Phys.*

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Chem. A, 108, 11600–11608, doi:10.1021/jp0462862, 2004.
- Pickett-Heaps, C. A., Jacob, D. J., Wecht, K. J., Drevet, J., Diskin, G. S., Wofsy, S. C., Worthy, D. E. J., Kort, E. A., Jimenez, R., Daube, B., and Park, S.: Methane Emissions from the Hudson Bay Lowlands: Interpretation of airborne and surface methane observations, in preparation, 2010.
- Plummer, D. A., McConnell, J. C., Shepson, P. B., Hastie, D. R., and Niki, H.: Modeling of ozone formation at a rural site in southern ontario, *Atmos. Environ.*, 30, 2195–2217, 1996.
- Prather, M. J. and Jacob, D. J.: A persistent imbalance in HO_x and NO_x photochemistry of the upper troposphere driven by deep tropical convection, *Geophys. Res. Lett.*, 24, 3189–3192, 1997.
- Quinn, P. K., Bates, T. S., Baum, E., Doubleday, N., Fiore, A. M., Flanner, M., Fridlind, A., Garrett, T. J., Koch, D., Menon, S., Shindell, D., Stohl, A., and Warren, S. G.: Short-lived pollutants in the Arctic: their climate impact and possible mitigation strategies, *Atmos. Chem. Phys.*, 8, 1723–1735, 2008, <http://www.atmos-chem-phys.net/8/1723/2008/>.
- Reid, J. S., Hyer, E. J., Prins, E. M., Westphal, D. L., Zhang, J. L., Wang, J., Christopher, S. A., Curtis, C. A., Schmidt, C. C., Eleuterio, D. P., Richardson, K. A., and Hoffman, J. P.: Global monitoring and forecasting of biomass-burning smoke: Description of and lessons from the Fire Locating and Modeling of Burning Emissions (FLAMBE) program, *IEEE J. Sel. Topics Appl. Earth Observations Remote Sens.*, 2, 144–162, doi:10.1109/jstars.2009.2027443, 2009.
- Remorov, R. G., Gershenzon, Y. M., Molina, L. T., and Molina, M. J.: Kinetics and mechanism of HO₂ uptake on solid NaCl, *J. Phys. Chem. A*, 106, 4558–4565, doi:10.1021/jp013179o, 2002.
- Ren, X. R., Olson, J. R., Crawford, J. H., Brune, W. H., Mao, J. Q., Long, R. B., Chen, Z., Chen, G., Avery, M. A., Sachse, G. W., Barrick, J. D., Diskin, G. S., Huey, L. G., Fried, A., Cohen, R. C., Heikes, B., Wennberg, P. O., Singh, H. B., Blake, D. R., and Shetter, R. E.: HO_x chemistry during inter-a 2004: Observation, model calculation, and comparison with previous studies, *J. Geophys. Res.-Atmos.*, 113(13), D05310, doi:10.1029/2007jd009166, 2008.
- Richter, D., Fried, A., and Weibring, P.: Difference frequency generation laser based spectrometers, *Laser and Photonics Reviews*, 3, 343–354, doi:10.1002/lpor.200810048, 2009.
- Ridley, B., Walega, J., Montzka, D., Grahek, F., Atlas, E., Flocke, F., Stroud, V., Deary, J., Gallant, A., Boudries, H., Bottenheim, J., Anlauf, K., Worthy, D., Sumner, A. L., Splawn, B.,

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- and Shepson, P.: Is the arctic surface layer a source and sink of NO_x in winter/spring?, *J. Atmos. Chem.*, 36, 1–22, 2000.
- Saathoff, H., Naumann, K. H., Riemer, N., Kamm, S., Mohler, O., Schurath, U., Vogel, H., and Vogel, B.: The loss of NO_2 , HNO_3 , $\text{NO}_3/\text{N}_2\text{O}_5$, and $\text{HO}_2/\text{HOONO}_2$ on soot aerosol: A chamber and modeling study, *Geophys. Res. Lett.*, 28, 1957–1960, 2001.
- Salawitch, R. J., Canty, T. P., Kurosu, T. P., Chance, K., Liang, Q., Pawson, S., Liu, X., Huey, L. G., Liao, J., Stickel, R. E., Tanner, D., Dibb, J. E., Weinheimer, A. J., Flocke, F. M., Knapp, D. J., Montzka, D. D., Neuman, J., Simpson, W. R., Donohoue, D., Carlson, D., Blake, D. R., Kinnison, D. E., Tilmes, S., Pan, L., Pierce, R., Hendrick, F., Kreher, K., Wang, Y., Choi, S., and Atlas, E. L.: Airborne, Ground-based, and Satellite Measurements of BrO during ARCTAS and ARCPAC, in preparation, 2010.
- Sander, S. P., Finlayson-Pitts, B. J., Friedl, R. R., Golden, D. M., Huie, R. E., Keller-Rudek, H., Kolb, C. E., Kurylo, M. J., Molina, M. J., Moortgat, G. K., Orkin, V. L., Ravishankara, A. R., and Wine, P. H.: Chemical Kinetic and Photochemical Data for Use in Atmospheric Studies, Evaluation Number 15, 2006.
- Sauvage, B., Martin, R. V., van Donkelaar, A., Liu, X., Chance, K., Jaeglé, L., Palmer, P. I., Wu, S., and Fu, T.-M.: Remote sensed and in situ constraints on processes affecting tropical tropospheric ozone, *Atmos. Chem. Phys.*, 7, 815–838, 2007, <http://www.atmos-chem-phys.net/7/815/2007/>.
- Scheuer, E., Talbot, R. W., Dibb, J. E., Seid, G. K., DeBell, L., and Lefer, B.: Seasonal distributions of fine aerosol sulfate in the north american arctic basin during TOPSE, *J. Geophys. Res.-Atmos.*, 108(D4), 8370, doi:10.1029/2001jd001364, 2003.
- Shindell, D. T., Faluvegi, G., Stevenson, D. S., Krol, M. C., Emmons, L. K., Lamarque, J. F., Petron, G., Dentener, F. J., Ellingsen, K., Schultz, M. G., Wild, O., Amann, M., Atherton, C. S., Bergmann, D. J., Bey, I., Butler, T., Cofala, J., Collins, W. J., Derwent, R. G., Doherty, R. M., Drevet, J., Eskes, H. J., Fiore, A. M., Gauss, M., Hauglustaine, D. A., Horowitz, L. W., Isaksen, I. S. A., Lawrence, M. G., Montanaro, V., Muller, J. F., Pitari, G., Prather, M. J., Pyle, J. A., Rast, S., Rodriguez, J. M., Sanderson, M. G., Savage, N. H., Strahan, S. E., Sudo, K., Szopa, S., Unger, N., van Noije, T. P. C., and Zeng, G.: Multimodel simulations of carbon monoxide: Comparison with observations and projected near-future changes, *J. Geophys. Res.-Atmos.*, 111, D19306, doi:10.1029/2006jd007100, 2006.
- Shindell, D. T., Chin, M., Dentener, F., Doherty, R. M., Faluvegi, G., Fiore, A. M., Hess, P., Koch, D. M., MacKenzie, I. A., Sanderson, M. G., Schultz, M. G., Schulz, M., Stevenson, D.

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

S., Teich, H., Textor, C., Wild, O., Bergmann, D. J., Bey, I., Bian, H., Cuvelier, C., Duncan, B. N., Folberth, G., Horowitz, L. W., Jonson, J., Kaminski, J. W., Marmer, E., Park, R., Pringle, K. J., Schroeder, S., Szopa, S., Takemura, T., Zeng, G., Keating, T. J., and Zuber, A.: A multi-model assessment of pollution transport to the Arctic, *Atmos. Chem. Phys.*, 8, 5353–5372, 2008,

<http://www.atmos-chem-phys.net/8/5353/2008/>.

Simpson, W. R., von Glasow, R., Riedel, K., Anderson, P., Ariya, P., Bottenheim, J., Burrows, J., Carpenter, L. J., Frie, U., Goodsite, M. E., Heard, D., Hutterli, M., Jacobi, H.-W., Kaleschke, L., Neff, B., Plane, J., Platt, U., Richter, A., Roscoe, H., Sander, R., Shepson, P., Sodeau, J., Steffen, A., Wagner, T., and Wolff, E.: Halogens and their role in polar boundary-layer ozone depletion, *Atmos. Chem. Phys.*, 7, 4375–4418, 2007, <http://www.atmos-chem-phys.net/7/4375/2007/>.

Singh, H., Chen, Y., Tabazadeh, A., Fukui, Y., Bey, I., Yantosca, R., Jacob, D., Arnold, F., Wohlfarth, K., Atlas, E., Flocke, F., Blake, D., Blake, N., Heikes, B., Snow, J., Talbot, R., Gregory, G., Sachse, G., Vay, S., and Kondo, Y.: Distribution and fate of selected oxygenated organic species in the troposphere and lower stratosphere over the Atlantic, *J. Geophys. Res.-Atmos.*, 105, 3795–3805, 2000.

Snow, J. A., Heikes, B. G., Merrill, J. T., Wimmers, A. J., Moody, J. L., and Cantrell, C. A.: Winter-spring evolution and variability of HO_x reservoir species, hydrogen peroxide, and methyl hydroperoxide, in the northern middle to high latitudes, *J. Geophys. Res.-Atmos.*, 108(D4), 8362, doi:10.1029/2002jd002172, 2003.

Snow, J. A., Heikes, B. G., Shen, H. W., O'Sullivan, D. W., Fried, A., and Walega, J.: Hydrogen peroxide, methyl hydroperoxide, and formaldehyde over North America and North Atlantic, *J. Geophys. Res.-Atmos.*, 112, D12S07, doi:10.1029/2006jd007746, 2007.

Sommariva, R., Haggerstone, A.-L., Carpenter, L. J., Carslaw, N., Creasey, D. J., Heard, D. E., Lee, J. D., Lewis, A. C., Pilling, M. J., and Zádor, J.: OH and HO₂ chemistry in clean marine air during SOAPEX-2, *Atmos. Chem. Phys.*, 4, 839–856, 2004, <http://www.atmos-chem-phys.net/4/839/2004/>.

Sommariva, R., Bloss, W. J., Brough, N., Carslaw, N., Flynn, M., Haggerstone, A.-L., Heard, D. E., Hopkins, J. R., Lee, J. D., Lewis, A. C., McFiggans, G., Monks, P. S., Penkett, S. A., Pilling, M. J., Plane, J. M. C., Read, K. A., Saiz-Lopez, A., Rickard, A. R., and Williams, P. I.: OH and HO₂ chemistry during NAMBLEX: roles of oxygenates, halogen oxides and heterogeneous uptake, *Atmos. Chem. Phys.*, 6, 1135–1153, 2006,

**Chemistry of
hydrogen oxide
radicals in the Arctic
troposphere in spring**

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

<http://www.atmos-chem-phys.net/6/1135/2006/>.

Sumner, A. L. and Shepson, P. B.: Snowpack production of formaldehyde and its effect on the arctic troposphere, *Nature*, 398, 230–233, 1999.

Sumner, A. L., Shepson, P. B., Grannas, A. M., Bottenheim, J. W., Anlauf, K. G., Worthy, D., Schroeder, W. H., Steffen, A., Domine, F., Perrier, S., and Houdier, S.: Atmospheric chemistry of formaldehyde in the arctic troposphere at polar sunrise, and the influence of the snowpack, *Atmos. Environ.*, 36, 2553–2562, 2002.

Taketani, F., Kanaya, Y., and Akimoto, H.: Kinetics of heterogeneous reactions of HO₂ radical at ambient concentration levels with (NH₄)₂SO₄ and NaCl aerosol particles, *J. Phys. Chem. A*, 112, 2370–2377, doi:10.1021/jp0769936, 2008.

Taketani, F., Kanaya, Y., and Akimoto, H.: Heterogeneous loss of HO₂ by KCl, synthetic sea salt, and natural seawater aerosol particles, *Atmos. Environ.*, 43, 1660–1665, doi:10.1016/j.atmosenv.2008.12.010, 2009.

Thornton, J. and Abbatt, J. P. D.: Measurements of HO₂ uptake to aqueous aerosol: Mass accommodation coefficients and net reactive loss, *J. Geophys. Res.-Atmos.*, 110, D08309, doi:10.1029/2004jd005402, 2005.

Thornton, J. A., Jaeglé, L., and McNeill, V. F.: Assessing known pathways for HO₂ loss in aqueous atmospheric aerosols: Regional and global impacts on tropospheric oxidants, *J. Geophys. Res.-Atmos.*, 113, D05303, doi:10.1029/2007jd009236, 2008.

van Donkelaar, A., Martin, R. V., Leaitch, W. R., Macdonald, A. M., Walker, T. W., Streets, D. G., Zhang, Q., Dunlea, E. J., Jimenez, J. L., Dibb, J. E., Huey, L. G., Weber, R., and Andreae, M. O.: Analysis of aircraft and satellite measurements from the Intercontinental Chemical Transport Experiment (INTEX-B) to quantify long-range transport of East Asian sulfur to Canada, *Atmos. Chem. Phys.*, 8, 2999–3014, 2008,

<http://www.atmos-chem-phys.net/8/2999/2008/>.

Wagner, T., Leue, C., Wenig, M., Pfeilsticker, K., and Platt, U.: Spatial and temporal distribution of enhanced boundary layer bromine concentrations measured by the GOME instrument aboard ERS-2, *J. Geophys. Res.-Atmos.*, 106, 24225–24235, 2001.

Wang, Y. H., Ridley, B., Fried, A., Cantrell, C., Davis, D., Chen, G., Snow, J., Heikes, B., Talbot, R., Dibb, J., Flocke, F., Weinheimer, A., Blake, N., Blake, D., Shetter, R., Lefer, B., Atlas, E., Coffey, M., Walega, J., and Wert, B.: Springtime photochemistry at northern mid and high latitudes, *J. Geophys. Res.-Atmos.*, 108(D4), 8358, doi:10.1029/2002jd002227, 2003.

Wang, Q., Jacob, D. J., Fisher, J. A., Mao, J., Le Sager, P., Lebensperger, E. M., Carouge, C.,

**Chemistry of
hydrogen oxide
radicals in the Arctic
troposphere in spring**

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Kondo, Y., Jimenez, J. L., Cubison, M., Howell, S. G., Freitag, S., Clarke, A. D., McNaughton, C. S., Weber, R., and Apel, E. C.: Sources and Sinks of Carbonaceous Aerosols in the Arctic in Spring, in preparation, 2010.
- Warneke, C., Bahreini, R., Brioude, J., Brock, C. A., de Gouw, J. A., Fahey, D. W., Froyd, K. D.,
 5 Holloway, J. S., Middlebrook, A., Miller, L., Montzka, S., Murphy, D. M., Peischl, J., Ryerson, T. B., Schwarz, J. P., Spackman, J. R., and Veres, P.: Biomass burning in Siberia and Kazakhstan as an important source for haze over the Alaskan Arctic in April 2008, *Geophys. Res. Lett.*, 36, L02813, doi:10.1029/2008gl036194, 2009.
- Weibring, P., Richter, D., Walega, J. G., and Fried, A.: First demonstration of a high performance
 10 difference frequency spectrometer on airborne platforms, *Optics Express*, 15, 13476–13495, 2007.
- Wesely, M. L.: Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical-models, *Atmos. Environ.*, 23, 1293–1304, 1989.
- Wild, O., Zhu, X., and Prather, M. J.: Fast-j: Accurate simulation of in- and below-cloud photolysis in tropospheric chemical models, *J. Atmos. Chem.*, 37, 245–282, 2000.
- 15 Yang, J., Honrath, R. E., Peterson, M. C., Dibb, J. E., Sumner, A. L., Shepson, P. B., Frey, M., Jacobi, H. W., Swanson, A., and Blake, N.: Impacts of snowpack emissions on deduced levels of OH and peroxy radicals at summit, greenland, *Atmos. Environ.*, 36, 2523–2534, 2002.
- 20 Zhang, L., Jacob, D. J., Boersma, K. F., Jaffe, D. A., Olson, J. R., Bowman, K. W., Worden, J. R., Thompson, A. M., Avery, M. A., Cohen, R. C., Dibb, J. E., Flock, F. M., Fuelberg, H. E., Huey, L. G., McMillan, W. W., Singh, H. B., and Weinheimer, A. J.: Transpacific transport of ozone pollution and the effect of recent Asian emission increases on air quality in North America: an integrated analysis using satellite, aircraft, ozonesonde, and surface observations, *Atmos. Chem. Phys.*, 8, 6117–6136, 2008,
 25 <http://www.atmos-chem-phys.net/8/6117/2008/>.
- Zhou, X. L., Beine, H. J., Honrath, R. E., Fuentes, J. D., Simpson, W., Shepson, P. B., and Bottenheim, J. W.: Snowpack photochemical production of HONO: A major source of OH in the arctic boundary layer in springtime, *Geophys. Res. Lett.*, 28, 4087–4090, 2001.

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.




Fig. 1. Median vertical profiles of HO_2 , OH , H_2O_2 , CH_3OOH , NO , NO_2 , HCHO , and HNO_4 concentrations during ARCTAS in the North American Arctic (1–21 April 2008). Observations from the DC-8 aircraft (black lines) are compared to three GEOS-Chem model simulations: (1) without HO_2 uptake by aerosol (green dashed line), referred to in the text as “standard GEOS-Chem”; (2) with HO_2 uptake yielding H_2O_2 (blue dashed line); (3) with HO_2 uptake not yielding H_2O_2 (solid red line). All concentrations are in unit of pptv except OH (ppqv). Most data were collected under sunlit conditions, between 08:00 and 18:00 local time. Stratospheric data were excluded as described in the text.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.




Fig. 2. Laboratory data reported in the literature for the reactive uptake coefficient $\gamma(\text{HO}_2)$ by different surfaces as a function of temperature. Vertical arrows indicate “greater than” or “less than”. Open symbols indicate aqueous surfaces, closed symbols indicate copper-doped aqueous surfaces, and other symbols indicate solid surfaces (noted as (s) in the legend). The solid red line is the median value of $\gamma(\text{HO}_2)$ computed in GEOS-Chem along the ARCTAS flight tracks using the Thornton et al. (2008) parameterization. Literature references are given by footnotes in legend: **(a)** Mozurkewich et al. (1987); **(b)** Hanson et al. (1992); **(c)** Gershenson et al. (1995); **(d)** Cooper and Abbatt (1996); **(e)** Saathoff et al. (2001); **(f)** Remorov et al. (2002); **(g)** Thornton and Abbatt (2005); **(h)** Taketani et al. (2008); **(i)** Taketani et al. (2009), **(j)** Loukhovitskaya et al. (2009).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Chemistry of
hydrogen oxide
radicals in the Arctic
troposphere in spring**

J. Mao et al.




Fig. 3. Scatterplot of observed ammonium vs. sulfate submicron aerosol concentrations measured aboard the DC-8 aircraft during ARCTAS-A (April 2008). The observed points are colored by altitude (km). The data from the first two flights (1 and 4 April) are excluded due to low quality of the ammonium data. In red is the linear least squares regression line for all altitudes.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Chemistry of
hydrogen oxide
radicals in the Arctic
troposphere in spring

J. Mao et al.




Fig. 4. Scatterplots of simulated vs. observed HO₂, OH, H₂O₂, and CH₃OOH concentrations, for the model simulation with $\gamma(\text{HO}_2 \rightarrow \text{products})$ and the ensemble of tropospheric observations during ARCTAS (April 2008). The red solid line is the reduced major axis regression line. Panel titles give the correlation coefficients and regression slopes.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Chemistry of
hydrogen oxide
radicals in the Arctic
troposphere in spring

J. Mao et al.




Fig. 5. Median vertical profiles of major HO_x and HO_y sources and sinks computed from observed tropospheric concentrations in ARCTAS (April 2008). Values are instantaneous rates. Gas-phase rate constants are from JPL06 and IUPAC06. HO_2 uptake by aerosol is computed with a reaction probability $\gamma(\text{HO}_2 \rightarrow \text{products})$ from Thornton et al. (2008). Stratospheric data are excluded as described in the text.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Chemistry of
hydrogen oxide
radicals in the Arctic
troposphere in spring

J. Mao et al.




Fig. 6. Circumpolar GEOS-Chem model budget of HO_y for the Arctic tropospheric column (60–90 N) during ARCTAS (1–21 April 2008). Mean production rates (P) and loss rates (L) are shown for three altitude bands. The transport term describes exchange with northern mid-latitudes south of 60 N.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Chemistry of hydrogen oxide radicals in the Arctic troposphere in spring

J. Mao et al.




Fig. 7. Schematic diagram of HO_x-HO_y chemistry in Arctic spring as constructed from the GEOS-Chem model simulation of the ARCTAS observations. Values are tropospheric column averages for April 2008 over the Arctic cap (60–90°N). Masses of chemicals within the domain (in parentheses) are in units of Mmol and rates are in units of Mmol d⁻¹. The dashed line for HCHO production indicates that it is not a HO_y sink.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion