

1 **Anthropogenic and natural constituents in particulate**  
2 **matter in the Netherlands**

3

4 **E.P. Weijers<sup>1</sup>, M. Schaap<sup>2</sup>, L. Nguyen<sup>3</sup>, J. Matthijsen<sup>4</sup>, H.A.C. Denier van der**  
5 **Gon<sup>2</sup>, H.M. ten Brink<sup>1</sup>, R. Hoogerbrugge<sup>3</sup>**

6

7 [1] {Department of Air Quality and Climate Change, Energy research Centre of the Nether-  
8 lands (ECN), Petten, the Netherlands}

9 [2] {TNO, Business unit Environment, Health and Safety, Utrecht, the Netherlands}

10 [3] {National Institute for Public Health and the Environment (RIVM), Bilthoven, the Neth-  
11 erlands}

12 [4] {Netherlands Environmental Assessment Agency (PBL), Bilthoven, the Netherlands}

13 Correspondence to: E.P. Weijers (weijers@ecn.nl)

14

1 **Abstract**

2 To develop mitigation strategies for reducing concentrations of both PM<sub>2.5</sub> (and PM<sub>10</sub>), the  
3 origin of particulate matter (PM) needs to be established. An intensive, one-year  
4 measurement campaign from August 2007 to August 2008 was carried out to determine the  
5 composition of PM<sub>10</sub> and PM<sub>2.5</sub> at five locations in the Netherlands, aiming at reducing the  
6 uncertainties on the origin of PM. Generally, a considerable conformity in the chemical  
7 composition of PM<sub>2.5</sub> (and PM<sub>10</sub>) was observed. From all the constituents present in PM<sub>2.5</sub>,  
8 the secondary inorganic aerosol is the most dominant (42-48%), followed by total  
9 carbonaceous matter (22-37%). Contributions due to sea salt (maximum 8%), mineral dust  
10 and metals (maximum 5%) are relatively low. For the first time, a detailed overview can be  
11 presented of the composition of the coarse fraction. Compared to the fine fraction,  
12 contributions of the sea salt, mineral dust and metals are now larger resulting in a more  
13 balanced distribution between the various constituents. Through mass closure a considerable  
14 part of the PM mass could be defined (PM<sub>2.5</sub>: 80-94%). The chemical distribution on days  
15 with high PM levels showed a distinct increase of nitrate as well as in the unaccounted mass  
16 ~~not accounted for~~. Contributions of the remaining constituents remained equal or f were lower  
17 (sea salt) when expressed in percentages. A correspondence between nitrate and the unknown  
18 mass was observed hinting at the presence of water on the filters. The contribution from  
19 natural sources in the Netherlands was estimated to be about 25% for PM<sub>10</sub> and 20% for  
20 PM<sub>2.5</sub>.

21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34

1

## 2 **1. Introduction**

3 Although air quality in Europe has improved substantially over the past decades, air pollution  
4 still poses a significant threat to human health (EEA, 2007). Health effects of air pollution are  
5 dominated by particulate matter, both  $PM_{2.5}$  and  $PM_{10}$ . Short-term exposure to PM has fre-  
6 quently been associated with increased human morbidity and mortality (e.g., Brunekreef and  
7 Holgate, 2002). Effects of long-term exposure to PM are much more uncertain than the short-  
8 term effects, but are believed to have a much greater effect on health loss (Dockery et al.,  
9 1993; Pope et al., 1995). Recent estimates arrive at an average loss of life expectancy of  
10 about 6-12 months in the Netherlands (EEA, 2007). So far, there is no conclusive evidence  
11 which component or components are most relevant for deteriorating human health, although  
12 some suggestions have been made, e.g. soot (Hoek et al., 2002). Therefore, the European air  
13 quality standards currently focus on  $PM_{10}$  and  $PM_{2.5}$ .

14 At present, many countries in Europe, including the Netherlands, have problems adhering to  
15 the daily limit value for  $PM_{10}$  at hotspot locations. In the Directive (2008/50/EC) on ambient  
16 air quality and cleaner air for Europe (EU, 2008) new air quality standards for fine particulate  
17 matter ( $PM_{2.5}$ ) have been established. An annual mean  $PM_{2.5}$  concentration of  $25 \mu\text{g}/\text{m}^3$  has  
18 been set as target value to be achieved in 2010 and as limit value to be adhered to by 2015.  
19 The new directive introduces additional  $PM_{2.5}$  objectives, that aim to a reduce human expo-  
20 sure to  $PM_{2.5}$ . The average exposure indicator (AEI) is a three-year averaged  $PM_{2.5}$  level  
21 measured at urban background locations throughout European Member States. A reduction  
22 target of 15% of the AEI between 2010 and 2020 for the Netherlands might be reached under  
23 current legislation, whereas a reduction of 20% is probably not even reached with outlined  
24 additional measures (Matthijsen et al., 2009). To develop mitigation strategies to meet these  
25 targets and to reduce the concentrations of both  $PM_{2.5}$  and  $PM_{10}$  in general one needs to es-  
26 tablish the origin of particulate matter. Of particular interest is the quantification of the an-  
27 thropogenic and natural fractions in PM as these provide an indication of the 'playing field'  
28 of policy makers.

29 Important information on the origin of particulate matter can be obtained through chemical  
30 analysis to establish its composition (Putaud et al., 2004, 2009; Viana et al., 2008; Sillanpää  
31 et al., 2006; Salvador et al., 2007; Mazzei et al., 2008; Almeida et al., 2007). Many compo-

1 nents can be used as tracers for specific sources. For example, sodium is a tracer that is al-  
2 most exclusively associated with sea salt. However, one is usually not able to analyze the full  
3 composition and a significant portion of the particulate mass is therefore unknown. Using as-  
4 sumptions on the relationships between (a group of) tracers and major constituents one can  
5 attribute the mass to these constituents, the ~~so-called~~ so-called practical chemical mass bal-  
6 ~~ance-closure (CMB)~~ approach (Harrison et al., 2003). In this way an estimate can be derived  
7 for the major source contributions to PM. Though this approach ~~CMB~~-significantly reduces  
8 the unknown mass a total mass closure is usually not attained. The (relative) magnitude of the  
9 unexplained mass is an indicator for the uncertainty in the mass attribution. To independently  
10 address such issues one can apply statistical techniques to the observation dataset to address  
11 the possible source categories for PM. Techniques such as positive matrix factorization  
12 (PMF) can be used to search for groups of components with the same behaviour in time,  
13 which are then assumed to derive from a single source or group of similar sources (Hopke et  
14 al., 2001). Hence, a mass closure approach as well as PMF will give us independent insights  
15 from an empirical point of view in the actual PM sources.

16 Here, a mass closure approach is used to study the chemical composition of  $PM_{10}$  and  $PM_{2.5}$   
17 filter samples that were collected in the Netherlands at five measurement sites for a period of  
18 one year. The study was executed within the framework of the Policy-Oriented Research  
19 Program (BOP) and described in detail by Schaap et al. (2010). The general aim of this na-  
20 tional program was to reduce knowledge gaps about composition, distribution and origin of  
21 particulate matter in the Netherlands. This study established the chemical composition meas-  
22 ured at Dutch rural and urban locations. Properties of various constituents in fine and coarse  
23 matter, respective mass deficits and the changing composition on days with higher PM  
24 masses are demonstrated. Also, the contribution of natural material in particulate matter and  
25 political implications are discussed. The usefulness of the PMF approach is demonstrated  
26 elsewhere (Mooibroek et al., 2010; Schaap et al., 2010) who applied this technique on the  
27 same data set.

28  
29

## 30 **2. Experimental methods**

### 31 **2.1 Monitoring locations**

32 Five sites were selected for the monitoring of the  $PM_{10}$  and  $PM_{2.5}$  composition (see Fig.1).  
33 These include three rural background sites: Hellendoorn (~150 km from Rotterdam), Cabauw

1 (~50 km from Rotterdam) and Vredepeel (~100 km from Rotterdam). Cabauw represents the  
2 rural background for the (industrialised) western part of the country whereas Hellendoorn and  
3 Vredepeel were selected to add information from the eastern and southern parts where agri-  
4 culture dominates. At Vredepeel intensive livestock activity exists in the direct surroundings.  
5 Like to the rural sites, the urban background location Schiedam belongs to the Dutch Na-  
6 tional Air Quality Monitoring Network (NAQMN) maintained by RIVM. The kerbside (traf-  
7 fic) site in Rotterdam is part of the ~~regional~~ air quality network operated by the regional envi-  
8 ronmental protection agency (DCMR). Further characteristics of the locations are given in  
9 Table 1.

10

## 11 **2.2 Measurement strategy**

12 The filter collection routine (PM<sub>10</sub> and PM<sub>2.5</sub>) took place every second day between 15 Au-  
13 gust 2007 and 4 September 2008. Chemical analysis was done every fourth day. For each of  
14 the five sites the following components in the filter samples were determined: ~~sodium, chlo-~~  
15 ~~ride,~~ sulphate, nitrate, ammonium, elemental and organic carbon, ~~elemental carbon, sodium~~  
16 ~~and chloride, and the elemental composition and metals (and silicon)~~. The filters were col-  
17 lected according to the reference method described in the European Standards (EN12341:  
18 1998 and EN14907: 2005). The measurement instrument was the SEQ47/50 (Leckel GmbH,  
19 Germany). In the BOP programme's experimental set-up this sequential low-volume system  
20 (LVS) used a flow of 2.3 m<sup>3</sup>/h. Four of these reference samplers were placed at every loca-  
21 tion for a paired measurement of PM<sub>10</sub> and PM<sub>2.5</sub>. The sampling period per filter is 24 hours  
22 and ran from midnight to midnight. All filters were protected during storage and transport.  
23 The logistic and operational (QA/QC) procedures were those used in the NAQMN network.  
24 Two types of filter media, quartz (QMA) and teflon (PTFE), were used. The teflon filters  
25 were chosen to quantify the presence of silicon. The quartz fibre filter media is the material  
26 commonly used in gravimetric studies. They were cut into two equal parts for the SIA and  
27 EC/OC analyses.

28

## 29 **2.3 Chemical analyses**

30 The soluble fraction was extracted from the quartz filters by 7 ml deionized distilled water  
31 and kept for 8 h in a sealed tube (10 ml). Ammonium ~~captured on the filters~~ was then ana-  
32 lysed on the AMFIA system (AMmonia Flow Injection Analyser, a flow injection system

1 built at ECN). The system is based on the selective diffusion of  $\text{NH}_4^+$  at reduced pH by con-  
2 ductivity. The detection level is 0.2  $\mu\text{g}$  per filter. Sulphate and nitrate were measured in  
3 aqueous solutions with the Dionex IC25 ion chromatograph. ~~This system combines a Ion~~  
4 ~~Chromatography System with a conductivity detector and a high performance dual piston~~  
5 ~~pump combined into one compact package. Samples were eluted through an ion exchange~~  
6 ~~column using a dilute sodium hydroxide solution. The eluent is passed through self-~~  
7 ~~regenerating suppressors that neutralise the eluent conductance, ensuring that electrochemical~~  
8 ~~detectors (conductivity or pulsed amperometric) can detect the ions down to sub ppm concen-~~  
9 ~~trations.~~ Detection level is 0.7  $\mu\text{g}$  per filter.

10 The analysis of carbon is based on thermography (Cachier-method). The filter-sample is  
11 heated in an oven resulting in the release of carbon. For quantitative determination of the  
12 amount of released carbon the evolved gases are oxidised to  $\text{CO}_2$  in a post-combustor oven.  
13 The discrimination of carbon in OC and EC is operationally defined. OC is the carbon re-  
14 leased during the 1<sup>st</sup> step (8 minutes at 340 C); the remainder gives an estimate for EC after  
15 combustion for 6 minutes at 750C. The analysis occurs in a Coulomat 702, manufactured by  
16 JUWE (Viersen, Germany). The total amount of carbon, TC, is directly proportional to the  
17 total amount of  $\text{CO}_2$  produced. The detection system, in principle, is an absolute method.  
18 Monthly calibrations with a standard were performed. Cleaning of the instrument occurred  
19 once a month (after a full month of operation). Several samples were in the oven for a longer  
20 period and also at a higher temperature to check for complete combustion and proper assess-  
21 ment of the EC. The lower detection limit of the method was around 5  $\mu\text{g}$  per filter corre-  
22 sponding to a concentration of 0.1  $\mu\text{g m}^{-3}$ . An average experimental blank of 1.2  $\mu\text{gOC/m}^3$   
23 was found during the campaign (ten Brink et al., 2009) consisting of a filter blank of 0.5  
24  $\mu\text{g/m}^3$ , and a field blank of 0.7  $\mu\text{g/m}^3$  (due to gas adsorption on the filter material). As the  
25 latter contribution ordinarily occurs in routine PM determination following the reference  
26 method, we only subtracted the filter blank from the measured OC values.

27 A selection of 28 metals was analysed with Inductively Coupled Plasma-Mass Spectrometry  
28 (ICP-MS) after microwave digestion of the Teflon filters with a nitric acid/H<sub>2</sub>O<sub>2</sub> mixture  
29 (Na, Mg, Al, Si, P, K, Ca, Ti, V, Cr, Mn, Fe, Co, Ni, Cu, Zn, Pb, As, Ba, Se, Sr, Y, Mo, Cd,  
30 Sn, Li, Be). ~~ICP-MS was employed to analyse the elemental composition of the collected~~  
31 ~~PM<sub>10</sub> and PM<sub>2.5</sub> masses. In short, ICP-MS (Inductively Coupled Plasma Mass Spectrometry)~~  
32 ~~is capable of determining a range of metal and non-metal concentrations with high sensitivity.~~  
33 ~~It is based on combining an inductively coupled plasma to produce ions with a mass spec-~~

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

1 ~~trometer as a method of separating and detecting the ions. ICP-MS ICP-MS is the a~~ standard  
2 operational technique in NAQMN. ~~To account for the non-complete dissolution~~In this study,  
3 ~~we use the (summated) elemental mass primarily to estimate the contributions of suspended~~  
4 ~~mineral dust and total metals~~ of several metals (e.g. Si, Al) using this digestion method. av-  
5 erage correction factors established in a comparison to digestion with HF were used (Schaap  
6 et al., 2010) . ~~The sum of these elements (with the exception of sodium) is denoted by “total~~  
7 ~~metals”.~~

Met opmaak: Lettertype: Vet

## 10 2.2 Data handling

11 For each site only filter samples were included with a complete characterization. Complete-  
12 ness was interpreted rather strict: if one of the components were missing the entire sample  
13 was excluded. As a consequence, the number of samples at a site available for interpretation  
14 differ from site to site varying from 26 (Cabauw) to 38 (Hellendoorn). The selected filter data  
15 lie within the period January-August 2008. The composition of the coarse fraction was ob-  
16 tained by subtracting the concentrations in the PM<sub>2.5</sub> sample from those in PM<sub>10</sub> followed by  
17 the calculation of the averages and standard deviations. Next, the mass concentrations of all  
18 the components were summated and compared with the weighted mass to establish the “not-  
19 analysed” part (further denoted by ‘NA’). ~~The sum of the measured elements (without so-~~  
20 ~~dium) is abbreviated as “total metals”.~~

## 23 3. Results

### 24 3.1 Chemical composition

25 The average PM concentrations in our data set are 22.1 µg/m<sup>3</sup> (PM<sub>10</sub>), 14.8 µg/m<sup>3</sup> (PM<sub>2.5</sub>)  
26 and 7.3 µg/m<sup>3</sup> (PM<sub>10-2.5</sub>) indicating that PM<sub>2.5</sub> is around two-third of PM<sub>10</sub> which is common  
27 in the Netherlands. As expected, levels appear highest at the Rotterdam kerbside and lowest  
28 at rural Hellendoorn. The mass difference for PM<sub>10</sub> between these sites is considerable (near  
29 9 µg/m<sup>3</sup>) and originating from nearly equal increments of the fine and coarse mass (4-5  
30 µg/m<sup>3</sup>).

31 The average composition of the daily PM<sub>10</sub> and PM<sub>2.5</sub> filter samples is presented in Fig. 2 and  
32 summarised in Table 2. An example of the compositional distribution can be seen in Fig. 3  
33 (Schiedam). Fig. 2 directly shows that similarity exists in the chemistry of PM at the various

1 locations. The largest contributions come from nitrate. The concentrations of the 'not-  
2 analysed' (NA) part in the various fractions is at least of similar level. The only exception is,  
3 not unexpectedly, observed at the kerbside location (Rotterdam) where comparable quantities  
4 of carbon in the fine fraction and (summed) metals in the coarse fraction were found. This  
5 is explained by the emissions from traffic passing nearby. At the other sites the next impor-  
6 tant component in PM<sub>2.5</sub> (and PM<sub>10</sub>) is sulphate, followed by elemental carbon, and, in vari-  
7 ous order, organic carbon, ammonium, total metals, chloride and sodium. The major parts of  
8 the inorganic components, elementary and organic carbon, and NA are in the fine fraction.  
9 Contributions of metals and sea-salt related components (sodium and chloride) mainly ar-  
10 rived from the coarser-sized particles. Notable is further the spatial similarity of the nitrate  
11 and ammonium levels with those of the not-analysed part at the five sites. In particular, this is  
12 seen in the PM<sub>2.5</sub> data and suggests a common origin.

13 At the two urban sites (Schiedam and Rotterdam) nitrate appeared as one of the more impor-  
14 tant components within PM<sub>10</sub> with levels near or above 4 µg/m<sup>3</sup>. Considerable part of this  
15 was in the coarse fraction (1.5 and 1.8 µg/m<sup>3</sup>, respectively). A similar pattern was seen for  
16 sulphate but with lower concentrations. Somewhat surprisingly, the highest nitrate, ammo-  
17 nium and sulphate concentrations occurred in the PM<sub>2.5</sub> mass collected at Cabauw. These  
18 were even higher than at Vredepeel which is a site known to be influenced by livestock activ-  
19 ity. Although the rural location of Cabauw was less suspect in this respect, local emissions of  
20 ammonia cannot be excluded and it would explain the elevated concentrations. As a conse-  
21 quence, the assumption that Cabauw could serve as a background site for the urban region  
22 (containing the Rotterdam and Schiedam locations) is not confirmed. In addition, due to a  
23 technical malfunctioning in the winter period at least 7 measurement days at Cabauw were  
24 missing. This explains the relatively low levels for sodium and chloride at this site.

25 The range of OC concentrations in PM<sub>10</sub> is between 1.6 µg/m<sup>3</sup> (Hellendoorn, Cabauw) and  
26 2.3 µg/m<sup>3</sup> (Rotterdam). A recent study by ten Brink et al. (2009) indicated that 70% of the  
27 OC in the Netherlands is contemporary (due to emissions from biogenic material and wood  
28 combustion). The remainder would originate from fossil-fuel sources (i.e. diesel traffic emis-  
29 sions). OC in the coarse fraction is highest at Vredepeel, Cabauw and Rotterdam (compared  
30 to Hellendoorn). At the first two sites this is probably related to nearby biogenic emissions  
31 nearby while at Rotterdam emissions are predominantly from vehicles.



### 3.2 Conversion formula used in the mass closure approach

Certain conversions were applied to the data set to identify the various constituents in PM. This procedure contained estimations of associated oxygen of metals, sea salt (SS), sea-salt sulphate (SS-sulphate), total carbonaceous matter (TCM) and mineral dust (MD). The outcome effectively reduces the 'not-analysed' part into a 'not-explained' or 'unknown' part in the final mass closure. We shortly discuss here the impacts on the mass closure (summarised in Table 3).

Elemental concentrations have been recalculated to their respective oxides to account for associated oxygen (Schaap et al., 2010). The summated "oxidised" contribution appeared 1.4 to 1.5 higher than the elemental contribution. Its incorporation had the largest effect on the coarse fraction: up to  $1.0 \mu\text{g}/\text{m}^3$  (Rotterdam). The change in  $\text{PM}_{2.5}$  was rather small ( $<0.4 \mu\text{g}/\text{m}^3$ ). Consequently, the mass closure improved most for the coarse fraction (9%, on average).

The sea-salt aerosol is generated by sea spray processes when waves break on the sea shore and out at sea. Here, sodium is used as a tracer because it is not affected by depletion and measurements are reliable. The presence of sea salt is then estimated by:  $\text{SS} = 3.26 \cdot \text{Na}$  (contribution of actually measured Cl is then neglected). The factor 3.26 follows from the composition of sea water (i.e., 1 l sea water contains 1 g material among which 0.308 g sodium).

~~The estimated SS contribution will differ from the actually measured sum of Na and Cl concentrations. The difference between this sum and the estimated SS contribution has to be discounted from the NA fraction. The change in the not analysed part is calculated with respect to the sum of Na and Cl (The where minor contributions of magnesium and other components present in seawater were neglected actually lead to a very small double counting of these components as they are part of the "metals" fraction.). The measured Na and Cl concentrations are eliminated from the mass closure total after the SS constituent has been calculated.~~

Generally, the application of the sea-salt conversion has a similar effect as the incorporation of oxidised elemental concentrations. Again the coarse fraction is affected most: the average sea-salt contribution rose with  $0.7 \mu\text{g}/\text{m}^3$  which corresponds with an improvement of 10% in the mass closure. ~~Double counting of sea-salt sulphate has been taken into account by subtraction of this contribution from the actually measured sulphate concentration. The sulphate concentrations have been corrected for a~~The sea-salt contribution which is estimated by  $\text{Na} \cdot (7.68/30.60)$ . The ratio is determined by the molar weights of sulphate and sodium. The sea salt contributions in relation to sulphate appear rather low ( $<0.3 \mu\text{g}/\text{m}^3$ ). The sea salt con-

Met opmaak: Spatiëring tussen Aziatische en Latijnse tekst niet aanpassen, Spatiëring tussen Aziatische tekst en en nummers niet aanpassen

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

1 tribution to the coarse fraction and the effect of Cl-depletion will be treated further in the  
2 dDiscussion.

3 Carbonaceous constituents contain elements other than carbon that contribute to the mass of  
4 PM (e.g., oxygen). The conversion from OC to organic material (OM) is even quite uncertain  
5 in the US where many more monitoring data of OC/EC are available than in Europe. This is  
6 the reason to take a minimalistic approach for a conversion factor via the parameter TC. The  
7 use of total carbon in turn is due to the uncertain OC-EC split in the various chemical analy-  
8 sis approaches in use in Europe (ten Brink et al., 2004). These analysis procedures corres-  
9 pond well for total carbon, but yield very different EC/OC ratios.

10 We follow here the approach of US-EPA (Frank, 2006) who compared total mass with sum-  
11 med contributions of the chemical components analysed for a very large number of filters.

12 Frank obtained an optimal mass closure with total carbonaceous matter (TCM) defined as 1.3  
13 times the amount of total carbon (TC). This is derived from measurements in the Speciation  
14 Trends Network which contains by definition urban background stations. Therefore, an aver-  
15 age factor of 1.3 seems a good compromise in the present study with three types of stations  
16 (rural, urban background and kerbside). Using this approach the largest relative change is  
17 calculated for PM<sub>2.5</sub> which is no surprise as TC is predominantly present in the fine fraction.  
18 The mass closure improved with some 7% (for PM<sub>2.5</sub>). The maximum change was observed  
19 at the kerbside site in Rotterdam (+1.4 µg/m<sup>3</sup> for PM<sub>2.5</sub>).

20 Carbonaceous constituents in PM contain other elements than carbon contributing to the  
21 mass of PM (e.g., oxygen). The conversion from OC to organic material (OM) is rather un-  
22 certain: in literature the factor varies between 1.2 and 2 and it may well be possible that the  
23 value decreases near emission sources and increases in aged air masses. US EPA includes  
24 the associated elements in their approach by following Frank (2006) who compared total  
25 mass with summated contributions of components analysed for a very large number of filters.

26 Frank calculated that total carbonaceous matter (TCM) defined by 1.3 times the amount of  
27 total carbon (TC) would have the best agreement. An additional reason to use total carbon  
28 and not convert OC to OM is due to the uncertain OC-EC split in the chemical analysis. Cur-  
29 rent analysis procedures correspond well for total carbon, but yield very different EC/OC ra-  
30 tios (Ten Brink et al., 2009). Using the approach of Frank the largest relative change is cal-  
31 culated for PM<sub>2.5</sub> which is no surprise as TC is predominantly present in the fine fraction.  
32 The mass closure improved with some 7%. The maximum change was observed at the kerb-  
33 side site in Rotterdam (+1.4 µg/m<sup>3</sup>).

Met opmaak: Lettertype: (Standaard)  
Times New Roman, 12 pt

Met opmaak: Uitsluitend, Regelafstand:  
1,5 regel

Met opmaak: Lettertype: (Standaard)  
Times New Roman, 12 pt

Met opmaak: Lettertype: (Standaard)  
Times New Roman, 12 pt

Met opmaak: Lettertype: (Standaard)  
Times New Roman, 12 pt

Met opmaak: Lettertype: (Standaard)  
Times New Roman, 12 pt

Met opmaak: Lettertype: (Standaard)  
Times New Roman, 12 pt

Met opmaak: Lettertype: (Standaard)  
Times New Roman, 12 pt

Met opmaak: Lettertype: (Standaard)  
Times New Roman, 12 pt

Met opmaak: Lettertype: (Standaard)  
Times New Roman, 12 pt

Met opmaak: Lettertype: (Standaard)  
Times New Roman, 12 pt

Met opmaak: Lettertype: (Standaard)  
Times New Roman, 12 pt

Met opmaak: Lettertype: (Standaard)  
Times New Roman, 12 pt

Met opmaak: Lettertype: (Standaard)  
Times New Roman, 12 pt

Met opmaak: Lettertype: (Standaard)  
Times New Roman, 12 pt

Met opmaak: Lettertype: (Standaard)  
Times New Roman, 12 pt

Met opmaak: Lettertype: (Standaard)  
Times New Roman, 12 pt

Met opmaak: Lettertype: (Standaard)  
Times New Roman, 12 pt

Met opmaak: Lettertype: (Standaard)  
Times New Roman, 12 pt

Met opmaak: Lettertype: (Standaard)  
Times New Roman, 12 pt, Engels (V.S.)

Met opmaak: Lettertype: Cursief

Met opmaak: Lettertype: 11 pt,  
Cursief

Met opmaak: Lettertype: Cursief

Met opmaak: Lettertype: Cursief,  
Engels (V.S.)

1 Mineral dust is defined here as fugitive wind-blown and mechanically resuspended soil dust  
2 with a composition comparable to the earth's crust containing elements like Si, Al, Ca, K, Fe  
3 and Ti. In our data set the correlation between Al and Si turned out to be very good ( $r^2=0.96$ ).  
4 This suggests that both originate from the same source and are present in similar ratios. The  
5 method to estimate the contribution of MD here is by focusing on these two abundant ele-  
6 ments according to  $MD = 0.49 * Si + (2.36*Si + 2.7*Al)$  (proposed by Denier van der Gon,  
7 2009). All elemental weights were recalculated first to their respective oxide concentrations.  
8 The calculated MD contribution is subtracted from the total elemental mass which leaves the  
9 final mass balance unaffected. The outcome is further denoted by 'metals'.

### 10 3.3 Mass closure

11 The resulting concentrations of the various constituents are given in Table 4. The resulting  
12 mass closure is again given for the Schiedam site in Fig. 4. The dominant constituent in the  
13  $PM_{2.5}$  filter samples is the secondary inorganic components (SIA). There is not much differ-  
14 ence between most measurement sites as concentrations range between 6 and 7  $\mu g/m^3$ . A  
15 maximum occurs at Cabauw (8.4  $\mu g/m^3$ ) due the enhanced levels of ammonium and nitrate  
16 noted earlier. The second important contribution comes from total carbonaceous matter  
17 (TCM) showing more variation. TCM in  $PM_{2.5}$  is higher at the urban sites (4.0-6.1  $\mu g/m^3$ )  
18 than at the rural sites (3.3-3.9  $\mu g/m^3$ ). The remaining constituents, i.e., sea salt (SS), mineral  
19 dust (MD), and metals contribute considerably less with concentrations always below 1  
20  $\mu g/m^3$ . The unexplained part in the  $PM_{2.5}$  mass fraction tends to be higher at the rural sites  
21 with a maximum at Cabauw of 3.5  $\mu g/m^3$ . At the other sites the missing part was always less  
22 than 2.5  $\mu g/m^3$ .

23 The presence of the various constituents in the coarse fraction is more evenly distributed than  
24 in the fine fraction. Relatively high contributions arise from sea salt ( $\leq 2.7 \mu g/m^3$ ; maximum  
25 at Schiedam) and SIA ( $\leq 2.3 \mu g/m^3$ ; higher levels at Rotterdam and Schiedam). The concen-  
26 trations of MD and the remaining metals are always less than 1.4  $\mu g/m^3$  (Cabauw) and 1.6  
27  $\mu g/m^3$  (Rotterdam), respectively. Remarkable is rather high TC levels at Vredepeel (1.3  
28  $\mu g/m^3$ ) and Cabauw (0.7  $\mu g/m^3$ ) compared to Rotterdam (1.2  $\mu g/m^3$ ) and Schiedam (0.7  
29  $\mu g/m^3$ ) The reason for these enhanced TC levels in the coarse fraction is not clear.

30 To determine which of the constituents are the main contributors when high PM levels (e.g.,  
31 on exceedence days) occur the data set has been partitioned into two subsets, containing the  
32 data for days at which the  $PM_{10}$  ( $PM_{2.5}$ ) concentrations were less or larger than 30 (25)  
33  $\mu g/m^3$ . Not surprisingly, the absolute contributions of most constituents rise when PM in-

1 creases. The difference between the  $PM_{10}$  levels when averaged over the two sets is 20  
2  $\mu\text{g}/\text{m}^3$ . The largest changes in absolute concentrations are observed for SIA (+11  $\mu\text{g}/\text{m}^3$  in  
3  $PM_{10}$ ) and the unknown part (+6  $\mu\text{g}/\text{m}^3$ ) followed by smaller or negligible changes for TCM  
4 (+3  $\mu\text{g}/\text{m}^3$ ), MD (+1.5  $\mu\text{g}/\text{m}^3$ ) and metals (<+0.5  $\mu\text{g}/\text{m}^3$ ). The only exception is sea salt  
5 which decreases at higher PM levels (-1.8  $\mu\text{g}/\text{m}^3$ ). The corresponding changes in chemical  
6 distributions of  $PM_{10}$  and  $PM_{2.5}$  are summarised in Fig. 5. It can be seen that when expressed  
7 in percentages both SIA and the unknown part become more dominant at days with more PM.  
8 Correspondingly, the relative contributions of the remaining constituents (MD, TCM and SS)  
9 decrease at higher PM concentrations.

10 The two measurement sites in the Rotterdam urban area allow investigating which compo-  
11 nents are emitted by traffic by comparing measured concentrations at the kerb site (Rotter-  
12 dam) to those at the urban background site (Schiedam). The difference in  $PM_{10}$  between the  
13 two sites appeared 3.5  $\mu\text{g}/\text{m}^3$  (based on 27 corresponding days) and can be attributed to car-  
14 bonaceous material (2.5  $\mu\text{g}/\text{m}^3$ ) and metals (1  $\mu\text{g}/\text{m}^3$ ). The carbonaceous material increment  
15 is largely due to elementary carbon. The metals that are enhanced include copper, iron, bar-  
16 ium and molybdenum and result from tail-pipe emissions ~~(TCM)~~, and brake and tire wear.  
17 Surprisingly, there is no similar increment in the concentrations of mineral dust. ~~This indi-~~  
18 ~~icates implying~~ that this is not an important source at Rotterdam ~~resuspension is not a signifi-~~  
19 ~~cant source~~. As this disagrees with the outcomes from other studies in the Netherlands  
20 (e.g. Visser et al., 2001) and elsewhere (e.g., Amato et al., 2009) further investigations on this  
21 issue the origin of mineral dust and the role of traffic are necessary.  
22

## 23 4. Discussion

### 24 4.1 Chemical composition

25 The data set gives a proper insight in the chemical characteristics of PM in the Netherlands.  
26 Generally, a considerable similarity in the chemical composition of PM is observed at the  
27 five measurement sites. Always, secondary inorganic components are the dominant constitu-  
28 ent with contributions responsible for nearly half of the total mass (42-48% of  $PM_{2.5}$ ) with  
29 nitrate as the most important component. The variation in SIA concentrations appears rather  
30 limited indicating that its behaviour is mainly driven by larger-scale meteorology and not by  
31 local sources. Nearby emissions of ammonia may incidentally lead to locally high levels  
32 (e.g., Cabauw). The next important contribution originates from total carbonaceous matter  
33

1 showing more variation over the sites studied here (22-37%). TCM in PM<sub>2.5</sub> tends to be  
2 higher in areas where more traffic prevails (Rotterdam, Schiedam). The remaining constitu-  
3 ents, i.e., sea salt, mineral dust and metals are relatively unimportant as they contribute each  
4 less than 8%.

5 The study allows the first detailed overview of the chemical composition of the coarse frac-  
6 tion in the Netherlands. Compared to the fine fraction both SIA and TCM are less dominant  
7 (around 29% and 25%, respectively) becoming comparable with mineral dust (<23%) and sea  
8 salt (<30%). The remaining metals sum up to 16% (Rotterdam). It was further noted that at  
9 the two urban sites considerable part of SIA (25%) was in the coarse fraction. It is assumed  
10 that in the marine urban environment (Schiedam and Rotterdam are within 30 km of the sea-  
11 shore) the coarse sea-salt particles act as reactive sinks for nitric and sulphuric acids. The im-  
12 plications of these mechanisms will be discussed further below.

13 As anticipated, absolute contributions of most constituents rise when PM concentrations go  
14 up. However, this does not occur at the same rate for each constituent. The largest increase is  
15 observed for SIA, most notably nitrate, and the missing mass part. By consequence, the  
16 chemical distribution rearranges at days with higher PM concentrations: SIA and the un-  
17 known part become more dominant, the contribution of mineral dust remains the same while  
18 the relative importance of TCM and sea salt decrease. Such behaviour can be explained by  
19 considering the atmospheric situation these days and the localisation of the major emissions.  
20 The high concentrations of PM<sub>2.5</sub> and PM<sub>10</sub> occur at days with stagnating conditions or slow  
21 transport from south, south-east or east. As the emissions are diluted less due to the lower  
22 wind speeds and removal is low, pollution starts to build up. During the winter period this ef-  
23 fect is strengthened due to the shallow boundary layers. In view of the dominant transport di-  
24 rection (continental) it is not surprising that the contribution of sea salt diminishes. In addi-  
25 tion to unfavourable meteorological conditions, considerable emissions for SO<sub>2</sub> and NO<sub>x</sub>  
26 take place east of the Netherlands (industrial Ruhr area); for NH<sub>3</sub> this is in the Southeast (a  
27 centre of intensive livestock industry). Model studies also indicate high contributions of na-  
28 tional sources to the concentrations of primary material and ammonium nitrate (Schaap and  
29 Denier van der Gon, 2007). Also some special events contribute to a (small) number of ex-  
30 ceedence or days with high concentrations. In the data set we have identified contributions  
31 from desert dust, fireworks at New Year's Eve, and bonfires at Easter.

32

#### 33 **4.2 Mass closure and the unknown part**

1 To correct for known underestimations a number of algorithms to estimate various constitu-  
2 ents has been applied. Most of these (oxide concentrations, sea salt, total carbon material)  
3 improve the mass closure. In the case of PM<sub>2.5</sub> the conversion from TC to TCM results in a  
4 more complete mass closure in the PM<sub>2.5</sub> samples at the urban sites (93-94%). The closure  
5 appears less perfect at the rural sites (80, 83 and 90%) demonstrating the sensitivity on site  
6 characteristics. In contrast to the fine fraction, the mass closure in the coarse fraction seems  
7 better at the rural sites. The unexplained parts are now less than 4% of the total mass (which  
8 is in absolute mass units negligible) while in the urban environment this is 12 (Rotterdam)  
9 and 16% (Schiedam) or, 1 and 1.5 µg/m<sup>3</sup>, respectively.

10 A similarity has been noted between SIA in PM<sub>2.5</sub> (more specific, nitrate and, to a lesser ex-  
11 tent, ammonium) and the unexplained mass. The largest mass deficit (measured at Cabauw)  
12 coincides with the highest SIA concentration; at Schiedam minimum SIA concentrations and  
13 mass deficit have been measured. To a lesser extent, similar behaviour is seen in the coarse  
14 mass fraction. Here, SIA concentrations at the urban sites are (somewhat) higher and corre-  
15 pond with larger mass deficits. By consequence, the 'contribution' of the unexplained mass  
16 is also important at days with higher PM concentrations since the presence of SIA (or more  
17 specific nitrate) is the main reason for these augmented levels.

18 A possible reason may be the amount of water attached to SIA. Hygroscopic salts on parti-  
19 cles, like ammonium nitrate and ammonium sulphate in the fine fraction, and sodium nitrate  
20 and sodium sulphate in the coarse fraction attract water and contributes to the PM mass  
21 measurements. It can be estimated that the amount of water on PM<sub>10</sub> due to the presence of  
22 SIA is about 10% of the SIA mass which is 4-5% of the PM<sub>10</sub> mass concentration. Such a  
23 contribution is possible as it is lower than the total unexplained mass. However, although im-  
24 portant it can not be the sole contributor to the unexplained mass.

25 ~~To correct for known underestimations we applied a number of algorithms to estimate vari-~~  
26 ~~ous constituents. Most of these recalculation (oxide concentrations, sea salt, total carbon ma-~~  
27 ~~terial) improve the mass closure. For example, in case of PM<sub>2.5</sub> the conversion from TC to~~  
28 ~~TCM results in a decrease in the unknown mass part of 6-8%. Not surprisingly, the more~~  
29 ~~complete mass closure in the PM<sub>2.5</sub> samples was obtained at the urban sites (93-94%). The~~  
30 ~~closure appears less perfect at the rural sites (80, 83 and 90%) demonstrating the sensitivity~~  
31 ~~on site characteristics. As noted earlier, SIA do not largely differ in PM<sub>2.5</sub> (and PM<sub>10</sub>) be-~~  
32 ~~tween the locations. As the contributions of the remaining components are relatively minor~~  
33 ~~in PM<sub>2.5</sub>, a larger contribution of TCM in PM<sub>2.5</sub> at urban locations results in a smaller part of~~  
34 ~~PM that is unexplained.~~

1 In contrast to the fine fraction, the mass closure in the coarse fraction seems better at the rural  
2 sites. Here, the unexplained parts are less than 4% of the total mass (which is in absolute  
3 mass units negligible) while in the urban environment it is 12% (Rotterdam) and 16%  
4 (Schiedam) or, in absolute value, 1 and 1.5  $\mu\text{g}/\text{m}^3$ , respectively.

5 A correlation is observed between the SIA presence in  $\text{PM}_{2.5}$  (more specific, nitrate and, to a  
6 lesser extent, ammonium) and the amount of unexplained mass at the five sites. For example,  
7 the largest mass deficit (measured at Cabauw) coincides with the highest SIA concentration;  
8 at Schiedam minimum SIA concentrations and mass deficit have been measured. To a lesser  
9 extent, a similar behaviour is noted for the coarse mass fraction. Here, the SIA concentrations  
10 at the urban sites are (somewhat) higher corresponding with larger mass deficits. By conse-  
11 quence, also the 'contribution' of the unexplained mass becomes more important at days with  
12 higher PM concentrations since the presence of SIA (or more specific nitrate) is the main rea-  
13 son for these augmented levels.

14 A possible reason may be the amount of water attached to SIA. Water is a natural component  
15 of particulate matter. Hygroscopic salts on particles, like ammonium nitrate and ammonium  
16 sulphate in the fine fraction, and sodium nitrate and sodium sulphate in the coarse fraction at-  
17 tract water. The water bound to particles contributes to the PM mass measurements. Ideally,  
18 this water is removed by pre heating of the air sample or by conditioning of the filters under  
19 dry (50% RH) circumstances following the reference method (NEN12341). However, heating  
20 or drying only partly removes the particle bound water due to hysteresis (e.g. Speer et al.,  
21 2003). They report molar ratios between 0.5 and 1.0 for water on ammonium nitrate and am-  
22 monium sulphate particles at a RH of 50% and 75%, respectively.  $\text{PM}_{10}$  reference measure-  
23 ments are to be performed at temperature of 20 °C and a relative humidity of 50%. With a  
24 molar ratio of 0.5 for water on ammonium nitrate and ammonium sulphate particles. As such,  
25 we estimate the amount of water on  $\text{PM}_{10}$  due to the presence of SIA is about 10% of the SIA  
26 mass. This results in an estimate of 4-5% of the  $\text{PM}_{10}$  mass concentration. Such a contribu-  
27 tion is possible as it is generally lower than the total unexplained mass. Concluding, water is  
28 expected to be an important, though not the sole, contributor to the unexplained mass.

29 A similar water uptake may be associated with sea salt. However, as the contribution of sea  
30 salt to the total mass is low, its water contribution is much less compared to that of SIA, at  
31 least in the case of  $\text{PM}_{2.5}$ . Likewise the amount of water on organic carbon (OC) is unlikely to  
32 play an important role. The contribution of hygroscopic OC to ambient PM levels in the  
33 Netherlands is much smaller than the contribution of SIA. Hygroscopic OC is a fraction of

1 the total OC amount ( $\sim 2 \mu\text{g}/\text{m}^3$ ) and thus much lower than SIA levels ( $\sim 10 \mu\text{g}/\text{m}^3$ ). More-  
2 over, the amount of water bound to a unit OC is less than on SIA (Speer et al., 2003).

### 4.3 Role of nitrate

5 According to the formation pathways ammonia gas reacts reversibly with  $\text{HNO}_3$  to form its  
6 ammonium salt,  $\text{NH}_4\text{NO}_3$ , which is semi-volatile and may release their gas phase precursors  
7 under conditions of low atmospheric  $\text{NH}_3$  levels or high temperature according to  $\text{NH}_3 +$   
8  $\text{HNO}_3 \leftrightarrow \text{NH}_4\text{NO}_3$ . In marine and coastal atmospheres, nitric acid is also converted into par-  
9 ticulate nitrate ( $\text{NaNO}_3$ ) through the reaction with sea salt particles resulting in the release of  
10  $\text{HCl}$ :  $\text{NaCl} + \text{HNO}_3 \rightarrow \text{NaNO}_3 + \text{HCl}$ . In contrast to  $\text{NH}_4\text{NO}_3$ ,  $\text{NaNO}_3$  is a non-volatile com-  
11 pound under atmospheric conditions therefore, partitioning of nitric acid into the sea salt is  
12 irreversible. In the Netherlands most of the nitrate is found in the fine mode as ammonium  
13 nitrate. However, the coarse nitrate fraction is still a substantial part of the analysed coarse  
14 mode (Table 2). In particular, the coarse-to-  $\text{PM}_{10}$  ratio in the case of nitrate is near 40% at  
15 the urban sites (and some 26% at the rural sites).

16 ~~From this formation pathway the problem arises how to correctly assign the coarse mode ni-~~  
17 ~~trate to the PM constituents SIA and SS, respectively. It is this formation pathway that poses~~  
18 ~~a complication in the chemical mass closure approach.~~ In the chemical mass balance sodium  
19 is used as a tracer of sea salt and multiply its concentration with a factor to obtain an estimate  
20 of fresh sea-salt aerosol, including all the chloride. Hence, ~~when one would put both the sea~~  
21 ~~salt estimate and the full coarse mode nitrate into the SS constituent, the sea salt~~  
22 ~~when one~~  
23 ~~would keep both the sea salt estimate and the full coarse mode nitrate in the chemical mass~~  
24 ~~balance, the sea salt~~ and nitrate contributions cannot simply be added as the nitrate replaces  
25 chloride. Consequently, one needs to make a correction in the chemical mass balance ap-  
26 proach for this double counting.

26 In principle, there are two options to remove the double counting. ~~The first is to replace (on~~  
27 ~~a molar basis) a part or all of the estimated chloride concentration in the SS constituent by the~~  
28 ~~measured coarse nitrate concentration. In case the coarse nitrate exceeds the estimated chlo-~~  
29 ~~ride the remaining part of nitrate is kept in the SIA constituent. The first is to remove the lost~~  
30 ~~chloride mass from the sea salt estimate and to keep the coarse nitrate as it is. The second op-~~  
31 ~~tion is to keep the estimated SS constituent as it is and to discount from the SIA constituent~~  
32 ~~the coarse nitrate fraction that potentially may replace the sea salt chloride. The second op-~~  
33 ~~tion is to assign only the net mass addition to the coarse nitrate part. The first option reflects~~  
34 ~~the substitution processes taking place on sea salt particles. The first option reflects the meas-~~

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Regelafstand: 1,5 regel

Met opmaak: Lettertype: 12 pt

Met opmaak: Spatiering tussen Aziatische en Latijnse tekst niet aanpassen, Spatiering tussen Aziatische tekst en en nummers niet aanpassen

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt

Met opmaak: Lettertype: 12 pt



~~ured chemical composition of the particles.~~ The second option reflects the sources and especially the associated reduction potential. Though both choices are valid and defensible, it can be argued that the second option suits better if we are after the source apportionment. Reduction of NO<sub>x</sub> emissions may cause a reduction of coarse mode nitrate, which is partly compensated by the fact that chloride is not lost anymore. Such a reduction would yield a net result of  $((\text{NO}_3\text{-Cl})/\text{NO}_3 = (62-35)/62 = 27/62)$  times the nitrate reduction (where the number involved are molar weights of the respective components). Hence, the same factor could be used to scale back the coarse nitrate fraction in the chemical mass balance. The motivation would be that when all anthropogenic sources would be non-existing one still has the sea salt left that is given now in the mass distributions. Doing so, the coarse nitrate would, on average, be lower with some  $0.6 \mu\text{g}/\text{m}^3$  which gives an increase in the NA part of the coarse fraction with some 4%. A similar reasoning may be valid for the anthropogenic sulphate in the coarse fraction. Corrections like these are uncommon in current mass closure studies and consequences will have to be explored in more detail in future.

#### 4.4 Natural fraction

For policy makers the extent to which particulate matter can be influenced by measures is of importance. It is the anthropogenic part that could be targeted by abatement strategies. Here, a first-order approximation of the natural and anthropogenic contribution to PM<sub>10</sub> and PM<sub>2.5</sub> is developed.

Sea spray emissions are the most important natural source of primary *sulphate*. Applying the sodium-to-sulphate ratio learns that roughly 5% of the particulate sulphate is marine. Though sulphate may be emitted directly into the atmosphere, its major source is the oxidation of sulfur dioxide. In the Netherlands anthropogenic SO<sub>2</sub> emissions derive mostly from the combustion of sulfur containing fuels for power generation and international shipping. In Europe, natural emissions for SO<sub>2</sub> include volcanoes but given location, height and distance to the Netherlands their contribution is probably low (i.e., not more than 1 percent to ground level sulphate). Contributions of other biogenic sources of SO<sub>2</sub> like oxidation of DMS, COS and H<sub>2</sub>S are also of minor importance (Bates et al., 1992). Wildfire emissions of SO<sub>2</sub> are commonly neglected in studies directed to wild land fire emissions. ~~In short, the natural contribution to sulphate in the Netherlands is estimated to be 5-10%. In the approximation 5% is used (as a conservative estimate).~~

To estimate how much of the *nitrate* is natural NO<sub>x</sub> emissions from soil, biomass burning and lightning should be considered. Studies on NO emissions from soil report emissions between

1 59 and 190 Kton in the EU15 (Simpson et al., 1999; Friedrich et al., 2008). Relative to the  
2 total inventoried emissions the central value is 4%. From these emissions about two third is  
3 attributed to agricultural soils and one third to forest soils. Soil NO<sub>x</sub> emissions largely depend  
4 on the nutrient input (through fertilization or atmospheric deposition) implying that the ab-  
5 ovementioned contribution is only partly natural. Lightning depends linearly on the amount  
6 of convective precipitation. In Europe corresponding emissions are estimated to be about 1%  
7 (65 Kton) of the total inventoried emissions It mainly takes place between 1 and 5 km alti-  
8 tude and mostly in southern Europe (Meijer et al., 2001; Friedrich et al., 2008). The available  
9 wild fires estimates range between 20-50 Kt (Friedrich et al., 2009; Hoelzemann et al., 2004)  
10 and also occur mostly in countries around the Mediterranean. Given the short life time of  
11 NO<sub>x</sub> and the rather short transport distance of nitrate we assume the impact of lightning and  
12 wild land fires can be negligible (<1%) in the Netherlands. ~~In short, the natural fraction of~~  
13 ~~NO<sub>x</sub> emissions and therewith nitrate is estimated to be 5% (absolute upper limit).~~In the ap-  
14 proximation, the natural fraction of NO<sub>x</sub> emissions and therewith nitrate is estimated to be  
15 between 0 and 5%.

16 *Ammonia* is emitted from, most notably, animal manure, traffic and fertilizer application  
17 (Olivier et al., 1998). Emissions in Europe are for the largest part (80-95 %) associated with  
18 agricultural activities (Van der Hoek, 1998). Recently, Erisman et al. (2009) estimated the  
19 natural emissions at about 10% of the total emission in Europe. This percentage includes con-  
20 tributions from wild animals and wetlands. A similar percentage is assumed for the Nether-  
21 lands.

22 *Elemental carbon* is emitted from combustion processes and a pure primary component. The  
23 main source is the (incomplete) combustion of fossil fuels. Wild land fires are occasional  
24 sources of vast amounts of carbonaceous particles. Though wild land fires may seem natural,  
25 the vast majority of present day fires are due to human behavior. Hence, elemental carbon is  
26 almost exclusively anthropogenic.

27 *Organic matter* is released from combustion processes, agricultural activities, as well as from  
28 nature. Anthropogenic combustion particles (as well as small contribution of stable emis-  
29 sions) are inventoried. About half of the observed concentrations of OC (Schaap et al., 2004;  
30 Simpson et al., 2007) can be explained. Other primary sources include biological material  
31 (fungal spores and plant debris) may contribute up to 0.4 µg/m<sup>3</sup> (about 6% of OM) at rural  
32 background sites in central Europe (Sanchez-Ochoa et al., 2008). Recently, <sup>14</sup>C analysis sug-  
33 gested that some 70% (Ten Brink et al., 2010) of the OC mass in the Netherlands was of liv-  
34 ing material where biogenic SOA, agriculture and other landscaping activities may have con-

1 tributed. The complicating factor in the estimation of the natural part of OM is the uncertainty  
2 related to its secondary component (oxidation of VOCs yields products with low vapor pres-  
3 sures that may condense on existing aerosol) as the major formation routes are not well  
4 known. Similarly, the ratio of natural to anthropogenic SOA is under large scientific debate.  
5 Note that the oxidation pathways that yield SOA are largely influenced through hydroxyl and  
6 ozone from anthropogenic emissions. Presently, it is postulated that biogenic sources are a  
7 major contributor to atmospheric SOA (e.g. Simpson et al., 2007). Assuming that the unex-  
8 plained OC using present day emission inventories is for the largest part SOA an upper limit  
9 of ~50% of natural OM is obtained. As a substantial part of the SOA may very well be anth-  
10 ropogenic we postulate a second guess with half of the SOA as natural, leading to 25% of  
11 natural OM.

12 | *Mineral dust* usually originates from different sources, e.g., wind erosion of bare soils, agri-  
13 cultural land management, re-suspension of road dust, road wear, driving on unpaved roads,  
14 handling of materials and building and construction activities. The contributions of these  
15 sources are still poorly understood. Here, only wind erosion is relevant as natural source.  
16 Globally the atmospheric input through wind erosion is dominated by a few areas exception-  
17 ally susceptible for wind erosion. Saharan dust is regularly transported to countries around  
18 the Mediterranean Sea. In central and northern Europe dust transport from the Sahara occurs  
19 once or twice a year and is not considered very significant. During this study we could only  
20 identify one day with a significant impact of Saharan dust to PM (contribution ~4% of the  
21 annual mean MD concentration). Korcz et al. (2009) and Schaap et al. (2009) show that wind  
22 blown dust at the European continent is a much smaller source for dust than traffic re-  
23 suspension and agricultural land management. More importantly, the wind blown dust source  
24 strength from soils other than arable land is low. Hence, wind blown dust emissions are  
25 strongly related to anthropogenic changes in surface vegetation cover and are regarded as  
26 mostly anthropogenic. Consequently, the total mineral dust concentration in air is expected to  
27 be anthropogenic for a large part. ~~To be on safe side an upper limit of 20% of the mineral~~  
28 ~~dust is assumed to be of natural origin. In the approximation we use 10% as a conservative~~  
29 ~~estimate for the natural contribution.~~

30 | *Metals* originate from a vast range of abrasion processes and metallurgy industry. Though  
31 soils may contribute to metal concentrations they are in principle accounted for in the mineral  
32 dust estimate. Here, all metals are assumed to be anthropogenic.

33 The natural fraction of the *unknown mass* is by definition unknown. As water partly explains  
34 the unknown fraction and most of the water appears associated with SIA it is likely that a

1 significant part of the unknown fraction can be considered to anthropogenic. We assume here  
2 that the anthropogenic/natural ratio within the unknown mass fraction resembles that of the  
3 explained mass.

4 The natural fraction in the Netherlands is then obtained by adding relative natural mass con-  
5 tributions per component. For instance, in Hellendoorn 11.5% of  $PM_{10}$  is associated with sea  
6 salt adding to 11.5% of natural  $PM_{10}$  (see Table 5). Note, that this fraction represents the

7 source attribution and therewith fresh sea salt as discussed in section 4.3 and would be lower  
8 when one reflects the chloride loss in this calculation. Nitrate contributes 21 % of which 0 or  
9 5% is assumed to be natural resulting in a contribution of 0 or 1% natural  $PM_{10}$  for the low  
10 and high cases, respectively. Hence, nitrate contributes very little to the natural fraction. -

11 Adding all contributions, the estimated natural fraction of  $PM_{10}$  is 245%. The low estimate  
12 for carbonaceous particles yields a natural fraction of 1924%. Hence, around one quarter of  
13  $PM_{10}$  is estimated to be natural. Performing the same exercise for  $PM_{2.5}$  yields estimates of  
14 178 and 134% for the two cases, respectively. Application of this methodology to all sites in  
15 this campaign yields natural fractions of about one fourth for  $PM_{10}$  and one fifth for  $PM_{2.5}$   
16 (Fig. 6). In case of the low natural OM fraction the natural estimates are about 5% lower than  
17 with the high case.

## 18 5. Conclusions

19 There is a considerable conformity in the chemical composition of PM over the Netherlands.  
20 Most dominant is the secondary inorganic aerosol – the sum of sulphate, nitrate and ammo-  
21 nium – showing slightly higher levels at the rural sites. Carbon containing components are  
22 generally the second largest contributor being enhanced in the traffic dominated urban area.

23 The most important natural constituent is sea salt, especially in the western part of the Neth-  
24 erlands near the North Sea. Contributions of mineral dust and metals appear relatively low.  
25 The distribution of species in  $PM_{10}$  resembles that of  $PM_{2.5}$  which is not surprising as the  
26 mass of  $PM_{2.5}$  is about two thirds of the total  $PM_{10}$  mass. The chemical composition observed  
27 in the coarse mass fraction appears more balanced: contributions of mineral dust and sea salt  
28 are of comparable magnitude with those of SIA and TCM.

29 On days with high PM concentrations, all constituents, except sea salt, show higher concen-  
30 trations. Sea salt maximises in marine air masses associated with clean air conditions. The ni-  
31 trate and unknown contributions increase most on these days suggesting a relationship by

Opmerking [s1]: Hier moet nog wat  
over min/mac nitraat

1 means of the associated water content. The relative contribution of the remaining components  
2 is mostly stable as function of PM concentration.

3 After analysing the filter samples on chemical content the sum of all the components ac-  
4 counts on average for 70-80% of the total mass. Hence, a full analysis is not obtained, but ex-  
5 isting knowledge can be used to narrow down partially the remaining gap. For example, met-  
6 als can be taken into account as their oxides and total carbonaceous matter (including non-C  
7 atoms) is estimated based on the total carbon content. In addition, the contributions of min-  
8 eral dust and sea salt were estimated. After this procedure, 80 to 94% of the total mass can be  
9 explained. Uncertainties occurring in the mass closure are conversion factors for total carbo-  
10 naceous matter and the displacement of chloride by nitrate in the coarse fraction.

11 Based on a literature review, it is concluded that the most important natural contributions ori-  
12 ginate from sea salt (100%) and organic material (with an upper limit of 50%). Including the  
13 natural parts of the remaining constituents, estimates for the entire natural contribution yield  
14 25% for PM<sub>10</sub> and 20% for PM<sub>2.5</sub>. This (fairly pragmatic) analysis illustrates that the majority  
15 of PM in the Netherlands is of anthropogenic origin.

16 A major gap in knowledge remains the characterisation of the unknown mass and the role of  
17 the water content of the filter. Another caveat is the chemical characterisation of the urban  
18 increment between (with respect to rural background concentrations). The technical inability  
19 in the differentiation between EC and OC influences results. For more traditional PM compo-  
20 nents, like SIA and primary material, a modelling may identify source contributions at sector  
21 level, and provide insight into national versus foreign sources. The origin of carbonaceous  
22 matter remains one of the biggest challenges, as knowledge in this respect is still moderate.  
23

1

2 **Acknowledgements**

3 This study is conducted under the auspices of the Netherlands Research Program on Particu-  
4 late Matter (BOP), a national program on PM<sub>10</sub> and PM<sub>2.5</sub>, funded by the Netherlands Minis-  
5 try of Housing, Spatial planning and the Environment (VROM). The program is a framework  
6 of cooperation, involving four Dutch institutes: the Energy research Centre of the Nether-  
7 lands (ECN), the Netherlands Environmental Assessment Agency (PBL), the National Insti-  
8 tute for Public Health and the Environment (RIVM), and TNO Built Environment and Geo-  
9 sciences.

10

11

1 **References**

2 Amato, F., Pandolfi, M., Escrig, A., Querol, X., Alastuey, A., Pey, J., Perez, N., and Hopke,  
3 P.K.: Quantifying road dust resuspension in urban environment by multilinear engine: A  
4 comparison with PMF2, *Atmos. Environ.*, 43, 2770-2780, 2009.

Met opmaak: Lettertype:

5  
6 Almeida, S.M., Pio, C.A., Freitas, M.C., Reis, M.A. and Trancoso, M.A.: Source apportionment of fine and coarse particulate matter in a sub-urban area at the Western European Coast. *Atmos. Environ.* 39, 3127–3138, 2005.

7  
8  
9  
10 Bates, T. S., Lamb, B. K., Guenther, A., Dignon, J. and Stoiber, R. E.: Sulfur emissions to the atmosphere from natural sources. *J. Atmos. Chem.*, 14, 315-337, 1992.

11  
12  
13 Brunekreef, B. and Holgate, S.T.: Air Pollution and Health, *Lancet*, 360, 1233-1244, 2002.

14  
15 Denier van der Gon, H., Jozwicka, M., Hendriks, E., Gondwe, M. and Schaap, M.: Mineral dust as a constituent of particulate matter. PBL report 500099003, Bilthoven, the Netherlands, 2010.

16  
17  
18  
19 Dockery, D., Pope, C. A., Xu, X., Spengler, J., Ware, J., Fay, M., Ferris, B., and Speizer, F.: An association between air pollution and mortality in six U.S. cities. *New Engl. J. Med.* 329,1753-1759, 1993.

20  
21  
22  
23 EU: Directive 2008/50/EC of the European Parliament and of the Council on ambient air quality and cleaner air for Europe, 2008.

24  
25  
26 EEA: Air pollution in Europe 1990-2004, EEA report 2/2007, Copenhagen, 2007.

27  
28 Erisman, J.W., Sutton, M.A., Galloway, J., Klimont, Z. and Winiwarter, W.: How a century of ammonia synthesis changed the world, *Nature Geoscience*, 1, 636-639, 2009.

29  
30

1 Frank, N.H.: Retained nitrate, hydrated sulfates, and carbonaceous mass in Federal Reference  
2 Method fine particulate matter for six eastern cities. *J. Air Waste Manage. Assoc.*, 56, 500-  
3 511, 2006.  
4  
5 Friedrich, R. :Improving and applying methods for the calculation of natural and biogenic  
6 emissions and assessment of impacts to the air quality. Final project activity report 2007.  
7 <http://natair.ier.uni-stuttgart.de/>  
8  
9 Harrison, R.M., Jones, A.M., and Lawrence, R.G.: A pragmatic mass closure model for air-  
10 borne particulate matter at urban background and roadside sites. *Atmos. Environ.* 37, 4927-  
11 4933, 2003.  
  
12 Hoek, G., Meliefste, K., Cyrys, J., Lewné, M., Bellander, T., Brauer, M., Fischer, P., Ge-  
13 hring, U., Heinrich, J., van Vliet, P., Brunekreef, B.: Spatial variability of fine particle con-  
14 centrations in three European areas, *Atmos. Environ.* 36, 4077-4088, 2002.  
  
15 Hoelzemann, J.; Schultz, M. G.; Brasseur, G. P.; Granier, C. and Simon, M.: Global Wildland  
16 Fire Emission Model (GWEM): Evaluating the use of global area burnt satellite data, *J. Geo-*  
17 *phys. Res.* 109, D14S04, 2004.  
18  
19 Hopke, P. K., Liu, C. and Rubin, D. B.:Multiple imputation for multivariate data with miss-  
20 ing and below-threshold measurements: time-series concentrations of pollutants in the arctic.  
21 *Biometrics* 57, 22–33, 2001.  
22  
23 Korcz, M., Fudała, J. and Kliś, C.: Estimation of wind blown dust emissions in Europe and its  
24 vicinity, *Atmos. Environ.* 43, 1410-1420, 2009.  
25  
26 Matthijsen, J., Jimmink, B., De Leeuw, F. and Smeets, W.: Attainability of PM<sub>2.5</sub> air quality  
27 standards, situation for the Netherlands in a European context. PBL report 500099015, Bil-  
28 thoven, the Netherlands, 2009.  
29  
30 Mazzei, F., D'Alessandro, A., Lucarelli, F., Nava, S., Prati, P., Valli, G. and Vecchi, R.):  
31 Characterization of particulate matter sources in an urban environment, *Sci. Total Environ.*  
32 41, 81-89, 2008.



1  
2 Meijer, E., van Velthoven, P., Brunner, D., Huntrieser, H. and Kelder, H.: Improvement and  
3 evaluation of the parameterization of nitrogen oxide production by lightning, *Phys. Chem.*  
4 *Earth*, 26, 557-583, 2001.  
5  
6 Mooibroek, D., Schaap, M., Weijers, E.P. and Hoogerbrugge, R.: Source apportionment and  
7 spatial variability of PM2.5 using measurements at five sites in the Netherlands, to be pub-  
8 lished.  
9  
10 Pope, C. A., Thun, M., Namboodiri, M., Dockery, D., Evans, J., Speizer, F., and Heath, C.:  
11 Particulate air pollution as a predictor of mortality in a prospective study of U.S. adults. *Am.*  
12 *J. Resp. Crit. Care*, 151, 669-674, 1995.  
13  
14 Putaud, J.-P., Raesa, F., Van Dingenen, R., Brüggemann, E., Facchini, M., Decesari, S., Fuzzi,  
15 S., Gehrig, R., Hueglin, C., Laj, P., Lorbeer, G., Maenhaut, W., Mihalopoulos, N., Muel-  
16 ler, K., Querol, X., Rodriguez, S., Schneider, J., Spindler, G., ten Brink, H., Tørseth, K., Wie-  
17 densöhler, A., 2004. A European aerosol phenomenology - 2: Chemical characteristics of  
18 particulate matter at kerbside, urban, rural and background locations in Europe, *Atmos. Envi-*  
19 *ron.*, 38, 2579–2595, 2004.  
20  
21 Putaud, J.-P., Van Dingenen, R., Alastuey, A., H. Bauer, Birmili, W. , Cyrus, J. , Flentje, H.,  
22 Fuzzi, S., Gehrig, R. , Hansson, H.C. , Harrison, R.M., Herrmann, H. , Hitznerberger, R. ,  
23 Hüglin, C., Jones, A.M., Kasper-Giebl, A., Kiss, G. , Kousa, A. , Kuhlbusch, T.A.J. ,  
24 Löschau, G. , Maenhaut, W., Molnar, A. , Moreno, T., Pekkanen, J. , Perrino, C. , Pitz, M. ,  
25 Puxbaum, H., Querol, X., Rodriguez, S. , Salma, I. , Schwarz, J. Smolik, J. Schneider, G.  
26 Spindler, H. ten Brink, Tursic, J. J., Viana, M., Wiedensöhler, A., Raes, F.: A European  
27 aerosol phenomenology - 3: Physical and chemical characteristics of particulate matter from  
28 60 rural, urban, and kerbside sites across Europe, *Atmos. Environ.* 44, 1308-1320, 2009.  
29  
30 Salvador, P., Arti, B., Querol, X., Alastuey, A., Costoya, M.: Characterisation of local and  
31 external contributions of atmospheric particulate matter at a background coastal site. *Atmos.*  
32 *Environ.* 41, 1–17, 2007.  
33

- 1 Schaap, M. and Denier van der Gon, H.A.C.: On the variability of Black Smoke and carbona-  
2 ceous aerosols in the Netherlands, *Atmos. Environ.*, 41, 5908-5920, 2007.
- 3
- 4 Schaap, M., Manders, A.M.M., Hendriks, E.C.J., Clossen, J.M., Segers, A.J.S., Denier van  
5 der Gon, H.A.C., Jozwicka, M., Sauter, F.J., Velders, G.J.M., Matthijsen, J., Builtjes, P.J.H.:  
6 Regional modelling of particulate matter for the Netherlands, PBL Report 500099006, Bilt-  
7 hoven, the Netherlands, 2009.
- 8
- 9 Schaap, M., Spindler, G., Schulz, M., Acker, K., Maenhaut, W., Berner, A., Wieprecht, W.,  
10 Streit, N., Mueller, K., Brüggemann, E., Putaud, J.-P., Puxbaum, H., Baltensperger, U., ten  
11 Brink H. M.: Artefacts in the sampling of nitrate studied in the "INTERCOMP" campaigns of  
12 EUROTRAC-AEROSOL, *Atmos. Environ.* 38, 6487-6496, 2004.
- 13
- 14 Schaap, M., Weijers, E.P., Mooibroek, D., Nguyen, L. and Hoogerbrugge, R.: Composition  
15 and origin of particulate matter in the Netherlands, PBL report 500099007, Bilthoven, the  
16 Netherlands, 2010.
- 17
- 18 Sillanpää, M., Hillama, R., Saarikoski, S., Frey, A., Pennanen, A., Makkonen, U., Spolnik, Z.,  
19 Van Grieken, R., Braniš, M., Brunekreef, B., Chalbot, M.-C., Kuhlbusch, Th., Sunyer, J.,  
20 Kerminen, V.-M., Kulmala M. and Salonen R. O.: Chemical composition and mass closure of  
21 particulate matter at six urban sites in Europe, *Atmos. Environ.*, 40, 212-223, 2006.
- 22
- 23 Simpson D., Winiwarter W., Börjesson G., Cinderby S., Ferreiro A., Guenther A., Hewitt N.,  
24 Janson R., Khalil M.A.K., Owen S., Pierce T., Puxbaum H., Shearer M., Skiba U., Stein-  
25 brecher R., Tarrason L. and Öquist, M.G.: Inventorying emissions from nature in Europe. *J.*  
26 *Geophys. Res.* 104, 8113-8152, 1999.
- 27
- 28 Simpson, D., Yttri, K.E., Klimont, Z., Kupiainen, K., Caseiro, A., Gelencsér, A., Pio, C.A.,  
29 Puxbaum, H., Legrand, M.: Modeling carbonaceous aerosol over Europe: Analysis of the  
30 CARBOSOL and EMEP EC/OC campaigns, *J. Geophys. Res.-Atmos.*, 112, D23S14, 2007.
- 31
- 32 ~~Speer, R.E., Edney, E.O., Kleindienst, T.E.: Impact of organic compounds on the concentra-~~  
33 ~~tions of liquid water in ambient PM<sub>2.5</sub>, *J. Aerosol Sci.*, 34, 63-77, 2003.~~

1  
2 Ten Brink, H.M., Weijers, E.P., Van Arkel, F. Th., De Jonge, D.: Carbon (EC/OC) concentra-  
3 tions as derived from routine PM measurements in the Netherlands', PBL Report 500099005,  
4 Bilthoven, the Netherlands, 2009.  
5  
6 Ten Brink, H.M., Weijers, E.P., Röckmann, T. and Dusek, U.: <sup>14</sup>C analysis of filter samples  
7 for source apportionment of PM in the Netherlands, ECN Report E--10-005, 2010.  
8  
9 Ten Brink, H.M., Maenhaut, W., Hitzenberger, R., Gnauk, T., Spindler, G., Even, A., Chi, X.,  
10 Bauer, H., Puxbaum, H., Putaud, J.-P., Tursic J. and Berner A.: INTERCOMP2000: the  
11 comparability of methods in use in Europe for measuring the carbon content of aerosol. *At-*  
12 *mos. Environ.*, 38, 6507–6519, 2004.  
13  
14  
15 Van der Hoek, K.W.: Estimating ammonia emission factors in Europe: Summary of the work  
16 of the UNECE ammonia expert panel, *Atmos. Environ.*, 32, 315-316, 1998.  
17  
18 Viana, M., Kuhlbusch, T.A.J., Querol, X, Alastuey, A., Harrison, R.M., Hopke, P.K., Wini-  
19 warter, W., Vallius, M., Szidat, S., Prévôt, A.S.H., Hueglin, C., Bloemen, H., Wählin, P.,  
20 Vecchi, R., Miranda, A.I., Kasper-Giebl, A., Maenhaut W., and Hitzenberger, R.: Source ap-  
21 portionment of particulate matter in Europe: A review of methods and results, *J. Aerosol Sci.*  
22 39, 827-849, 2008.  
23  
24 Visser et al., 2001 Visser, H., Buringh, E., van Breugel, P.B., 2001. Composition and origin  
25 of airborne particulate matter in The Netherlands. National Institute for Public Health and the  
26 Environment, RIVM. The Netherlands, Report 650010029. 104pp.  
27  
28  
29 Weijers, E.P., Sahan, E., Brink, H.M. ten, Schaap, M., Matthijsen, J., Otjes, R.P., van Arkel,  
30 F. van (2010): Contribution of secondary inorganic aerosols to PM<sub>10</sub> and PM<sub>2.5</sub> in the Neth-  
31 erlands; measurement and modeling results. PBL report 500099006, Bilthoven, 2010.  
32  
33  
34

- Met opmaak: Nederlands (standaard)
- Met opmaak: Lettertype: (Standaard) Times New Roman, 12 pt, Engels (V.S.)
- Met opmaak: Engels (V.S.)
- Met opmaak: Lettertype: (Standaard) Times New Roman, 12 pt, Engels (V.S.)
- Met opmaak: Engels (V.S.)
- Met opmaak: Lettertype: (Standaard) Times New Roman, 12 pt, Engels (V.S.)
- Met opmaak: Engels (V.S.)
- Met opmaak: Lettertype: (Standaard) Times New Roman, 12 pt, Engels (V.S.)
- Met opmaak: Engels (V.S.)
- Met opmaak: Lettertype: (Standaard) Times New Roman, 12 pt, Engels (V.S.)
- Met opmaak: Engels (V.S.)
- Met opmaak: Lettertype: (Standaard) Times New Roman, 12 pt, Engels (V.S.)
- Met opmaak: Lettertype: (Standaard) Times New Roman, 12 pt
- Met opmaak: Lettertype: (Standaard) Times New Roman, 12 pt
- Met opmaak: Lettertype: (Standaard) Times New Roman, 12 pt
- Met opmaak: Lettertype: (Standaard) Times New Roman, 12 pt
- Met opmaak: Lettertype: (Standaard) Times New Roman, 12 pt
- Met opmaak: Lettertype: (Standaard) Times New Roman, 12 pt, Niet Vet
- Met opmaak: Lettertype: (Standaard) Times New Roman, 12 pt
- Met opmaak: Lettertype: (Standaard) Times New Roman, 12 pt, Engels (V.S.)

1 Table 1. Properties of the measurement locations

<i>Location</i>	<i>Characteristics</i>
Cabauw	Background location in the centre of the Netherlands Rural surrounding Some agricultural emissions in the direct surroundings Urban areas in 20-40 km distance
Hellendoorn	Background location in the east of the Netherlands Rural surrounding No known anthropogenic emissions nearby
Vredepeel	Background location in the southeast of the Netherlands Rural surrounding Intensive live stock (pigs) breeding nearby
Schiedam	Urban background location in the west of the Netherlands Part of Rotterdam urban agglomeration (appr. 600,000 inhabitants) Residential areas surrounded by motorways Close (<5 km) to industrial activity (refineries, oil-industry) and harbour
Rotterdam	Traffic location ('kerbside') in the west of the Netherlands Part of the Rotterdam agglomeration Less than 5 m from the kerb; traffic junction at 80 m <u>with traffic lights.</u> Heavy traffic.

1 | Table 2 Average concentrations with standard deviations of the main components as measured at the five sites

number of samples	Hellendoorn		Vredepeel		Cabauw		Schiedam		Rotterdam	
	38		35		26		33		41	
PM <sub>10</sub>	17,7 ±	6,4	21,2 ±	7,3	23,6 ±	11,9	21,8 ±	9,6	26,3 ±	11,6
NO <sub>3</sub>	3,6 ±	2,1	4,3 ±	2,2	5,4 ±	3,7	4,0 ±	3,6	4,8 ±	4,2
SO <sub>4</sub>	2,3 ±	1,0	2,5 ±	1,1	3,0 ±	1,4	3,1 ±	1,6	3,0 ±	1,6
NH <sub>4</sub>	1,2 ±	0,7	1,5 ±	1,1	2,0 ±	1,7	1,2 ±	1,6	1,5 ±	1,7
EC	1,7 ±	1,0	2,0 ±	0,9	2,1 ±	1,0	2,0 ±	1,3	3,7 ±	1,5
OC	1,6 ±	1,0	1,8 ±	0,7	1,6 ±	0,9	1,8 ±	1,3	2,3 ±	1,0
Cl	0,6 ±	0,9	1,1 ±	1,2	0,4 ±	0,4	1,2 ±	1,3	0,9 ±	0,9
Na	0,6 ±	0,6	0,9 ±	0,8	0,5 ±	0,4	1,1 ±	0,8	0,9 ±	0,6
total metals	1,4 ±	0,8	1,7 ±	0,7	2,0 ±	1,2	2,0 ±	1,0	2,9 ±	1,1
not-analysed	4,6 ±	3,5	5,5 ±	2,5	6,7 ±	4,5	5,3 ±	2,5	6,4 ±	3,3
PM <sub>2.5</sub>	12,5 ±	5,4	14,5 ±	6,3	17,5 ±	11,0	13,0 ±	7,6	16,4 ±	10,0
NO <sub>3</sub>	2,6 ±	1,7	3,2 ±	2,3	4,1 ±	3,7	2,5 ±	2,5	3,0 ±	3,0
SO <sub>4</sub>	2,1 ±	0,9	2,2 ±	1,1	2,7 ±	1,4	2,5 ±	1,3	2,4 ±	1,1
NH <sub>4</sub>	1,3 ±	0,9	1,3 ±	1,0	1,6 ±	1,4	1,1 ±	1,2	1,4 ±	1,5
EC	1,5 ±	0,8	1,4 ±	0,7	1,7 ±	0,9	1,9 ±	1,0	3,2 ±	1,4
OC	1,4 ±	0,7	1,2 ±	0,6	1,3 ±	0,6	1,2 ±	0,7	1,5 ±	0,8
Cl	0,2 ±	0,2	0,3 ±	0,4	0,2 ±	0,1	0,2 ±	0,2	0,2 ±	0,2
Na	0,2 ±	0,2	0,3 ±	0,3	0,1 ±	0,1	0,3 ±	0,2	0,3 ±	0,3
total metals	0,7 ±	0,4	0,7 ±	0,2	0,8 ±	0,3	0,9 ±	0,3	1,0 ±	0,3
not-analysed	2,6 ±	2,0	3,8 ±	2,7	4,9 ±	5,7	2,4 ±	2,3	3,3 ±	3,6
PM <sub>10-2.5</sub>	5,1 ±	2,1	6,8 ±	2,9	6,1 ±	2,5	8,8 ±	3,6	9,9 ±	3,4
NO <sub>3</sub>	1,0 ±	0,9	1,1 ±	1,0	1,3 ±	2,7	1,5 ±	1,6	1,8 ±	1,7
SO <sub>4</sub>	0,2 ±	0,4	0,3 ±	0,5	0,2 ±	1,1	0,6 ±	0,8	0,6 ±	1,0
NH <sub>4</sub>	-0,1 ±	0,6	0,2 ±	0,4	0,4 ±	1,0	0,1 ±	0,6	0,1 ±	0,5
EC	0,2 ±	0,9	0,6 ±	0,7	0,4 ±	0,3	0,2 ±	0,9	0,4 ±	0,6
OC	0,2 ±	0,8	0,7 ±	0,6	0,3 ±	0,4	0,5 ±	1,0	0,8 ±	0,6
Cl	0,4 ±	0,7	0,7 ±	0,9	0,3 ±	0,3	1,0 ±	1,2	0,7 ±	0,7
Na	0,4 ±	0,5	0,6 ±	0,5	0,4 ±	0,3	0,8 ±	0,6	0,6 ±	0,5
total metals	0,7 ±	0,6	0,9 ±	0,5	1,2 ±	0,9	1,2 ±	0,8	1,9 ±	0,9
not-analysed	2,1 ±	2,6	1,6 ±	2,5	1,7 ±	5,0	2,9 ±	1,6	3,0 ±	2,3

1 Table 3 Absolute and percentual change in mass concentrtrion after applying converson  
 2 formula

3

	<b>PM<sub>10</sub></b>		<b>PM<sub>2.5</sub></b>		<b>PM<sub>10-2.5</sub></b>	
	rural	urban	rural	urban	rural	urban
<b>associated oxygen</b>						
increase (µg/m <sup>3</sup> )	0,9	1,2	0,3	0,4	0,5	0,8
%-decrease of 'NA'	4%	5%	2%	3%	9%	8%
<b>sea salt</b>						
increase (µg/m <sup>3</sup> )	0,8	1,3	0,3	0,4	0,6	0,8
%-decrease of 'NA'	4%	5%	2%	3%	10%	9%
<b>nss sulphate</b>						
increase (µg/m <sup>3</sup> )	-0,2	-0,3	-0,1	-0,1	-0,1	-0,2
%-decrease of 'NA'	-1%	-1%	0%	0%	-2%	-2%
<b>total carb. material</b>						
increase (µg/m <sup>3</sup> )	1,1	1,5	0,9	1,2	0,3	0,3
%-decrease of 'NA'	5%	6%	6%	8%	4%	3%

4

5

1 Table 4. Concentrations of the constituents as determined in the PM<sub>10</sub> and PM<sub>2.5</sub> data set  
 2 for the five BOP-locations

3

	- - Hellendoorn		- - Vredepeel		- - Cabauw		- - Schiedam		- - Rotterdam	
number of samples	38 - -		35 - -		26 - -		33 - -		41 - -	
PM <sub>10</sub>	17,7 ±	6,4	21,2 ±	7,3	23,6 ±	11,9	21,8 ±	9,6	26,3 ±	11,6
SLA	7,0 ±	3,4	8,0 ±	4,2	10,2 ±	6,4	8,0 ±	6,6	9,1 ±	7,3
TCM	4,3 ±	2,4	5,0 ±	2,0	5,0 ±	2,3	4,9 ±	2,5	7,7 ±	3,1
SS	2,0 ±	2,0	3,0 ±	2,5	1,6 ±	1,2	3,7 ±	2,7	2,9 ±	2,1
MD	1,2 ±	1,4	1,2 ±	1,0	2,1 ±	1,9	1,6 ±	1,3	1,8 ±	1,3
metals	0,8 ±	0,3	1,2 ±	0,4	1,0 ±	0,4	1,4 ±	0,5	2,4 ±	0,8
unknown	2,3 ±	3,5	2,7 ±	2,0	3,7 ±	4,2	2,1 ±	2,7	2,3 ±	2,8
PM <sub>2.5</sub>	12,5 ±	5,4	14,5 ±	6,3	17,5 ±	11,0	13,0 ±	7,6	16,4 ±	10,0
SLA	6,0 ±	2,8	6,6 ±	4,1	8,4 ±	6,1	6,0 ±	4,6	6,8 ±	5,1
TCM	3,8 ±	1,9	3,3 ±	1,5	3,9 ±	2,0	4,0 ±	2,2	6,1 ±	2,7
SS	0,6 ±	0,8	1,0 ±	0,9	0,5 ±	0,3	1,0 ±	0,8	0,9 ±	1,0
MD	0,5 ±	0,4	0,5 ±	0,3	0,7 ±	0,4	0,6 ±	0,5	0,6 ±	0,3
metals	0,4 ±	0,2	0,6 ±	0,1	0,5 ±	0,2	0,6 ±	0,2	0,8 ±	0,3
unknown	1,2 ±	1,7	2,5 ±	2,6	3,5 ±	5,5	0,7 ±	2,2	1,2 ±	3,5
PM <sub>10-2.5</sub>	5,1 ±	2,1	6,8 ±	2,9	6,1 ±	2,5	8,8 ±	3,6	9,9 ±	3,4
SLA	1,0 ±	1,5	1,4 ±	1,6	1,8 ±	4,5	2,0 ±	2,8	2,3 ±	2,9
TCM	0,5 ±	2,0	1,7 ±	1,4	1,1 ±	0,9	0,9 ±	0,9	1,6 ±	1,1
SS	1,4 ±	1,5	2,0 ±	1,7	1,2 ±	0,9	2,7 ±	2,0	2,0 ±	1,5
MD	0,7 ±	1,0	0,8 ±	0,7	1,4 ±	1,5	1,0 ±	1,2	1,2 ±	1,0
metals	0,4 ±	0,2	0,6 ±	0,3	0,5 ±	0,3	0,8 ±	0,4	1,6 ±	0,7
unknown	1,1 ±	3,0	0,2 ±	2,2	0,2 ±	5,0	1,4 ±	2,0	1,1 ±	2,3

4

5

1

	Hellendoorn			Vredepeel			Cabauw			Schiedam			Rotterdam		
number of samples	38			35			26			33			41		
	conc	sd	%	conc	sd	%	conc	sd	%	conc	sd	%	conc	sd	%
PM10	17.7 ±	6.4	-	21.2 ±	7.3	-	23.6 ±	11.9	-	21.8 ±	9.6	-	26.3 ±	11.6	-
SIA	7.0 ±	3.4	40	8.0 ±	4.2	38	10.2 ±	6.4	43	8.0 ±	6.6	37	9.1 ±	7.3	35
TCM	4.3 ±	2.4	28	5.0 ±	2.0	27	5.0 ±	2.3	24	4.9 ±	2.5	26	7.7 ±	3.1	32
SS	2.0 ±	2.0	11	3.0 ±	2.5	14	1.6 ±	1.2	7	3.7 ±	2.7	17	2.9 ±	2.1	11
MD	1.2 ±	1.4	7	1.2 ±	1.0	6	2.1 ±	1.9	9	1.6 ±	1.3	7	1.8 ±	1.3	7
metals	0.8 ±	0.3	5	1.2 ±	0.4	6	1.0 ±	0.4	4	1.4 ±	0.5	6	2.4 ±	0.8	9
unknown	2.3 ±	3.5	9	2.7 ±	2.0	10	3.7 ±	4.2	13	2.1 ±	2.7	7	2.3 ±	2.8	6
PM2.5	12.5 ±	5.4	-	14.5 ±	6.3	-	17.5 ±	11.0	-	13.0 ±	7.6	-	16.4 ±	10.0	-
SIA	6.0 ±	2.8	48	6.6 ±	4.1	46	8.4 ±	6.1	48	6.0 ±	4.6	46	6.8 ±	5.1	42
TCM	3.8 ±	1.9	35	3.3 ±	1.5	28	3.9 ±	2.0	26	4.0 ±	2.2	36	6.1 ±	2.7	41
SS	0.6 ±	0.8	5	1.0 ±	0.9	7	0.5 ±	0.3	3	1.0 ±	0.8	8	0.9 ±	1.0	5
MD	0.5 ±	0.4	4	0.5 ±	0.3	3	0.7 ±	0.4	4	0.6 ±	0.5	5	0.6 ±	0.3	4
metals	0.4 ±	0.2	4	0.6 ±	0.1	4	0.5 ±	0.2	3	0.6 ±	0.2	5	0.8 ±	0.3	5
unknown	1.2 ±	1.7	5	2.5 ±	2.6	13	3.5 ±	5.5	16	0.7 ±	2.2	1	1.2 ±	3.5	3
PM10-2.5	5.1 ±	2.1	-	6.8 ±	2.9	-	6.1 ±	2.5	-	8.8 ±	3.6	-	9.9 ±	3.4	-
SIA	1.0 ±	1.5	14	1.4 ±	1.6	21	1.8 ±	4.5	29	2.0 ±	2.8	23	2.3 ±	2.9	23
TCM	0.5 ±	2.0	9	1.7 ±	1.4	25	1.1 ±	0.9	18	0.9 ±	0.9	10	1.6 ±	1.1	16
SS	1.4 ±	1.5	28	2.0 ±	1.7	30	1.2 ±	0.9	19	2.7 ±	2.0	30	2.0 ±	1.5	21
MD	0.7 ±	1.0	14	0.8 ±	0.7	11	1.4 ±	1.5	23	1.0 ±	1.2	11	1.2 ±	1.0	12
metals	0.4 ±	0.2	8	0.6 ±	0.3	9	0.5 ±	0.3	8	0.8 ±	0.4	9	1.6 ±	0.7	16
unknown	1.1 ±	3.0	21	0.2 ±	2.2	4	0.2 ±	5.0	3	1.4 ±	2.0	16	1.1 ±	2.3	12

2

3



1

2 Table 5 Illustration of natural contribution (in %) for all components in the case of Hel-  
 3 lendoorn. For each component the relative contribution to PM<sub>2.5</sub> and PM<sub>10</sub> is listed as  
 4 well as the estimated natural contribution. The two columns on the right provide the re-  
 5 sulting natural contribution to PM<sub>10</sub> and PM<sub>2.5</sub> for each component for a high and low  
 6 estimate of the natural OC content. All numbers are percentages.

7

<i>Comp</i>	<i>PM<sub>10</sub></i>	<i>PM<sub>2.5</sub></i>	<i>Natural</i>	<i>Hellendoorn</i>	<i>Hellendoorn</i>
	<i>Contrib.</i>	<i>Contrib.</i>	<i>%</i>	<i>PM<sub>10</sub></i>	<i>PM<sub>2.5</sub></i>
				Low / high	Low / high
NO3	21	21	<u>0-5</u>	<u>0-1</u>	<u>0-1</u>
SO4	13	17	5	1	1
NH4	7	10	10	0.5	1
MD	7	4	<u>1-20</u>	<u>1-5</u>	<u>0.5-1</u>
SS	11.5	5	100	11.5	5
Metals	5	4	0	0	0
EC*1.3	12	15	0	0	0
OC*1.3	12	14	25 / 50	3 / 6	3.5 / 7
Sum known				<u>16.58 / 212</u>	<u>112.5 / 15.56</u>
Unknown	13	10	#	2.5 / 3	1.5 / 1.5
Total				<u>1924 / 245</u>	<u>1413 / 187</u>

8 # Natural contribution is assumed to be the same as the total natural percentage of the explained  
 9 mass.

10

- 1
- 2
- 3 Figure 1. Localisation of the five BOP measurement sites in the Netherlands
- 4
- 5 Figure 2. Absolute concentrations at the five sites
- 6
- 7 Figure 3 Chemical distributions measured at the urban background site Schiedam
- 8
- 9 Figure 4 Mass closures for the different fractions estimated at the urban background site
- 10 Schiedam
- 11
- 12 Figure 5 Average chemical distributions in data subsets with high and low PM concen-
- 13 trations. Above:  $PM_{10} > 30 \mu\text{g}/\text{m}^3$  versus  $PM \leq 30 \mu\text{g}/\text{m}^3$ ; below  $PM_{2.5} \leq 25 \mu\text{g}/\text{m}^3$  ver-
- 14 sus  $PM_{2.5} > 25 \mu\text{g}/\text{m}^3$ . Averages over the five BOP stations.
- 15
- 16 Figure 6 Natural and anthropogenic contributions to  $PM_{10}$  and  $PM_{2.5}$  as estimated from
- 17 the BOP data set.