

IUPAC Subcommittee on Gas Kinetic Data Evaluation – Data Sheet V.A1.19

Datasheets can be downloaded for personal use only and must not be retransmitted or disseminated either electronically or in hardcopy without explicit written permission. The citation for this datasheet is: IUPAC Subcommittee for Gas Kinetic Data Evaluation, <http://www.iupac-kinetic.ch.cam.ac.uk>

This datasheet last evaluated: April 2010; last change to preferred values: April 2010.

CH₃C(O)OH + ice

Experimental data

Parameter	Temp./K	Reference	Technique/ Comments
<i>Partition coefficients: K_{linC}</i>			
$1.9 \times 10^{-11} \exp(6800/T)$	220-245	Sokolov and Abbatt, 2002	CWFT-MS (a)
320	193	Picaud et al., 2005	CWFT-MS (b)
258	203		
153	213		
73	223		
$2.3 \times 10^{-10} \exp(6530/T)$	197-227	von Hessberg et al., 2007	CWFT-MS (c)
$1.9 \times 10^{-12} \exp(7400/T)$	213-243	Kerbrat et al., 2010	PBFT-CIMS (d)
1040	218	Symington et al, 2010	CWFT-MS (e)
497	228		
179	238		

Comments

(a) Ice film made by freezing distilled water. Uptake was found to be reversible and equilibrium surface coverages were calculated using the geometric ice surface area. Equilibrium uptake of CH₃C(O)OH to ice at various temperatures was analysed using the Langmuir isotherm. The values for K_{linC} at individual temperatures given in the Table uses the reported values of $K_{LangP}(T)$ and $N_{max}(T)$. No errors were reported. The temperature dependent expression of K_{linC} was derived by fitting to these five data points. A value of $\Delta H_{ads} = -73 \pm 12$ kJ mol⁻¹ was reported. CH₃C(O)OH concentration varied between $\approx 4 \times 10^{10}$ and 2×10^{13} molecule cm⁻³, with between 2 and 20 % present as dimers.

(b) Ice film, 30-80 μ m thick was made by freezing distilled water. Uptake was found to be reversible and equilibrium surface coverages were calculated using the geometric ice surface area. Equilibrium uptake of CH₃C(O)OH to ice at various temperatures was analysed using the BET isotherm to derive a value of $\Delta H_{ads} = -33.5 \pm 4.2$ kJmol⁻¹. The parameterised BET isotherms were used to calculate values of K_{linC} at the four temperatures where reversible uptake was observed. The authors suggest that most of the acetic acid was in the form of dimers in their experiments.

(c) Ice film made by freezing distilled water. Uptake was found to be reversible and equilibrium surface coverages were calculated using the geometric ice surface area. Values of $N_{max} = 2.4 \times 10^{14}$ molecule cm⁻² (independent of temperature) and $\Delta H_{ads} = -55 \pm 9$ kJ mol⁻¹ were reported. CH₃C(O)OH concentration varied between 3×10^9 and 2×10^{11} molecule cm⁻³ and the fraction of dimers present was calculated to be less than 15 % for all temperatures and concentrations except for 197 K and $[HC(O)OH] > 2 \times 10^{10}$ molecule cm⁻³.

(d) Packed ice bed flow tube at atmospheric pressure. Partition coefficients derived from analysis of breakthrough curves and using the geometric ice surface area.

(e) Ice film made by freezing distilled water. Uptake was found to be reversible and equilibrium surface coverages were calculated using the geometric ice surface area. Equilibrium uptake of CH₃C(O)OH to ice at various temperatures was analysed using the Langmuir isotherm. Table uses the reported values of $K_{\text{LangP}}(T)$ and $N_{\text{max}}(T)$. The temperature dependent expression of K_{linC} was derived by fitting to these three data points. A value of $\Delta H_{\text{ads}} = -52.6 \text{ kJ mol}^{-1}$, was reported. CH₃C(O)OH concentration varied between $\approx 3 \times 10^{10}$ and $4 \times 10^{12} \text{ molecule cm}^{-3}$ and were corrected for the presence of dimers.

Preferred Values

Parameter	Value	T/K
$K_{\text{linC}} / \text{cm}$	$1.5 \times 10^{-14} \exp(8500/T)$	195 - 240
$N_{\text{max}} / \text{molecule cm}^{-2}$	2.5×10^{14}	
<i>Reliability</i>		
$\Delta(E/R) / \text{K}$	± 1000	195 - 240
$\Delta \log N_{\text{max}}$	0.1	

Comments on Preferred Values

There are five experimental studies of the reversible uptake of CH₃C(O)OH to pure ice surfaces, the results differing somewhat, possibly due to the presence of varying amounts of the acetic acid dimer. Sokolov et al. (2002), von Hessberg et al. (2007), Kerbrat et al. (2010) and Symington et al. (2010) suggest that the fraction of dimers present in their samples was of the order of percent, whereas Picaud et al. (2005) calculated that almost 100 % of their sample was dimerised. It is not clear if this assumption was partially responsible for the much lower partitioning coefficients found by Picaud et al. The published data of Sokolov et al. (2002), von Hessberg et al. (2007) and Kerbrat et al., (2010) are in good agreement and were used to derive the preferred expression, which indicates an enthalpy of adsorption of $70 (\pm 10) \text{ kJ mol}^{-1}$. The values of N_{max} returned by the various studies is variable, probably due to extrapolations from low coverages and also due to lateral interactions at high coverages, which make this parameter generally difficult to access experimentally (Jedlovszky et al., 2006).


Theoretical investigations (Compoint et al. 2002; Picaud et al. 2005) have shown that the hydroxyl and carbonyl groups of acetic acid are bound to two surface water molecules, with the CH₃ group directed away from the ice surface. This would tend to suggest that acetic acid dimers, for which OH and CO bonds are no longer available, would undergo significantly weaker interaction with the ice surface.

References

- Compoint, M., Toubin, C., Picaud, S., Hoang, P. N. M. and Girardet, C.: Chem. Phys. Lett. 365, 1-7, 2002.
- Kerbrat, M., Huthwelker, T., Bartels-Rausch, T., Gäggeler, H. W., and Ammann, M., Phys. Chem. Chem. Phys., in press, 2010.
- Jedlovszky, P., Partay, L., Hoang, P. N. M., Picaud, S., von Hessberg, P. and Crowley, J. N.: J. Am. Chem. Soc. 128, 15300-15309, 2006.
- Picaud, S., Hoang, P. N. M., Peybernes, N., Le Calve, S. and Mirabel, P.: J. Chem. Phys. 122, article 194707, 2005.
- Sokolov, O. and Abbatt, J. P. D.: J. Phys. Chem. 106, 775-782, 2002.

Symington, A. PhD thesis, University of Cambridge, 2010.

von Hessberg, P., Pouvesle, N., Winkler, A. K., Schuster, G. and Crowley, J. N.: Phys. Chem. Chem. Phys. 10, 2345-2355, 2008.



Partitioning coefficients (K_{linC}) for acetic acid uptake to ice.