

# **Climatology and comparison of ozone from ENVISAT/GOMOS and SHADOZ/balloon-sonde observations in the southern tropics**

**N. Mze<sup>1</sup>, A. Hauchecorne<sup>2</sup>, H. Bencherif<sup>1</sup>, F. Dalaudier<sup>2</sup> and J.-L. Bertaux<sup>2</sup>**

<sup>1</sup>Laboratoire de l'Atmosphère et des Cyclones, Université de la Réunion, CNRS-INSU, UMR 8105, St-Denis, Reunion Island, France

<sup>2</sup>Laboratoire Atmosphères, Milieux, Observations Spatiales, Université Versailles St-Quentin, CNRS-INSU, Verrières-le-Buisson, France

Correspondence to: N. Mze (nahoudha.mze@univ-reunion.fr)

## **Abstract**

In this paper, the stellar occultation instrument GOMOS is compared with ozonesondes from the SHADOZ network. We only used nighttime O<sub>3</sub> profiles and a requirement selection at 8 southern hemisphere stations. 7 years of GOMOS datasets (GOPR 6.0cf and IPF 5.0) and 11 years of balloon-sondes are used in this study. A monthly distribution of GOMOS O<sub>3</sub> mixing ratios is performed in the upper-troposphere and in the stratosphere (15-50 km). A comparison with SHADOZ is done in the altitude range from 15 km to 30 km.

In the 21-30km altitude range, a satisfactory agreement is observed between GOMOS and SHADOZ although some differences are observed depending on the station. The range for monthly differences is generally decreasing with increasing height and is within  $\pm 15\%$ . It is found that the agreement between GOMOS and SHADOZ degrades below  $\sim 20$  km. The median differences are nearly within  $\pm 5\%$  particularly above 23 km. But a large positive bias is found below 21 km compared to SHADOZ.

## 1 Introduction

The long-term evolution of stratospheric ozone concentrations depends not only on changes of many stratospheric constituents (including ozone-depleting substances (ODSs), greenhouse gases (GHGs), water vapor, and aerosols), but also on changes in the troposphere and in the stratosphere caused by natural variability and anthropogenic forcing (WMO, 2006). Besides, the decrease of ODSs and the associated context of ozone recovery make that stratospheric ozone is a thematic of importance for ongoing researches. Air enters into the stratosphere from the troposphere primarily in the ozone photochemical source region: the tropics (Shepherd et al., 2000). It is indeed important to follow and investigate the evolution of stratospheric ozone in the tropics. Ozone-sonde networks such as SHADOZ (Southern Hemisphere Additional Ozone-sondes) provide continuous and accurate measurements at different selected locations. On the other hand, satellites provide a global coverage. But these measurements have to be validated through comparisons with ground-based observations like balloon-sondes in order to give confidence to the results. Furthermore, the role of satellites for intercomparing ground-based instruments between them is also essential to detect possible station to station biases.

Global Ozone Monitoring by Occultation of Stars (GOMOS), on board the European satellite ENVISAT (ENVironmental SATellite), is the first instrument dedicated to the study of atmosphere by the technique of stellar occultation (Bertaux et al., 2004). The instrument has the advantage of self calibrating method and a good vertical resolution with a global coverage (Kyrölä et al., 2004). The key objective of GOMOS is the long-term ozone monitoring with a high vertical resolution, a high accuracy at global coverage and consequently assessment of ozone trends in the stratosphere (Bertaux et al., 2000).

The purpose of this paper is to give an overview of GOMOS ozone climatology in the southern tropics and a comparison with ground-based ozone-sondes at different stations operating in the framework of the SHADOZ network. Contrary to the northern hemisphere, ground based stations in the southern hemisphere are very sparse. Moreover, the tropical stratosphere is a region where ozone is created by sunlight and where significant changes are expected to occur. The tropical stratosphere is however a region where it is difficult to measure ozone by satellite experiments because of increased Rayleigh atmospheric attenuation, high altitude clouds, low temperature, high humidity and dense aerosols (Borchi et al., 2007). It is therefore essential to compare the few stations operating in the SHADOZ project with the performances of the stellar occultation instrument GOMOS. The study

1 focuses on the validation of the GOMOS level 2 data processing version GOPR 6.0cf (and  
2 IPF 5.0). In that regard, we use GOMOS data from August 2002 to December 2008 and focus  
3 on the southern hemisphere sites based on SHADOZ ozonesonde network (Thompson et al.,  
4 2003). SHADOZ datasets are well known and increasingly used for climatological studies  
5 (Lamsal et al., 2004, McPeters et al., 2007), variability studies (Logan et al., 2003; Witte et  
6 al., 2008) or comparison studies (Liu et al., 2006, Sivakumar et al., 2007), for case studies  
7 such as ozone isentropic transport in the lower stratosphere (Semane et al., 2006; Bencherif et  
8 al., 2007), and for trend analyses (Clain et al., 2009).

9 The first results for multi-year GOMOS ozone climatology were performed by Kyrölä et al.  
10 (2007). They concluded that, even if GOMOS data includes gaps due to instrumental  
11 difficulties or nighttime requirement, the available time series allow one to investigate ozone  
12 variations in the stratosphere and in the mesosphere. Meijer et al. (2004) validated the ozone  
13 data GOPR 5.4b derived from GOMOS observations under dark limb condition with  
14 correlative data based on balloon-sondes and ground based instruments. They reported an  
15 insignificant negative bias from 2.5% to 7.5% between 14 km and 64 km with standard  
16 deviations of 11-16% between 19 km and 64 km. Furthermore, this result was demonstrated to  
17 be independent of the star temperature, magnitude and the latitudinal region, except in polar  
18 regions in the altitude range from 35 km to 45 km where a slightly larger bias was observed.

19 Kyrölä et al. (2006) provided a nighttime stratospheric ozone climatology and a comparison  
20 of GOMOS 2003 data (GOPR 6.0a). In ~~Faet~~ fact, they compared GOMOS stratospheric ozone  
21 with the Fortuin-Kelder ozone climatology (Fortuin and Kelder, 1998) and found large  
22 differences in polar areas correlated with large increases of NO<sub>2</sub>. They also observed that  
23 GOMOS values are systematically larger in the upper stratosphere due to the diurnal variation  
24 of ozone above 45 km. They added that GOMOS finds a few percent less ozone than Fortuin-  
25 Kelder in the middle and lower stratosphere. On the opposite, they reported that in the  
26 equatorial areas GOMOS values are much lower than Fortuin-Kelder climatological values in  
27 the upper troposphere-lower stratosphere.

28 Moreover, GOMOS dark occultations were compared with ozone soundings at high-latitudes  
29 stations: Marambio (64.3°S, 56.7°W) and Sodankylä (67.4°N, 23.6°E) by Tamminen et al.  
30 (2006). They found a good agreement in the 15-30 km altitude range. The differences between  
31 the averages were within  $\pm 5\%$  for Marambio and somewhat worse for Sodankylä (up to -  
32 10%). In addition, a good agreement in the middle stratosphere between GOMOS and  
33 various balloon-borne instruments at mid- and high-latitudes stations was reported by Renard

1 et al. ~~(2006)~~ (2008). Liu et al. (2008) performed a comparison of GOMOS and MIPAS  
2 (Michelson Interferometer for Passive Atmospheric Sounding) with ozonesonde profiles at  
3 Beijing (39.48°N, 116.28°E). They observed a good agreement above 15 km. The differences  
4 between GOMOS and balloon-sonde are found to be positive and significant within  $\pm 10\%$   
5 above 15 km, particularly between 19 km and 30 km where biases are found below 5%. A  
6 cross validation was also performed with the ground based microwave radiometer SOMORA  
7 (Stratospheric Ozone Monitoring Radiometer) by Hocke et al. (2007). The relative differences  
8 were within 10% at altitudes below 45 km.

9 Comparisons were also performed versus satellite data. Verronen et al. (2005) compared  
10 nighttime ozone profiles from MIPAS and GOMOS (version 5.4b and 6.0a). They reported a  
11 good agreement between the two instruments. They agree within 10-15% in the middle  
12 atmosphere. Bracher et al. (2005) compared three ENVISAT instruments: MIPAS,  
13 SCIAMACHY (SCanning Imaging Absorption spectroMeter for Atmospheric CHartographY)  
14 and GOMOS (dark observations version 6.0a). Cross comparisons have shown an agreement  
15 for ozone within 15% in the 21-40km height range between the instruments. Dupuy et al.  
16 (2009) observed a good agreement in the stratosphere between GOMOS and ACE-FTS  
17 (Atmospheric Chemistry Experiment-Fourier Transform Spectrometer). The median relative  
18 differences were within  $\pm 10\%$ .

19  
20 This work investigates 7 years of GOMOS measurements in order to evaluate its potential for  
21 establishing a climatology of stratospheric ozone in comparison with ground-based  
22 observations at different tropical/subtropical locations in the southern hemisphere operated in  
23 the framework of the SHADOZ program. Section 2 describes data measurements and methods  
24 of analysis. Results are reported and discussed in Sect. 3. Summary is presented in Sect. 4.

## 2 Datasets and analysis

The Global Ozone Monitoring by Occultation of Stars (GOMOS) on board the European Space Agency's ENVISAT satellite was launched on 1<sup>st</sup> ~~march~~ March 2002. GOMOS is a medium resolution spectrometer covering the wavelength range from 250 nm to 950 nm. The four spectrometers of the instrument cover the spectral ranges: 248-387 nm, 387-693 nm, 750-776 nm and 915-956nm. This coverage allows monitoring O<sub>3</sub> and other species, i.e., NO<sub>2</sub>, NO<sub>3</sub>, neutral density, aerosols, H<sub>2</sub>O and O<sub>2</sub> from the upper troposphere up to the mesosphere. The altitude sampling resolution is better than 1.7 km (Bertaux et al., 2000, 2004; Kyrölä et al., 2004).

The retrieval method from GOMOS occultations is based on the transmission spectra and divided into two processes: a spectral inversion allows retrieving the line densities and a vertical inversion permits to retrieve the vertical distribution of the local densities using the onion-peeling method (Bertaux et al., 2004, Kyrölä et al., 2004). The vertical resolution of the retrieved ozone profiles is 2 km below 30 km and then it increases linearly to 3 km up to 40 km and remains the same above, due to the inversion algorithm which is based on Tikhonov regularization (Sofieva et al., 2004; Kyrölä et al., 2006). The height range covered by GOMOS is typically from 15 km to 100 km. For more details about the instrument parameters, measurements characteristics and data processing the reader can refer to Bertaux et al. (2000, 2004) and Kyrölä et al. (2004, 2006) papers.

Ground based datasets used for the present study are obtained from balloon-soundes launched at 8 southern hemisphere stations. An overview is presented in Table 1 which summarizes, site by site, geographical and size-data characteristics. Figure 1 displays their geographical localisation on the map. Ozonesondes are used for measuring height profiles of ozone from ground to about 30 km (balloon burst) with a precision of about 5% (Thompson et al., 2003). The present study uses 11 years of ozone-radiosonde datasets from 1998 to 2008. All balloon-sonde profiles are available on the SHADOZ website: <http://croc.gsfc.nasa.gov/shadoz/>. Details about quality or sonde parameters are described by Johnson et al. (2002) and by Thompson et al. (2003, 2007).

With regard to ozone profiles derived from GOMOS occultations, level 2 data processing version 6.0cf is used. Reprocessed data are available from August 2002 to June 2006. And

1 data from July 2006 to December 2008 are processed by the operational version IPF 5.00.  
2 Level 2 data products were analyzed with the Basic Envisat Atmospheric Toolbox software  
3 (BEAT). It provides tools for ingesting, processing, and analyzing atmospheric remote sensing  
4 data (<http://www.stcorp.nl/beat/>).  
5

6 Moreover, it should be noted that only nighttime ozone profiles are used in the altitude range  
7 between 15 km and 50 km. Only occultations with solar zenith angle (SZA) larger than  $108^\circ$   
8 (dark limb limit) at the geolocation of the tangent point are usable. Daytime retrieval suffers  
9 from additional noise caused by the stray light from the Sun (Meijer et al., 2004; Kyrölä et al.,  
10 2006). In addition, we considered dark limb occultations with latitudinal and longitudinal  
11 differences of  $\pm 5^\circ$  and  $\pm 10^\circ$  over each station. With data product GPR 5.4b and GPR 6.0a,  
12 Meijer et al. (2004) and Bracher et al. (2005) used a quality filter and took into account only  
13 measurements with a reported error below 20%. But in the version 6.0cf of the algorithm,  
14 error is overestimated in the 25-40 km altitude range (Johanna Tamminen, personal  
15 communication 2009). We therefore included GOMOS measurements recorded with a  
16 reported error below 30% and only  $O_3$  concentrations between 0 and  $10^{19}$  mol/m<sup>3</sup> are taken  
17 into account.  
18

19 Furthermore, ozone mixing ratios are derived by dividing the GOMOS ozone number density  
20 by the atmospheric neutral density included in the data product and calculated from ECMWF  
21 fields (Kyrölä et al., 2006).  
22

23 ~~Additionally~~ Additionally, following the method of analysis used by Kyrölä et al. (2006), we  
24 used the median and the interquartile range (*iqr*) as a robust estimator and statistic tools on  
25 GOMOS profiles. The interquartile range is defined as:  $q_3 - q_1$  where  $q_3$  and  $q_1$  are the 75<sup>th</sup> and  
26 25<sup>th</sup> percentiles, respectively (or third and first quartile, respectively). In this way, the study of  
27 GOMOS ozone climatology and variability is less affected by outliers.  
28

29 Due to height limitation in balloon-sonde data, comparison between ground-based and  
30 satellite observations, as studied in this issue, can be performed within a limited altitude  
31 range, i.e., in the 15-30 km altitude range. In that regard, ozone profiles from GOMOS and

1 balloon-sonde have been interpolated into a 1 km vertical grid. In addiditon, all GOMOS and  
2 SHADOZ data have been monthly averaged. As mentioned above, balloon-sonde profiles are  
3 recorded from 1998 to 2008, while GOMOS observations cover the period from August 2002  
4 to December 2008. It should be noted that there is no GOMOS data in February, April and  
5 May 2005 due to technical problems on the pointing system that occurred on January 2005  
6 (Kyrölä et al., 2006). Figure 2 shows the monthly numbers of dark limb occultations and of  
7 balloon-sonde profiles for each station. For the most part the number of GOMOS occultations  
8 is larger than the number of ground based profiles. In fact, the number of observations used in  
9 this work is 2836 profiles from SHADOZ sites and 6708 profiles from GOMOS occultations.  
10 It can be seen from the figure that there are less GOMOS dark limb occultations by August in  
11 comparison with the other months of the year.

### 3 Results

The monthly distributions of ozone mixing ratio derived from GOMOS observations are illustrated in Fig. 3 for each station in the height range between 15 km and 50 km. The altitude range for the maximum of mixing ratio is between 29 km and 37 km. The maximum mixing ratios is in the range from 7.5 ppmv to 10.5 ppmv. Furthermore, Fig. 3 illustrates the semi-annual variation of ozone for equatorial stations (Nairobi, Natal, Java and Ascension) with maxima in February-March and September-October and the annual variation in tropical stations (Samoa, Fiji, Reunion and Irene) with a minimum during austral winter (June-July). A lack of data is generally observed in August between 15 km and 20 km. The mean altitude of the maximum mixing ratios for equatorial and tropical stations is ~31.3 km and ~32.2 km, respectively. The mean of maximum mixing ratios for equatorial and tropical stations is 9.8 ppmv and 8.9 ppmv, respectively. The range for maximum mixing ratios at the equatorial sites is from 9.1 ppmv to 10.5 ppmv, while the range at the tropical stations is from 7.5 ppmv to 10.1 ppmv. These results are in agreement with Kyrölä et al. (2006) who provided a nighttime stratospheric ozone climatology measured by GOMOS in 2003.

Figure 4 displays the monthly distributions of ozone variability in the 15-50 km altitude range as computed from GOMOS occultations above SHADOZ stations. Variability is defined as the ratio between the quartile deviation (interquartile range divided by 2) and the median, in percent:

$$V = \frac{iqr}{2 \cdot median} \times 100 \quad (1)$$

In the height range from 20 km to 50 km, the variability values vary between 3% and 10% over all the months and all the sites.

For all stations, the variability is larger during austral winter (5 to 10%) than during austral summer (less than 5%) in the 27-35km altitude range. Furthermore, Fig. 4 shows much larger variability in the altitude range between 15 km and 20 km heights regardless of seasons or sites. It displays strong deviations (>20%), partially due to larger error bars in the lower part of the profiles.



Taking into account the altitude range of overlapping between GOMOS and balloon-sonde ozone observations, their monthly climatological profiles are compared in the height range from 15 km to 30 km. This comparison can be used for two purposes: first to evaluate the quality of GOMOS data with SHADOZ profiles at different locations and second to inter-compare SHADOZ stations using the same space instrument assuming that if a bias is present in GOMOS profiles, it will not be location dependent. Figure 5 displays the relative percentage of differences with respect to monthly climatological values obtained from balloon-sonde observations. It is defined as:

$$\Delta O_3(z) = \frac{[O_3]_{GOMOS} - [O_3]_{SHADOZ}}{[O_3]_{SHADOZ}} \times 100 \quad (2)$$

where  $[O_3]_{GOMOS}$  represents the median values for GOMOS and  $[O_3]_{SHADOZ}$  represents the averaged values for SHADOZ.

One can observe from Fig. 5 that differences between the median ozone mixing ratios from GOMOS dark occultations and the mean of SHADOZ profiles depend on station (vary from station to station) and altitude.

It can be noted however that for all sites, GOMOS finds much larger values than SHADOZ for heights between 15 km and ~20 km. Indeed, GOMOS seems to overestimate ozone mixing ratios in this altitude range over all the studied sites whatever the season (> 50%). Above 20 km, a better agreement is found for each site. The range in percent for the differences between ~20 km and 30 km is generally decreasing with increasing height and is within  $\pm 15\%$  for almost all the stations. However it is found that over Nairobi and Irene ozone climatological values derived from GOMOS observations are less than the values obtained from balloon-sondes above 25 km and 22 km, respectively. At Nairobi SHADOZ is 2-10% larger than GOMOS and at Irene SHADOZ is 2-15% larger than GOMOS. On the opposite, GOMOS values obtained over Natal, Java and Ascension are larger than SHADOZ values, except during the period from September to December in the 22.5-30km altitude range (25-30 km for Java). The agreement between GOMOS and SHADOZ is the poorest at Ascension (within 20-45%) between ~20 km and 27 km for the period from April to June. GOMOS is larger than SHADOZ at Samoa, except in May, July and between September and November in the 26-30km heights range. At Fiji, GOMOS is larger than SHADOZ except during the period from May to September above 27 km. And GOMOS values are larger than SHADOZ

values at Reunion except in August in the 20-25 km heights range and between April to August above 27 km and in November and December above 29 km. Further studies are needed to understand the causes of the station dependent observed differences. On the average, GOMOS is ~5-10 % larger than SHADOZ between 20 km and 30 km and we will come back later to these point. Despite the observed differences,~~the agreement between GOMOS and SHADOZ in this altitude range can be considered as for almost all the months and the stations~~ there is a satisfactory agreement between GOMOS and SHADOZ in this altitude range.

The ~~global~~ median and mean over all monthly comparisons ~~differences~~ between GOMOS and SHADOZ ozone mixing ratios are presented in Fig. 6 for each station. The outcome from analyzing median and mean of the differences allows examining if the statistical distribution of the differences is well represented by a standard Gaussian distribution (mean and median should be quite similar) or if it is strongly influenced by outliers (mean and median are very different). We note that the mean values nearly follow the median ones. In the 21-30km altitude range GOMOS is generally larger than SHADOZ at all sites, except at Nairobi and Irene, where we find that SHADOZ is larger than GOMOS above 23 km and 21.5 km, respectively, confirming the analysis on monthly mean differences. The observed differences between GOMOS and SHADOZ may be partly caused by ~~due to~~ an underestimation of ozone by sondes below the ozone maximum (around 26 km in the tropics) due to the 50-60s time constant of electrochemical sondes, creating a shift up of 300 m in the altitude registration as proposed by Borchini et al. (2005) who found a 5% negative bias in SHADOZ profiles compared to SAOZ (Système d'Analyse par Observation Zénithale). Johnson et al. (2002) and Thompson et al. (2003) also suggested that errors on sonde measurements could be introduced by instrumental uncertainties (pump efficiency corrections, sensing solutions changes) at higher altitudes, i.e., at lower ambient pressure. Moreover, GOMOS errors contribute to discrepancies. As reported by Tamminen et al. (2010), the main sources of GOMOS errors are due to random effects: measurement noise and scintillations (10% around 15 km and 0.5% to 4% in the stratosphere, values correspond to nighttime measurements). And the largest part of the systematic error is due to imperfect aerosol modeling which impacts mainly the O<sub>3</sub> retrievals (others sources of systematic errors are due to uncertainties in cross sections of the trace gases and in the atmospheric temperature). Scintillation caused by air density irregularities is a nuisance for retrievals of atmospheric composition. In GOMOS retrievals,

the scintillation effect is corrected using scintillation measurements by the fast photometer (Sofieva et al., 2009). The remaining perturbations, due to the incomplete scintillation correction, are not negligible. It induces an error of 0.5–1.5% in ozone retrieval at altitudes 20–40 km; see recent studies of Sofieva et al. (2009) and Tamminen et al. (2010) for more details. The coarse spatial sampling of GOMOS could also be considered as a minor source of error. The displacement of the sondes from 0 to 30 km is most of the time smaller than 120 km (assuming a wind speed  $< 20\text{ms}^{-1}$  and an ascent time of 6000s,  $5\text{ ms}^{-1}$  ascent speed) which is small compared to the area of GOMOS observations around the station ( $\pm 5^\circ$ ,  $\pm 10^\circ$ ). The meteorology may represent a source of the differences between GOMOS and SHADOZ, but it does not have a significant effect.

On average, the percentage of differences between GOMOS and SHADOZ is nearly within  $\pm 10\%$  in the height range from 21 km to 23 km and decrease to nearly  $\pm 5\%$  above 23km, except for Ascension, where it reaches about 8%. The overall results of the comparison between 21 km and 30 km are summarized in Table 2. In the lower stratosphere (below 21 km), it comes that GOMOS systematically and increasingly overestimates ozone mixing ratios with decreasing height. This is in agreement with Meijer et al. (2004) who found an overestimation of GOMOS between 18 km and 21 km compared to ozonesondes in the tropical region using GPR v5.4b. And this is consistent with Borchers et al. (2007), who found an altitude limitation of GOMOS measurements at 22 km in the tropics. They compared satellites and sondes with SAOZ onboard long duration balloons during the southern hemisphere summer in 2003 and 2004. They observed a consistent agreement between GOMOS and SAOZ above 22 km in the stratosphere, showing no altitude shift and a small low ozone bias of GOMOS of 1–2.5%. And they reported a degradation of GOMOS performances in the stratosphere below 22 km. Above 21 km, we observe a good agreement between ozone measurements derived from GOMOS occultations and ground-based balloon-sonde, similar to results reported in previous studies at tropical station (Meijer et al., 2004). And these results in the tropics complete the good agreement observed at mid- and high-latitude stations between GOMOS and ozonesondes (Tamminen et al., 2006; Renard et al., 2006 2008; Liu et al., 2008). Besides, the usage of the presented ozone climatology could lead to the GOMOS primary objective, i.e. study and assessment of trends in the stratosphere. Furthermore, as reported by Bertaux et al. (2000), it could be essential for improving atmospheric models for prediction of future changes.

## 4 Summary

In this work we provided ozone climatology based on 7 years of GOMOS data at southern hemisphere stations from 15 km to 50 km. We required a quality selection in order to have good data measurements. The range for maximum mixing ratios at the equatorial stations is from 9.1 ppmv to 10.5 ppmv, while the range at the tropical stations is from 7.5 ppmv to 10.1 ppmv. In the altitude range from 20 km to 50 km, ozone variability varies between 3% and 10% over all the months and it is larger during austral winter in the 27-35km altitude range.

We compared GOMOS dark limb occultations with 8 southern ozonesondes stations based on SHADOZ network. GOMOS is in satisfactory agreement with SHADOZ in the altitude range from ~20 km to 30 km. On most of the stations, in that altitude range, GOMOS is larger than SHADOZ by ~5-10%. A possible explanation is a bias in the altitude registration of ozonesondes due to their time constant and the impact of GOMOS errors (imperfect scintillation correction, inaccurate aerosol modeling and uncertainties in cross sections of trace gases and in the atmospheric temperature) on ozone data quality. On the contrary at Nairobi and Irene GOMOS values are generally lower than SHADOZ ones. Further studies are needed to understand the causes of the station dependent observed differences. In the altitude range from 21 km to 30 km the monthly differences are generally within  $\pm 15\%$ . And the median of the relative percentage of differences between GOMOS and SHADOZ ozone mixing ratios in the 21-23km height range is nearly within  $\pm 10\%$  for almost all the stations and decrease to nearly  $\pm 5\%$  above 23 km. Below 21 km, GOMOS data show an increasing variability with decreasing altitude and an increasing positive bias compared to SHADOZ in the tropics. We recommend using with caution GOMOS measurements below 21 km in the tropics.

## Acknowledgements

The LACy (Laboratoire de l'Atmosphère et des Cyclones) is supported by INSU (Institut National des Sciences de l'Univers), a CNRS institut, and by the Regional Council of Reunion (Conseil Régional de La Réunion). The present work is part of the Regional COMPTRAST programme. ENVISAT is an ESA mission and GOMOS is an ESA funded instrument.

The authors thank the GOMOS team and SHADOZ network for providing the ozone profiles used in this work. We also acknowledge the European Space Agency (ESA), Centre National d'Etudes Spatiales (CNES) and the European Commission within the SCOUT-O3 project (contract

- 1 505390-GOCECT-2004) for their support. The development of the Basic ENVISAT Atmospheric
- 2 Toolbox is primarily funded by the European Space Agency (ESA).
- 3

## References

- Bencherif H., El Amraoui, L., Semane, N., Massart, S., Vidyarabya Charyulu, D., Hauchecorne, A., and Peuch, V. H.: Examination of the 2002 major warming in the southern hemisphere using ground based and Odin/SMR assimilated data: stratospheric ozone distributions and tropic midlatitude exchange, *Canadian Journal Physics*, 85: 1287-1300, 2007.
- Bertaux, J. L., Kyrölä, E., and Wehr, T.: Stellar occultation technique for atmospheric ozone monitoring: GOMOS on Envisat, *Earth Observation Quarterly*, 67, 17– 20, 2000.
- Bertaux, J.L., Hauchecorne, A., Dalaudier, F., Cot, C., Kyrölä, E., Fussen, D., Tamminen, J., Leppelmeier, G.W., Sofieva, V., Hassinen, S., d’Andon, O.F., Barrot, G., Mangin, A., Theodore, B., Guirlet, M., Korablev, O., Snoeij, P., Koopman, R., and Fraisse, R.: First results on GOMOS/Envisat, *Adv. Space Res.* 33, 1029–1035, 2004.
- Borchi, F., Pommereau, J.-P., Garnier, A., and Nunes-Pinhanrada, M.: Evaluation of SHADOZ sondes, HALOE and SAGE II ozone profiles at the tropics from SAOZ UV-Vis remote measurements on board long duration balloons, *Atmos. Chem. Phys.*, 5, 1381–1397, 2005, <http://www.atmos-chem-phys.net/5/1381/2005/>.
- Borchi, F. and Pommereau, J.P.: Evaluation of ozonesondes, HALOE, SAGE II and III, Odin-OSIRIS and -SMR, and ENVISAT-GOMOS, -SCIAMACHY and-MIPAS ozone profiles in the tropics from SAOZ long duration balloon measurements in 2003 and 2004, *Atmos. Chem. Phys.*, 7, 2671–2690, 2007, <http://www.atmos-chem-phys.net/7/2671/2007/>.
- Bracher, A., Bovensmann, H., Bramstedt, K., Burrows, J. P., von Clarmann, T., Eichmann, K.-U., Fischer, H., Funke, B., Gil-Lopez, S., Glatthor, N., Grabowski, U., Höpfner, M., Kaufmann, M., Kellmann, S., Kiefer, M., Koukouli, M. E., Linden, A., Lopez-Puertas, M., Mengistu Tsidu, G., Milz, M., Noël, S., Rohen, G., Rozanov, A., Rozanov, V. V., von Savigny, C., Sinnhuber, M., Skupin, J., Steck, T., Stiller, G. P., Wang, D.-Y., Weber, M., and Wuttke, M. W.: Cross comparisons of O<sub>3</sub> and NO<sub>2</sub> measured by the atmospheric ENVISAT instruments GOMOS, MIPAS, and SCIAMACHY, *Adv. Space Res.*, 36, 855–867, doi: 10.1016/j.asr.2005.04.005, 2005.
- Clain, G., Baray, J. L., Delmas, R., Diab, R., Leclair de Bellevue, J., Keckhut, P., Posny, F., Metzger, J. M., and Cammas, J. P.: Tropospheric ozone climatology at two Southern Hemisphere tropical/subtropical sites, (Reunion Island and Irene, South Africa) from

ozonesondes, LIDAR, and in situ aircraft measurements, *Atmos. Chem. Phys.*, 9, 1723-1734, 2009, <http://www.atmos-chem-phys.net/9/1723/2009/>.

Dupuy, E., Walker, K. A., Kar, J., Boone, C. D., McElroy, C. T., Bernath, P. F., Drummond, J. R., Skelton, R., McLeod, S. D., Hughes, R. C., Nowlan, C. R., Dufour, D. G., Zou, J., Nichitiu, F., Strong, K., Baron, P., Bevilacqua, R. M., Blumenstock, T., Bodeker, G. E., Borsdorff, T., Bourassa, A. E., Bovensmann, H., Boyd, I. S., Bracher, A., Brogniez, C., Burrows, J. P., Catoire, V., Ceccherini, S., Chabrillat, S., Christensen, T., Coffey, M. T., Cortesi, U., Davies, J., De Clercq, C., Degenstein, D. A., De Mazire, M., Demoulin, P., Dodion, J., Firanski, B., Fischer, H., Forbes, G., Froidevaux, L., Fussen, D., Gerard, P., Godin-Beekmann, S., Goutail, F., Granville, J., Griffith, D., Haley, C. S., Hannigan, J. W., Höpfner, M., Jin, J. J., Jones, A., Jones, N. B., Jucks, K., Kagawa, A., Kasai, Y., Kerzenmacher, T. E., Kleinböhl, A., Klekociuk, A. R., Kramer, I., Kullmann, H., Kuttippurath, J., Kyrölä, E., Lambert, J.-C., Livesey, N. J., Llewellyn, E. J., Lloyd, N. D., Mahieu, E., Manney, G. L., Marshall, B. T., McConnell, J. C., Mc-Cormick, M. P., McDermid, I. S., McHugh, M., McLinden, C. A., Mellqvist, J., Mizutani, K., Murayama, Y., Murtagh, D. P., Oelhaf, H., Parrish, A., Petelina, S. V., Piccolo, C., Pommereau, J.-P., Randall, C. E., Robert, C., Roth, C., Schneider, M., Senten, C., Steck, T., Strandberg, A., Strawbridge, K. B., Sussmann, R., Swart, D. P. J., Tarasick, D. W., Taylor, J. R., Tard, C., Thomason, L. W., Thompson, A. M., Tully, M. B., Urban, J., Vanhellemont, F., Vigouroux, C., von Clarmann, T., von der Gathen, P., von Savigny, C., Waters, J. W., Witte, J. C., Wolff, M., and Zawodny, J. M.: Validation of ozone measurements from the Atmospheric Chemistry Experiment (ACE), *Atmos. Chem. Phys.*, 9, 287–343, 2009, <http://www.atmos-chem-phys.net/9/287/2009/>.

Fortuin, J. P. F., and Kelder, H.: An ozone climatology based on ozonesonde and satellite measurements, *J. Geophys. Res.*, 103(D24), 31,709–31,734, 1998.

Hocke, K., Kämpfer, N., Ruffieux, D., Froidevaux, L., Parrish, A., Boyd, I., von Clarmann, T., Steck, T., Timofeyev, Y. M., Polyakov, A. V., and Kyrölä, E.: Comparison and synergy of stratospheric ozone measurements by satellite limb sounders and the ground-based microwave radiometer SOMORA, *Atmos. Chem. Phys.*, 7, 4117-4131, 2007, <http://www.atmos-chem-phys.net/7/4117/2007/>.

Johnson, B. J., Oltmans, S. J., Vömel, H., Deshler, T., Kroger, C., and Smit, H. G. J.: ECC ozonesonde pump efficiency measurements and sensitivity tests of buffered and unbuffered sensor solutions, *J. Geophys. Res.*, 107(D19), 4393, doi:10.1029/2001JD000557, 2002.

1 Kyrölä, E., Tamminen, J., Leppelmeier, G. W., Sofieva, V., Hassinen, S., Bertaux, J.-L.,  
2 Hauchecorne, A., Dalaudier, F., Cot, C., Korablev, O., Hembise, O., Barrot, G., Mangin, A.,  
3 Théodore, B., Guirlet, M., Etanchaud, F., Snoeij, P., Koopman, R., Saavedra, L., Fraisse, R.,  
4 Fussen, D., and Vanhellemont, F.: GOMOS on Envisat: An overview, *Adv. Space Res.*, 33,  
5 1020–1028, doi:10.1016/S0273-1177(03)00590-8, 2004.

6 Kyrölä, E., Tamminen, J., Leppelmeier, G. W., Sofieva, V., Hassinen, S., Bertaux, J.-L.,  
7 Hauchecorne, A., Dalaudier, F., Fussen, D., Vanhellemont, F., Fanton d’Andon, O., Barrot,  
8 G., Mangin, A., Théodore, B., Guirlet, M., Snoeij, P., Koopman, R., Saavedra, L., and Meijer,  
9 Y. J.: Nighttime ozone profiles in the stratosphere and mesosphere by the Global Ozone  
10 Monitoring by Occultation of Stars on Envisat, *J. Geophys. Res.*, 111, D24306,  
11 doi:10.1029/2006JD00577193, 2006.

12 Kyrölä, E., Tamminen, J., Sofieva, V., Bertaux, J.-L., Hauchecorne, A., Dalaudier, F., Fussen,  
13 D., Vanhellemont, F., Fanton d’Andon, O., Barrot, G., Mangin, A., Guirlet, M., Saavedra, L.,  
14 Snoeij, P., Fehr, T., Meijer, Y., and Fraisse R.: Global distribution of ozone vertical profiles  
15 from GOMOS measurements 2002–2006, in: *Proceedings of the ‘Envisat Symposium 2007’*,  
16 Montreux, Switzerland 23–27 April 2007 (ESA SP-636, July 2007).

17 Lamsal, L.N., Weber, M., Tellmann, S., and Burrows, J. P.: Ozone column classified  
18 climatology of ozone and temperature profiles based on ozonesonde and satellite data, *J.*  
19 *Geophys. Res.*, vol. 109, D20304, doi: 10.1029/2004JD004680, 2004.

20 Logan J. A., Jones, D. B. A., Megretskaia, I. A., Oltmans, S. J., Johnson, B. J. , Vomel, H.,  
21 Randel, W. J. , Kimani, W. , and Schmidlin, F. J.: Quasi-biennial oscillation in tropical ozone  
22 as revealed by ozonesonde and satellite data, *J. Geophys. Res.*, 108, NO. D8, 4244,  
23 doi:10.1029/2002JD002170, 2003.

24 Liu, X., Chance, K., Sioris, C. E., Kurosu, T. P., and Newchurch, M. J.: Intercomparison of  
25 GOME, ozonesonde, and SAGE II measurements of ozone: Demonstration of the need to  
26 homogenize available ozonesonde data sets, *J. Geophys. Res.*, 111, D14305,  
27 doi:10.1029/2005JD006718, 2006.

28 Liu Y., Cai, Z., Kyrölä, E.: Comparison of ENVISAT GOMOS and MIPAS ozone profiles  
29 with balloon sonde measurements from Beijing, in: *Proceedings of the Dragon 1 Programme*  
30 *Final Results 2004–2007*, Beijing, P.R. China 21– 25 April 2008 (ESA SP-655, April 2008).

31 McPeters, R. D., Labow, G. J., and Logan, J. A.: Ozone climatological profiles for satellite  
32 retrieval algorithms, *J. Geophys. Res.*, 112, D05308, doi:10.1029/2005JD006823, 2007.



1 Meijer, Y. J., Swart, D. P. J., Allaart, M., Andersen, S. B., Bodeker, G., Boyd, I., Braathen,  
2 G., Calisesi, Y., Claude, H., Dorokhov, V., von der Gathen, P., Gil, M., Godin-Beekmann, S.,  
3 Goutail, F., Hansen, G., Karpetchko, A., Keckhut, P., Kelder, H. M., Koelemeijer, R., Kois,  
4 B., Koopman, R. M., Kopp, G., Lambert, J.-C., Leblanc, T., McDermid, I. S., Pal, S., Schets,  
5 H., Stubi, R., Suortti, T., Viscont G., and Yela, M.: Pole-to-pole validation of Envisat  
6 GOMOS ozone profiles using data from groundbased and balloon sonde measurements, J.  
7 Geophys. Res., 109, D23305, doi:10.1029/2004JD004834, 2004.

8 ~~Renard, J. B., Berthet, G., Brogniez, C., Catoire, V., Chartier, M., Gaubicher, B., Goutail, F.,~~  
9 ~~Oelhaf, H., and Robert, C.: Overview of the validation of GOMOS measurements using~~  
10 ~~various stratospheric balloon borne instruments, in: Proceedings of the Third Workshop on~~  
11 ~~the Atmospheric Chemistry Validation of Envisat (ACVE 3) 4-7 December 2006, ESA-~~  
12 ~~ESRIN, Frascati, Italy, (ESA SP 642, February 2007).~~

13 Renard, J., Berthet, G., Brogniez, C., Catoire, V., Fussen, D., Goutail, F., Oelhaf, H.,  
14 Pommereau, J., Roscoe, H. K., Wetzol, G., Chartier, M., Robert, C., Balois, J., Verwaerde, C.,  
15 Auriol, F., Francois, P., Gaubicher, B., and Wursteisen, P.: Validation of GOMOS-Envisat  
16 vertical profiles of O<sub>3</sub>, NO<sub>2</sub>, NO<sub>3</sub>, and aerosol extinction using balloon-borne instruments  
17 and analysis of the retrievals, J. Geophys. Res. (Space Physics), 113, A02302,  
18 doi:10.1029/2007JA012345, 2008. 2172

19 Semane, N., Bencherif, H., Morel, B., Hauchecorne, A., and Diab, R. D.: An unusual  
20 stratospheric ozone decrease in the Southern Hemisphere subtropics linked to isentropic air-  
21 mass transport as observed over Irene (25.5° S, 28.1° E) in mid-May 2002, Atmospheric  
22 Chemistry and Physics, 6, 1927–1936, 2006, <http://www.atmos-chem-phys.net/6/1927/2006/>.

23 Shepherd, T. G.: The middle atmosphere, Journal of Atmospheric and Solar-Terrestrial  
24 Physics, 62, 1587-1601, 2000.

25 Sivakumar, V., Portafaix, T., Bencherif, H., Godin-Beekmann, S., and Baldy, S.:  
26 Stratospheric ozone climatology and variability over a southern subtropical site: Reunion  
27 Island (21°S; 55°E), Ann. Geophys., 25, 2321–2334, 2007, [http://www.ann-](http://www.ann-geophys.net/25/2321/2007/)  
28 [geophys.net/25/2321/2007/](http://www.ann-geophys.net/25/2321/2007/).

29 Sofieva, V. F., Tamminen, J., Haario, H., Kyrölä, E., and Lehtinen, M.: Ozone profile  
30 smoothness as a priori information in the inversion of limb measurements, Annales  
31 Geophysicae, 22(10), 3411–3420, 2004, <http://www.ann-geophys.net/22/3411/2004/>.

1 [Sofieva, V. F., Kan, V., Dalaudier, F., Kyrölä, E., Tamminen, J., Bertaux, J.-L., Hauchecorne,](#)  
2 [A., Fussen, D., and Vanhellemont, F.: Influence of scintillation on quality of ozone](#)  
3 [monitoring by GOMOS, Atmos. Chem. Phys., 9, 9197–9207, 2009,](#)  
4 [<http://www.atmos-chem-phys.net/9/9197/2009/>. 6767, 6768](#)

5 Tamminen, J., Karhu, J. A., Kyrölä, E., Hassinen, S., Kyrö, E., Karpechko, A. Y., and  
6 Piacentini, E.: GOMOS Ozone Profiles at High Latitudes: Comparison with Marambio and  
7 Sodankylä Sonde Measurements, in: Atmosphere and Climate, Studies by Occultation  
8 Methods, Springer Berlin Heidelberg, doi: 10.1007/3-540-34121-8, 2006.

9 [Tamminen, J., Kyrölä, E., Sofieva, V. F., Laine, M., Bertaux, J.-L., Hauchecorne, A.,](#)  
10 [Dalaudier, F., Fussen, D., Vanhellemont, F., Fanton-d'Andon, O., Barrot, G., Mangin, A.,](#)  
11 [Guirlet, M., Blanot, L., Fehr, T., Saavedra de Miguel, L., and Fraisse, R.: GOMOS data](#)  
12 [characterization and error estimation, Atmos. Chem. Phys. Discuss., 10, 6755-6796, 2010.](#)

13 Thompson, A. M., Witte, J. C., McPeters, R. D., Oltmans, S. J., Schmidlin, F. J., Logan, J. A.,  
14 Fujiwara, M., Kirchhoff, V. W. J. H., Posny, F., Coetzee, G. J. R., Hoegger, B., Kawakami,  
15 S., Ogawa, T., Johnson, J. B., Vömel, H., and Labow, G.: Southern Hemisphere Additional  
16 Ozonesondes (SHADOZ) 1998–2000 tropical ozone climatology 1. Comparison with Total  
17 Ozone Mapping Spectrometer (TOMS) and ground-based measurements, J. Geophys. Res.,  
18 108, 8238, doi:10.1029/2001JD000967, 2003a.

19 Thompson, A. M., Witte, J. C., Smit, H. G. J., Oltmans, S. J., Johnson, J. B., Kirchhoff, V. W. J.  
20 H., and Schmidlin, F. J.: Southern Hemisphere Additional Ozonesondes (SHADOZ) 1998–  
21 2004 tropical ozone climatology: 3. Instrumentation, station-to-station variability, and  
22 evaluation with simulated flight profiles, J. Geophys. Res., 112, D03304, doi:  
23 10.1029/2005JD007042, 2007.

24 Verronen, P. T., Kyrölä, E., Tamminen, J., Funke, B., Gil-Lopez, S., Kaufmann, M., Lopez-  
25 Puertas, M., Clarmann, T. v., Stiller, G., Grabowski, U., and Höpfner, M.: A comparison of  
26 nighttime GOMOS and MIPAS ozone profiles in the stratosphere and mesosphere, Adv.  
27 Space Res., 36, 958–966, 2005.

28 Witte, J. C., Schoeberl, M. R., Douglass, A. R., and Thompson, A. M.: The Quasi-biennial  
29 Oscillation and annual variations in tropical ozone from SHADOZ and HALOE. Atmos.  
30 Chem. Phys., 8, 3929–3936, 2008, [<http://www.atmos-chem-phys.net/8/3929/2008/>](#).

- 1 WMO, Scientific Assessment of Ozone Depletion: 2006, World Meteorological Organization,
- 2 Global Ozone Research and Monitoring Project, Report N°50, Switzerland, 2007.

Table 1. Overview of SHADOZ stations used in this study

Location	Latitude	Longitude	Period	Profiles	Elevation	Sonde Info*
<i>Equatorial stations</i>						
Nairobi	1.27°S	36.8°E	1998-2008	459	1795 m	EnSci 2Z
Natal	5.42°S	35.38°W	1998-2008	394	42 m	EnSci Z, Science Pump 6A
Java, Watukosek	7.57°S	112.65°E	1998-2008	282	50 m	MEISEI RSII-KC79D, EnSci 2Z
Ascension	7.98°S	14.42°W	1998-2008	492	91 m	Science Pump 6A
<i>Tropical stations</i>						
Samoa	14.23 S	170.56°W	1998-2008	383	77 m	Science Pump 6A
Fiji	18.13°S	178.40°E	1998-2008	275	6 m	Science Pump ECC6A
Reunion	21.06°S	55.48°E	1998-2008	319	24 m	EnSci Z & SPC 6A
Irene	25.90°S	28.22°E	1998-2008	232	1524 m	Science Pump ECC6A

\* See more details for instruments and solution strengths in Thompson et al. (2003, 2007)

Table 2. Summary of results for the GOMOS comparison with SHADOZ. For each station: the vertical range used to calculate the median difference, mean value and minimum/maximum values in this range are indicated.

Location	Range (km)	Difference (%)	
		Mean	Range
<i>Equatorial stations</i>			
Nairobi	21-30	-0.89	-4.57 to +6.42
Natal	21-30	+1.75	-0.60 to +8.35
Java, Watukosek	21-30	+4.13	-1.48 to +15.66
Ascension	21-30	+8.00	+4.65 to +10.94
<i>Tropical stations</i>			
Samoa	21-30	+1.64	-1.58 to +5.53
Fiji	21-30	+2.68	+0.20 to +6.85
Reunion	21-30	+4.47	-0.74 to +10.71
Irene	21-30	-1.72	-3.85 to +5.05




Figure 1. Map of SHADOZ stations in the southern hemisphere. Ozone profiles obtained by radiosonde experiments at 8 stations (black circles) are used for the purpose at the present issue, from west to east: Samoa, Natal, Ascension Island, Irene, Nairobi, Reunion, Java and Fiji. The regions where GOMOS profiles are selected within  $\pm 5^\circ$  latitude and  $\pm 10^\circ$  longitude differences over each station are also represented (dashed boxes).





Figure 2. Monthly distributions of the number of the GOMOS dark occultations (black bars) and SHADOZ profiles (white bars). The GOMOS dataset has a global coverage for the time period from 2002 to 2008, while the SHADOZ dataset is covers the years from 1998 to 2008 and includes 8 southern tropical/subtropical sites.




1

2 Figure 3. Monthly median distributions of ozone mixing ratio (in ppmv) as derived from  
 3 GOMOS observations per station. The contours are separated by 1 ppm.


4






1  
2 Figure 4. Contour plots of ozone variability (in %) from GOMOS observations. Variability is  
3 defined as the ratio between the interquartile range (*iqr*) and the median. Contours are  
4 separated by 5 %.

5



1  
2 Figure 5. Monthly distributions of the relative percentage of differences between GOMOS  
3 and SHADOZ observations (these differences are based on comparisons of monthly means).  
4 It is calculated with respect to ground based measurements in the altitude range from 15 km to  
5 30 km. The contours are separated by 5%.

6



1

2 Figure 6. Median (thick continuous lines) and mean (dashed lines) ~~of the global differences~~  
3 over all monthly comparisons between GOMOS and SHADOZ ozone mixing ratios, for  
4 heights between 15 km and 30 km. The thin continuous lines refer to the 25<sup>th</sup> and 75<sup>th</sup>  
5 percentiles for the median of the percentage of differences.