

IUPAC Subcommittee on Gas Kinetic Data Evaluation – Data Sheet V.A1.33

Datasheets can be downloaded for personal use only and must not be retransmitted or disseminated either electronically or in hardcopy without explicit written permission. The citation for this datasheet is: IUPAC Subcommittee for Gas Kinetic Data Evaluation, <http://www.iupac-kinetic.ch.cam.ac.uk>

This datasheet last evaluated: May 2010; last change to preferred values: May 2010.

HOI (Ice) → Products

Experimental Data

Parameter	Temp./K	Reference	Technique/Comments
<i>Experimental uptake coefficients: γ, γ_0</i>			
$\gamma = (8 \pm 2) \times 10^{-2}$	180	Allanic and Rossi, 1999	Knudsen-MS (a)
$\gamma = (5.6 \pm 2) \times 10^{-2}$	190		
$\gamma = (3.1 \pm 0.7) \times 10^{-2}$	200		
$\gamma_0 > 10^{-2}$	243	Holmes et al., 2001	CWFT-MS (b)

Comments

- (a) HOI (at concentrations of $\sim 10^{10}$ molecule cm $^{-3}$) was formed in the reaction of O(3P) with C₂H₅I. The ice film was formed by vapour deposition. The uptake did not saturate (time independent) under these conditions and displayed a negative temperature dependence. It was found to be independent of HOI concentration (varied over a factor of four). The uncertainty in γ reflects complexities in the HOI kinetics. The only gas-phase product detected was I₂ with no recoverable HOI after the evaporation of the 20 μ m thick ice film.
- (b) HOI (at concentrations $< 10^{10}$ molecule cm $^{-3}$) was formed in the reaction of O(3P) with C₃H₇I. The ice film was formed by freezing liquid water at 258 K. The uptake coefficient decreased with exposure time and HOI was observed to desorb from the ice film after exposure stopped, indicating a (partially) reversible process. Diffusion limitation prevented precise measurement of the initial uptake coefficient.

Preferred Values

none


Comments on Preferred Values

The results of Allanic and Rossi (1999) and Holmes et al (2001) were obtained in different temperature ranges and concur that the uptake coefficient is large. A parameterisation, $\gamma = 5 \times 10^{-5} \exp(1300/T)$ generates uptake coefficients that are consistent with both datasets. At the lower temperatures covered by Allanic and Rossi (1999), the uptake is observed to be irreversible, whereas Holmes et al. (2001) observe partially reversible adsorption at 243 K. From the experimental datasets it is not obvious how HOI reacts with the ice surface. Both studies observed I₂ as product and Holmes et al (2001) suggest that this may arise from self-reaction of HOI (or reaction of HOI with IONO₂ impurity) on the ice surface. It is possible

that on a pure ice surface, HOI will adsorb reversibly unless high concentrations and low temperatures result in formation of a thermodynamically stable phase (e.g. formation of hydrates). For this reason we make no recommendation for uptake to pure ice, whilst noting that HOI reacts readily on ice surfaces containing reactive species (Allanic and Rossi, 1999; Holmes et al., 2001).

References

Allanic, A. and Rossi, M.J.: J. Geophys. Res. 104, 18689-18696, 1999.
 Holmes, N. S., Adams, J. W., and Crowley, J. N.: Phys. Chem. Chem. Phys., 3, 1679-1786, 2001.



Experimental uptake coefficients for HOI interaction with ice.