

## ***Interactive comment on “Measurements of HONO during BAQS-Met” by J. J. B. Wentzell et al.***

**T. Johnson**

timothy.johnson@pnl.gov

Received and published: 12 July 2010

The authors may wish to refer to the recently reinforced importance of biomass burning as a source of HONO. This was first noted by Keene et al (2006) using MC-IC and then more quantitatively using infrared spectroscopy by Yokelson et al (2007, 2009). Very recently, HONO has been confirmed as a significant product in biomass burning with a very high degree of correlation between FTIR and Negative-ion Proton-Transfer Chemical-Ionization Mass Spectrometry (NI-PT-CIMS) measurements (Roberts et al, 2010, Burling et al, 2010)

William C. Keene, et al. “Emissions of major gaseous and particulate species during experimental burns of southern African biomass,” *Journal of Geophysical Research*, Vol. 111, D04301.

R. J. Yokelson, et al. “The Tropical Forest and Fire Emissions Experiment: overview  
C5178

and airborne fire emission factor measurements,” *Atmos. Chem. Phys.*, 7, 5175–5196, 2007

R.J. Yokelson, et al. “Emissions from biomass burning in the Yucatan,” *Atmos. Chem. Phys.*, 9, 5785–5812, 2009

J.M. Roberts et al. “Measurement of HONO, HNCO, and other inorganic acids by negative-ion proton-transfer chemical-ionization mass spectrometry (NI-PT-CIMS): application to biomass burning emissions” *Atmos. Meas. Tech. Discuss.*, 3, 301–331, 2010

I.R. Burling, et al. “Laboratory measurements of trace gas emissions from biomass burning of fuel types from the Southeastern and Southwestern United States” *Atmos. Chem. Phys. Discuss.*, 10, 16425–16473, 2010.

---

Interactive comment on *Atmos. Chem. Phys. Discuss.*, 10, 15295, 2010.