

Interactive
Comment

Interactive comment on “Role of aldehyde chemistry and NO_x concentrations in secondary organic aerosol formation” by A. W. H. Chan et al.

A. W. H. Chan et al.

achan@caltech.edu

Received and published: 13 July 2010

We thank the reviewer for the helpful comments. Please see responses below:

- *Page 10240, lines 20–25: Do you mean specifically alpha-beta unsaturated aldehydes may be important in SOA formation or all aldehydes?*

Since only α , β -unsaturated aldehydes are studied in detail here, we expect these aldehydes are important SOA precursors through PAN chemistry. As shown in this work, other aldehydes (e.g. 4-pentenal) can be a source of SOA, but the formation pathways are different than those shown for α , β -unsaturated aldehydes. At this point, we cannot establish that the chemistry elucidated in this work can be applied to other aldehydes.

C5261

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



- *Page 10237, lines 11–15 and Page 10263, Figure 10: It looks like the stoichiometry of the reactions for forming the dihydroxyacid and hydroxynitrooxyacid by reaction of the dioxketone intermediate with H_2O and HNO_3 is incorrect. To form these products the dioxketone would need to react with H_2 and HNO_2 , or if the reactions occurred with H_2O and HNO_3 then O atoms would have to be formed as products. This probably makes the proposed reactions less likely.*

We agree with the reviewer that reaction with H_2 and HONO is unlikely. Reaction with H_2O or HNO_3 is far more likely, but the stoichiometry would result in an additional O atom. This pathway cannot be ruled out from our experimental results. Another possible pathway is an analogous cyclization of the alkoxy radical (following reaction with NO), leading to a C_4 -hydroxy-epoxyketone (with one fewer O atom than the dioxketone proposed here). Reaction of this intermediate with H_2O or HNO_3 will yield the observed oligoesters. However, decomposition of the alkoxy radical is likely favored over cyclization, and, in addition, such an intermediate is highly strained, making this pathway less likely. At this stage, we propose the dioxketone pathway as the most viable pathway, and more work needs to be done to elucidate the exact chemistry.

- *Page 10252, Table 1: I think the MBO232 and MBO231 structures in the table are reversed.*

Table 1 has been corrected.

Interactive comment on Atmos. Chem. Phys. Discuss., 10, 10219, 2010.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)