

Interactive comment on “Does acetone react with HO₂ in the upper-troposphere?” by T. J. Dillon et al.

J. Peeters (Referee)

jozef.peeters@chem.kuleuven.be

Received and published: 23 August 2010

Referee Comments on ACPD manuscript acp-2010-442 by T. J. Dillon et al.

Referee: Prof. Emeritus Jozef Peeters, Department of Chemistry, University of Leuven
e-mail: jozef.peeters@chem.kuleuven.be

General Comments

The authors have performed an experimental investigation of the reaction between HO₂ radicals and acetone at low temperatures of 207 - 228 K and reduced pressures of 25 - 85 Torr, essentially based on measuring the OH resulting from the competition between (i) the sequence HO₂ + CH₃C(O)CH₃ ↔ (CH₃)₂C(OH)OO (R1, reversible; k₁ and k₋₁) followed by (CH₃)₂C(OH)OO + NO → products (k₂), and (ii) the well known re-

C6746

action HO₂ + NO → OH + NO₂ (k₈). The OH was monitored by LIF and calibration was achieved on basis of the kinetically well characterized reaction system in absence of acetone. The experimental results, obtained using proven methodologies, are of high quality, and the lower-limit k₁ data that could be derived directly are by themselves of much value, showing that the reaction does indeed occur at these low temperatures at rates of at least 1 × 10⁻¹² cm³ s⁻¹ at T ≤ 210 K and 25 Torr. However, the more involved interpretation of the experimental data in terms of both k₁ and k₋₁ (or K₁) is not straightforward (see Specific Comments, below), and more importantly, two aspects of reaction (R1), with a large impact on the chemistry as detailed in the Specific Comments below, were not taken into account. Thus, though the paper shows convincingly that reaction (R1) does occur, the derived k₋₁(T) and equilibrium constant K₁(T) data are both questionable, and the conclusion of the authors that the HO₂ initiated oxidation of acetone and formation of acetic acid thereby is not significant in the tropopause appears therefore insufficiently substantiated. The interpretation of the data should therefore be re-addressed, in the light of the particular aspects of reaction (R1) detailed below.

Specific Comments

1. First, it should be noted that the k₋₁(T) and equilibrium constant K₁(T) results obtained by the authors appear rather unusual in the light of the experimental data for the analogous reactions of HO₂ with HCHO and CH₃CHO and the reverse reactions. The reported k₋₁(T) for (CH₃)₂C(OH)OO decomposition imply a very high k₋₁(300 K) of around 2 × 10⁶ s⁻¹. When comparing this result to the experimental values of 175 s⁻¹ for H₂C(OH)OO (Veyret et al., 1989) and 1900 s⁻¹ for CH₃CH(OH)OO (Tomas et al., 2001), the carbonyl-stabilizing inductive effect of the second CH₃ group seems a hundred times stronger than that of the first CH₃ substituent, which is definitely not in keeping with group additivity concepts. Further, from the reported K₁(T) results, the stability of (CH₃)₂C(OH)OO with respect to the reactants appears to be 11 kJ mol⁻¹ less than the 59 ± 2 kJ mol⁻¹ computed at various high-levels of theory by Hermans

C6747

et al., 2005, whereas the stabilities computed in that same paper at the same levels of theory for the two analogous α -OH-peroxys $\text{CH}_3\text{CH}(\text{OH})\text{OO}$ and $\text{H}_2\text{C}(\text{OH})\text{OO}$ were shown to agree with experiment within 2 kJ mol⁻¹. This gives reason to question the more involved interpretation of the [OH] data in presence/absence of acetone in terms of both k_1 and k_{-1} .

2. The apparent rate constant that can be derived from the measured reduction of [OH] upon acetone addition is essentially $k_1/(1 + k_{-1}/k_2[\text{NO}])$, or $K_1/(1/k_{-1} + 1/k_2[\text{NO}])$, and hence k_1 and k_{-1} (or k_1 and K_1) could not be obtained independently from these experiments. The authors opted for adopting the theoretically predicted k_1 of Hermans et al. 2004, and so deriving k_{-1} (and hence K_1) from the observed [OH] reduction upon acetone addition. This by itself was not the best choice, since the theoretical k_1 of Hermans et al. 2004 (as all the k in that paper and likewise in Hermans et al. 2005) are TST-based and therefore by definition high-pressure limit values, whereas the reaction in question, through a stable pre-reactive complex (PRC) and over a submerged barrier, will necessarily show pressure-dependence, while the experimental pressures are quite low. Moreover, equilibrium constants, besides being independent of pressure, can in general be predicted (much) more accurately than rate coefficients; therefore, a better choice would have been to adopt the theoretical K_1 of Hermans et al. 2005, in which paper the stability of the α -OH-peroxy at hand was verified at higher levels of theory compared to the 2004 paper, and the rate coefficients at issue here were accordingly upgraded.

3. As to the pressure-dependence of reaction (R1), the rate of prompt redissociation of the nascent chemically activated PRC^\ddagger formed at 210-230 K can be estimated at roughly $k_{\text{red}}(E_{\text{av}} = 8 \text{ kJ/mol}) \approx 3 \times 10^9 \text{ s}^{-1}$, such that at 25 and 85 Torr, collisional stabilization, with $\beta_{\text{scZLJ}}[\text{M}]$ of $\approx 5 \times 10^7 \text{ s}^{-1}$ and $\approx 1.5 \times 10^8 \text{ s}^{-1}$, respectively, is too slow and hence product formation occurs mainly through the nascent, activated PRC^\ddagger , meaning that the reaction is in its low-P limit. The forward $k_f(E_{\text{av}} = 8 \text{ kJ/mol})$ rate to form the product is estimated at about $2 \times 10^7 \text{ s}^{-1}$, i.e. 100 times slower than redis-

C6748

sociation; both these rates are estimated here from the data in Hermans et al. 2005 and their available Supporting Information. (See Note on rates estimated in these comments, below). That higher pressures should increase the product formation rate (i.e. the effective k_1) substantially, is readily seen: $\text{PRC}(\ddagger)$ that are brought just below the reactants level by collisions can no longer redissociate promptly, but still promptly yield the product as the submerged transition state lies 11 kJ mol⁻¹ below the reactants (see Hermans et al. 2005 for the energetics). The pressure effect at these low T should be pronounced; assuming an HO₂-acetone capture rate $k_{\text{capt}} = 10^{(-10)} \text{ cm}^3 \text{ s}^{-1}$, the low-P k_1 is estimated at barely $10^{(-12)} \text{ cm}^3 \text{ s}^{-1}$, i.e. substantially lower than the upgraded high-P limit $k_{1,\infty}(207 \text{ K}) = 5.8 \times 10^{(-12)} \text{ cm}^3 \text{ s}^{-1}$ and $k_{1,\infty}(228 \text{ K}) = 3.0 \times 10^{(-12)} \text{ cm}^3 \text{ s}^{-1}$ predicted by Hermans et al. 2005.

4. Equally important, but not considered by the authors, is that at these low T and high [acetone] around $10^{(15)} \text{ cm}^{-3}$, the bulk of the HO₂ can be expected to be sequestered in the very stable H-bonded $\text{OOH}-\text{O}=\text{C}(\text{CH}_3)_2$ PRC complex, in thermal equilibrium with acetone and free HO₂. The $\text{HO}_2 + \text{CH}_3\text{C}(\text{O})\text{CH}_3 \leftrightarrow \text{OOH}-\text{O}=\text{C}(\text{CH}_3)_2$ complexation (K_{compl}) was first described by Aloisio and Francisco 2000 at B3LYP level, while Hermans et al. 2005 computed an even slightly higher complex stability of $41 \pm 4 \text{ kJ mol}^{-1}$ at several higher levels of theory, giving also the vibration/rotation parameters in Supporting Information (SI). Of much impact on the issue at hand are the magnitude of $K_{\text{compl}}(T)$ and the timescale to attain the equilibrium. $K_{\text{compl}}(200-230 \text{ K})$ is large because the complex is stable but at the same time quite "loose" and hence exhibits a high density of states and high partition function on account of the quasi-free internal rotation of the terminal hydroperoxy-O about the O-H—O= axis. The harmonic-oscillation-approximation of Gaussian attributes a "vibration frequency" of 60 cm⁻¹ to this mode (see SI Hermans et al. 2005), but this results merely from a small dip in the (otherwise flat) $V(\theta)$ internal rotation potential, at a dihedral angle where the terminal hydroperoxy-O is close to the nearest methyl-H (but still 2.6 Å distant); the 207 K partition function for this quasi-free internal rotation mode is estimated at about 20, i.e. much higher than the HOA value of 2.9. As earlier reported by Aloisio and Fran-

C6749

cisco, two other new modes of the complex (Gaussian "frequencies 83 and 157 cm⁻¹" of Hermans et al.) are hindered internal rotations, which together can put the partition function another factor of ≈ 3 above the HOA approximation. Thus, $K_{\text{compl}}(T)$ is roughly estimated at about 3×10^{-14} cm³ at 207 K and 5×10^{-15} cm³ at 228 K, with a probable error of a factor of 10. Given the experimental [acetone] of order 10^{15} cm⁻³, this implies that at equilibrium the bulk of the HO₂ (some 80% to well over 90%) can be tied up in the thermal OOH—O=C(CH₃)₂ complex. As the terminal O is quasi-free, this complex should still react with NO, to yield an OH that is more weakly H-bonded to the carbonyl-O (see Aloisio and Francisco, 2000) and thus easily released. On the other hand, reaction of the complex with another acetone to form the α -OH-peroxy is not expected to contribute, as it is both energetically (by 41 kJ mol⁻¹) and entropically strongly disfavored compared to (R1). Besides thermal redissociation, the complex can thermally convert, about 20 times slower, to the α -OH-peroxy, which brings the effective k_1 somewhat above the low-P limit. The rate of approach to the equilibrium can be approximated as $k_{\text{capt}} \times [\text{acetone}] \times \beta_{\text{scZLJ}}[M] / (k_{\text{red}}(E_{\text{av}}) + \beta_{\text{scZLJ}}[M])$ (given that the term for the thermal redissociation is sizably smaller, at least at 207 K and 25 Torr). This gives an equilibration time constant of roughly 1000 μ s (with a large uncertainty) for the 207 K, 25 Torr, 7.5×10^{14} cm⁻³ acetone experiments, such that at [NO] = 1.5×10^{14} cm⁻³ much of the HO₂ will already have reacted before the complex attains equilibrium. At 228 K, 85 Torr and 1.7×10^{15} cm⁻³ acetone however, where also the thermal redissociation term becomes important, the time constant is nearer 50 μ s, such that HO₂ complexation is quasi-immediate. It should be stressed however that the K_{compl} estimates above may be off by a factor of 10, given the ± 4 kJ mol⁻¹ uncertainty on the complex stability, as stated for most relative energies in the Hermans et al. 2005 paper.

Items 3 and 4 above can explain the strong apparent disagreement between the k_1 and K_1 data derived in this manuscript and the predictions of Hermans et al. For instance, at 207 K, 25 Torr: a same effective $k_1 / (1 + k_1/k_2[\text{NO}])$ rate as concluded to in the paper is obtained with k_1 negligible and $k_1 = 1.6 \times 10^{-12}$ cm³ s⁻¹ (i.e. 0.27 C6750

times the $k_{1,\infty}$ prediction of Hermans et al. 2005); and at 228 K, 85 Torr: likewise, with k_1 negligible and $k_1 \approx 1 \times 10^{-12}$ cm³ s⁻¹ (i.e. $0.33 \times k_{1,\infty}$) but with 80-90% of the HO₂ quickly complexed.

5. According to the authors' interpretation of the observed [OH] behavior in Fig. 2, with a large k_1 of 30000 s⁻¹ at T = 228 K, the second term in the denominator of the apparent rate for HO₂ removal by acetone, $k_1[\text{acetone}] / (1 + k_1/k_2[\text{NO}])$ (see sub 1., above), would be much larger than 1, such that the apparent rate would simplify to $K_1 k_2 [\text{acetone}] [\text{NO}]$, and thus show the same dependence on [NO] as the rate $k_8 [\text{NO}]$ of the competing HO₂ + NO \rightarrow OH + NO₂ reaction. As a result, the competition between HO₂ removal by acetone and by NO to yield OH would be independent of [NO]. If on the other hand k_1 is negligible at these T as predicted by Hermans et al., the apparent rate of HO₂ removal by NO is independent of [NO], very different from the linear [NO]-dependence of the competing HO₂ + NO reaction. This might have been easily checked by the authors. However, the tell-tale dependence of the competition at issue on [NO] was not investigated, all experiments at given T having having been conducted at a single [NO].

6. In the light of the above, the section on atmospheric modelling in the manuscript appears rather premature.

Technical and minor Comments

Page 16755, line 24: this should be : $k_1 = 3 \times 10^{-12}$ cm³ molecule⁻¹ s⁻¹ (minus sign in exponent).

Table 3: the fast OH + HO₂ reaction is missing as a (minor) OH sink.

Table 3: A minus sign is missing in the exponent of 10 of the second term.

Table 3: OCH₂C(O)CH₃[†] decomposes promptly into CH₃CO + HCHO instead of reacting with O₂ (Orlando et al., 2000).

Table 3: The expression given for $k(\text{CH}_3 + \text{O}_2)$ would result at the experimental pres- C6751

tures in values well above the high-P limit.

Table 3: The reaction of $\text{CH}_3\text{CO} + \text{O}_2$ generates about 30% OH at 207 K and 25 Torr, and about 8% at 228 K and 85 Torr.

Table 3: Equilibrium formation of the $\approx 68 \text{ kJ mol}^{-1}$ stable HO_2OOH tetroxide from $\text{HO}_2 + \text{HO}_2$ (see Denis et al., *J. Phys. Chem. A*, 2009, 113, 499) is missing, but is only minor at the fairly low experimental $[\text{HO}_2]$ of this work, though quite important at the much higher $[\text{HO}_2]$ in another recent study.

Page 16761, line 24: twice "in line with (the) theoretical predictions".

NOTE on rates and equilibria estimated for these comments.

Chemically activated rates estimated for these comments by the referee are generally Whitten-Rabinovitch-type RRKM-estimates, and should not be considered as rigorous RRKM derivations, from which they may deviate by a factor of 3 or more. Estimated thermal rates in these comments are likewise not rigorous MC-TST derivations and can be off by a factor of 3 to 10; similar for the thermal-equilibrium constants.

REFERENCES

Aloisio, S. and Francisco, J. S., *J. Phys. Chem. A*, 104, 3211-3224, 2000.

Hermans, I., Nguyen, T. L., Jacobs, P. A. and Peeters, J., *J. Am. Chem. Soc.*, 126, 9908-9909, 2004.

Hermans, I., Müller, J. F., Nguyen, T. L., Jacobs, P. A. and Peeters, J., *J. Phys. Chem. A*, 109, 4303-4311, 2005.

Orlando, J. J., Tyndall, G. S., Vereecken, L. and Peeters, J., *J. Phys. Chem. A*, 104, 11578-11588, 2000.

Tomas, B., Villenave, E. and Lesclaux, R., *J. Phys. Chem. A*, 105, 3505, 2001.

Veyret, B., Lesclaux, R., Rayez, M.-T., Rayez, J.-C., Cox, R. A. and Moortgat, G. K.: *J.*

C6752

Phys. Chem., 93, 2368, 1989.

Interactive comment on *Atmos. Chem. Phys. Discuss.*, 10, 16747, 2010.

C6753