

Interactive  
Comment

## ***Interactive comment on “Anthropogenic imprints on nitrogen and oxygen isotopic composition of precipitation nitrate in a nitrogen-polluted city in southern China” by Y. T. Fang et al.***

**S. Morin (Referee)**

morin.samuel@gmail.com

Received and published: 6 October 2010

### **1 General comments**

Fang et al. present a new set of isotopic measurements of rainwater nitrate from an area subjected to significant anthropogenic influence. Concentration,  $\delta^{15}\text{N}$  and  $\delta^{18}\text{O}$  data are provided for two consecutive years (2008 and 2009), allowing the authors to i) attempt to derive seasonal patterns in the variations of  $\delta^{15}\text{N}$  and  $\delta^{18}\text{O}$  and ii) attempt to explore reasons for year-to-year variability. The paper is well written and the methods for chemical, isotopic analyses and their interpretation seem sound and wisely

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



used. My main concerns with this work, detailed below, pertains to the comparability of  $\delta^{15}\text{N}$  from various forms of "airborne reactive nitrogen", i.e.,  $\text{NO}_x$ , particulate or gas-phase nitrate, and nitrate found in precipitation, and the interpretation of  $\delta^{18}\text{O}$  variations through simplified mass-balance concepts.

## 2 Specific comments

### 2.1 $\delta^{15}\text{N}$ from $\text{NO}_x$ , particulate or gas-phase nitrate, and rainwater nitrate

Several publications have shown that  $\delta^{15}\text{N}$  of aerosol nitrate is very different from  $\delta^{15}\text{N}$  of rainwater nitrate at the same site. See e.g. a plot generated from the data presented in Freyer (1991), showing an offset on the order of 9 ‰ between rainwater and aerosol nitrate  $\delta^{15}\text{N}$ . In this case, aerosol  $\delta^{15}\text{N}$  is much higher than rainwater nitrate  $\delta^{15}\text{N}$ . Baker et al. (2007) reached opposite results from remote marine locations. This variable discrepancy was reviewed by Morin et al. (2009) (see in particular section 4.1.3). Caution is thus warranted when comparing  $\delta^{15}\text{N}$  data from different atmospheric matrices. In the ms, the authors often compare their  $\delta^{15}\text{N}$  values obtained from rainwater samples, to aerosol nitrate  $\delta^{15}\text{N}$  values. Before delving further into the interpretation of the data, the authors should make sure the reader is aware of this issue, which is not solved at the moment. This may have a significant impact on the interpretation of the results.

### 2.2 Interpretation of $\delta^{18}\text{O}$ of nitrate

$\delta^{18}\text{O}$  is not an isotopic tracer that is conserved during chemical reactions, in contrast to  $\Delta^{17}\text{O}$  (the isotopic anomaly). Thus interpreting  $\delta^{18}\text{O}$  of nitrate from the contribution of various  $\text{NO}_x$  oxidation pathways featuring different " $\delta^{18}\text{O}$  signatures" is not entirely correct. The reason is that, unlike  $\Delta^{17}\text{O}$ , isotopic fractionation occurring at each step

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

of any chemical mechanism can induce large variations between  $\delta^{18}\text{O}$  values of the reactants and the products (see e.g. Chakraborty and Chakraborty, 2003). The interpretation of seasonal variations of  $\delta^{18}\text{O}$  can therefore not be as detailed as could be the case with  $\Delta^{17}\text{O}$ . This must clearly be realized by the authors, and this point must be made clear to the reader. Along this line, references should be given to support the  $\delta^{18}\text{O}$  values provided for  $\text{O}_3$  and OH (e.g., page 21456, line 5). This may have a significant impact on the interpretation of the results.

### 3 Technical comments

page 21142, line 3 – 5 : to support statements relevant to the chemical oxidation pathways of  $\text{NO}_x$ , one would expect references to standard atmospheric chemistry textbooks, such as Finlayson-Pitts and Pitts or Seinfeld and Pandis, rather than references to studies discussing isotopic measurements, as currently quoted.

page 21442, line 6 : "equilibrates" → "equilibrates during the daytime"

Page 21444, line 27 : "lightning" should not be included in "biogenic emissions"

Page 21446, line 16 : "mass-independent  $^{17}\text{O}$  anomaly" needs reformulation: mass-independent fractionation (i.e., as a process) leads to a  $^{17}\text{O}$  anomaly, but the anomaly itself is not mass-independent, it is just an anomaly.

Page 21446, line 20 : "lowing" → "lowering"

Page 21447, line 11 : the stated lifetimes (1.2 and 0.27) should be supported by a reference. Also, such detailed numbers must refer to very specific conditions. I recommend relaxing a little the accuracy of these numbers to make more general statements (or maybe rather give a range of accepted lifetime values).

Page 21452, line 1 : "(with  $^{15}\text{N}/^{14}\text{N}$  ratio)" needs reformulation. It is not understood

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

what is meant in this parenthesis.

Page 21454, line 15 : the denitrifier method has currently been compared to other methods and showed no systematic bias, contrary to the tentative suspicion of such bias which was presented in Kendall et al. (2007). The relevant references are Chmura et al. (2009) and Xue et al. (2010).

Page 21455, line 29 : the large seasonal difference in  $\delta^{18}\text{O}$  was interpreted by Savarino et al. (2007) as a consequence of the role of stratospheric nitrate injection into the troposphere in late austral winter. Comparing this very different context to the seasonal variations found by the authors should thus be avoided.

Page 21456, line 15 : "This pathway accounts for 4% of the annual inorganic  $\text{NO}_3^-$  on the global scale" : this statement deserves more explanation (reference ? origin of this assessment ? uncertainty ?)

Page 21457, line 9 : Hastings et al., 2003, is probably not an adequate reference to support the  $\delta^{18}\text{O}$  of atmosphere  $\text{O}_2$  (rather : Barkan and Luz, 2003).

Page 21465, Table 1 : Why are arithmetic means of  $\text{NO}_3^-$  presented ? What is the value of such statistics ? The same applies to isotopic ratios. The authors should identify the most relevant kind of averaging method and use it consistently throughout the manuscript. For example, why are arithmetic averages presented in Table 2, rather than mass-weighted averages (which I understand the authors refer to "flux-weighted means", although I may be wrong here) ? To me mass-weighted averaged would make the most sense, especially when dealing with seasonal averages. This dampens the impact of isotopic outliers associated with low concentration levels.

Page 21468, Table 4 : a few typos ("Aersol" → "Aerosol", "Savario" → "Savarino"). Please mention in the captions that isotopic ratios are expressed in ‰.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

## References

Baker, A. R., Westin, K., Kelly, S. D., Voss, M., Streu, P., and Cape, J. N.: Dry and wet deposition of nutrients from the tropical Atlantic atmosphere: links to primary productivity and nitrogen fixation, *Deep-Sea Res. I*, 54, 1704 – 1720, doi:10.1016/j.dsr.2007.07.001, 2007.

Barkan, E. and Luz, B.: High-precision measurements of  $^{17}\text{O}/^{16}\text{O}$  and  $^{18}\text{O}/^{16}\text{O}$  of  $\text{O}_2$  and  $\text{O}_2/\text{Ar}$  ratios in air, *Rapid Commun. Mass Spectrom.*, 17(24), 2809–2814, 2003.

Chakraborty, S. and Chakraborty, S.: Isotopic fractionation of the  $\text{O}_3$ -nitric oxide reaction, *Curr. Sci.*, 85, 1210 – 1212, 2003.

Chmura, W. M., Rozanski, K., Kuc, T., and Gorczyca, Z., Comparison of two methods for the determination of nitrogen and oxygen isotope composition of dissolved nitrates, *Nukleonika*, 54, 17-23, 2009.

Finlayson-Pitts, B. J. and Pitts, J. N.: Chemistry of the upper and lower atmosphere : theory, experiments and applications, Academic Press, San Diego CA, 2000.

Freyer, H. D.: Seasonal variation of  $^{15}\text{N}/^{14}\text{N}$  ratios in atmospheric nitrate species, *Tellus*, 43B, 30–44, 1991.

Kendall, C., Elliott, E. M., and Wankel, S. D.: Stable Isotopes in Ecology and Environmental Science, 2nd edition, chap. 12 : Tracing anthropogenic inputs of nitrogen to ecosystems, Blackwell Publishing, 2007.

Morin, S., Savarino, J., Frey, M. M., Domine, F., Jacobi, H. W., Kaleschke, L., and Martins, J. M. F.: Comprehensive isotopic composition of atmospheric nitrate in the Atlantic Ocean boundary layer from 65°S to 79°N, *J. Geophys. Res.*, 114, D05303, doi:10.1029/2008JD010696, 2009.

Savarino, J., Kaiser, J., Morin, S., Sigman, D. M., and Thiemens, M. H.: Nitrogen and

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



oxygen isotopic constraints on the origin of atmospheric nitrate in coastal Antarctica, *Atmos. Chem. Phys.*, 7, 1925 – 1945, 2007.

Seinfeld, J. and Pandis, S.: *Atmospheric Chemistry and Physics*, Wiley Interscience, 1998.

Xue, D. M., De Baets, B., Botte, J., Vermeulen, J., Van Cleemput, O., and Boeckx, P., Comparison of the silver nitrate and bacterial denitrification methods for the determination of nitrogen and oxygen isotope ratios of nitrate in surface water, *Rapid Communications in Mass Spectrometry*, 24, 833-840, 10.1002/rcm.4445, 2010.

---


[Interactive comment on Atmos. Chem. Phys. Discuss.](#), 10, 21439, 2010.

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

[Discussion Paper](#)



**Fig. 1.** Rain ("pluie") and aerosol  $\delta^{15}\text{N}$  values from locations in Europe, from Freyer (1991), showing the large difference between  $\delta^{15}\text{N}$  of aerosol and rainwater nitrate.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)