

Interactive comment on “Investigating the sensitivity of high-resolution mesoscale models to microphysical parameters by the use of polarimetric radar observations” by R. Ferretti et al.

Anonymous Referee #1

Received and published: 14 October 2010

Review of: "Investigating the sensitivity of high-resolution mesoscale models to microphysical parameters by the use of polarimetric radar observations"

Authors: Ferretti et al.

GENERAL COMMENTS

The manuscript describes a series of sensitivity tests, pertaining to the graupel pa-
C8700

rameters in one-moment bulk microphysics schemes, that have been done with two mesoscale models for 1-km simulations of a hailstorm. Comparison are made to radar reflectivity and hydrometeor type and content derived from polarimetric radar observations from two radars. A considerable amount of work has obviously been done for this study and several different approaches to model-to-radar comparisons have been attempted. Overall, I find the depth of the analysis to be too insufficient to provide much useful insight into the understanding of model sensitivity to microphysics parameters. I believe that the observations, interesting though they are, are not at all useful for this study given that the model simulations are so different from the observations. This is not a criticism of the models or the modellers – it is very difficult to simulate an individual hailstorm. But given this, the comparisons to observations are thus intrinsically problematic for this type of examination of model microphysics.

My primary criticism, however, is the general lack of depth in any of the analysis. A series of sensitivity tests has been performed (following very closely to the study of Gilmore et al. 2004) and the results have been examined in a few different ways, but nearly all of the discussion is essentially just a description of the differences between the results. There is very little discussion, examination, or even speculation as to the reasons for the differences. One of the primary goals of such a study should be to provided recommendations to modellers regarding the best choice of parameter settings. But all this manuscript shows is that the model simulations are sensitive to graupel parameters (which is already known from previous studies). Even with major revisions, I do not believe this manuscript is sufficient for publication.

SPECIFIC COMMENTS

1. It is very unclear why two different mesoscale models have been used. The models give very different solutions for this case and different microphysics schemes have been used. So it is unclear to me how the use of two models helps answer the questions posed regarding the sensitivity to parameter settings for graupel. I suppose one could be looking to see if, given different models with different schemes, the same general

changes in the model solutions occur when the same changes are made to the graupel parameters, but it is not presented this way in the manuscript, at least not clearly. The authors need to do a much better job at explaining why two models are used and showing that this is useful.

2. Although comparison to observations is generally desirable when examining model performance and sensitivities, in this particular study I believe the inclusion of observations does not serve a useful purpose. The simulations using both models are very different from the radar observations, so one cannot make any claims that a given parameter change to the microphysics which leads to a solution that is closer to the observations is thus a change for the better. For example, we have no idea if the vertical motion in the model is accurate, so a change to the microphysics which improves the hydrometeor mass content, compared to the radar estimates, may or may not be due to genuine improvement to the microphysics – but that is what is argued. Unless the control simulations are very close to the observations, which is not the case here, use of radar data for this type of sensitivity study is intrinsically problematic.

3. A comparison of the model storm cell positions from the various sensitivity runs is done, where the cell location is defined as the location of the peak updraft. But what does this show? For one thing, the COSMO storm systems are broad and multicellular, so a change that slightly switches which updraft is stronger does not indicate a change in location of the storm system. More importantly, what does this show? If the location of the peak updraft changes when the assumed bulk graupel density is changed, what are we to take from this?

4. The spectral analysis of the hydrometeor fields confuses me. I have never seen this done before – if it has been, references should be given. I do not understand what is to be learned from this (admittedly, this may be my own fault, but I think it needs to be either better explained or referenced). Perhaps if the model storms were actually similar to the observed storms – which they are not – then we could conclude that the spectra which better match the radar-deduced (not "observed") hydrometeor energy

C8702

spectra are better. But as is, I fail to see how this section addresses the goals of the study.

5. One thing that strikes me as being conspicuously absent from the conclusions of this study is the following. The radar particle type retrieval indicates that there is both hail and large amounts of graupel in this storm. The authors argue that their sensitivity tests, and those of other studies, indicate that better solutions are obtained with more hail-like parameter settings for the graupel category. Surely the obvious conclusion is that a bulk microphysics scheme should include separate categories for graupel and hail.

Interactive comment on Atmos. Chem. Phys. Discuss., 10, 20461, 2010.

C8703