

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Reanalysis of tropospheric sulphate aerosol and ozone for the period 1980–2005 using the aerosol-chemistry-climate model ECHAM5-HAMMOZ

L. Pozzoli^{1,*}, G. Janssens-Maenhout¹, T. Diehl^{2,6}, I. Bey³, M. G. Schultz⁴, J. Feichter^{5,**}, E. Vignati¹, and F. Dentener¹

¹European Commission, Joint Research Centre, Institute for Environment and Sustainability, Ispra, Italy

²NASA Goddard Space Flight Center, Greenbelt, Maryland, USA

³Center for Climate Systems Modeling and Institute of Atmospheric and Climate Science, ETH Zurich, Zurich, Switzerland

⁴Forschungszentrum Jülich, Germany

⁵Max Planck Institute for Meteorology, Hamburg, Germany

⁶University of Maryland Baltimore County, Baltimore, Maryland, USA

* now at: Eurasia Institute of Earth Sciences, Istanbul Technical University, Turkey

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



** now at: ETH, Institute of Atmospheric and Climate Science, Zurich, Switzerland

Received: 10 March 2011 – Accepted: 14 March 2011 – Published: 30 March 2011

Correspondence to: F. Dentener (frank.dentener@jrc.ec.europa.eu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

**Reanalysis
1980–2005
ECHAM5-HAMMOZ**

L. Pozzoli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Understanding historical trends of trace gas and aerosol distributions in the troposphere is essential to evaluate the efficiency of the existing strategies to reduce air pollution and to design more efficient future air quality and climate policies. We performed coupled photochemistry and aerosol microphysics simulations for the period 1980–2005 using the aerosol-chemistry-climate model ECHAM5-HAMMOZ, to assess our understanding of long term changes and inter-annual variability of the chemical composition of the troposphere, and in particular of O₃ and sulphate concentrations, for which long-term surface observations are available. In order to separate the impact of the anthropogenic emissions and meteorology on atmospheric chemistry, we compare two model experiments, driven by the same ECMWF re-analysis data, but with varying and constant anthropogenic emissions, respectively. Our model analysis indicates an average increase of 1 ppbv (corresponding to 0.04 ppbv yr⁻¹) in global average surface O₃ concentrations due to anthropogenic emissions, but this trend is largely masked by natural variability (0.63 ppbv), corresponding to 75% of the total variability (0.83 ppbv). Regionally, annual mean surface O₃ concentrations increased by 1.3 and 1.6 ppbv over Europe and North America, respectively, despite the large anthropogenic emission reductions between 1980 and 2005. A comparison of winter and summer O₃ trends with measurements shows a qualitative agreement, except in North America, where our model erroneously computed a positive trend. O₃ increases of more than 4 ppbv in East Asia and 5 ppbv in South Asia can not be corroborated with long-term observations. Global average sulphate surface concentrations are largely controlled by anthropogenic emissions. Globally natural emissions are an important driver determining AOD variations, regionally AOD decreased by 28% over Europe, while it increased by 19% and 26% in East and South Asia. The global radiative perturbation calculated in our model for the period 1980–2005 was rather small (0.05 W m⁻² for O₃ and 0.02 W m⁻² for total aerosol direct effect), but larger perturbations ranging from -0.54 to 1.26 W m⁻² are estimated in those regions where anthropogenic emissions largely varied.

Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Air quality is determined by the emission of primary pollutants into the atmosphere, by chemical production of secondary pollutants and by meteorological conditions. The two air pollutants of most concern for public health, ozone (O₃) and particulate matter (PM), have also strong impacts on climate. In the fourth Assessment Report of the Intergovernmental Panel on Climate Change (IPCC, AR4), Solomon et al. (2007) estimate a Radiative Forcing (RF) from tropospheric ozone of +0.35 [−0.1, +0.3] W m^{−2}, which corresponds to the third largest contribution to the total RF after carbon dioxide (CO₂) and methane (CH₄). IPCC-AR4 also provides estimates for radiative forcing from aerosols. The direct RF (scattering and absorption of solar and infrared radiation) amounts to −0.5 [±0.4] W m^{−2}, and the RF through indirect changes in cloud properties is estimated to −0.70 [−1.1, +0.4] W m^{−2}.

Trends in global radiation and visibility measurements indeed suggest an important role for aerosol. Solar radiation measurements showed a consistent and worldwide decrease at the Earth's surface (an effect dubbed "dimming") from the 1960s. This trend reversed into "brightening" in the late 1990s in the US, Europe and parts of Korea (Wild, 2009). Similarly, an analysis of visibility measurements from 1973–2007 by Wang et al. (2009) suggests a global increase of AOD worldwide, except in Europe. Since, SO₄^{2−} is one of the main aerosol components that determine the aerosol optical depth (Streets et al., 2009), the SO₄^{2−} concentration reductions over Europe and US after the implementation of air quality policies may partly explain the dimming-brightening transition observed in the 1990s in Europe, whereas in emerging economies such as China and India the emission of air pollutants rapidly increased since 1990. To our knowledge, a consistent global re-analysis of the role of changes in emissions and meteorology has not yet been performed.

Inter-annual meteorological variability in the last decades also strongly determined the variations of the concentrations and geographical distribution of air pollutants. For example, the El Niño event in 1997–1998 and the period after the Mt. Pinatubo volcanic eruption in 1991, can explain much of the past chemical inter-annual variability

**Reanalysis
1980–2005
ECHAM5-HAMMOZ**

L. Pozzoli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of tropospheric O₃, CH₄ and OH (Fiore et al., 2009; Dentener et al., 2003; Hess and Mahowald, 2009). In Europe, the infamous summer of 2003, led to a strong positive anomaly of solar surface radiation (Wild, 2009) and exacerbated ozone pollution at ground-level and throughout the troposphere (Solberg et al., 2008; Tressol et al., 2008).

Meteorological variability and changes in the precursor emissions of O₃ and SO₄²⁻ are often antagonistic processes, and their impact on surface concentrations is difficult to understand from measurements alone (Vautard et al., 2006; Berglen et al., 2007). Therefore, there have been several efforts to re-analyze these trends using tropospheric chemistry and transport models. For example, the European project RE-analysis of the TROpospheric chemical composition (RETRO, Schultz et al., 2007) employed three different global models to simulate tropospheric ozone changes between 1960 and 2000. Recently Hess and Mahowald (2009) analyzed the role of meteorology in inter-annual variability of tropospheric ozone chemistry.

In this paper, we extend these analyses by using a coupled aerosol-chemistry-climate simulation, and discuss our results in the light of the previous studies. We analyze the chemical variability due to changes in meteorology (i.e. transport, chemistry) and natural emissions, and separate them from anthropogenic emissions induced variability. The period 1980–2005 was chosen because the advent of satellite observations in the 1970s introduced a discontinuity in the re-analysis meteorological datasets (Hess and Mahowald, 2009; van Noije et al., 2006) and, as mentioned above, the considered period includes large meteorological anomalies. Particular attention will be given to four regions of the world (North America, Europe, East Asia, and South Asia) where significant changes in terms of the absolute amount of emitted trace gases and aerosol precursors occurred in the last decades. We will focus on past changes of O₃ and SO₄²⁻ because of their importance for air quality and climate. Long measurement records are available since the 1980s and they will be used to evaluate our model results and the simulated trends. We further analyze changes in AOD, radiative perturbation (RP) and OH radical associated with changes in emissions and meteorology.

**Reanalysis
1980–2005
ECHAM5-HAMMOZ**

L. Pozzoli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The paper is organized as follows. In Sect. 2 the model description and experiment setup are outlined. In Sect. 3 anthropogenic and natural emissions are described. Sections 4 and 5 present an analysis of global and regional variability and trends of O₃ and SO₄²⁻ from 1980 to 2005. The regional analysis focuses on, Europe (EU), North America (NA), East Asia (EA), and South Asia (SA) (Fig. 1). In Sect. 6 we will describe the anthropogenic radiative perturbation due to changes in O₃ and SO₄²⁻. In Sects. 7 and 8 we will summarize the main findings and further discuss implication of our study for air quality-climate interactions.

2 Model and simulation descriptions

ECHAM5-HAMMOZ is a fully coupled aerosol-chemistry-climate model, composed of the general circulation model (GCM) ECHAM5, Luca Pozzoli 27 March 2011 10:39 a.m. The ECHAM5-HAMMOZ model is described in detail in Pozzoli et al. (2008a). The model has been extensively evaluated in previous studies (Stier et al., 2005; Pozzoli et al., 2008a,b; Auvray et al., 2007; Rast et al., 2011) with comparisons to several measurements and within model intercomparison studies.

In this study a triangular truncation at wavenumber 42 (T42) resolution was used for the computation of the general circulation. Physical variables are computed on an associated Gaussian grid with ca. 2.8° × 2.8° degrees. The model has 31 vertical levels from the surface up to 10 hPa and a time resolution for dynamics and chemistry of 20 min. We simulated the period 1979–2005 (the first year is discarded from the analysis as spin-up). Meteorology was taken from the ECMWF ERA40 re-analysis (Uppala et al., 2005) until 2000 and from operational analyses (IFS cycle-32r2) for the remaining period (2001–2005). ECHAM5 vorticity, divergence, sea surface temperature, and surface pressure are relaxed towards the re-analysis data every time step with a relaxation time scale of 1 day for surface pressure and temperature, 2 days for divergence, and 6 h for vorticity (Jeuken et al., 1996). The relaxation technique is advantageous to retain the ECHAM5 specific descriptions of clouds and aerosol. The concentrations

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of CO₂ and other GHGs, used to calculate the radiative budget, were set according to the specifications given in Appendix II of the IPCC-TAR report (Nakicenovic et al., 2000). In Appendix A we provide a detailed description of the chemical and micro-physical parameterisations included in ECHAM5-HAMMOZ. A detailed description of the ECHAM5 model can be found in Roeckner et al. (2003).

Two transient simulations were conducted:

- SREF: reference simulation for 1980–2005 where meteorology and emissions are changing on a hourly-to-monthly basis;
- SFIX: simulation for 1980–2005 with anthropogenic emissions fixed at year 1980, while meteorology, natural and wildfire emissions change as in SREF;

3 Emissions

3.1 Anthropogenic emissions

The anthropogenic emissions of CO, NO_x, and VOCs for the period 1980–2000 are taken from the RETRO inventory (<http://retro.enes.org/>) (Schultz et al., 2007; Endresen et al., 2003; Schultz et al., 2008) which provides monthly average emission fields interpolated to the model resolution of 2.8° × 2.8°. In order to prevent a possible drift in CH₄ concentrations with consequences for the simulation of OH and O₃, we prescribed monthly zonal mean CH₄ concentrations in the boundary layer obtained from the interpolation of surface measurements (Schultz et al., 2007). The prescribed CH₄ concentrations vary annually and range from 1520–1650 ppbv (Southern and Northern Hemisphere, respectively SH and NH) in 1980 to 1720–1860 ppbv in 2005 (SH and NH). NO_x aircraft emissions are based on Grewe et al. (2001) and distributed according to prescribed height profiles. The AeroCom hindcast aerosol emission inventory (<http://dataipsl.ipsl.jussieu.fr/AEROCOM/emissions.html>) was used for the annual total anthropogenic emissions of primary black carbon (BC), organic carbon (OC) aerosols

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and sulphur dioxide (SO₂). Specifically, the detailed global inventory of primary BC and OC emissions by Bond et al. (2004) was modified by Streets et al. (2004, 2006) to include additional technologies and new fuel attributes to calculate SO₂ emissions using the same energy drivers as for BC and OC, and extended to the period 1980–2006 (Streets et al., 2009) using annual fuel-use trends and economic growth parameters included in the IMAGE model (National Institute for Public Health and the Environment, 2001). Except for biomass burning, the BC, OC, and SO₂ anthropogenic emissions were provided as annual averages. Primary emissions of SO₄²⁻ are calculated as constant fraction (2.5%) of the anthropogenic sulphur emissions. The SO₂ and primary SO₄²⁻ emissions from international ship traffic for the years 1970, 1980, 1995, and 2001, were taken from the EDGAR 2000 FT inventory (Van Aardenne et al., 2001) and linearly interpolated in time. The emissions of CO, NO_x, VOCs, and SO₂ were available only until the year 2000. Therefore, we used for 2001–2005 the 2000 emissions, except for the regions where significant changes were expected between 2000 and 2005, such as US, Europe, East Asia and South East Asia, for which derived emission trends of CO, NO_x, VOCs, and SO₂ were applied to year 2000. These trends were derived from the USEPA (<http://www.epa.gov/ttn/chief/trends/>), EMEP (<http://www.emep.int/>), and REAS (<http://www.jamstec.go.jp/frcgc/research/p3/emission.htm>) emission inventories over the US, Europe, and Asia, respectively.

Table 1 summarizes the total annual anthropogenic emissions for the years 1980, 1985, 1990, 1995, 2000, and 2005. Global emissions of CO, VOCs and BC were relatively constant during the last decades, with small increases between 1980 and 1990, decreases in the 1990s and renewed increase between 2000 and 2005. During 25 yr, global NO_x and OC emissions increased up to 10%, while sulfur emissions decreased by 10%. However, these global numbers mask that the global distribution of the emission largely changed, with reductions over North America and Europe, balanced by strong increases in the economically emerging countries, such as China and India. Figure 2 shows the relative trends of anthropogenic emissions used during this study over the 4 selected world regions. In Europe and North America there is a general

**Reanalysis
1980–2005
ECHAM5-HAMMOZ**

L. Pozzoli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



decrease of emissions of all pollutants from 1990 on. In East Asia and South Asia anthropogenic emissions generally increase although for EA there is a decrease around 2000.

3.2 Natural and biomass burning emissions

5 Some O₃ and SO₄²⁻ precursors are emitted by natural processes, which exhibit inter-annual variability due to changing meteorological parameters (temperature, wind, solar radiation, clouds and precipitation). For example, VOC emissions from vegetation are influenced by surface temperature and short wavelength radiation, NO_x is produced by lightning and associated with convective activity, dimethyl sulphide (DMS) emissions
10 depend on the phytoplankton blooms in the oceans and wind speed, and some aerosol species, such as mineral dust and sea salt, are strongly dependent on surface wind speed. Inter-annual variability of weather will therefore influence the concentrations of tropospheric O₃ and SO₄²⁻ and may also affect the radiative budget. The emissions from vegetation of CO and VOCs (isoprene and terpenes) were calculated interactively
15 using the Model of Emissions of Gases and Aerosols from Nature (MEGAN) (Guenther et al., 2006). The total annual natural emissions in Tg(C) of CO and biogenic VOCs (BVOCs) for the period 1980–2005 range from 840 to 960 Tg(C) yr⁻¹ with a standard deviation of 26 Tg(C) yr⁻¹ (Fig. 3a), which corresponds to the 3% of annual mean natural VOC emissions for the considered period. 80% of these total biogenic emissions
20 occur in the tropics.

Lightning NO_x emissions (Fig. 3b) are calculated following the parameterization of Grewe et al. (2001). We calculated a 5% variability for NO_x emissions from lightning, from 3.55 to 4.25 Tg(N) yr⁻¹, with a decreasing trend of 0.017 Tg(N) yr⁻¹ (*R*² of 0.53; 95% confidence bounds ±0.007). We note here that there are considerable uncertainties in the parameterization for NO_x emissions (e.g., Grewe et al., 2001; Tost et al.,
25 2007; Schumann and Huntrieser, 2007), and different parameterizations simulated opposite trends over the same period (Schultz et al., 2007).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



DMS, predominantly emitted from the oceans, is a SO_4^{2-} aerosol precursor. Its emissions depend on seawater DMS concentrations associated with phytoplankton blooms (Kettle and Andreae, 2000) and model surface wind speed, which determines the DMS sea-air exchange Nightingale et al. (2000). Terrestrial biogenic DMS emissions follow Pham et al. (1995). The total DMS emissions are ranging from 22.7 to 24.4 Tg(S) yr^{-1} , with a standard deviation of 0.3 Tg(S) yr^{-1} or 1% of annual mean emissions (Fig. 3c). Again the highest DMS emissions correspond to the 1997–1998 ENSO event. Since we have used a climatology of seawater DMS concentrations instead of annually varying concentrations, the variability may be misrepresented.

The emissions of sea salt are based on Schulz et al. (2004). The strong function of wind speed dependency results in a range from 5000–5550 Tg yr^{-1} (Fig. 3e) with a standard deviation of 2%.

Mineral dust emissions are calculated online using the ECHAM5 wind speed, hydrological parameters (e.g. soil moisture), and soil properties, following the work of Tegen et al. (2002) and Cheng et al. (2008). The total mineral dust emissions have a large inter-annual variability (Fig. 3d), ranging from 620 to 930 Tg yr , with a standard deviation of 72 Tg yr , almost 10% of the annual mean average. Mineral dust originating from the Sahara and over Asia contribute on average 58% and 34%, of the global mineral dust emissions, respectively.

Biomass burning, from tropical savannah burning, deforestation fires, and mid-and high latitude forest fires are largely linked to anthropogenic activities but fire severity (and hence emissions) are also controlled by meteorological factors such as temperature, precipitations, and wind. We use the compilation of inter-annual varying biomass burning emissions published by (Schultz et al., 2008), which used literature data, satellite observations and a dynamical vegetation model in order to obtain continental-scale emission estimates and a geographical distribution of fire occurrence. We consider emissions of the components CO , NO_x , BC , OC , and SO_2 , and apply a time-invariant vertical profile of the plume injection height for forest and savannah fire emissions. Figure 3f shows the inter-annual variability for CO , NO_x , and OC biomass burning

**Reanalysis
1980–2005
ECHAM5-HAMMOZ**

L. Pozzoli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



emissions, with different peaks, such as in year 1998 during the strong ENSO episode.

Other natural emissions are kept constant during the entire simulation period. CO emissions from soil and ocean are based on the Global Emission Inventory Activity (GEIA) as in Horowitz et al. (2003), amounting to 160 and 20 Tg yr⁻¹, respectively. Natural soil NO_x emissions are taken from the ORCHIDEE model (Lathiere et al., 2006), resulting in 9 Tg(N) yr⁻¹. SO₂ volcanic emissions of 14 Tg(S) yr⁻¹ from Andres and Kasgnoc (1998); Halmer et al. (2002). Since in the current model version secondary organic aerosol (SOA) formation is not calculated online, we applied a monthly varying OC emission (19 Tg yr) to take into account the SOA production from biogenic monoterpenes (a factor of 0.15 is applied to monoterpene emissions of Guenther et al., 1995), as recommended in the AEROCOM project (Dentener et al., 2006).

4 Variability and trends of O₃ and OH during 1980–2005 and their relationship to meteorological variables

In this section we analyze the global and regional variability of O₃ and OH in relation to selected modeled chemical and meteorological variables. Section 4.1 will focus on global surface ozone, Sect. 4.2 will look into more detail to regional differences in O₃, and for Europe and the US, compare the data to observations. Section 4.3 will describe the variability of the global ozone budget, and Sect. 4.4 will focus on the related changes in OH. As explained earlier, we will separate the influence of meteorological and natural emission variability from anthropogenic emissions by differencing the SREF and SFIX simulations.

4.1 Global surface ozone and relation to meteorological variability

In Fig. 4 we show the ECHAM5-HAMMOZ SREF inter-annual monthly anomalies (difference between the monthly value and the 25-yr monthly average) of global mean surface temperature, water vapor, and surface concentrations of O₃, SO₄²⁻, total column

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



AOD and methane-weighted OH tropospheric concentrations will be discussed in later sections. To better visualize the inter-annual variability over 1980–2005, in Fig. 4, we also display the 12-months running average of monthly mean anomalies for both SREF (blue) and SFIX (red) simulations.

5 Surface temperature evolution showed large anomalies in the last decades, associated with major natural events. For example, large volcanic eruptions (El Chichon, 1982; Pinatubo, 1991) generated cooler temperatures due to the emission into the stratosphere of sulphate aerosols. The 1997–1998 ENSO caused ca. 0.4 K elevated temperatures. The strong coupling of the hydrological cycle and temperature is reflected in a correlation of 0.83 between the monthly anomalies of global surface temperature and water vapour content. These two meteorological variables influence O₃ and OH concentrations. For example the large 1997–1998 ENSO event corresponds with a positive ozone anomaly of more than 2 ppbv. There is also an evident correspondence between lower ozone and cooler periods in 1984 and 1988. However, the correlation between monthly temperature and O₃ anomalies (in SFIX) is only moderate (R = 0.43).

10 The 25-yr global surface O₃ average is 36.45 ppbv (Table 2) and increased by 0.48 ppbv compared to the SFIX simulation, with year 1980 constant anthropogenic emissions. About half of this increase is associated with anthropogenic emission changes (Fig. 4c (grey area)). The inter-annual monthly surface ozone concentrations varied by up to ±2.17 ppbv (1σ = 0.83 ppbv), of which 75% (0.63 ppbv in SFIX) was related to natural variations- especially the 1997–1998 ENSO event.

4.2 Regional differences in surface ozone trends, variability, and comparison to measurements

25 Global trends and variability may mask contrasting regional trends. Therefore we also perform a regional analysis for North America (NA), Europe (EU), East Asia (EA) and South Asia (SA) (see Fig. 1). To quantify the impact on surface concentrations after 25 yr of changing anthropogenic emission, we compare the averages of two 5-yr

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



periods, 1981–1985 and 2001–2005. We considered 5-yr averages to reduce the noise due to meteorological variability in these two periods.

In Fig. 5 we provide maps for the globe, Europe, North America, East Asia and South Asia (rows), showing in the first column (a) the reference surface concentrations, and in the other 3 columns relative to the period 1981–1985, the isolated effect of anthropogenic emission changes (b), meteorological changes (c), and combined meteorological and emissions changes (d). The global maps provide insight into inter-regional influences of concentrations.

A comparison to observed trends provides additional insight into the accuracy of our calculations (Fig. 6). This comparison, however, is hampered by the lack of observations before 1990, and the lack of long-term observations outside of Europe and North America. We will therefore limit the comparison to the period 1990–2005, separately for winter (DJF) and summer (JJA), acknowledging that some of the larger changes may have happened before. Figure 1 displays the measurement locations: we used 53 stations in North America and 98 stations in Europe. To allow a realistic comparison with our coarse-resolution model, we grouped the measurement in 5 subregions for EU and 5 subregions for NA. For each subregion we calculated the trend of the median winter and summer anomalies. In Appendix B we give an extensive description of the data used for these summary figures, and their statistical comparison with model results.

We note here that O_3 and SO_4^{2-} in ECHAM5-HAMMOZ were extensively evaluated in previous studies (Stier et al., 2005; Pozzoli et al., 2008a,b; Rast et al., 2011), showing in general a good agreement between calculated and observed SO_4^{2-} and an overestimation of surface O_3 concentrations in some regions. We implicitly assume that this model bias does not influence the calculated variability and trends, an assumption that will be discussed further in the discussion section.

**Reanalysis
1980–2005
ECHAM5-HAMMOZ**L. Pozzoli et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

4.2.1 Europe

In Fig. 5e we show the SREF 1981–1985 annual mean EU surface O_3 (i.e. before large emission changes) corresponding to an EU average concentration of 46.9 ppbv. High annual average O_3 concentrations up to 70 ppbv are found over the Mediterranean basin, and lower concentrations between 20 to 40 ppbv in Central and Eastern Europe. Figure 5f shows the difference in mean O_3 concentrations between the SREF and SFIX simulation for the period 2001–2005. The decline of NO_x and VOC anthropogenic emissions was 20% and 25%, respectively (Fig. 2a). Annual averaged surface O_3 concentration augments by 0.81 ppbv between 1981–1985 and 2001–2005. The spatial distribution of the calculated trends is very different over Europe, computed O_3 increased between 1 and 5 ppbv over Northern and Central Europe, while it decreased by up to 1–5 ppbv over Southern Europe. The O_3 responses to emission reductions is driven by complex nonlinear photochemistry of the O_3 , NO_x and VOC system. Indeed, we can see very different O_3 winter and summer sensitivities in Figs. 7a and 8. The increase (7% compared to year 1980) in annual O_3 surface concentration is mainly driven by winter (DJF) values, while in summer (JJA) there is a small 2% decrease between SREF and SFIX. This winter NO_x titration effect on O_3 is particularly strong over Europe (and less over other regions), as was also shown in e.g. Fig. 4 of Fiore et al. (2009), due to the relatively high NO_x emission density, and the mid-to-high latitude location of Europe. The impact of changing meteorology and natural emissions over Europe is shown in Fig. 5g, showing an increase by 0.37 ppbv between 1981–1985 and 2001–2005. Winter-time variability drives much of the inter-annual variability of surface O_3 concentrations (see Figs. 7a and 8a). While it is difficult to attribute the relationship between O_3 and meteorological conditions to a single process, we speculate that the European surface temperature increase of 0.7 K 1981–1985 to 2001–2005 could play a significant role. Figure 5d, 5h, and 5l show that North Atlantic ozone increased during the same period, contributing to the increase of the baseline O_3 concentrations at the western border of EU. Figure 5f and 5g show that both emissions and meteorological

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



variability may have synergistically caused upward trends in Northern and Central Europe, and downward trends in Southern Europe.

The regional responses of surface ozone can be compared to the HTAP multi-model emission perturbation study of Fiore et al. (2009), which considered identical regions and are thus directly comparable. The combined impact of the HTAP 20% emission reduction were resulting in an increase of O_3 by 0.2 ppbv in winter, and an O_3 decrease by -1.7 ppbv in summer (average of 21 models), to a large extent driven by NO_x emissions. Considering similar annual mean reductions in NO_x and VOC emissions over Europe from 1980–2005, we found a stronger emission driven increase of up to 3 ppbv in winter, and a decrease of up to 1 ppbv in summer. An important difference between the two studies may be the use of spatially homogeneous emission reduction in HTAP, while the emissions used here, generally included larger emission reductions in Northern Europe while emissions in Mediterranean countries remained more or less constant.

The calculated and observed O_3 trends for the period 1990–2005 are relatively small compared to the inter-annual variability. In winter (DJF) (Fig. 6) measured trends confirm increasing ozone in most parts of Europe. The observed trends are substantially larger (0.3 – 0.5 ppbv yr^{-1}) than the model results (0 – 0.2 ppbv yr^{-1}), except in Western Europe (WEU). In summer (JJA) the agreement of calculated and observed trends is small: in the observations they are close to zero for all European regions with large 95% confidence intervals, while calculated trends show significant decreases (0.1 – 0.45 ppbv yr^{-1}) of O_3 . Despite seasonal O_3 trends are not well captured by the model, the seasonally averaged modeled and measured surface ozone concentrations are generally reasonably well correlated (Appendix B, $0.5 < R < 0.9$). In winter the simulated inter-annual variability seemed to be somewhat underestimated, pointing to missing variability coming from e.g. stratosphere-troposphere or long-range transport. In summer the correlations are generally increasing for Central Europe and decreasing for Western Europe.

**Reanalysis
1980–2005
ECHAM5-HAMMOZ**L. Pozzoli et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

4.2.2 North America

5 Computed annual mean surface O_3 over North America (Fig. 5i), for 1981–1985 was 48.3 ppbv. Higher O_3 concentrations are found over California and in the continental outflow regions, Atlantic and Pacific Oceans, and the Gulf of Mexico, and lower concentrations north of 45° N. Anthropogenic NO_x in NA decreased by 17% between 1980 and 2005, particularly in the 1990s (-22%) (Fig. 2b). These emission reductions produced an annual mean O_3 concentration decrease up to 1 ppbv over all Eastern US, 1–2 ppbv over the Southern US, and 1 ppbv in the western US. Changes in anthropogenic emissions from 1980 to 2005 (Fig. 5j) resulted in a small average increase of ozone by 0.28 ppbv over NA, where effects on O_3 of emission reductions in the US and Canada were balanced by higher O_3 concentrations mainly over the tropics (below 25° N). Changes in meteorology (Fig. 5k) increased O_3 between 1 and 5 ppbv (average 0.89 ppbv) over the continent, and reductions in West Mexico and the Atlantic Ocean O_3 by up to 5 ppbv. Thus meteorological variability was the largest driver of the overall average regional increase of 1.58 ppbv in SREF (Fig. 5l). Over NA meteorological variability is the main driver of summer (JJA) O_3 fluctuations from -7% to 5% (Fig. 8b). In winter (Fig. 7b), the contribution of emissions and chemistry is strongly influencing these relative changes. Computed and measured summer concentrations were better correlated than those in winter (Appendix B). However, while in winter an analysis of the observed trends seems to suggest $0\text{--}0.2\text{ ppbv yr}^{-1}$ O_3 increases, the model rather predicts small O_3 decreases. However, often these differences are not significant (see also Appendix B). In summer modelled upward trends (SREF) are not confirmed by measurements, except for Western US (WUS). These computed upward trends were strongly determined by the large-scale meteorological variability (SFIX), and the model trends solely based on anthropogenic emission changes (SREF-SFIX) would be more consistent with observations. We speculate that the role of, and the meteorological feedbacks on, natural emissions may be too strongly represented in our model in North America, but process study would be needed to confirm this hypothesis.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4.2.3 East Asia

In East Asia, we calculated an annual mean surface O_3 concentration of 43.6 ppbv for 1981–1985. Surface O_3 is less than 30 ppbv over north-eastern China, and influenced by continental outflow conditions, up to 50 ppbv concentrations are computed over the Northern Pacific Ocean and Japan (Fig. 5m). Over EA, anthropogenic NO_x and VOC emissions increased by 125% and 50%, respectively, from 1980 to 2005. Between 1981–1985 and 2001–2005 O_3 is reduced by 10 ppbv in North Eastern China, due to reaction with freshly emitted NO. In contrast, O_3 concentrations increase close to the China coast by up to 10 ppbv, and up to 5 ppbv over the entire north Pacific, reaching North America (Fig. 5b). For the entire EA region (Fig. 5n) we found an increase of annual mean O_3 concentrations of 2.43 ppbv. The effect of meteorology and natural emissions is generally significantly positive, with an EA-wide increase of 1.6 ppbv, up to 5 ppbv in northern and southern continental EA (Fig. 5o). The combined effect of anthropogenic emissions and meteorology is an increase of 4.13 ppbv in O_3 concentrations (Fig. 5p). During this period, the seasonal mean O_3 concentrations were increasing by 3% and 9% in winter and summer, respectively (Figs. 7c and 8c). The effect of meteorology on the seasonal mean O_3 concentrations shows opposite effects in winter and summer: a reduction between 0 and 5% in winter, and an increase between 0 and 10% in summer. The 3 long-term measurement datasets at our disposal (not shown) indicate large inter-annual variability of O_3 and no significant trend in the time period from 1990 to 2005, therefore not plotted in Fig. 6.

4.2.4 South Asia

Of all 4 regions, the largest relative change in anthropogenic emissions occurred over SA: NO_x emissions increased by 150%, VOC 60%, and sulphur 220%. South Asian O_3 inter-annual variability is rather different from EA, NA, and EU, because the SA region is almost completely situated in the tropics. Meteorology is highly influenced by the Asian monsoon circulation, with the wet season in June–August. We calculate

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



an annual mean surface O_3 concentration of 48.2 ppbv, with values between 45 and 60 ppbv over the continent (Fig. 5q). Note that the high concentrations at the northern edge of the region may be influenced by the orography of the Himalaya. The increasing anthropogenic emissions enhanced annual mean surface O_3 concentrations by on average 4.24 ppbv (Fig. 5r), and more than 5 ppbv over India and the Gulf of Bengal. In the NH winter (dry season) the increase in O_3 concentrations of up to 10% due to anthropogenic emissions is more pronounced than in summer (wet season), 5% in JJA (Figs. 7d and 8d). The effect of meteorology produced an annual mean O_3 increase of 1.15 ppbv over the region and more than 2 ppbv in the Ganges valley and in the southern Gulf of Bengal (Fig. 5s). The total (emissions and meteorology) variability in seasonal mean O_3 concentrations is in the order of 5%, both in winter and summer (Figs. 7d and 8d). In 25 yr, the computed annual mean O_3 concentrations increased by 5.12 ppbv over SA, approximately 75% of which are related to increasing anthropogenic emissions (Fig. 5t). Unfortunately, to our knowledge no such long-term data of sufficient quality exist in India. We further remark the substantial overestimate of our computed ozone compared to measurements, a problem that ECHAM shares with many other global models, we refer for a further discussion to Ellingsen et al. (2008).

4.3 Variability of the global ozone budget

We will now discuss the changes in global tropospheric O_3 . To put our model results in a multi-model context, we show in Fig. 9 the global tropospheric O_3 budget along with budget terms derived from Stevenson et al. (2006). O_3 budget terms were calculated using an assumed chemical tropopause, with a threshold of 150 ppbv of O_3 . The annual globally integrated chemical production (P), loss (L), surface deposition (D), and stratospheric influx terms are well in the range of those reported by Stevenson et al. (2006), though O_3 burden and lifetime are at the high. In our study, the variability in production and loss are clearly determined by meteorological variability, with the 1997–1998 ENSO event standing out. The increasing turnover of tropospheric ozone manifests in gradually decreasing ozone lifetimes (–1 day from 1980 to 2005), while

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



total tropospheric ozone burden increases from 370 to 380 Tg, caused by increasing production and stratospheric influx.

Further, we compare our work to a re-analysis study by Hess and Mahowald (2009), which focused on the relationship between meteorological variability and ozone. Hess and Mahowald (2009) used the chemical transport model (CTM) MOZART2 to conduct two ozone simulations from 1979 to 1999 without considering the inter-annual changes in emissions (except for lightning emissions) and is thus very comparable to our SFIX simulation. The simulations were driven by two different re-analysis methodologies: the National Center for Environmental Prediction/National Center for Atmospheric Research (NCEP/NCAR) re-analysis; the output of the Community Atmosphere Model (CAM3, Collins et al., 2006), driven by observed sea surface temperatures (SNCEP and SCAM in Hess and Mahowald (2009), respectively). Comparison of our model (in particular the SFIX simulation) driven by the ECMWF ERA40 reanalysis with Hess and Mahowald (2009) provides insight in the extent to which these different approaches impact the inter-annual variability of ozone. In Table 3 we compare the results of SFIX with the two Hess and Mahowald (2009) model results. We excluded the last 5 yr of our SFIX simulation in order to allow direct statistical comparison with the period 1980–2000.

4.3.1 Hydrological cycle and lightning

The variability of photolysis frequencies of NO_2 (J_{NO_2}) at the surface is an indicator for overhead cloud cover fluctuations, with lower values corresponding to larger cloud cover. J_{NO_2} values are ca. 10% lower in our SFIX (ECHAM5) compared to the two simulations reported by Hess and Mahowald (2009). This may be due to a different representation of the cloud impact on photolysis frequencies (in presence of a cloud layer lower rates at surface and higher rates above) as calculated in our model using Fast-J.2 (see Appendix A) compared to the look-up-tables used by Hess and Mahowald (2009). Furthermore, we found 22% higher average precipitation and 37% higher tropospheric water vapor in ECHAM5, than reported for the NCEP and CAM

**Reanalysis
1980–2005
ECHAM5-HAMMOZ**

L. Pozzoli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



re-analyses, respectively. As discussed by Hagemann et al. (2006), ECHAM5 humidity may be biased high regarding processes involving the hydrological cycle, especially in the NH summer in the tropics. The computed production of NO_x from lightning (LNO) of 3.91 Tg yr in our SFIX simulation resides between the values found in the NCEP- and CAM-driven simulations of Hess and Mahowald (2009), despite the large uncertainties of lightning parameterizations (see Sect. 3.2).

4.3.2 O_3 , CO and OH

The global multi annual averages of tropospheric O_3 are very similar in the 3 simulations, ranging from 46 to 48.4 ppbv, while 20% higher values are found for surface O_3 in our SFIX simulation. Our global tropospheric average of CO concentrations is 15% higher than in Hess and Mahowald (2009), probably due to different biogenic CO and VOCs emissions. Despite higher water vapor, and lower surface J_{NO_2} in SFIX, our calculated OH tropospheric concentrations are smaller by 15%. Since a multitude of factors can influence tropospheric OH abundances, interpreting the differences among the three re-analysis approaches is beyond the subject of this paper.

4.3.3 Variability, re-analysis and nudging methods

A remarkable difference with Hess and Mahowald (2009) is the higher inter-annual variability of global ozone, CO, OH, and HNO_3 as simulated in our model, compared to that found in the CAM-driven simulation analysis. This likely indicates that the “mild” nudging (only forcing through monthly averaged sea-surface temperatures) incorporates only partially the processes that govern inter-annual variations in the chemical composition of the troposphere. The variability of OH (calculated as the relative standard deviation, RSD) in our SFIX simulation is higher by a factor of 2 than those in SCAM and SNCEP, and CO, HNO_3 up to a factor of 10. We speculate that these differences point to differences in the hydrological cycle among the models, which influence OH through changes in cloud cover and HNO_3 through different washout rates.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A similar conclusion was reached by Auvray et al. (2007), who analyzed ozone formation and loss rates from the ECHAM5-MOZ and GEOS-CHEM models for different pollution conditions over the Atlantic Ocean. Since the methodology used in our SFIX simulation should be rather comparable to that used for the NCEP-driven MOZART2 simulation, we speculate that in addition to the differences in re-analysis (NCEP/NCAR and ECMWF/ERA40), also different nudging methodologies may strongly impact the calculated inter-annual variabilities.

4.4 OH variability

To estimate the changes in the global oxidation capacity, we calculated the global mean tropospheric OH concentration weighted by the reaction coefficient of CH_4 , following Lawrence et al. (2001). We found a mean value of $1.2 \pm 0.016 \times 10^6$ molecules cm^{-3} in the SREF simulation (Table 2, within the $1.06\text{--}1.39 \times 10^6$ molecules cm^{-3} range calculated by Lawrence et al., 2001). In Fig. 4d we show the global tropospheric average monthly mean anomalies of OH. During the period 1980–2005 we found a decreasing OH trend of -0.33×10^4 molecules cm^{-3} ($R^2 = 0.79$ due to natural variability), balanced by an opposite trend of 0.30×10^4 molecules cm^{-3} ($R^2 = 0.95$) due to anthropogenic emission changes. In agreement with an earlier study by Fiore et al. (2006) we found a strong relationship between global lightning and OH inter-annual variability with a correlation of 0.78. This correlation drops to 0.32 for SREF, which additionally includes the effect of changing anthropogenic CO, VOC and NO_x emissions. The resulting OH inter-annual variability of 2% for the impact of meteorology (SFIX) was close to the estimate of Hess and Mahowald (2009) (for the period 1980–2000, Table 3), and much smaller than the 10% variability estimated by Prinn et al. (2005), but somewhat higher than the global inter-annual variability of 1.5% analyzed by Dentener et al. (2003). Latter authors however did not include inter-annual varying biomass burning emissions. Dentener et al. (2003) with a different model computed an increasing trend of $0.24 \pm 0.06\% \text{yr}^{-1}$ in OH global mean concentrations for the period 1979–1993, which was mainly caused by meteorological variability. For a slightly

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



shorter period (1980–1993), we found a decreasing trend of $-0.27\% \text{ yr}^{-1}$ in our SFIX run. Montzka et al. (2011) estimate an inter-annual variability of OH in the order of $2 \pm 1.8\%$ for the period 1985–2008, which compares reasonably well with our results of 1.6% and 2.4% for the SREF and SFIX runs (1980–2005, Table 2), respectively. The calculated OH decline from 2001 to 2005, which was not strongly correlated to global surface temperature, humidity or lightning, could have implications for the understanding of the stagnation of atmospheric methane growth during the first part of 2000s. However, we do not want to over-interpret this decline, since in this period we used meteorological data from the operational ECMWF analysis instead of ERA40 (Sect. 2). On the other hand we have no evidence of other discontinuities in our analysis, and the magnitude of the OH changes was similar to earlier changes in the period 1980–1995.

5 Surface and column SO_4^{2-}

In this section we analyze global and regional surface sulphate (Sect. 5.1) and the global sulphate budget (Sect. 5.2) including their variability. The regional analysis and comparison with measurements follows the approach of the ozone analysis above.

5.1 Global and regional surface sulphate

Global average surface SO_4^{2-} concentrations are 1.12 and $1.18 \mu\text{g m}^{-3}$ for the SREF and SFIX runs, respectively (Table 2). Anthropogenic emission changes induce a decrease of ca. $0.1 \mu\text{g m}^{-3}$ SO_4^{2-} between 1980 and 2005 in SREF (Fig. 4e). Monthly anomalies of SO_4^{2-} surface concentrations range from -0.1 , to $0.2 \mu\text{g m}^{-3}$. The 12-month running averages of monthly anomalies are in the range of $\pm 0.1 \mu\text{g m}^{-3}$ for SREF, and about half of this in the SFIX simulation. The anomalies in global average SO_4^{2-} surface concentrations do not show a significant correlation with meteorological variables on the global scale.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5.1.1 Europe

For 1981–1985 we compute an annual average SO_4^{2-} surface concentration of $2.57 \mu\text{g(S)} \text{m}^{-3}$ (Fig. 10e), with the largest values over the Mediterranean and Eastern Europe. Emission controls (Fig. 2a) reduced SO_4^{2-} surface concentrations (Figs. 10f, 11a, and 12a) by almost 50% in 2001–2005. Figure 10f and g show that emission reductions and meteorological variability contribute ca. 85% and 15% respectively, to the overall differences between 2001–2005 and 1981–1985, indicating a small but significant role for meteorological variability in the SO_4^{2-} signal. Figure 10h shows that the largest SO_4^{2-} decreases occurred in South and North East Europe. In Figs. 11a and 12a we display seasonal differences in the SO_4^{2-} response to emissions and meteorological changes. SFIX European winter (DJF) surface sulphate varies $\pm 30\%$ compared to 1980, while in summer (JJA) concentrations are between 0 and 20% larger than in 1980. The decline of European surface sulphur concentrations in SREF is larger in winter (50%) than in summer (37%).

Measurements mostly confirm these model findings. Indeed, inter-annual seasonal anomalies of SO_4^{2-} in winter (Appendix B) generally correlate well ($R > 0.5$) at most stations in Europe. In summer, the modelled inter-annual variability is always underestimated (normalized standard deviation of 0.3–0.9), most likely indicating an underestimate in the variability of precipitation scavenging in the model over Europe. Modeled and measured SO_4^{2-} trends are in good agreement in most European regions (Fig. 6). In winter the observed declines of $0.02\text{--}0.07 \mu\text{g(S)} \text{m}^{-3} \text{yr}^{-1}$ are underestimated in NEU and EEU, and overestimated in the other regions. In summer the observed declines of $0.02\text{--}0.08 \mu\text{g(S)} \text{m}^{-3} \text{yr}^{-1}$ are overestimated in SEU, underestimated in CEU, WEU, and EEU.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5.1.2 North America

The calculated annual mean surface concentration of sulphate over the NA region for the period 1981–1985 is $0.65 \mu\text{g}(\text{S})\text{m}^{-3}$. Highest concentrations are found over the Eastern and Southern US (Fig. 10i). NA emissions reductions of 35% (Fig. 2b) reduced SO_4^{2-} concentrations on average by $0.18 \mu\text{g}(\text{S})\text{m}^{-3}$, and up to $1 \mu\text{g}(\text{S})\text{m}^{-3}$ over the Eastern US (Fig. 10j). Meteorological variability results in a small overall increase of $0.05 \mu\text{g}(\text{S})\text{m}^{-3}$ (Fig. 10k). Changes in emissions and meteorology can almost be combined linearly (Fig. 10l). The total decline is thus $0.11 \mu\text{g}(\text{S})\text{m}^{-3}$: -20% in winter and -25% in summer between 1980–2005, indicating a fairly low seasonal dependency.

Like in Europe, also in North America measured inter-annual variability is smaller than in our calculations in winter and larger in summer. Observed winter downward trends are in the range of $0\text{--}0.03 \mu\text{g}(\text{S})\text{m}^{-3}\text{yr}^{-1}$ in reasonable agreement with the range of $0.01\text{--}0.06 \mu\text{g}(\text{S})\text{m}^{-3}\text{yr}^{-1}$ calculated in the SREF simulation. In summer, except for Western US (WUS), observed SO_4^{2-} trends range between $0.05\text{--}0.08 \mu\text{g}(\text{S})\text{m}^{-3}\text{yr}^{-1}$, the calculated trends decline only between $0.03\text{--}0.04 \mu\text{g}(\text{S})\text{m}^{-3}\text{yr}^{-1}$ (Fig. 6). We suspect that a poor representation of the seasonality of anthropogenic sulfur emissions contributes to both the winter overestimate and the summer underestimate of the SO_4^{2-} trends.

5.1.3 East Asia

Annual mean SO_4^{2-} during 1981–1985 was $0.9 \mu\text{g}(\text{S})\text{m}^{-3}$ with higher concentrations over eastern China, Korea, and in the continental outflow over the Yellow Sea (Fig. 10m). Growing anthropogenic sulfur emissions (60% over EA) produced an increase in regional annual mean SO_4^{2-} concentrations of $0.24 \mu\text{g}(\text{S})\text{m}^{-3}$ in the 2001–2005 period. Largest increases (practically a doubling of concentrations) were found over eastern China and Korea (Fig. 10n). The contribution of changing meteorology and natural emissions is relatively small ($0.01 \mu\text{g}(\text{S})\text{m}^{-3}$ or 15%, Fig. 10o), and the

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



coupled effect of changing emissions and meteorology is dominated by the emissions perturbation ($0.19 \mu\text{g(S)} \text{m}^{-3}$, Fig. 10p).

5.1.4 South Asia

Annual and regional average SO_4^{2-} surface concentrations of $0.70 \mu\text{g(S)} \text{m}^{-3}$ (1981–1985) increased by on average $0.38 \mu\text{g(S)} \text{m}^{-3}$, and up to $1 \mu\text{g(S)} \text{m}^{-3}$ over India, due to the 220% increase in anthropogenic sulfur emissions in 25 yr. The alternation of wet and dry seasons, is greatly influencing the seasonal SO_4^{2-} concentration changes. In winter (dry season), following the emissions, the SO_4^{2-} concentrations increase two-fold. In the wet season (JJA) we do not see a significant increase in SO_4^{2-} concentrations and the variability is almost completely dominated by the meteorology (Figs. 11d and 12d). This indicates that frequent rainfall in the monsoon circulation keeps SO_4^{2-} low regardless of increasing emissions. In winter we calculated a ratio of 0.45 between SO_4^{2-} wet deposition and total SO_4^{2-} production (both gas and liquid phases), while during summer months about 1.24 times more sulphur is deposited in the SA region than is produced.

5.2 Variability of the global SO_4^{2-} budget

We now analyze in more detail the processes that contribute to the variability of surface and column sulphate. Figure 13 shows that inter-annual variability of the global SO_4^{2-} burden is largely determined by meteorology in contrast to the surface SO_4^{2-} changes discussed above. In-cloud SO_2 oxidation processes with O_3 and H_2O_2 do not change significantly over 1980–2005 in the SFIX simulation ($47.2 \pm 0.4 \text{Tg(S)} \text{yr}^{-1}$), while in the SREF case the trend is very similar to those of the emissions. Interestingly, the H_2SO_4 (and aerosol) production resulting from the SO_2 reaction with OH (Fig. 13c), responds differently to meteorological and emission variability than in-cloud oxidation. Globally it increased by 1Tg(S) between 1980 and late 1990s (SFIX), and

**Reanalysis
1980–2005
ECHAM5-HAMMOZ**

L. Pozzoli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



then decreased again after year 2000 (see also Fig. 4e). The increase of gas phase SO_4^{2-} production (Fig. 13c) in the SREF simulation is even more striking in the context of overall declining emissions. These contrasting temporal trends can be explained by the changes in the geographical distribution of the global emissions and the variation of the relative efficiency of the oxidation pathways of SO_2 in SREF, which are given in Table 4. Thus the global increase in SO_4^{2-} gaseous phase production is disproportionately depending on the sulphur emissions over Asia, as noted earlier by e.g. Unger et al. (2009). For instance in the EU region the SO_4^{2-} burden increases by 2.2×10^{-3} Tg(S) per Tg(S) emitted, while this response is more than a factor of two higher in SA. Consequently, despite a global decrease in sulphur emissions of 8%, the global burden is not significantly changing and the lifetime of SO_4^{2-} is slightly increasing by 5%.

6 Variability of AOD and anthropogenic radiative perturbation of aerosol and O_3

The global annual average total aerosol optical depth (AOD) ranges between 0.151 and 0.167 during the period 1980–2005 (SREF); slightly higher than the range of model/measurement values (0.127–0.151) reported by Kinne et al. (2006). The monthly mean anomalies of total AOD (Fig. 4f, $1\sigma = 0.007$ or 4.3%) are determined by variations of natural aerosol emissions, including biomass burning; the changes in anthropogenic SO_2 emissions discussed above only cause little differences in global AOD anomalies. The effect of anthropogenic emissions is more evident at the regional scale. Figure 14a shows the AOD 5-yr average calculated for the period 1981–1985, with a global average of 0.155. The changes in anthropogenic emissions (Fig. 14b) decrease AOD over large part of the Northern Hemisphere, in particular over Eastern Europe, and they largely increase AOD over East and South Asia. The effect of meteorology and natural emissions is smaller, ranging between -0.05 and 0.1 (Fig. 14c), and it is almost linearly adding to the effect of anthropogenic emissions (Fig. 14d).

In Fig. 15 and Table 5 we see that over EU the reductions in anthropogenic emissions produced a 28% decrease in AOD, and 14% over NA. In EA and SA the increasing

emissions, and particularly sulphur emissions, produced an increase of AOD of 19% and 26%, respectively. The variability in AOD due to natural aerosol emissions and meteorology is significant. In SFIX the natural variability of AOD is up to 10% over EU, 17% over NA, 8% over EA, and 13% over SA. Interestingly, over NA the resulting AOD in the SREF simulation does not show a large signal. The same results were qualitatively found also from satellite observations (Wang et al., 2009), AOD decreased only over Europe, no significant trend was found for North America, and it increased in Asia.

In Table 5 we present an analysis of the radiative perturbations due to aerosol and ozone comparing the periods 1981–1985 and 2001–2005. We define the difference between the instantaneous clear-sky total aerosol and all sky O_3 RF of the SREF and SFIX simulations, as the total aerosol and O_3 short-wave radiative perturbation due to anthropogenic emissions, and we will refer to them as RP_{aer} and RP_{O_3} , respectively.

6.1 Aerosol radiative perturbation

The instantaneous aerosol radiative forcing (RF) in ECHAM5-HAMMOZ is diagnostically calculated from the difference in the net radiative fluxes including and excluding aerosol (Stier et al., 2007). For aerosol we focus on clear sky radiative forcing, since unfortunately a coding error, prevents us to evaluate all-sky forcing. In Fig. 15 we show, together with the AOD (see before), the evolution of the normalized RP_{aer} , at the top-of-the-atmosphere (RP_{aer}^{TOA}) and at the TOA the surface (and RP_{aer}^{Surf}).

For Europe, the AOD change by -28% related to the removal of mainly SO_4^{2-} aerosol corresponds to an increase of RP_{aer}^{TOA} by $1.26 W m^{-2}$ and RP_{aer}^{Surf} by 2.05 , respectively. The 14% AOD reduction in NA corresponds to a RP_{aer}^{TOA} of $0.39 W m^{-2}$ and RP_{aer}^{Surf} of $0.71 W m^{-2}$. In EA and SA, AOD increased by 19% and 26% , corresponding to a RP_{aer}^{TOA} of -0.53 and $-0.54 W m^{-2}$, respectively, while at surface we found RP_{aer} of $-1.19 W m^{-2}$ and $-1.83 W m^{-2}$. The larger difference between TOA and surface forcing in South Asia compared to the other regions indicates a much larger contribution of BC absorption in South Asia.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Globally, there is significant spatial correlation of the aerosol radiative perturbation (SREF-SFIX) $R^2 = 0.85$ for RP_{aer}^{TOA} , while there is no correlation between atmospheric and surface aerosol radiative perturbation and AOD or BC. This is the manifestation of the more global dispersion of SO_4^{2-} , and the more local character of BC dispersion. Within the four selected regions the spatial correlation between emission induced changes in AOD and RP_{aer} at the top-of-the-atmosphere, surface and the atmosphere is much higher ($R^2 > 0.93$ for all regions except for RP_{aer}^{ATM} over NA; Table 5).

We calculate a relatively constant RP_{aer}^{TOA} between -13 to $-17 W m^{-2}$ per unit AOD around the world; and a larger range of -26 to $-48 W m^{-2}$ per unit AOD for RP_{aer} at the surface. The atmospheric absorption by aerosol RP_{aer} (calculated from the difference of RP at TOA and RP at the surface) is around $10 W m^{-2}$ in EU and NA, $15 W m^{-2}$ in EA, and $34 W m^{-2}$ in SA, showing the importance of absorbing BC aerosols in determining surface and atmospheric forcing, as confirmed by the high correlations of RP_{aer}^{ATM} with surface black carbon levels.

6.2 Ozone radiative perturbation

For convenience and completeness, we also present in this section a calculation of O_3 RP diagnosed using ECHAM5-HAMMOZ O_3 columns, in combination with all-sky radiative forcing efficiencies provided by D. Stevenson (personal communication, 2008; for a further discussion, Gauss et al., 2006). Table 5 and Fig. 5 show that O_3 total column and surface concentrations increased by 1.54 DU (1.58 ppbv) over NA, 1.35 DU (1.28 ppbv) over EU, 2.85 (4.13 ppbv) over EA, and 2.99 DU (5.12 ppbv) over SA in the period 1980–2005. The RP_{O_3} over the different regions reflect the total column O_3 changes, $0.05 W m^{-2}$ over EU, $0.06 W m^{-2}$ over NA, $0.12 W m^{-2}$ over EA, and $0.15 W m^{-2}$ over SA. As expected, the spatial correlation of O_3 columns and radiative perturbations is nearly 1, also the surface O_3 concentrations in EA and SA correlate nearly as well with the radiative perturbation. The lower correlations in EU and NA suggest that a substantial fraction of the ozone production from emissions in NA and EU takes place above the boundary layer (Table 5).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



7 Summary and conclusions

We used the coupled aerosol-chemistry-general circulation model ECHAM5-HAMMOZ, constrained with 25 yr of meteorological data from ECMWF, and a compilation of recent emission inventories, to evaluate the response of atmospheric concentrations, aerosol optical depth and radiative perturbations to anthropogenic emission changes and natural variability over the period 1980–2005. The focus of our study was on O₃ and SO₄²⁻, for which most long-term surface observations in the period 1980–2005 were available. The main findings are summarized in the following points.

- We compiled a gridded database of anthropogenic CO, VOC, NO_x, SO₂, BC, and OC emissions, utilizing reported regional emission trends. Globally, anthropogenic NO_x and OC emissions increased by 10%, while sulphur emissions decreased by 10% from 1980 to 2005. Regional emission changes were larger, e.g. all components decreased by 10–50% in North America and Europe, but increased between 40–220% in East and South Asia.
- Natural emissions were calculated on-line and dependent on inter-annual changes in meteorology. We found a rather small global inter-annual variability for biogenic VOCs emissions (3%), DMS (1%), and sea salt aerosols (2%). A larger variability was found for lightning NO_x emissions (5%) and mineral dust (10%). Generally we could not identify a clear trend for natural emissions, except for a small decreasing trend of lightning NO_x emissions ($0.017 \pm 0.007 \text{ Tg(N) yr}^{-1}$).
- A large part of the global meteorological inter-annual variability during 1980–2005 can be attributed to major natural events, such as the volcanic eruptions of the El Chichon and Pinatubo, and the 1997–1998 ENSO event. Two important drivers for atmospheric composition change – humidity and temperature – are strongly correlated. The moderate correlation ($R = 0.43$) of global inter-annual surface ozone and surface temperature, suggests important contribution to variability of other processes.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Global surface O_3 increased in 25 yr on average by 0.48 ppbv due to anthropogenic emissions, but 75% of the inter-annual variability of the multi-annual monthly surface ozone was related to natural variations.
- A regional analysis suggests that changing anthropogenic emissions increased O_3 on average by 0.8 ppbv in Europe, with large differences between southern and other parts of Europe, which is generally a larger response compared to the HTAP study (Fiore et al., 2009). Measurements qualitatively confirm these trends in Europe, but especially in winter the observed trends are up to a factor of 3 ($0.3\text{--}0.5\text{ ppbv yr}^{-1}$) larger than calculated, while in summer the small negative calculated trends were not confirmed by absence of trend in the measurements. In North America anthropogenic emissions on average slightly increased O_3 by 0.3 ppbv, nevertheless in large parts of the US decreases between 1 and 2 ppbv were calculated. Annual averages hid some seasonal model discrepancies with observed trends. In East Asia, we computed an increase of surface O_3 by 4.1 ppbv, 2.4 ppbv from anthropogenic emissions and 1.6 ppbv contribution from meteorological changes. The scarce long-term observational datasets (mostly Japanese stations) do not contradict these computed trends. In 25 yr, annual mean O_3 concentrations increased of 5.1 ppbv over SA, with approximately 75% related to increasing anthropogenic emissions. Confidence in the calculations of O_3 and O_3 trends is low, since the few available measurements suggest much lower O_3 over India.
- The tropospheric O_3 budget and variability agrees well with earlier studies by Stevenson et al. (2006) and Hess and Mahowald (2009). During 1980–2005 we calculate an intensification of tropospheric O_3 chemistry, leading to an increase of global tropospheric ozone by 3% and a decreasing O_3 lifetime by 4%. In re-analysis studies, the choice of re-analysis product and nudging method was also shown to have strong impacts on variability of O_3 and other components. For instance the agreement of our study with an alternative data assimilation

**Reanalysis
1980–2005
ECHAM5-HAMMOZ**

L. Pozzoli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Reanalysis
1980–2005
ECHAM5-HAMMOZ**

L. Pozzoli et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

technique presented by Hess and Mahowald (2009), i.e. nudging of sea-surface-temperatures (often used in climate modeling time slice experiments) resulted in substantially less agreement, and casts doubts on the applicability of such techniques for future climate experiments.

- 5 – Global OH, which determines the oxidation capacity of the atmosphere, decreased by $-0.27\% \text{ yr}^{-1}$ due to natural variability, of which lightning was the most important contributor. Anthropogenic emissions changes caused an opposite trend of $0.25\% \text{ yr}^{-1}$ thus nearly balancing the natural emission trend. Calculated inter-annual variability is in the order of 1.6%, in disagreement with the earlier study of Prinn et al. (2005) of large inter-annual fluctuations in the order of 10%, but closer to the estimates of Dentener et al. (2003) (1.8%) and Montzka et al. (2011) (2%).
- 10 – The global inter-annual variability of surface SO_4^{2-} of 10% is strongly determined by regional variations of emissions. Comparison of computed trends with measurements in Europe and North America showed in general good agreement. Seasonal trend analysis gave additional information. For instance, in Europe, measurements suggest equal downward trends of $0.05\text{--}0.1 \mu\text{g(S)} \text{ m}^{-3} \text{ yr}^{-1}$ in both summer and winter, while computed surface SO_4^{2-} declined somewhat stronger in winter than in summer. In North America, in winter the model reproduces the observed SO_4^{2-} declines well in some, but not all regions. In summer computed trends are generally underestimated by up to 50%. We expect that a misrepresentation of temporal variations of emissions, together with non-linear oxidation chemistry, could play a role in these winter-summer differences. In East and South Asia the model results suggest increases of surface SO_4^{2-} by ca. 30%, however to our knowledge no datasets are available that could corroborate these results.
- 15
- 20
- 25 – Trend and variability of sulphate columns are very different from surface SO_4^{2-} . Despite a global decrease of SO_4^{2-} emissions from 1980 to 2005, global sulphate

burdens were not significantly changing, due to a southward shift of SO₂ emissions, which determines a more efficient production and longer lifetime of SO₄²⁻.

- Globally surface SO₄²⁻ concentration decreases by ca. 0.1 μg(S) m⁻³, while the global AOD increases by ca. 0.01 (or ca. 5%), the latter driven by variability of dust and sea salt emissions. Regionally anthropogenic emissions changes are more visible: we calculate significant decline of AOD over Europe (28%), a relatively constant AOD over North America (decreased of 14% only in the last 5 yr), and strongly increasing AOD over East (19%) and South Asia (26%). These results differ substantially from Streets et al. (2009). Since the emission inventory used in this study and the one by Streets et al. (2009) are very similar, we expect that our explicit treatment of aerosol chemistry and microphysics lead to very different results than the scaling of AOD with emission trends used by Streets et al. (2009). Our analysis suggests that the impact of anthropogenic emission changes on radiative perturbations is typically larger and more regional at the surface than at the top-of-the atmosphere, with especially over South Asia a strong atmospheric warming by BC aerosol. The global top-of-the-atmosphere radiative perturbation follows more closely aerosol optical depth, reflecting large-scale SO₄²⁻ dispersion patterns. Nevertheless, our study corroborates an important role for aerosol in explaining the observed changes in surface radiation over Europe (Wild, 2009; Wang et al., 2009). Our study is also qualitatively consistent with the reported worldwide visibility decline (Wang et al., 2009). In Europe our calculated AOD reductions are consistent with improving visibility Wang (2009) and increasing surface radiation Wild (2009). O₃ radiative perturbations (not including feedbacks of CH₄) are regionally much smaller than aerosol RPs, but globally equal to or larger than aerosol RPs.

**Reanalysis
1980–2005
ECHAM5-HAMMOZ**

L. Pozzoli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



8 Outlook

Our re-analysis study showed that several of the overall processes determining the variability and trend of O₃ and aerosols are qualitatively understood- but also that many of the details are not well included. As such it gives some trust in our ability to predict the future impacts of aerosol and reactive gases on climate, but also that many model parameterisation need further improvement for more reliable predictions. It is our feeling that comparisons focussing on 1 or 2 yr of data, while useful by itself, may mask issues with compensating errors, and wrong sensitivities. Re-analysis studies are useful tools to unmask these model deficiencies. The analysis of summer and winter differences in trends and variability was particularly insightful in our study, since it highlights our level of understanding of the relative importance of chemical and meteorological processes. The separate analysis of the influence of meteorology and anthropogenic emissions changes is of direct importance for the understanding and attribution of observed trends to emission controls. The analysis of differences in regional patterns again highlights our understanding of different processes.

The ECHAM5-HAMMOZ model is, like most other climate models, continuously being improved. For instance, the overestimate of surface O₃ in many world regions or the poor representation in a tropospheric model of the stratosphere-troposphere exchange (STE) fluxes (as reported by this study, Rast et al., 2011 and Schultz et al., 2007) , reduces our trust in our trend analysis, and should be urgently addressed. Participation in model inter-comparisons, and comparison of model results to intensive measurement campaigns of multiple components may help to identify deficiencies in the model process descriptions. Improvement of parameterizations and model resolution will in the long run improve the model performance. Continued efforts are needed to improve our knowledge on anthropogenic and natural emissions in the past decades will help to understand better recent trends and variability of ozone and aerosols. New re-analysis products, such as the re-analyses from ECMWF and NCEP are frequently becoming available, and should give improved constraints for the meteorological conditions. It is

**Reanalysis
1980–2005
ECHAM5-HAMMOZ**

L. Pozzoli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of outmost importance that the few long-term measurement datasets are being continued, and that these long-term commitments are also implemented in regions outside of Europe and North America. Other datasets such as AOD from AERONET, and various quality controlled satellite datasets may in future become useful for trend analysis.

5 Chemical re-analyses is computationally and time consuming, and cannot be easily performed for every new model version. However, an updated chemical re-analysis every couple of years, following major model and re-analysis product upgrades seems highly recommendable. These studies should preferentially be performed in close collaboration with other modeling groups, which allow sharing data and analysis methods.
10 A better understanding of the chemical climate of the past is particularly relevant in the light of the continued effort to abate the negative impacts of air pollution, and the expected impacts of these controls on climate (Arneth et al., 2009; Raes and Seinfeld, 2009).

Appendix A

15 ECHAM5-HAMMOZ: model description

A1 The ECHAM5 GCM

ECHAM5 is a spectral GCM developed at the Max Planck Institute for Meteorology (Roegner et al., 2003, 2006; Hagemann et al., 2006) based on the numerical weather prediction model of the European Center for Medium-Range Weather Forecast (ECMWF). The prognostic variables of the model are vorticity, divergence, temperature, and surface pressure and are represented in the spectral space. The multi-dimensional flux-form semi-Lagrangian transport scheme from Lin and Rood (1996) is used for water vapor, cloud related variables, and chemical tracers. Stratiform clouds are described by a microphysical cloud scheme (Lohmann and Roegner, 1996) with
25 a prognostic statistical cloud cover scheme (Tompkins, 2002). Cumulus convection is

**Reanalysis
1980–2005
ECHAM5-HAMMOZ**

L. Pozzoli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



parameterized with the mass flux scheme of Tiedtke (1989) with modifications from Nordeng (1994). The radiative transfer calculation considers vertical profiles of the greenhouse gases (e.g., CO₂, O₃, CH₄), aerosols, as well as the cloud water and ice. The shortwave radiative transfer follows Cagnazzo et al. (2007) considering 6 spectral bands. For this part of the spectrum, cloud optical properties are calculated on the basis of Mie calculations using idealized size distributions for both cloud droplets and ice crystals (Rockel et al., 1991). The long-wave radiative transfer scheme is implemented according to Mlawer et al. (1997) and Morcrette et al. (1998) and considers 16 spectral bands. The cloud optical properties in the long-wave spectrum are parameterized as a function of the effective radius (Roeckner et al., 2003; Ebert and Curry, 1992).

A2 Gas-phase chemistry module MOZ

The MOZ chemical scheme has been adopted from the MOZART-2 model (Horowitz et al., 2003), and includes 63 transported tracers and 168 reactions to represent the NO_x-HO_x-hydrocarbons chemistry. The sulfur chemistry includes oxidation of SO₂ by OH and DMS oxidation by OH and NO₃ Feichter et al. (1996). Stratospheric O₃ concentrations are prescribed as monthly mean zonal climatology derived from observations (Logan, 1999; Randel et al., 1998). These concentrations are fixed at the topmost two model levels (pressures of 30 hPa and above). At other model levels above the tropopause, the concentrations are relaxed towards these values with a relaxation time of 10 days following Horowitz et al. (2003). The photolysis frequencies are calculated with the algorithm Fast-J.2 (Bian and Prather, 2002) considering the calculated optical properties of aerosols and clouds. The rates of heterogeneous reactions involving N₂O₅, NO₃, NO₂, HO₂, SO₂, HNO₃, and O₃ are calculated based on the model calculated aerosol surface area. A more detailed description of the tropospheric chemistry module MOZ and the coupling between the gas phase chemistry and the aerosols is given in Pozzoli et al. (2008a).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A3 Aerosol module HAM

The tropospheric aerosol module HAM (Stier et al., 2005) predicts the size distribution and composition of internally- and externally-mixed aerosol particles. The microphysical core of HAM, M7 (Vignati et al., 2004), treats the aerosol dynamics and thermodynamics in the framework of modal particle size distribution; the 7 log-normal modes are characterized by three moments including median radius, number of particles, and a fixed standard deviation (1.59 for fine particles and 2.00 for coarse particles. Four modes are considered as hydrophilic aerosols composed of sulfate (SU), organic (OC) and black carbon (BC), mineral dust (DU), and sea salt (SS): nucleation (NS) ($r \leq 0.005 \mu\text{m}$), Aitken (KS) ($0.005 \mu\text{m} < r \leq 0.05 \mu\text{m}$), accumulation (AS) ($0.05 \mu\text{m} < r \leq 0.5 \mu\text{m}$) and coarse (CS) ($r > 0.5 \mu\text{m}$) (where r is the number median radius). Note that in HAM the nucleation mode is entirely constituted of sulfate aerosols. Three additional modes are considered as hydrophobic aerosols composed of BC and OC in the Aitken mode (KI), and of mineral dust in the accumulation (AI) and coarse (CI) modes. Wavelength-dependent aerosol optical properties (single scattering albedo, extinction cross section, and asymmetry factor) were pre-calculated explicitly using Mie theory (Toon and Ackerman, 1981) and archived in a look-up-table for a wide range of aerosol size distributions and refractive indices. HAM is directly coupled to the cloud microphysics scheme, allowing consistent calculations of the aerosol indirect effects Lohmann et al. (2007).

A4 Gas and aerosol deposition

Gas and aerosol dry deposition follows the scheme of Ganzeveld and Lelieveld (1995); Ganzeveld et al. (1998, 2006), coupling the Wesely resistance approach with land-cover data from ECHAM5. Wet deposition is based on Stier et al. (2005), including scavenging of aerosol particles by stratiform and convective clouds and below cloud scavenging. The scavenging parameters for aerosol particles are mode-specific with lower values for hydrophobic (externally-mixed) modes. For gases, the partitioning

ACPD

11, 10191–10263, 2011

Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



between the air and the cloud water is calculated based on Henry's law and cloud water content.

Appendix B

5 O_3 and SO_4^{2-} measurement comparisons

10 Long measurement records of O_3 and SO_4^{2-} are mainly available in Europe, North America, and few stations in East Asia from the following networks: the European Monitoring and Evaluation Programme (EMEP, <http://www.emep.int/>); the Clean Air Status and Trends Network (CASTNET, <http://www.epa.gov/castnet/>); the World Data Centre for Greenhouse Gases (WDCGG, <http://gaw.kishou.go.jp/wdcgg/>). O_3 measures are available starting from year 1990 for EMEP, and few stations back to 1987 in the CASTNET and WDCGG networks. For this reason we selected only the stations that have at least 10 yr records in the period 1990–2005 for both winter and summer. A total of 81 stations were selected over Europe from EMEP, 48 stations over North America from CASTNET, and 25 stations from WDCGG. The average record length among all selected stations is of 14 yr. For SO_4^{2-} measures, we selected 62 stations from EMEP, and 43 stations from CASTNET. The average record length among all selected stations is of 13 yr.

20 The location of each selected station is plotted in Fig. 1 for both O_3 and SO_4^{2-} measurements. Each symbol represents a measuring network (EMEP: triangle; CASTNET: diamond; WDCGG: square) and each color a geographical subregion. Similarly to Fiore et al. (2009), we grouped stations in Central Europe (CEU, which includes mainly the stations of Germany, Austria, Switzerland, The Netherlands and Belgium), and South Europe (SEU, stations below 45° N and in the Mediterranean basin), but we also included in our study a group of stations for North Europe (NEU, which includes the Scandinavian countries), Eastern Europe (EEU, which includes stations east of 17° E), Western Europe (WEU, UK and Ireland). Over North America we grouped the

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sites in 4 regions, Northeast (NEUS), Great Lakes (GLUS), Mid-Atlantic (MAUS) and Southwest (SUS), based on Lehman et al. (2004) representing chemically coherent receptor regions for O₃ air pollution, and an additional region for Western US (WUS).

For each station we calculated the seasonal mean anomalies (DJF and JJA) by subtracting the multi-year average seasonal means from each annual seasonal mean. The same calculations were applied to the values extracted from ECHAM5-HAMMOZ SREF simulation, and the observed and calculated records were compared. The correlation coefficients between observed and calculated O₃ DJF anomalies is larger than 0.5 for the 44%, 39%, and 36% of the selected EMEP, CASTNET, and WDCGG stations, respectively. The agreement ($R \geq 0.5$) between observed and calculated O₃ seasonal anomalies is improving in summer months, with 52% for EMEP, 66% for CASTNET, and 52% for WDCGG. For SO₄²⁻ the correlation between observed and calculated anomalies is large than 0.5 for the 56% (DJF) and 74% (JJA) of the EMEP selected stations. In the 65% of the selected CASTNET stations the correlation between observed and calculated anomalies is larger than 0.5, both in winter and summer.

The comparisons between model results and observations are synthesized in so-called Taylor 2001 diagrams, displaying the inter-annual correlation and normalized standard deviation (ratio between the standard deviations of the calculated values and of the observations). It can be shown that the distance of each point to the black dot (1;1) is a measure of the RMS error (Fig. A1). Winter (DJF) O₃ inter-annual anomalies are relatively well represented by the model in Western Europe (WEU), Central Europe (CEU), and Northern Europe (NEU), with most correlation coefficient between 0.5–0.9, but generally underestimated standard deviations. A lower agreement ($R < 0.5$; standard deviation < 0.5) is in general found for the stations in North America, except for the stations over GLUS and MAUS. These winter differences indicate that some drivers of wintertime anomalies (such as long-range transport- or stratosphere-troposphere exchange of O₃) may not be sufficiently strongly included in the model. The summer (JJA) O₃ anomalies are in general better represented by the model. The agreement of the modeled standard deviation with measurements is increasing compared to winter

**Reanalysis
1980–2005
ECHAM5-HAMMOZ**L. Pozzoli et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

anomalies. A significant improvement compared to winter anomalies is found over North American stations (SEUS, NEUS, and MAUS), while the CEU stations have low normalized standard deviation, even if correlation coefficients are above 0.6. Interestingly, while the European inter-annual variability of the summertime ozone remains underestimated (normalized standard deviation around 0.5); the opposite is true for North American variability, indicating a too large inter-annual variability of chemical O₃ production, perhaps caused by too large contributions of natural O₃ precursors. SO₄²⁻ winter anomalies are in general reasonably well captured in winter, with correlation $R > 0.5$ at most stations, and the magnitude of the inter-annual variations. In general we found a much better agreement between calculated and observed SO₄²⁻, with inter-annual coefficients generally larger than 0.5. variability in summer than in winter. In strong contrast with the winter season, now the inter-annual variability is always underestimated (normalized standard deviation of 0.3–0.9) indicating an underestimate in the model of chemical production variability, or removal efficiency. It is unlikely that anthropogenic emissions (the dominant emissions) variability was causing this lack of variability.

Tables A1 and A2 list the observed and calculated seasonal trends of O₃ and SO₄²⁻ surface concentrations in the European and North American regions as defined before. In winter we found statistically significant increasing O₃ trends (p-value < 0.05) in all European regions and in 3 North American regions (WUS, NEUS, and MAUS), see Table A1. The SREF model simulation could capture significant trends only over Central Europe (CEU) and Western Europe (WEU). We did not find significant trends for the SFIX simulation, which may indicate no O₃ trends over Europe due to natural variability. In 2 North American regions we found significant decreasing trends (WUS and NEUS), in contrast with the observed increasing trends. We must note that in these 2 regions we also observed a decreasing trend due to natural variability (T_{SFIX}), which may be too strongly represented in our model. In summer, the observed decreasing O₃ trends over Europe are not significant, while the calculated trends show a significant decrease (T_{SREF} : NEU, WEU, and EEU), which is partially due to a natural

**Reanalysis
1980–2005
ECHAM5-HAMMOZ**

L. Pozzoli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



trend (T_{SFIX} : NEU and EEU). In North America the observations show an increasing trend in WUS, and decreasing trends in all other regions. All the observed trends are statistically significant. The model could reproduce a significant positive trend only over WUS (which seems to be determined by natural variability, $T_{\text{SFIX}} > T_{\text{SREF}}$). In all other North American regions we found increasing trends, but not significant. Nevertheless the correlation coefficients between observed and simulated anomalies are better in summer and for both Europe and North America.

SO_4^{2-} trends are in general better represented by the model. Both in Europe and North America the observations show decreasing SO_4^{2-} trends (Table A2), both in winter and summer. The trends are statistically significant in all European and North American subregions, both in winter and summer. In North America (except WUS) the observed decreasing trends are slightly higher in summer (from -0.05 and $-0.08 \mu\text{g(S)} \text{m}^{-3} \text{yr}^{-1}$) than in winter (from -0.02 and $-0.03 \mu\text{g(S)} \text{m}^{-3} \text{yr}^{-1}$). The calculated trends (T_{SREF}) are in general overestimated in winter and underestimated in summer. We must note that a seasonality in anthropogenic sulfur emissions was introduced only over Europe (30% higher in winter and 30% lower in summer compared to annual mean), while in the rest of the world, annual mean sulfur emissions were provided.

In Fig. A2 we show a comparison between the observed and calculated (SREF) annual trends of O_3 and SO_4^{2-} for the single European (EMEP) and North American (CASTNET) measuring stations. The grey areas represent the grid boxes of the model where trends are not statistically significant. We also excluded from the plot the stations where trends are not significant. Over Europe the observed O_3 annual trends are increasing, while the model does not show a significant O_3 trends for almost all Europe. In the model statistically significant decreasing trends are found over the Mediterranean and in part of Scandinavia and Baltic Sea. In North America both the observed and modeled trends are mainly not statistically significant. Decreasing SO_4^{2-} annual trends are found over Europe and North America, with a general good agreement between the observations from single stations and the model results.

**Reanalysis
1980–2005
ECHAM5-HAMMOZ**

L. Pozzoli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Acknowledgements. We greatly acknowledge Sebastian Rast at Max Planck Institute for Meteorology, Hamburg, for the scientific and technical support. We would like also to thank the Deutsches Klimarechenzentrum (DKRZ) and the Forschungszentrum Jülich for the computing resources and technical support. We would like to thank the EMEP, WDCGG, and CASTNET networks for providing ozone and sulfate measurements over Europe and North America.

References

- Andres, R. and Kasgnoc, A.: A time-averaged inventory of subaerial volcanic sulfur emissions, *J. Geophys. Res.-Atmos.*, 103, 25251–25261, 1998. 10201
- Arneth, A., Unger, N., Kulmala, M., and Andreae, M. O.: Clean the Air, Heat the Planet?, *Science*, 326, 672–673, <http://www.sciencemag.org/content/326/5953/672.short>, 2009. 10224
- Auvray, M., Bey, I., Lull, E., Schultz, M. G., and Rast, S.: A model investigation of tropospheric ozone chemical tendencies in long-range transported pollution plumes, *J. Geophys. Res.*, 112, D05304, doi:10.1029/2006JD007137, 2007. 10196, 10211
- Berglen, T., Myhre, G., Isaksen, I., Vestreng, V., and Smith, S.: Sulphate trends in Europe: Are we able to model the recent observed decrease?, *Tellus B*, 59, 773–786, <http://www.scopus.com/inward/record.url?eid=2-s2.0-3454791924%7&partnerID=40&md5=b70fa1dd6e13861b2282403bf2239728>, 2007. 10195
- Bian, H. and Prather, M.: Fast-J2: Accurate simulation of stratospheric photolysis in global chemical models, *J. Atmos. Chem.*, 41, 281–296, 2002. 10225
- Bond, T. C., Streets, D. G., Yarber, K. F., Nelson, S. M., Woo, J.-H., and Klimont, Z.: A technology-based global inventory of black and organic carbon emissions from combustion, *J. Geophys. Res.*, 109, D14203, doi:10.1029/2003JD003697, 2004. 10198
- Cagnazzo, C., Manzini, E., Giorgetta, M. A., Forster, P. M. De F., and Morcrette, J. J.: Impact of an improved shortwave radiation scheme in the MAECHAM5 General Circulation Model, *Atmos. Chem. Phys.*, 7, 2503–2515, doi:10.5194/acp-7-2503-2007, 2007. 10225
- Cheng, T., Peng, Y., Feichter, J., and Tegen, I.: An improvement on the dust emission scheme in the global aerosol-climate model ECHAM5-HAM, *Atmos. Chem. Phys.*, 8, 1105–1117, doi:10.5194/acp-8-1105-2008, 2008. 10200
- Collins, W. D., Bitz, C. M., Blackmon, M. L., Bonan, G. B., Bretherton, C. S., Carton, J. A., Chang, P., Doney, S. C., Hack, J. J., Henderson, T. B., Kiehl, J. T., Large, W. G., McKenna,

**Reanalysis
1980–2005
ECHAM5-HAMMOZ**

L. Pozzoli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


D. S., Santer, B. D., and Smith, R. D.: The Community Climate System Model Version 3 (CCSM3), *J. Climate*, 19, 2122–2143, doi:10.1175/JCLI3761.1, 2006. 10209

Dentener, F., Peters, W., Krol, M., van Weele, M., Bergamaschi, P., and Lelieveld, J.: Inter-annual variability and trend of CH₄ lifetime as a measure for OH changes in the 1979-1993 time period, *J. Geophys. Res.-Atmos.*, 108, 4442, doi:10.1029/2002JD002916, 2003. 10195, 10211, 10221

Dentener, F., Kinne, S., Bond, T., Boucher, O., Cofala, J., Generoso, S., Ginoux, P., Gong, S., Hoelzemann, J. J., Ito, A., Marelli, L., Penner, J. E., Putaud, J.-P., Textor, C., Schulz, M., van der Werf, G. R., and Wilson, J.: Emissions of primary aerosol and precursor gases in the years 2000 and 1750 prescribed data-sets for AeroCom, *Atmos. Chem. Phys.*, 6, 4321–4344, doi:10.5194/acp-6-4321-2006, 2006. 10201

Ebert, E. and Curry, J.: A parameterization of ice-cloud optical properties for climate models, *J. Geophys. Res.-Atmos.*, 97, 3831–3836, 1992. 10225

Ellingsen, K., Gauss, M., Van Dingenen, R., Dentener, F. J., Emberson, L., Fiore, A. M., Schultz, M. G., Stevenson, D. S., Ashmore, M. R., Atherton, C. S., Bergmann, D. J., Bey, I., Butler, T., Drevet, J., Eskes, H., Hauglustaine, D. A., Isaksen, I. S. A., Horowitz, L. W., Krol, M., Lamarque, J. F., Lawrence, M. G., van Noije, T., Pyle, J., Rast, S., Rodriguez, J., Savage, N., Strahan, S., Sudo, K., Szopa, S., and Wild, O.: Global ozone and air quality: a multi-model assessment of risks to human health and crops, *Atmos. Chem. Phys. Discuss.*, 8, 2163–2223, doi:10.5194/acpd-8-2163-2008, 2008. 10208

Endresen, A., Sorgard, E., Sundet, J. K., Dalsoren, S. B., Isaksen, I. S. A., Berglen, T. F., and Gravir, G.: Emission from international sea transportation and environmental impact, *J. Geophys. Res.*, 108, 4560, doi:10.1029/2002JD002898, 2003. 10197

Feichter, J., Kjellstrom, E., Rodhe, H., Dentener, F., Lelieveld, J., and Roelofs, G.: Simulation of the tropospheric sulfur cycle in a global climate model, *Atmos. Environ.*, 30, 1693–1707, 1996. 10225

Fiore, A., Horowitz, L., Dlugokencky, E., and West, J.: Impact of meteorology and emissions on methane trends, 1990–2004, *Geophys. Res. Lett.*, 33, L12809, doi:10.1029/2006GL026199, 2006. 10211

Fiore, A. M., Dentener, F. J., Wild, O., Cuvelier, C., Schultz, M. G., Hess, P., Textor, C., Schulz, M., Doherty, R. M., Horowitz, L. W., MacKenzie, I. A., Sanderson, M. G., Shindell, D. T., Stevenson, D. S., Szopa, S., Van Dingenen, R., Zeng, G., Atherton, C., Bergmann, D., Bey, I., Carmichael, G., Collins, W. J., Duncan, B. N., Faluvegi, G., Folberth, G., Gauss, M., Gong,

**Reanalysis
1980–2005
ECHAM5-HAMMOZ**

L. Pozzoli et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



S., Hauglustaine, D., Holloway, T., Isaksen, I. S. A., Jacob, D. J., Jonson, J. E., Kaminski, J. W., Keating, T. J., Lupu, A., Marmer, E., Montanaro, V., Park, R. J., Pitari, G., Pringle, K. J., Pyle, J. A., Schroeder, S., Vivanco, M. G., Wind, P., Wojcik, G., Wu, S., and Zuber, A.: Multimodel estimates of intercontinental source-receptor relationships for ozone pollution, *J. Geophys. Res.*, 114, D04301, doi:10.1029/2008JD010816, 2009. 10195, 10204, 10205, 10220, 10227

Ganzeveld, L. and Lelieveld, J.: Dry deposition parameterization in a chemistry general circulation model and its influence on the distribution of reactive trace gases, *J. Geophys. Res.-Atmos.*, 100, 20999–21012, 1995. 10226

Ganzeveld, L., Lelieveld, J., and Roelofs, G.: A dry deposition parameterization for sulfur oxides in a chemistry and general circulation model, *J. Geophys. Res.-Atmos.*, 103, 5679–5694, 1998. 10226

Ganzeveld, L. N., van Aardenne, J. A., Butler, T. M., Lawrence, M. G., Metzger, S. M., Stier, P., Zimmermann, P., and Lelieveld, J.: Technical Note: Anthropogenic and natural offline emissions and the online EMISSIONS and dry DEPOSITION submodel EMDEP of the Modular Earth Submodel system (MESSy), *Atmos. Chem. Phys. Discuss.*, 6, 5457–5483, doi:10.5194/acpd-6-5457-2006, 2006. 10226

Gauss, M., Myhre, G., Isaksen, I. S. A., Grewe, V., Pitari, G., Wild, O., Collins, W. J., Dentener, F. J., Ellingsen, K., Gohar, L. K., Hauglustaine, D. A., Iachetti, D., Lamarque, F., Mancini, E., Mickley, L. J., Prather, M. J., Pyle, J. A., Sanderson, M. G., Shine, K. P., Stevenson, D. S., Sudo, K., Szopa, S., and Zeng, G.: Radiative forcing since preindustrial times due to ozone change in the troposphere and the lower stratosphere, *Atmos. Chem. Phys.*, 6, 575–599, doi:10.5194/acp-6-575-2006, 2006. 10218

Grewe, V., Brunner, D., Dameris, M., Grenfell, J. L., Hein, R., Shindell, D., and Staehelin, J.: Origin and variability of upper tropospheric nitrogen oxides and ozone at northern mid-latitudes, *Atmos. Environ.*, 35, 3421–3433, 2001. 10197, 10199

Guenther, A., Hewitt, C., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau, M., McKay, W., Pierce, T., Scholes, B., Steinbrecher, R., Tallamraju, R., Taylor, J., and Zimmerman, P.: A global-model of natural volatile organic-compound emissions, *J. Geophys. Res.-Atmos.*, 100, 8873–8892, 1995. 10201

Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, doi:10.5194/acp-6-3181-2006,

Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


2006. 10199

Hagemann, S., Arpe, K., and Roeckner, E.: Evaluation of the Hydrological Cycle in the ECHAM5 Model, *J. Climate*, 19, 3810–3827, 2006. 10210, 10224

Halmer, M., Schmincke, H., and Graf, H.: The annual volcanic gas input into the atmosphere, in particular into the stratosphere: a global data set for the past 100 years, *J. Volcanol. Geothermal Res.*, 115, 511–528, 2002. 10201

Hess, P. and Mahowald, N.: Interannual variability in hindcasts of atmospheric chemistry: the role of meteorology, *Atmos. Chem. Phys.*, 9, 5261–5280, doi:10.5194/acp-9-5261-2009, 2009. 10195, 10209, 10210, 10211, 10220, 10221, 10242

Horowitz, L., Walters, S., Mauzerall, D., Emmons, L., Rasch, P., Granier, C., Tie, X., Lamarque, J., Schultz, M., Tyndall, G., Orlando, J., and Brasseur, G.: A global simulation of tropospheric ozone and related tracers: Description and evaluation of MOZART, version 2, *J. Geophys. Res.-Atmos.*, 108, 4784, doi:10.1029/2002JD002853, 2003. 10201, 10225

Jeuken, A., Siegmund, P., Heijboer, L., Feichter, J., and Bengtsson, L.: On the potential of assimilating meteorological analyses in a global climate model for the purpose of model validation, *Journal of Geophysical Research-Atmospheres*, 101, 16 939–16 950, 1996. 10196

Kettle, A. and Andreae, M.: Flux of dimethylsulfide from the oceans: A comparison of updated data seas and flux models, *J. Geophys. Res.-Atmos.*, 105, 26793–26808, 2000. 10200

Kinne, S., Schulz, M., Textor, C., Guibert, S., Balkanski, Y., Bauer, S. E., Bernsten, T., Berglen, T. F., Boucher, O., Chin, M., Collins, W., Dentener, F., Diehl, T., Easter, R., Feichter, J., Fillmore, D., Ghan, S., Ginoux, P., Gong, S., Grini, A., Hendricks, J., Herzog, M., Horowitz, L., Isaksen, I., Iversen, T., Kirkevåg, A., Kloster, S., Koch, D., Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Lesins, G., Liu, X., Lohmann, U., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S., Seland, O., Stier, P., Takemura, T., and Tie, X.: An AeroCom initial assessment - optical properties in aerosol component modules of global models, *Atmos. Chem. Phys.*, 6, 1815–1834, doi:10.5194/acp-6-1815-2006, 2006. 10216

Lathièrre, J., Hauglustaine, D. A., Friend, A. D., De Noblet-Ducoudré, N., Viovy, N., and Folberth, G. A.: Impact of climate variability and land use changes on global biogenic volatile organic compound emissions, *Atmos. Chem. Phys.*, 6, 2129–2146, doi:10.5194/acp-6-2129-2006, 2006. 10201

Lawrence, M. G., Jöckel, P., and von Kuhlmann, R.: What does the global mean OH concentration tell us?, *Atmos. Chem. Phys.*, 1, 37–49, doi:10.5194/acp-1-37-2001, 2001. 10211

Lehman, J., Swinton, K., Bortnick, S., Hamilton, C., Baldrige, E., Eder, B., and Cox, B.: Spatio-

Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


temporal characterization of tropospheric ozone across the eastern United States, *Atmos. Environ.*, 38, 4357–4369, 2004. 10228

Lin, S. and Rood, R.: Multidimensional flux-form semi-Lagrangian transport schemes, *Mon. Weather Rev.*, 124, 2046–2070, 1996. 10224

5 Logan, J.: An analysis of ozonesonde data for the troposphere: Recommendations for testing 3-D models and development of a gridded climatology for tropospheric ozone, *J. Geophys. Res.-Atmos.*, 104, 16115–16149, 1999. 10225

Lohmann, U. and Roeckner, E.: Design and performance of a new cloud microphysics scheme developed for the ECHAM general circulation model, *Climate Dynamics*, 12, 557–572, 1996. 10224

10 Lohmann, U., Stier, P., Hoose, C., Ferrachat, S., Kloster, S., Roeckner, E., and Zhang, J.: Cloud microphysics and aerosol indirect effects in the global climate model ECHAM5-HAM, *Atmos. Chem. Phys.*, 7, 3425–3446, doi:10.5194/acp-7-3425-2007, 2007. 10226

Mlawer, E., Taubman, S., Brown, P., Iacono, M., and Clough, S.: Radiative transfer for inhomogeneous atmospheres: RRTM, a validated correlated-k model for the longwave, *J. Geophys. Res.-Atmos.*, 102, 16663–16682, 1997. 10225

15 Montzka, S. A., Krol, M., Dlugokencky, E., Hall, B., Jockel, P., and Lelieveld, J.: Small Interannual Variability of Global Atmospheric Hydroxyl, *Science*, 331, 67–69, <http://www.sciencemag.org/content/331/6013/67.abstract>, 2011. 10212, 10221

20 Morcrette, J., Clough, S., Mlawer, E., and Iacono, M.: Impact of a validated radiative transfer scheme, RRTM, on the ECMWF model climate and 10-day forecasts, *Tech. Rep. 252*, ECMWF, Reading, UK, 1998. 10225

Nakicenovic, N., Alcamo, J., Davis, G., de Vries, H., Fenhann, J., Gaffin, S., Gregory, K., Grubler, A., Jung, T., Kram, T., Rovere, E. L., Michaelis, L., Mori, S., Morita, T., Papper, W., Pitcher, H., Price, L., Riahi, K., Roehrl, A., Rogner, H.-H., Sankovski, A., Schlesinger, M., Shukla, P., Smith, S., Swart, R., van Rooijen, S., Victor, N., and Dadi, Z.: *Special Report on Emissions Scenarios*. Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, 2000. 10197

25 Nightingale, P., Malin, G., Law, C., Watson, A., Liss, P., Liddicoat, M., Boutin, J., and Upstill-Goddard, R.: In situ evaluation of air-sea gas exchange parameterizations using novel conservative and volatile tracers, *Global Biogeochem. Cy.*, 14, 373–387, 2000. 10200

Nordeng, T.: Extended versions of the convective parameterization scheme at ECMWF and their impact on the mean and transient activity of the model in the tropics, *Tech. Rep. 206*,

**Reanalysis
1980–2005
ECHAM5-HAMMOZ**

L. Pozzoli et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


ECMWF, Reading, UK, 1994. 10225

Pham, M., Muller, J., Brasseur, G., Granier, C., and Megie, G.: A three-dimensional study of the tropospheric sulfur cycle, *J. Geophys. Res.-Atmos.*, 100, 26061–26092, 1995. 10200

Pozzoli, L., Bey, I., Rast, S., Schultz, M. G., Stier, P., and Feichter, J.: Trace gas and aerosol interactions in the fully coupled model of aerosol-chemistry-climate ECHAM5-HAMMOZ: 1. Model description and insights from the spring 2001 TRACE-P experiment, *J. Geophys. Res.*, 113, D07308, doi:10.1029/2007JD009007, 2008a. 10196, 10203, 10225

Pozzoli, L., Bey, I., Rast, S., Schultz, M. G., Stier, P., and Feichter, J.: Trace gas and aerosol interactions in the fully coupled model of aerosol-chemistry-climate ECHAM5-HAMMOZ: 2. Impact of heterogeneous chemistry on the global aerosol distributions, *J. Geophys. Res.*, 113, D07309, doi:10.1029/2007JD009008, 2008b. 10196, 10203

Prinn, R. G., Huang, J., Weiss, R. F., Cunnold, D. M., Fraser, P. J., Simmonds, P. G., McCulloch, A., Harth, C., Reimann, S., Salameh, P., O'Doherty, S., Wang, R. H. J., Porter, L. W., Miller, B. R., and Krummel, P. B.: Evidence for variability of atmospheric hydroxyl radicals over the past quarter century, *Geophys. Res. Lett.*, 32, L07809, doi:10.1029/2004GL022228, 2005. 10211, 10221

Rast, J., Schultz, M., Aghedo, A., Bey, I., Brasseur, G., Diehl, T., Esch, M., Ganzeveld, L., Kirchner, I., Kornblueh, L., Rhodin, A., Roeckner, E., Schmidt, H., Schroede, S., Schulzweida, U., Stier, P., and van Noije, T.: Interannual variability in tropospheric ozone over the 1980–2000 period: Results from the global chemistry climate model ECHAM5MOZ, *J. Geophys. Res.*, in review, 2011. 10196, 10203, 10223

Raes, F. and Seinfeld, J.: Climate Change and Air Pollution Abatement: A Bumpy Road., *Atmos. Environ.*, 43, 5132–5133, 2009. 10224

Randel, W., Wu, F., Russell, J., Roche, A., and Waters, J.: Seasonal cycles and QBO variations in stratospheric CH₄ and H₂O observed in UARS HALOE data, *J. Atmos. Sci.*, 55, 163–185, 1998. 10225

Rockel, B., Raschke, E., and Weyres, B.: A parameterization of broad band radiative transfer properties of water, ice and mixed clouds, *Beitr. Phys. Atmos.*, 64, 1–12, 1991. 10225

Roeckner, E., Bauml, G., Bonaventura, L., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kirchner, I., Kornblueh, L., Manzini, E., Rhodin, A., Schlese, U., Schulzweida, U., and Tompkins, A.: The atmospheric general circulation model ECHAM5: Part 1, *Tech. Rep. 349*, Max Planck Institute for Meteorology, Hamburg, 2003. 10197, 10224, 10225

Roeckner, E., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kornblueh, L., Manzini, E.,

Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Schlese, U., and Schulzweida, U.: Sensitivity of Simulated Climate to Horizontal and Vertical Resolution in the ECHAM5 Atmosphere Model, *J. Climate*, 19, 3771–3791, 2006. 10224

Schultz, M., Backman, L., Balkanski, Y., Bjoerndalsaeter, S., Brand, R., Burrows, J., Dalsorenen, S., de Vasconcelos, M., Grodtmann, B., Hauglustaine, D., Heil, A., Hoelzemann, J., Isaksen, I., Kaurola, J., Knorr, W., Ladstaetter-Weienmayer, A., Mota, B., Oom, D., Pacyna, J., Panasiuk, D., Pereira, J., Pulles, T., Pyle, J., Rast, S., Richter, A., Savage, N., Schnadt, C., Schulz, M., Spessa, A., Staehelin, J., Sundet, J., Szopa, S., Thonicke, K., van het Bolscher, M., van Noije, T., van Velthoven, P., Vik, A., and Wittrock, F.: REanalysis of the TROpospheric chemical composition over the past 40 years (RETRO). A long-term global modeling study of tropospheric chemistry. Final Report, Tech. rep., Max Planck Institute for Meteorology, Hamburg, Germany, 2007. 10195, 10197, 10199, 10223

Schultz, M. G., Heil, A., Hoelzemann, J. J., Spessa, A., Thonicke, K., Goldammer, J. G., Held, A. C., Pereira, J. M. C., and van het Bolscher, M.: Global wildland fire emissions from 1960 to 2000, *Global Biogeochem. Cy.*, 22, GB2002, doi:10.1029/2007GB003031, 2008. 10197, 10200

Schulz, M., de Leeuw, G., and Balkanski, Y.: Emission Of Atmospheric Trace Compounds, chap. Sea-salt aerosol source functions and emissions, 333–359, Kluwer, 2004. 10200

Schumann, U. and Huntrieser, H.: The global lightning-induced nitrogen oxides source, *Atmos. Chem. Phys.*, 7, 3823–3907, doi:10.5194/acp-7-3823-2007, 2007. 10199

Solberg, S., Hov, O., Sovde, A., Isaksen, I. S. A., Coddeville, P., De Backer, H., Forster, C., Orsolini, Y., and Uhse, K.: European surface ozone in the extreme summer 2003, *J. Geophys. Res.*, 113, D07307, doi:10.1029/2007JD009098, 2008. 10195

Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K., Tignor, M., and Miller, H. L.: Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, 2007, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2007. 10194

Stevenson, D., Dentener, F., Schultz, M., Ellingsen, K., van Noije, T., Wild, O., Zeng, G., Amann, M., Atherton, C., Bell, N., Bergmann, D., Bey, I., Butler, T., Cofala, J., Collins, W., Derwent, R., Doherty, R., Drevet, J., Eskes, H., Fiore, A., Gauss, M., Hauglustaine, D., Horowitz, L., Isaksen, I., Krol, M., Lamarque, J., Lawrence, M., Montanaro, V., Muller, J., Pitari, G., Prather, M., Pyle, J., Rast, S., Rodriguez, J., Sanderson, M., Savage, N., Shindell, D., Strahan, S., Sudo, K., and Szopa, S.: Multimodel ensemble simulations of present-day and near-future tropospheric ozone, *J. Geophys. Res.-Atmos.*, 111, D08301, doi:10.1029/2005JD006338,

**Reanalysis
1980–2005
ECHAM5-HAMMOZ**

L. Pozzoli et al.

[Title Page](#)
[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2006. 10208, 10220, 10255

Stier, P., Feichter, J., Kinne, S., Kloster, S., Vignati, E., Wilson, J., Ganzeveld, L., Tegen, I., Werner, M., Balkanski, Y., Schulz, M., Boucher, O., Minikin, A., and Petzold, A.: The aerosol-climate model ECHAM5-HAM, *Atmos. Chem. Phys.*, 5, 1125–1156, doi:10.5194/acp-5-1125-2005, 2005. 10196, 10203, 10226

Stier, P., Seinfeld, J. H., Kinne, S., and Boucher, O.: Aerosol absorption and radiative forcing, *Atmos. Chem. Phys.*, 7, 5237–5261, doi:10.5194/acp-7-5237-2007, 2007. 10217

Streets, D. G., Bond, T. C., Lee, T., and Jang, C.: On the future of carbonaceous aerosol emissions, *J. Geophys. Res.*, 109, D24212, doi:10.1029/2004JD004902, 2004. 10198

Streets, D. G., Wu, Y., and Chin, M.: Two-decadal aerosol trends as a likely explanation of the global dimming/brightening transition, *Geophys. Res. Lett.*, 33, L15806, doi:10.1029/2006GL026471, 2006. 10198

Streets, D. G., Yan, F., Chin, M., Diehl, T., Mahowald, N., Schultz, M., Wild, M., Wu, Y., and Yu, C.: Anthropogenic and natural contributions to regional trends in aerosol optical depth, 1980–2006, *J. Geophys. Res.*, 114, D00D18, doi:10.1029/2008JD011624, 2009. 10194, 10198, 10222

Taylor, K.: Summarizing multiple aspects of model performance in a single diagram, *J. Geophys. Res.-Atmos.*, 106, 7183–7192, 2001. 10228

Tegen, I., Harrison, S., Kohfeld, K., Prentice, I., Coe, M., and Heimann, M.: Impact of vegetation and preferential source areas on global dust aerosol: Results from a model study, *J. Geophys. Res.-Atmos.*, 107, 4576, doi:10.1029/2001JD000963, 2002. 10200

Tiedtke, M.: A comprehensive mass flux scheme for cumulus parameterization in large-scale models, *Mon. Weather Rev.*, 117, 1779–1800, 1989. 10225

Tompkins, A.: A prognostic parameterization for the subgrid-scale variability of water vapor and clouds in large-scale models and its use to diagnose cloud cover, *J. Atmos. Sci.*, 59, 1917–1942, 2002. 10224

Toon, O. and Ackerman, T.: Algorithms for the calculation of scattering by stratified spheres, *Appl. Optics*, 20, 3657–3660, 1981. 10226

Tost, H., Jöckel, P., and Lelieveld, J.: Lightning and convection parameterisations - uncertainties in global modelling, *Atmos. Chem. Phys.*, 7, 4553–4568, doi:10.5194/acp-7-4553-2007, 2007. 10199

Tressol, M., Ordonez, C., Zbinden, R., Brioude, J., Thouret, V., Mari, C., Nedelec, P., Cammas, J.-P., Smit, H., Patz, H.-W., and Volz-Thomas, A.: Air pollution during the 2003 European heat

Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


wave as seen by MOZAIC airliners, *Atmos. Chem. Phys.*, 8, 2133–2150, doi:10.5194/acp-8-2133-2008, 2008. 10195

Unger, N., Menon, S., Koch, D. M., and Shindell, D. T.: Impacts of aerosol-cloud interactions on past and future changes in tropospheric composition, *Atmos. Chem. Phys.*, 9, 4115–4129, doi:10.5194/acp-9-4115-2009, 2009. 10216

Uppala, S. M., KÅllberg, P. W., Simmons, A. J., Andrae, U., Bechtold, V. D. C., Fiorino, M., Gibson, J. K., Haseler, J., Hernandez, A., Kelly, G. A., Li, X., Onogi, K., Saarinen, S., Sokka, N., Allan, R. P., Andersson, E., Arpe, K., Balmaseda, M. A., Beljaars, A. C. M., Berg, L. V. D., Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M., Fisher, M., Fuentes, M., Hagemann, S., Hlm, E., Hoskins, B. J., Isaksen, L., Janssen, P. A. E. M., Jenne, R., McNally, A. P., Mahfouf, J.-F., Morcrette, J.-J., Rayner, N. A., Saunders, R. W., Simon, P., Sterl, A., Trenberth, K. E., Untch, A., Vasiljevic, D., Viterbo, P., and Woollen, J.: The ERA-40 re-analysis, *Q. J. Roy. Meteorol. Soc.*, 131, 2961–3012, doi:10.1256/qj.04.176, 2005. 10196

Van Aardenne, J., Dentener, F., Olivier, J., Goldewijk, C., and Lelieveld, J.: A 1 1 resolution data set of historical anthropogenic trace gas emissions for the period 1890–1990, *Global Biogeochem. Cy.*, 15, 909–928, 2001. 10198

van Noije, T. P. C., Segers, A. J., and van Velthoven, P. F. J.: Time series of the stratosphere-troposphere exchange of ozone simulated with reanalyzed and operational forecast data, *J. Geophys. Res.*, 111, D03301, doi:10.1029/2005JD006081, 2006. 10195

Vautard, R., Szopa, S., Beekmann, M., Menut, L., Hauglustaine, D. A., Rouil, L., and Roemer, M.: Are decadal anthropogenic emission reductions in Europe consistent with surface ozone observations?, *Geophys. Res. Lett.*, 33, L13810, doi:10.1029/2006GL026080, 2006. 10195

Vignati, E., Wilson, J., and Stier, P.: M7: An efficient size-resolved aerosol microphysics module for large-scale aerosol transport models, *J. Geophys. Res.-Atmos.*, 109, D22202, doi:10.1029/2003JD004485, 2004. 10226

Wang, K., Dickinson, R., and Liang, S.: Clear sky visibility has decreased over land globally from 1973 to 2007, *Science*, 323, 1468–1470, 2009. 10194, 10217, 10222

Wild, M.: Global dimming and brightening: A review, *J. Geophys. Res.*, 114, D00D16, doi:10.1029/2008JD011470, 2009. 10194, 10195, 10222

Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.

Table 1. Global anthropogenic emissions of CO [Tg yr⁻¹], NO_x [Tg(N) yr⁻¹], VOCs [Tg(C) yr⁻¹], SO₂ [Tg(S) yr⁻¹], SO₄²⁻ [Tg(S) yr⁻¹], OC [Tg yr⁻¹], and BC [Tg yr⁻¹].

	1980	1985	1990	1995	2000	2005
CO	673.7	680.1	713.8	685.3	655.4	678.6
NO _x	34.2	33.7	36.1	36.4	36.7	37.2
VOCs	84.6	85.0	87.9	84.8	80.5	84.3
SO ₂	67.1	67.2	66.2	61.0	58.8	59.0
SO ₄ ²⁻	1.8	1.8	1.8	1.7	1.6	1.6
OC	7.8	8.2	8.3	8.7	8.5	8.7
BC	4.9	4.9	4.9	4.8	4.7	4.9

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.

Table 2. Average, standard deviation (SD) and relative standard deviation (RSD, standard deviation divided by the mean) of globally averaged variables in this work for the simulation with changing anthropogenic emission and with fixed anthropogenic emissions (1980–2005).

	SREF			SFIX		
	Average	SD	RSD	Average	SD	RSD
Sfc O ₃ (ppbv)	36.45	0.826	0.0227	35.97	0.627	0.0174
O ₃ (ppbv)	48.37	1.1020	0.0228	47.64	0.8851	0.0186
CO (ppbv)	0.103	0.000416	0.0402	0.101	0.000529	0.0519
OH (molecules cm ⁻³ × 10 ⁶)	1.20	0.016	0.013	1.18	0.029	0.024
HNO ₃ (pptv)	129.11	14.70	0.1139	125.73	13.91	0.1106
Emi S (Tg yr ⁻¹)	104.2	3.49	0.0336	108.09	0.47	0.0043
SO ₂ (pptv)	231.7	15.02	0.0648	246.9	8.70	0.0352
Sfc SO ₄ ⁻² (μg m ⁻³)	1.12	0.071	0.0640	1.18	0.061	0.0515
SO ₄ ⁻² (μg m ⁻³)	0.69	0.028	0.0406	0.72	0.025	0.0352

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.

Table 3. Average, standard deviation (SD) and relative standard deviation (RSD, standard deviation divided by the mean) of globally averaged variables in this work, SCAM and SNCEP (Hess and Mahowald, 2009) (1980–2000). Three dimension variables are density weighted and averaged between the surface and 280 hPa. Three dimension quantities evaluated at the surface are prefixed with Sfc. The standard deviation is calculated as the standard deviation of the monthly anomalies (the monthly value minus the mean of all years for that month).

	ERA40 (this work SFIX)			SCAM (Hess, 2009)			SNCEP (Hess, 2009)		
	Average	SD	RSD	Average	SD	RSD	Average	SD	RSD
Sfc T (K)	287	0.112	0.000391	287	0.116	0.000403	287	0.121	0.00042
Sfc JNO_2 ($s^{-1} \times 10^{-3}$)	2.13	0.0121	0.00567	2.43	0.00454	0.00187	2.39	0.00814	0.00341
LNO ($TgN yr^{-1}$)	3.91	0.153	0.0387	4.71	0.118	0.0251	2.79	0.211	0.0759
PRECIP ($mm day^{-1}$)	2.95	0.0331	0.011	2.42	0.0145	0.006	2.4	0.0389	0.0162
Q ($g kg^{-1}$)	4.72	0.060	0.0127	3.46	0.0411	0.0119	3.38	0.0361	0.0107
O_3 (ppbv)	47.79	0.819	0.01714	46	0.192	0.00418	48.4	0.752	0.0155
Sfc O_3 (ppbv)	36.1	0.595	0.0165	29.8	0.122	0.0041	31.2	0.468	0.015
CO (ppbv)	0.100	0.000450	0.04482	0.083	0.000449	0.00542	0.0847	0.000388	0.00458
OH ($mole/mole \times 10^{15}$)	63.1	1.269	0.02012	73.5	0.707	0.00962	70.4	0.847	0.012
HNO_3 (pptv)	127	13.95	0.1096	121	1.22	0.0101	121	1.49	0.0123

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.

Table 4. Relationships, in Tg(S) per Tg(S) emitted, calculated for the period 1980–2005 between sulfur emissions and SO_4^{2-} burden (B), SO_4^{2-} production from SO_2 in-cloud oxidation (In-cloud), and SO_4^{2-} gaseous phase production (Cond) over Europe (EU), North America (NA), East Asia (EA), and South Asia (SA).

	EU		NA		EA		SA	
	slope	R^2	slope	R^2	slope	R^2	slope	R^2
B ($\times 10^{-3}$)	2.21	0.86	0.79	0.12	2.86	0.79	5.36	0.90
In-cloud	0.31	0.97	0.38	0.93	0.25	0.79	0.18	0.90
Cond	0.08	0.93	0.09	0.70	0.17	0.85	0.37	0.98

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I ◀

▶ I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.

Table 5. Globally and regionally (EU, NA, EA, and SA) averaged effect of changing anthropogenic emissions (SREF-SFIX) during the 5-yr periods 1981–1985 and 2001–2005 on: surface concentrations of O_3 (ppbv), SO_4^{2-} (%), and BC (%); total aerosol optical depth (AOD) (%) and total column O_3 (DU); The total anthropogenic aerosol radiative perturbation at top of the atmosphere (RP_{aer}^{TOA}), at surface (RP_{aer}^{SURF}) ($W m^{-2}$), and in the atmosphere ($RP_{aer}^{ATM} = (RP_{aer}^{TOA} - RP_{aer}^{SURF})$); the anthropogenic radiative perturbation of O_3 (RP_{O_3}); correlations calculated over the entire period 1980–2005 between anthropogenic RP_{aer}^{TOA} and ΔAOD ; between anthropogenic RP_{aer}^{SURF} and ΔAOD ; between RP_{aer}^{ATM} and ΔAOD ; between RP_{aer}^{ATM} and ΔBC ; between anthropogenic RP_{O_3} and ΔO_3 at surface; between anthropogenic RP_{O_3} and ΔO_3 column.

	GLOBAL		EU		NA		EA		SA	
ΔO_3 [ppbv]	0.98		0.81		0.27		2.44		4.25	
ΔSO_4^{2-} [%]	-10		-36		-25		27		59	
ΔBC [%]	0		-43		-34		30		70	
ΔAOD [%]	0		-28		-14		19		26	
ΔO_3 [DU]	1.18		1.35		1.54		2.85		2.99	
RP_{aer}^{TOA} [$W m^{-2}$]	0.02		1.26		0.39		-0.53		-0.54	
RP_{aer}^{SURF} [$W m^{-2}$]	-0.03		2.05		0.71		-1.19		-1.83	
RP_{aer}^{ATM} [$W m^{-2}$]	0.05		-0.79		-0.32		0.66		1.29	
RP_{O_3} [$W m^{-2}$]	0.05		0.05		0.06		0.12		0.15	
	slope	R^2	slope	R^2	slope	R^2	slope	R^2	slope	R^2
RP_{aer}^{TOA} [$W m^{-2}$] vs. ΔAOD	-17.86	0.85	-16.1	0.99	-16.99	0.97	-13.57	0.99	-13.84	0.99
RP_{aer}^{SURF} [$W m^{-2}$] vs. ΔAOD	-1.32	0.24	-26.71	0.99	-28.10	0.97	-28.95	0.99	-47.57	0.99
RP_{aer}^{ATM} [$W m^{-2}$] vs. ΔAOD	-4.62	0.03	10.61	0.93	11.11	0.68	15.38	0.97	33.73	0.98
RP_{aer}^{ATM} [$W m^{-2}$] vs. ΔBC [%]	0.35	0.11	1.73	0.99	0.95	0.99	1.92	0.97	1.74	0.99
RP_{O_3} [$mW m^{-2}$] vs. ΔO_3 [ppbv]	42.8	0.91	35.2	0.65	51.3	0.49	46.7	0.94	36.0	0.97
RP_{O_3} [$mW m^{-2}$] vs. ΔO_3 [DU]	40.8	0.99	36.4	0.99	44.2	0.99	44.1	0.99	49.1	0.99

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.

Table A1. Observed and simulated trends of surface O₃ seasonal anomalies for European (EU) and North American (NA) stations, grouped as in Fig. 1 (North Europe (NEU); Central Europe (CEU); West Europe (WEU); East Europe (EEU); West US (WUS); North East US (NEUS); Mid-Atlantic US (MAUS); Great lakes US (GLUS); South US (SUS)). The number of stations (NSTA) for each regions is listed in the second column. The seasonal trends are listed separately for winter (DJF) and summer (JJA). Seasonal trends (ppbvyr⁻¹) are calculated for observations (T_{OBS}), SREF and SFIX simulations (T_{SREF} and T_{SFIX}) as linear fitting of the median surface O₃ anomalies of each group of stations (the 95% confidence interval is also shown for each calculated trend). The trends statistically significant (p-value < 0.05) are highlighted in bold. The correlation coefficient between median observed and simulated (SREF) anomalies are listed ($R_{\text{OBS/SREF}}$). The correlation coefficients statistically significant (p-value < 0.05) are highlighted in bold.

O ₃ Stations		Winter anomalies (DJF)				Summer anomalies (JJA)			
REGION	NSTA	T_{OBS}	T_{SREF}	T_{SFIX}	$R_{\text{OBS/SREF}}$	T_{OBS}	T_{SREF}	T_{SFIX}	$R_{\text{OBS/SREF}}$
NEU	20	0.27±0.18	0.05±0.23	-0.08±0.26	0.49	-0.00±0.22	-0.44±0.26	-0.29±0.27	0.36
CEU	54	0.44±0.15	0.21±0.40	0.06±0.40	0.46	0.04±0.32	-0.11±0.30	-0.00±0.29	0.73
WEU	16	0.42±0.36	0.40±0.55	0.21±0.56	0.93	-0.10±0.23	-0.15±0.28	-0.11±0.28	0.62
EEU	8	0.34±0.38	0.06±0.27	-0.09±0.28	0.07	-0.02±0.25	-0.36±0.36	-0.22±0.34	0.71
WUS	7	0.12±0.21	-0.08±0.11	-0.12±0.12	0.26	0.41±0.30	0.11±0.21	0.21±0.22	0.80
NEUS	11	0.22±0.13	-0.10±0.17	-0.17±0.15	0.03	-0.27±0.28	0.03±0.47	0.14±0.49	0.55
MAUS	17	0.11±0.18	-0.02±0.23	-0.07±0.26	0.66	-0.40±0.37	0.09±0.81	0.19±0.83	0.68
GLUS	14	0.04±0.18	0.05±0.19	-0.02±0.20	0.82	-0.19±0.29	0.20±0.47	0.34±0.49	0.43
SUS	4	0.08±0.20	-0.08±0.26	-0.06±0.27	0.50	-0.25±0.48	0.29±0.84	0.40±0.83	0.64

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.

Table A2. Observed and simulated trends of surface SO_4^{2-} seasonal anomalies for European (EU) and North American (NA) stations, grouped as in Fig. 1 (North Europe (NEU); Central Europe (CEU); West Europe (WEU); East Europe (EEU); South Europe (SEU); West US (WUS); North East US (NEUS); Mid-Atlantic US (MAUS); Great lakes US (GLUS); South US (SUS)). The number of stations (NSTA) for each regions is listed in the second column. The seasonal trends are listed separately for winter (DJF) and summer (JJA). Seasonal trends ($\mu\text{g(S)} \text{ m}^{-3} \text{ yr}^{-1}$) are calculated for observations (T_{OBS}), SREF and SFIX simulations (T_{SREF} and T_{SFIX}) as linear fitting of the median surface SO_4^{2-} anomalies of each group of stations (the 95% confidence interval is also shown for each calculated trend). The trends statistically significant (p-value < 0.05) are highlighted in bold. The correlation coefficient between median observed and simulated (SREF) anomalies are listed ($R_{\text{OBS/SREF}}$). The correlation coefficients statistically significant (p-value < 0.05) are highlighted in bold.

SO ₄ ²⁻ Stations		Winter anomalies (DJF)				Summer anomalies (JJA)			
REGION	NSTA	T_{OBS}	T_{SREF}	T_{SFIX}	$R_{\text{OBS/SREF}}$	T_{OBS}	T_{SREF}	T_{SFIX}	$R_{\text{OBS/SREF}}$
NEU	18	-0.03±0.02	-0.01±0.04	0.02±0.08	0.59	-0.03±0.01	-0.03±0.01	-0.01±0.02	0.88
CEU	18	-0.04±0.03	-0.10±0.08	-0.06±0.14	0.67	-0.06±0.02	-0.04±0.02	0.00±0.02	0.79
WEU	10	-0.05±0.03	-0.08±0.05	-0.08±0.10	0.83	-0.04±0.02	-0.02±0.02	0.01±0.03	0.75
EEU	11	-0.07±0.04	-0.01±0.12	0.05±0.18	0.28	-0.08±0.02	-0.04±0.02	-0.01±0.02	0.78
SEU	5	-0.02±0.02	-0.06±0.05	-0.03±0.08	0.78	-0.02±0.02	-0.05±0.02	-0.02±0.03	0.75
WUS	6	-0.00±0.00	-0.01±0.01	-0.00±0.01	0.52	-0.00±0.00	-0.00±0.01	0.01±0.01	-0.11
NEUS	9	-0.03±0.01	-0.04±0.03	-0.01±0.05	0.29	-0.08±0.03	-0.03±0.02	0.02±0.03	0.52
MAUS	14	-0.02±0.01	-0.06±0.02	-0.02±0.04	0.57	-0.08±0.03	-0.04±0.02	0.00±0.02	0.73
GLUS	10	-0.02±0.02	-0.04±0.03	-0.01±0.06	0.11	-0.08±0.04	-0.03±0.02	0.01±0.02	0.41
SUS	4	-0.02±0.02	-0.03±0.02	-0.00±0.02	-0.05	-0.05±0.04	-0.03±0.03	0.00±0.04	0.37

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.




Fig. 1. Map of the selected regions for the analysis and measurement stations with long records of O_3 and SO_4^{2-} surface concentrations. North America (NA) [15°N – 55°N ; 60°W – 125°W], Europe (EU) [25°N – 65°N ; 10°W – 50°E], East Asia (EA) [15°N – 50°N ; 95°E – 160°E], and South Asia (SA) [5°N – 35°N ; 50°E – 95°E]. Triangles show the location of EMEP stations, squares of WDCGG stations, and diamonds of CASTNET stations. The stations are grouped in sub-regions: North Europe (NEU); Central Europe (CEU); West Europe (WEU); East Europe (EEU); South Europe (SEU); West US (WUS); North East US (NEUS); Mid-Atlantic US (MAUS); Great lakes US (GLUS); South US (SUS).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.




Fig. 2. Percentage changes in total anthropogenic emissions (CO, VOC, NO_x, SO₂, BC and OC) from 1980 to 2005 over the four selected regions as shown in Fig. 1: **(a)** Europe, EU; **(b)** North America, NA; **(c)** East Asia, EA; **(d)** South Asia, SA. In the legend of each regional plot, the total annual emissions for each species (Tg yr⁻¹) are reported for year 1980.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.




Fig. 3. Total annual natural and biomass burning emissions for the period 1980–2005: **(a)** Biogenic CO and VOCs emissions from vegetation; **(b)** NO_x emissions from lightning; **(c)** DMS emissions from oceans; **(d)** mineral dust aerosol emissions; **(e)** marine sea salt aerosol emissions; **(f)** CO, NO_x, and OC aerosol biomass burning emissions.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 4. Monthly mean anomalies for the period 1980–2005 of globally averaged fields for the SREF and SFIX ECHAM5-HAMMOZ simulations. Light blue lines are monthly mean anomalies for the SREF simulation, with overlaying dark blue giving the 12 month running averages. Red lines are the 12 month running averages of monthly mean anomalies for the SFIX simulation. The grey area represents the difference between SREF and SFIX. The global fields are respectively: **(a)** surface temperature (K); **(b)** tropospheric specific humidity (g Kg^{-1}); **(c)** O₃ surface concentrations (ppbv); **(d)** OH tropospheric concentration weighted by CH₄ reaction (molecules cm^{-3}); **(e)** SO₄²⁻ surface concentrations ($\mu\text{g m}^{-3}$); **(f)** total aerosol optical depth.

Reanalysis
1980–2005
ECHAM5-HAMMOZ

L. Pozzoli et al.




Fig. 5. Maps of surface O_3 concentrations and the changes due to anthropogenic emissions and natural variability. In the first column we show 5 yr averages (1981–1985) of surface O_3 concentrations over the selected regions, Europe, North America, East Asia, and South Asia. In the second column (**b**) we show the effect of anthropogenic emission changes in the period 2001–2005 on surface O_3 concentrations, calculated as the difference between SREF and SFIX simulations. In the third column (**c**) the natural variability of O_3 concentrations, effect of natural emissions and meteorology in the simulated 25 yr, calculated as the difference between 5 yr average periods (2001–2005) and (1981–1985) in the SFIX simulation. The combined effect of anthropogenic emissions and natural variability is shown in column (**d**) and it is expressed as the difference between 5 yr average period (2001–2005)–(1981–1985) in the SREF simulation.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 6. Trends of the observed (OBS) and calculated (SREF and SIFX) O_3 and SO_4^{2-} seasonal anomalies (DJF and JJA) averaged over each group of stations as shown in Fig. 1: North Europe (NEU); Central Europe (CEU); West Europe (WEU); East Europe (EEU); South Europe (SEU); West US (WUS); North East US (NEUS); Mid-Atlantic US (MAUS); Great lakes US (GLUS); South US (SUS). The vertical bars represent the 95% confidence interval of the trends. The number of stations used to calculate the average seasonal anomalies for each subregion is shown in parenthesis. Further details in Appendix B.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.





Fig. 7. Winter (DJF) anomalies of surface O_3 concentrations averaged over the selected regions (shown in Fig. 1). On the left y-axis anomalies of O_3 surface concentrations for the period 1980–2005 are expressed as the ratios between seasonal means for each year and year 1980. The blue line represents the SREF simulation (changing meteorology and changing anthropogenic emissions), the red line the SFIX simulation (changing meteorology and fixed anthropogenic emissions at the level of year 1980), while the gray area indicates the SREF-SFIX difference. On the right y-axis green and pink dashed lines represent the changes, ratio between each year and year 1980, of total annual VOC and NO_x emissions, respectively.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


**Reanalysis
1980–2005
ECHAM5-HAMMOZ**

L. Pozzoli et al.

**Fig. 8.** As Fig. 7 for summer (JJA) anomalies of surface O₃ concentrations.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.




Fig. 9. Global tropospheric O_3 budget calculated for the period 1980–2005 for the SREF (blue) and SFIX (red) ECHAM5-HAMMOZ simulations: **(a)** Chemical production (P); **(b)** chemical loss (L); **(c)** surface deposition (D); **(d)** stratospheric influx ($S_{\text{inf}} = L + D - P$); **(e)** tropospheric burden (B_{O_3}); **(f)** lifetime ($\tau_{\text{O}_3} = B_{\text{O}_3}/(L + D)$). The black points for year 2000 represent the mean \pm standard deviation budgets as found in the multi model study of Stevenson et al. (2006).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.




Fig. 10. As Fig. 5 but for SO_4^{2-} surface concentrations.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


**Reanalysis
1980–2005
ECHAM5-HAMMOZ**

L. Pozzoli et al.




Fig. 11. Winter (DJF) anomalies of surface SO_4^{2-} concentrations averaged over the selected regions (shown in Fig. 1). On the left y-axis anomalies of SO_4^{2-} surface concentrations for the period 1980–2005 are expressed as the ratios between seasonal means for each year and year 1980. The blue line represents the SREF simulation (changing meteorology and changing anthropogenic emissions), the red line the SFIX simulation (changing meteorology and fixed anthropogenic emissions at the level of year 1980), while the gray area indicates the SREF-SFIX difference. On the right y-axis pink dashed line represents the changes, ratio between each year and year 1980, of total annual sulfur emissions.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.




Fig. 12. As Fig. 11 for summer (JJA) anomalies of surface SO_4^{2-} concentrations.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.




Fig. 13. Global tropospheric SO_4^{2-} budget calculated for the period 1980–2005 for the SREF (blue) and SFIX (red) ECHAM5-HAMMOZ simulations: **(a)** total sulfur emissions; **(b)** SO_4^{2-} liquid phase production; **(c)** SO_4^{2-} gaseous phase production; **(d)** surface deposition; **(e)** SO_4^{2-} burden; **(f)** lifetime.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)





Fig. 14. Maps of total aerosol optical depth (AOD) and the changes due to anthropogenic emissions and natural variability. We show **(a)** the 5-yr averages (1981–1985) of global AOD; **(b)** the effect of anthropogenic emission changes in the period 2001–2005 on AOD, calculated as the difference between SREF and SFIX simulations; **(c)** the natural variability of AOD, effect of natural emissions and meteorology in the simulated 25 years, calculated as the difference between 5-yr average periods (2001–2005) and (1981–1985) in the SFIX simulation; **(d)** the combined effect of anthropogenic emissions and natural variability expressed as the difference between 5-yr average period (2001–2005)–(1981–1985) in the SREF simulation.

Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.




Fig. 15. Annual anomalies of total aerosol optical depth (AOD) averaged over the selected regions (shown in Fig. 1). On the left y-axis anomalies of AOD for the period 1980–2005 are expressed as the ratios between annual means for each year and year 1980. The blue line represents the SREF simulation (changing meteorology and changing anthropogenic emissions), the red line the SFIX simulation (changing meteorology and fixed anthropogenic emissions at the level of year 1980), while the gray area indicates the SREF-SFIX difference. On the right y-axis pink and green dashed lines represent the clear-sky aerosol anthropogenic radiative perturbation at the top of the atmosphere ($RP_{\text{aer}}^{\text{TOA}}$) and at surface ($RP_{\text{aer}}^{\text{Surf}}$), respectively.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Reanalysis 1980–2005 ECHAM5-HAMMOZ

L. Pozzoli et al.




Fig. A1. Taylor diagrams comparing the ECHAM5-HAMMOZ SREF simulation with EMEP, CASTNET, and WDCGG observations of O_3 seasonal means **(a)** DJF and **(b)** JJA and SO_4^{2-} seasonal means **(c)** DJF and **(d)** JJA. The black dot is used as reference to which simulated fields are compared. Continuous grey lines show iso-contours of skill score. Stations are grouped by regions as in Fig. 1: North Europe (NEU); Central Europe (CEU); West Europe (WEU); East Europe (EEU); South Europe (SEU); West US (WUS); North East US (NEUS); Mid-Atlantic US (MAUS); Great lakes US (GLUS); South US (SUS).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. A2. Annual trends of O_3 and SO_4^{2-} surface concentrations. The trends are calculated as linear fitting of the annual mean surface O_3 and SO_4^{2-} anomalies for each grid box of the model simulation SREF over Europe and North America. The grid boxes with not statistically significant (p -value > 0.05) trends are displayed in grey. The colored circles represent the observed trends for EMEP and CASTNET measuring stations, over Europe and North America, respectively. Only the stations with statistically significant (p -value < 0.05) trends are plotted.