

**Spatiotemporal  
distribution of  
light-absorbing  
carbon**

P. Krecl et al.

# Spatiotemporal distribution of light-absorbing carbon and its relationship to other atmospheric pollutants in Stockholm

P. Krecl<sup>1,\*</sup>, A. C. Targino<sup>2</sup>, and C. Johansson<sup>1,3</sup>

<sup>1</sup>Department of Applied Environmental Science, Atmospheric Science Unit, Stockholm University, Stockholm, Sweden

<sup>2</sup>Universidade Tecnológica Federal do Paraná, Londrina, Brazil

<sup>3</sup>Stockholm Environment and Health Administration, Stockholm, Sweden

\*now at: The School of Earth and Environment, University of Leeds, Leeds, UK

Received: 19 February 2011 – Accepted: 20 April 2011 – Published: 29 April 2011

Correspondence to: P. Krecl (p.krecl@leeds.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

Carbon-containing particles have deleterious effects on both Earth's climate and human health. In Europe, the main sources of light-absorbing carbon (LAC) emissions are the transport (67%) and residential (25%) sectors. Information on the spatiotemporal variability of LAC particles in urban areas is relevant for air quality management and to better diagnose the population exposure to these particles. This study reports on results of an intensive field campaign conducted at four sites (two kerbside stations, one urban background site and a rural station) in Stockholm, Sweden, during the spring 2006. Light-absorbing carbon mass concentrations ( $M_{LAC}$ ) were measured with custom-built Particle Soot Absorption Photometers (PSAP). The spatiotemporal variability of  $M_{LAC}$  concentrations was explored by examining correlation coefficients ( $R$ ), coefficients of divergence (COD), and diurnal patterns at all sites. Simultaneous measurements of  $NO_x$ ,  $PM_{10}$ ,  $PM_{2.5}$ , and meteorological variables were also carried out at the same locations to help characterize the LAC emission sources.

Hourly mean and standard deviation  $M_{LAC}$  concentrations ranged from 0.36 (rural) to  $5.39 \mu g m^{-3}$  (street canyon) and from 0.50 to  $3.60 \mu g m^{-3}$ , respectively. Concentrations of LAC between urban sites were poorly correlated even for daily averages ( $R < 0.70$ ), combined with highly heterogeneously distributed concentrations ( $COD > 0.30$ ) even at spatial scales of few kilometers. This high variability is connected to the distribution of emission sources and processes contributing to the LAC fraction at these sites. At urban sites,  $M_{LAC}$  tracked  $NO_x$  levels and traffic density well and mean  $M_{LAC}/PM_{2.5}$  ratios were larger (26–38%) than at the background sites (4–10%). The results suggest that vehicle exhaust emissions are the main responsible for the high  $M_{LAC}$  concentrations found at the urban locations whereas long-range transport (LRT) episodes of combustion-derived particles can generate a strong increase of levels at background sites.

### Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



To decrease pollution levels at kerbside and urban background locations in Stockholm, we recommend abatement strategies that target reductions of vehicle exhaust emissions, which are the main contributors to  $M_{LAC}$  and  $NO_x$  concentrations.

## 1 Introduction

5 Primary carbonaceous particles are mostly emitted by combustion processes (transport, residential sector, industry, and power generation) and represent a large fraction of the submicron aerosol. Regarding the optical properties of the carbonaceous material, it is usually described as two major constituents: light-absorbing carbon (LAC), with strong absorption of visible light, and organic carbon (OC), with weak absorption in the  
10 visible electromagnetic spectrum. Despite the detrimental effect of carbon-containing particles on human health (Lighty et al., 2000; Franco Suglia et al., 2007) and the potential of the LAC fraction to modify the radiative budget regionally (Ramanathan et al., 2007) and to have other adverse effects on climate (e.g., Ramanathan et al., 2001; Ackerman et al., 2000; Menon et al., 2002), their emission and ambient concentrations  
15 are still not addressed by local regulations worldwide.

The main emission sources of LAC in Europe are road transport (49%), residential combustion of solid fuels (25%), and off-road transport (18%) according to estimations by Kupiainen and Klimont (2007) for the year 2000. In Sweden, the main sources contributing to LAC emissions are road transport (37%), residential combustion of biofuels  
20 (25%), and off-road transport (23%) as extracted from the Bond et al. (2004) global inventory based on 1996 fuel-use data. Streets et al. (2004) projected a reduction of LAC emissions in Europe in 2030 and 2050 (36–64% depending on the IPCC scenario), mainly due to the introduction of advanced vehicle technology. A 40% decrease of LAC emissions is predicted in Sweden for the year 2020 compared to 2007, associated to more stringent particulate emission standards for diesel engines and other vehicles (R. Abrahamsson, personal communication, Swedish Environmental Protection  
25

### Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Agency, 2011). However, advances in reducing atmospheric emissions by the development of cleaner vehicle technologies could be balanced or even exceeded by increasing traffic density in many cities.

Compared to rural zones, urban areas are characterized by higher population density, larger number of pollutant emission sources, higher atmospheric pollutants concentrations, and a higher LAC content in the fine particle ( $PM_{2.5}$ ) fraction (Putaud et al., 2004). Simultaneous and continuous time-resolved LAC measurements are required in different parts of the city to (1) characterize the carbon carbon-containing particle sources which can be used to streamline successful air quality control strategies to safeguard public health, (2) understand the particles fate and transport in the urban atmosphere, (3) study the spatiotemporal variability of LAC particles to better diagnose the population exposure to these particles, and (4) determine to what extent the limit values for  $PM_{2.5}$  and  $PM_{10}$  regulate the LAC concentrations, since LAC would be a good candidate substance to be regulated if both climate and health effects of emissions are to be considered.

Stockholm is the largest metropolitan region in Scandinavia (1.25 million inhabitants in the urban area) and has an extensive air quality monitoring program created by the Stockholm Environment and Health Administration and operated by SLB analys. For several atmospheric pollutants (carbon monoxide, sulphur dioxide, benzene, lead, and benzo[a]pyrene) Stockholm concentrations are below the national environmental quality standards (which are the same as the EU directives) even on the busiest streets in the inner city (Johansson et al., 2009). However,  $NO_2$  and  $PM_{10}$  mass concentrations still exceed the air quality standards in Stockholm city in connection to high traffic emissions (Johansson et al., 2009). High  $PM_{10}$  concentrations are found in the inner city in springtime mostly due to particle resuspension, as a result of the use of studded tires when roads are free from snow (Norman and Johansson, 2006). While the spatial variability of particle mass concentrations (Johansson et al., 2007a) and particle number concentrations (Gidhagen et al., 2004a,b, 2005) over Stockholm has already been assessed in previous studies, considerably less is known about the distribution

## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of LAC particles within the city (Johansson et al., 2007b) and the relationship between the LAC particle fraction and other atmospheric pollutants.

This work investigates the spatiotemporal variation of hourly light-absorbing carbon concentrations in Stockholm city in springtime. First, the performance of the custom-built instruments is evaluated in ambient air by comparing their measurements with those from commercial instruments. Then, descriptive statistics are presented for different monitoring sites and the spatial and temporal variation of the LAC mass concentrations ( $M_{LAC}$ ) is also examined. To help characterize the LAC emission sources, we also report the relationships between LAC and other atmospheric pollutants ( $PM_{2.5}$ ,  $PM_{10}$ , and  $NO_x$ ), the diurnal cycles of these pollutants, and the wind direction dependence of the concentrations. Finally, the urban and rural contributions to the atmospheric pollutants load are discussed.

## 2 Description of measurement sites

An intensive field study was conducted simultaneously at three monitoring sites in central Stockholm (Hornsgatan, Torkel Knutssonsgatan (referred to as “Torkel”), and Essingeleden) and at a rural station (Aspvreten) in spring 2006 (Figs. 1 and 2). Hornsgatan is a four-lane 24 m wide street canyon with 24 m high buildings. Instruments were housed in a trailer parked on the street with a traffic volume of  $\sim 28\,000$  vehicles per day. On average, the composition of the vehicle fleet is: passenger cars 77%, light-duty vehicles 12%, buses 2%, and heavy-duty vehicles 1%. Half of the vehicle fleet runs on gasoline, 30% uses diesel (21% of the passenger cars, most of the taxis and heavy-duty vehicles), and 14% are ethanol vehicles (80% of the buses and 15% of the passenger cars) (SLB, 2010). Torkel is located on a rooftop platform (25 m height) and thereby represents urban background concentrations for the Stockholm region, since it is not directly affected by nearby atmospheric emissions. Measurements were also carried out near Essingeleden, on the island of Lilla Essingen, by the kerb of the eight-lane E4 highway with a traffic flow of  $\sim 128\,000$  vehicles per day. The distance

### Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



between Hornsgatan and Torkel sites is 450 m, Hornsgatan-Essingeleden distance is 2.7 km, and Torkel and Essingeleden sites are 3.1 km apart. Aspvreten is a rural background station operated by ITM and is part of two European-scale aerosol monitoring networks: the European Monitoring and Evaluation Programme (EMEP) and the European Supersites for Atmospheric Aerosol Research (EUSAAR). The station is located 80 km southwest of Stockholm and 2 km inland from the Baltic Sea, and surrounded by deciduous and coniferous forest and grasslands. Measurements conducted at Norr Malma rural background station, located in a rural area ~ 70 km northeast of Stockholm and 1 km south of Lake Erken (Fig. 1), were used to complete or complement Aspvreten data.

### 3 Instrumentation and sampling program

#### 3.1 Carbonaceous aerosol

Measurements of light-absorbing carbon were performed using custom-built Particle Soot Absorption Photometers (PSAP) at the four sites (central Stockholm and Aspvreten) and two commercial Aethalometers installed at Hornsgatan and Torkel. The measurement principle of these instruments is the same and relies on the light absorbing properties of the carbonaceous aerosols. The method is based on the integrating plate technique (Lin et al., 1973) that measures the attenuation of light transmitted through particles that are continuously collected on a filter. The LAC mass absorption cross section  $\sigma_a$  relates the absorption coefficient of the aerosol to the LAC mass concentration  $M_{LAC}$ . The calculation of  $\sigma_a$  requires the determination of  $M_{LAC}$  by an independent method (e.g., thermo-optical measurements).

The Aethalometers series 8100 (Magee Scientific, USA) were operated with PM<sub>2.5</sub> sample inlets (installed at 3.5 m a.g.l.) and a sample flow rate of 2 l min<sup>-1</sup>. The instruments reported 1-min values, calculated from 5-min moving averages, and 15-min average values were logged into the acquiring system. Aethalometer concentrations

## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Spatiotemporal  
distribution of  
light-absorbing  
carbon**

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

were computed using the mass specific attenuation cross section provided by the manufacturer ( $16.6 \text{ m}^2 \text{ g}^{-1}$ ). In this study, mean offset values of  $0.5 \mu\text{g m}^{-3}$  and  $0.21 \mu\text{g m}^{-3}$  were subtracted from Aethalometer  $M_{\text{LAC}}$  concentrations measured at Hornsgatan and Torkel, respectively. These offset values were found when operating the Aethalometer in the field with particle-free air drawn through the filter for several days. Similar Aethalometer offset values were observed by the instrument supplier when running baseline measurements (Krecl et al., 2007).

The custom-built PSAPs were constructed at ITM, Stockholm University (herein after referred to as PSAP-ITM) and a detailed description can be found in Krecl et al. (2007, 2010). For this campaign, sample inlets were installed at  $\sim 2 \text{ m a.g.l.}$  and no pre-cut size devices were used. The PSAP particle sampling was conducted through  $1/4''$  stainless steel tubing with the flow rate controlled by a needle valve ( $\sim 0.06 \text{ l min}^{-1}$  at Central Stockholm sites, and  $0.2 \text{ l min}^{-1}$  at the rural site). Data were logged every 30 s and a 4-min moving average was computed for all voltages prior to calculating the attenuation coefficients. Several corrections were applied to PSAP measurements, as described in Krecl et al. (2007, 2010), to obtain the absorption coefficient of the airborne particles: filter property correction, filter loading correction, scattering correction and additional absorption correction. To correct for aerosol light scattering aerosol deposited onto the PSAP filter, two approaches were used depending on the data availability at each site: (1) aerosol scattering coefficient measurements performed at Hornsgatan in the period 23 May–6 July 2006 using an integrating nephelometer (520 nm wavelength, model M9003, Ecotech, Australia), (2) aerosol light-scattering coefficients estimated from  $\text{PM}_{2.5}$  mass measurements multiplied by the aerosol mass scattering coefficient (Krecl et al., 2007). For the urban sites, the mass scattering coefficient was obtained when linearly correlating daily mean  $\text{PM}_{2.5}$  and aerosol light scattering measurements at Hornsgatan in the period 23 May–30 August 2006 (slope of  $4.0 \pm 0.37$  (95% confidence interval), correlation coefficient  $R = 0.92$ , number of samples: 99). Essingeleden  $\text{PM}_{2.5}$  measurements were used in these calculations when Hornsgatan data were missing. Finally, a mass scattering coefficient of  $4.0 \text{ m}^2 \text{ g}^{-1}$  was

## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



used to correct all Stockholm PSAP calculations which agrees well with reported values for polluted continental air ( $3.5 \pm 1.2 \text{ m}^2 \text{ g}^{-1}$ , IPCC, 2001). In the case of Aspvreten, a mass scattering coefficient of  $2.0 \text{ m}^2 \text{ g}^{-1}$  was taken from the literature for rural background conditions (IPCC, 2001; Pereira et al., 2008) and  $\text{PM}_{2.5}$  concentrations were estimated as 80% of  $\text{PM}_{10}$  levels (Areskoug et al., 2004) when  $\text{PM}_{2.5}$  data were missing.

Finally, to convert the aerosol light absorption coefficient into  $M_{\text{LAC}}$  we assumed that all light absorption was from light-absorbing carbon and that all LAC had the same mass absorption cross section at each site over the sampling period. In situ LAC mass absorption cross sections  $\sigma_a$  were determined by linearly correlating aerosol light-absorption coefficients (after applying all corrections) with elemental carbon (EC) mass concentrations. For EC/OC analysis, quartz fiber filters were collected at Torkel and Aspvreten using a  $10 \mu\text{m}$  cut-off size sequential sampler (model SEQ 47/50, Leckel GmbH, Germany) installed at 1.6 m above rooftop level operated at a standard flow rate of  $38.3 \text{ l m}^{-1}$ . Filters were pre-baked in a furnace at  $800^\circ\text{C}$  for 150 min and placed in clean glass Petri dishes and kept refrigerated before and after the filter sampling. The EC/OC analysis was carried out using a Thermal/Optical Carbon Aerosol Analyzer (Sunset Laboratory Inc., Forest Grove, USA) operating on the NIOSH method 5040 (Birch and Cary, 1996). Mass specific absorption cross sections of  $7.6 \text{ m}^2 \text{ g}^{-1}$  ( $R = 0.80$ , number of samples: 9) and  $10 \text{ m}^2 \text{ g}^{-1}$  ( $R = 0.90$ , number of samples: 10) were obtained at Torkel and Aspvreten, respectively. The  $\sigma_a$  value for Torkel is in agreement with freshly generated light-absorbing carbon values ( $7.5 \pm 1.2 \text{ m}^2 \text{ g}^{-1}$ ) summarized by Bond and Bergstrom (2005) at a wavelength of 550 nm and was used for all Stockholm sites. The Aspvreten value is typical of aged LAC aerosol at background sites (Cozic et al., 2008).

### 3.2 Other measurements

$\text{PM}_{10}$  and  $\text{PM}_{2.5}$  mass measurements were performed at all sites using automatic Tapered Element Oscillating Microbalance (TEOM 1400a, Rupprecht & Patashnick Inc.,

USA) instruments with heated inlets (50 °C) to avoid condensing water. To account for losses of volatile material on the particles, all TEOM data were corrected following Areskoug (2007). In short, built-in TEOM corrections were removed, mass concentrations were expressed at ambient pressure and temperature and, finally, concentrations were multiplied by 1.19 and an offset of 1.15 was added.

Because of the short PM<sub>2.5</sub> time series at Aspvreten (started on 22 June 2006), Norr Malma PM<sub>2.5</sub> measurements were used as a replacement of Aspvreten data only for calculations of average values. NO<sub>x</sub> concentrations were measured at the three monitoring sites in central Stockholm and at Norr Malma station using commercial chemiluminescence analyzers (model 31 M LCD, Environment SA, France). Total traffic counts were recorded at Hornsgatan and Essingeleden using automatic counters based on magnetic impedance. Meteorological measurements were conducted at Torkel and Aspvreten including air temperature, relative humidity, wind speed (WS), wind direction (WD), precipitation, atmospheric pressure, and global solar radiation. Sensors were installed at 2 m above roof at Torkel and 2 m above ground at Aspvreten, except for the anemometer installed at 10 m height.

## 4 Results and discussion

### 4.1 Comparison of aerosol light-absorption measurements

Krecl et al. (2007), when comparing Aethalometer and PSAP-ITM measurements in an area impacted by woodsmoke, showed that the custom-built PSAP was a reliable alternative for the commercial Aethalometer. Because of the different physicochemical characteristics of combustion particles from vehicle exhaust and residential woodsmoke (Kocbach et al., 2005), we evaluate here the performance of the custom-built PSAP at a kerbside site (Hornsgatan) and at an urban background monitoring station (Torkel). The PSAP-ITM instruments used in this study were previously intercompared in the laboratory when sampling air from outdoors and showed a very high correlation for

## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



15-min averages ( $R = 0.99$ ) (Krecl et al., 2010). Figure 3 displays the linear regression between hourly PSAP-ITM  $b_{ap}$  and Aethalometer  $M_{LAC}$  for measurements carried out at Hornsgatan (left panel) and at Torkel (right panel) in the period 20 April–5 July 2006. The two methods were found to be highly correlated ( $R > 0.85$ ) for both sites, but a large  $y$ -intercept ( $-7.2 \text{ M m}^{-1}$ ) was observed when correlating the Aethalometer and PSAP-ITM measurements at Hornsgatan, even after subtracting an offset value as described previously. If instead the PSAP-ITM  $M_{LAC}$  concentrations are linearly correlated with the Aethalometer mass concentrations, the slope of the regression line is  $1.4 \pm 0.03$  (95% confidence interval) for both sites (not shown). The slope being larger than 1 might partly result from (a) the use of an Aethalometer mass specific attenuation cross that is not site and season specific for this study (Sharma et al., 2002), and (b) biases of the thermo-optical measurements of EC (Bond and Bergstrom, 2005) used to determine the custom-built PSAP $\sigma_a$ .

## 4.2 General overview

During the selected sampling period (20 April–5 July 2006), daily mean temperatures ranged between  $3.7^\circ\text{C}$  and  $23.6^\circ\text{C}$  at Torkel, and between  $3.1^\circ\text{C}$  and  $19.6^\circ\text{C}$  at Aspvreten. The mean wind speed was  $3.5 \text{ m s}^{-1}$  at Torkel (rooftop) and the prevailing wind directions were south (22%), southwest (18%), and west (15%) whereas Aspvreten mean WS was  $1.8 \text{ m s}^{-1}$  and the dominant winds were from south (22%), southeast (20%), and east (15%).

Descriptive statistics of hourly mean  $M_{LAC}$  concentrations measured with the PSAP-ITM are displayed in Table 1 for the period of simultaneous measurements. Missing data corresponded to time periods when the instruments were not operating or were known to be operating improperly (e.g., filter transmittance below the recommended threshold, voltage signal higher than the maximum data logger input voltage, maintenance for filter change or flow rate measurement).

Mean  $M_{LAC}$  concentrations ranged from 0.36 to  $5.39 \mu\text{g m}^{-3}$ , and standard deviation (SD) varied between 0.50 and  $3.60 \mu\text{g m}^{-3}$ . As expected, higher values of  $M_{LAC}$

### Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

concentrations were found at the street canyon site whereas lower levels always corresponded to the rural station. The 95th percentile at Hornsgatan was 12 times higher than the 95th percentile at Aspvreten even for daily averages (not shown), and the lowest values at Hornsgatan (5th percentile) were higher than the average at Aspvreten (1.44  $\mu\text{g m}^{-3}$  versus 0.36  $\mu\text{g m}^{-3}$ ). The other two urban sites presented intermediate concentrations, with Essingeleden (close to the highway) showing on average 2.8 times higher concentrations than the urban rooftop site. Street level concentrations of  $M_{\text{LAC}}$  in Stockholm (Hornsgatan and Essingeleden) were higher than concentrations measured in a residential area impacted by woodsmoke in Northern Sweden during wintertime (Krecl et al., 2007, 2010). Compared to other European kerbside sampling sites, mean  $M_{\text{LAC}}$  concentrations at Hornsgatan and Essingeleden were within the 3.11–10.68  $\mu\text{g m}^{-3}$  annual average range reported by Putaud et al. (2004). Average values at Aspvreten were similar to other European rural background sites (Putaud et al., 2004), with concentrations lower than 1  $\mu\text{g m}^{-3}$ . The Torkel median  $M_{\text{LAC}}$  concentration is comparable to the median value observed at a Tokyo rooftop site (1.18  $\mu\text{g m}^{-3}$ , ~20 m height) in the period 2003–2005 (Kondo et al., 2006). Mean  $M_{\text{LAC}}$  concentrations at two New York city sites were 1.01 and 1.38  $\mu\text{g m}^{-3}$  for wintertime (Venkatachari et al., 2006), being comparable to Torkel levels but lower than Hornsgatan and Essingeleden mean values.

A large hour-to-hour and day-to-day variability of  $M_{\text{LAC}}$  concentrations was observed at the kerbside sites whereas concentrations at the rural background station showed very low levels. Figure 4 illustrates the spatiotemporal variability for one example week from 14 May through 20 May 2006 when no long-range transport (LRT) from polluted areas was detected in Stockholm (as observed in HYSPLIT4 backward trajectory analysis and corroborated by the low levels at the rural site Aspvreten). More tellingly, the maximum hourly concentration at Hornsgatan reached 15.84  $\mu\text{g m}^{-3}$  at 11:00 LT whereas a minimum value of 2.15  $\mu\text{g m}^{-3}$  was observed at 21:00 LT on Friday 19 May. At Aspvreten,  $M_{\text{LAC}}$  concentrations ranged between 0.20 and 0.32  $\mu\text{g m}^{-3}$  for the same day. During that week, the daily mean temperature at Torkel varied between 7.9 °C

## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

and 10.4 °C with low precipitation records on 17 May (0.6 mm), 18 May (3.8 mm), and 20 May (5.6 mm). Aspvreten weather was colder than at Torkel with air temperatures between 5.1 °C and 9.0 °C and precipitations occurred on 17 May (1.0 mm), 18 May (2.6 mm), and 20 May (3.2 mm).

During the present field campaign, a long-range transport of pollutants emitted by agricultural wildfires in Russia and the Baltic region had a strong impact on aerosol concentrations in Sweden in the period 24 April–9 May 2006, mainly for background sites (Targino et al., 2011). On average,  $M_{\text{LAC}}$  concentrations were 4 times higher at Torkel and 7 times higher at Aspvreten during this LRT episode compared to the rest of the study period.

### 4.3 Spatiotemporal variability of $M_{\text{LAC}}$ concentrations

Ambient particle concentrations vary spatially and temporally to different degrees at the intraurban scale and two statistical tools are commonly used to determine whether a pollutant concentration is homogeneously distributed in time and space. Correlation coefficients accurately track the temporal similarity of paired measurements conducted at two sites, while coefficients of divergence (COD) are often used to analyze the spatial uniformity of the observations (Wilson et al., 2005).

The Pearson correlation coefficients between  $M_{\text{LAC}}$  time series were calculated for hourly and daily (starting at midnight) mean concentrations during the period of simultaneous measurements and the results are displayed in Table 2. All correlation coefficients were significant at the 99% level (not shown). Daily correlations were higher than hourly correlations for all site pairs. However, daily  $M_{\text{LAC}}$  time series were still poorly correlated except for Aspvreten and Torkel measurements (hourly  $R = 0.78$ ; daily  $R = 0.88$ ). The high correlation between Aspvreten and Torkel could be explained by the long-range transport origin of the  $M_{\text{LAC}}$  observed at the two sites since these sites were not affected by local emission sources.

The coefficients of divergence were calculated according to Wongphatarakul et al. (1998) and results from collocated hourly and daily  $M_{\text{LAC}}$  concentrations in the period

## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



23.0 at the kerbside locations, which is consistent with ratios observed close to major roads (distance of 5–10 m) in Japanese urban areas (Naser et al., 2009). The mean  $M_{\text{LAC}}/\text{PM}_{2.5}$  ratio is larger at the kerbside stations (38% at Hornsgatan, and 26% at Essingeleden) than at the background sites (10% at Torkel, and 4% at Aspvreten) as observed by Putaud et al. (2004) when studying the chemical characteristics of the particulate matter across Europe. The strongly decreasing  $\text{PM}_{2.5}/\text{PM}_{10}$  ratio from rural (0.69) to kerbside sites (0.26), indicates a higher emission of coarse particles in the urban area (e.g. road dust emissions, Norman and Johansson, 2006).

Linear correlations between hourly concentrations of atmospheric pollutants ( $M_{\text{LAC}}$ ,  $\text{PM}_{10}$ ,  $\text{PM}_{2.5}$ , and  $\text{NO}_x$ ) measured at each station were calculated in the period 20 April–5 July 2006: (1) excluding data during the LRT event (24 April–9 May 2006), (2) including all days. Table 5 displays the inner city correlation coefficients only when local sources dominated the pollutants emissions (period 1). The number of valid samples ranged from 171 to 1451 because not all instruments were always operating correctly, and this variation has to be considered when comparing  $R$  values. High  $R$  values ( $R > 0.75$ ) were only observed for the variable combination  $M_{\text{LAC}}\text{-NO}_x$  at all urban stations, suggesting that both compounds originated mainly from vehicle traffic emissions. For period 2, the correlation between  $M_{\text{LAC}}$  and  $\text{NO}_x$  concentrations decreased to 0.47 at Torkel (not shown) and remained constant for the kerbside sites. This finding indicates a strong contribution of the LRT episode to  $M_{\text{LAC}}$  concentrations at Torkel whereas  $\text{NO}_x$  was emitted by local pollution sources. The correlation between  $M_{\text{LAC}}$  and  $\text{PM}_{10}$  time series increased from 0.45 to 0.85 at Aspvreten (not shown) when including the LRT episode, suggesting a common aerosol origin. However, the correlation between  $M_{\text{LAC}}$  and  $\text{PM}_{2.5}$  at Aspvreten was weak and might be probably caused by the relative small number of valid measurements.

### 4.5 Diurnal patterns

The diurnal variation of mean  $M_{\text{LAC}}$ ,  $\text{PM}_{10}$ ,  $\text{PM}_{2.5}$ , and  $\text{NO}_x$  concentrations for the 4 sites is shown in Fig. 5, together with the daily pattern of traffic density at Hornsgatan

and Essingeleden. Measurements were classified as weekdays (Monday–Friday) and weekends (Saturday/Sunday) and mean hourly values were calculated for the period 20 April–5 July 2006. PM<sub>2.5</sub> data from Norr Malma replaced Aspvreten measurements for this calculation due to the low data availability (only 10%). Because of missing PM<sub>2.5</sub> observations at Hornsgatan (36% data coverage), PM<sub>2.5</sub> weekend daily patterns at these sites were not fully synchronized with the other observations. Time displayed in Fig. 5 corresponds to LT.

It can be observed that pollutants concentrations were generally higher on weekdays than at weekends between early morning and evening at the urban sites. To test the significance of the weekday concentrations being higher than the weekend levels, an unpaired *t*-test was performed on the mean difference concentration for each hour of the day at 95% confidence level. Measurements at Aspvreten showed no significant difference between weekend and weekdays for all variables analyzed, indicating no evidence of local anthropogenic activities. Weekday concentrations were statistically significantly higher than weekend levels for: (1)  $M_{LAC}$  concentrations only at the kerbside stations during daytime (07:00–17:00 at Hornsgatan, and 06:00–20:00 at Essingeleden), (2) PM<sub>10</sub> concentrations in the morning (08:00–11:00) at Hornsgatan and Torkel, and in the morning and afternoon at Essingeleden (07:00–20:00), (3) NO<sub>x</sub> levels at the three urban sites during daytime (06:00–20:00), (4) PM<sub>2.5</sub> concentrations only at Hornsgatan in the period 07:00–20:00 (this result should be considered with caution as previously mentioned).

On weekdays, the pollutants concentration rise observed at the kerbside stations in the early morning (05:00–06:00) was consistent with an increase in vehicle traffic, and maximum values were reached at ~09:00. Peak values were followed by a gradual decrease in  $M_{LAC}$  and NO<sub>x</sub> concentrations at both sites through the afternoon, partially explained by the growth of the mixing layer depth and strong turbulent mixing during the afternoon. A similar pattern for NO<sub>x</sub> was observed at Hornsgatan by Olivares et al. (2007) when analyzing diurnal variations on weekdays and at weekends in the period June 2002–September 2004. Hornsgatan  $M_{LAC}$  diurnal cycle was similar to the pattern

## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

observed at Vallila kerbside station in Helsinki during three campaigns conducted between 1996 and 2005 (Järvi et al., 2008). However, Hornsgatan median concentrations were much higher than Valilla levels (4.47 versus 1.11  $\mu\text{g m}^{-3}$ ) as well as morning peak values on weekdays (9.69 versus 2.40  $\mu\text{g m}^{-3}$ ).

Weekday and weekend daily patterns of traffic rate were distinct for Hornsgatan and Essingeleden as depicted in Fig. 5i–j. On average, the traffic volume at Essingeleden was 4.5 times higher than at Hornsgatan. On weekdays, two peaks represented the morning (06:00–09:00) and afternoon (14:00–18:00) traffic rush hours at both sites with moderate traffic volume between these periods. In the early morning (00:00–05:00) at weekends, the traffic rate was higher at weekends than on weekdays, particularly at Hornsgatan station.

Even though the vehicle traffic was more intensive at Essingeleden than at Hornsgatan, pollutants concentrations were usually higher at Hornsgatan. This is most likely mainly connected to poor ventilation conditions at Hornsgatan because of the street canyon geometry. Higher  $\text{PM}_{10}$  concentrations were observed at Essingeleden than at Hornsgatan during the afternoon on weekdays, which might be explained by higher road dust emissions due to the higher vehicle speed at Essingeleden than at Hornsgatan. We found a positive wind speed dependence of  $\text{PM}_{10}$  concentrations for  $\text{WS} > 5 \text{ m s}^{-1}$  at these urban sites (an especially important for Essingeleden), in contrast to the dilution of  $M_{\text{LAC}}$  observed when the WS increased (not shown). This is consistent with results by Vogt et al. (2010) who reported a WS dependence of the coarse particle fluxes over Stockholm in the period April 2008–April 2009. These findings confirm that  $\text{PM}_{10}$  to a large degree is due to non-exhaust particles, whereas  $M_{\text{LAC}}$  is mainly due to vehicle exhaust particles. At weekends, a reduced number of vehicles circulated and the contribution of road resuspended particles to the  $\text{PM}_{10}$  fraction was smaller than on weekdays. Thus, Hornsgatan  $\text{PM}_{10}$  concentrations were slightly higher than at Essingeleden on Saturday/Sunday.

During early morning hours (00:00–05:00), Hornsgatan  $M_{\text{LAC}}$  and  $\text{NO}_x$  concentrations were found statistically significantly higher at weekends than on weekdays

## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

(unpaired  $t$ -test, 95% confidence level). This rise beyond the evening rush hour at Hornsgatan might be due to the higher early morning traffic at weekends than on weekends, mainly composed of diesel taxis (SLB, 2010) and possibly associated to leisure-time activities in Stockholm downtown area.

#### 4.6 Wind direction dependence

Wind direction is a key variable to determine the relative position of the measuring sites to the main pollutants emission sources. To study the pollutants concentration dependence on the wind direction, the concentration roses of the pollutants time series were calculated for each site. Figure 6 displays the  $M_{LAC}$ ,  $PM_{10}$ , and  $NO_x$  concentration roses at the four sites for the whole period of simultaneous measurements (Fig. 6a–c), and excluding measurements during the LRT event (Fig. 6d–f). Due to the poor data coverage of  $PM_{2.5}$  measurements at Hornsgatan and Aspvreten,  $PM_{2.5}$  concentration roses were calculated only for Essingeleden and Torkel (not shown). The average of hourly concentrations was calculated for each WD sector including only data records with valid measurements from all sites.

Pollutants concentrations at the kerbside stations are higher than backgrounds levels for all wind directions.  $M_{LAC}$  and  $NO_x$  concentrations roses showed the same pattern for the kerbside stations during both periods (i.e. including and excluding the LRT event). This suggests that local emission sources (i.e. traffic exhaust) were the main contributors to  $M_{LAC}$  and  $NO_x$  measurements at Hornsgatan and Essingeleden. However, Torkel and Aspvreten concentration roses for  $M_{LAC}$  showed a distinct behavior. When the LRT episode was included,  $M_{LAC}$  concentrations were higher at both sites for E and SE winds (Fig. 6a), matching the predominant WD at Torkel and Aspvreten during the agricultural wildfires episode in Russia.  $M_{LAC}$  levels were lower at the background sites when the LRT episode was excluded and the WD dependence was negligible (Fig. 6d).  $PM_{10}$  concentrations showed a positive WD dependence at all sites for SE and E winds when the LRT event was present in the analysis (Fig. 6b).

## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Hornsgatan concentration rises for  $M_{LAC}$  and  $NO_x$  presented the highest values associated with the NE and E sectors matching the street canyon orientation of  $75^\circ$  from the north towards the east. Gidhagen et al. (2004b) showed that modeled  $NO_x$  levels at the Hornsgatan site were improved when traffic emissions from the surrounding streets, especially in the sector  $40^\circ$ – $140^\circ$ , were included in the simulations. This suggests that  $M_{LAC}$  and  $NO_x$  emitted by traffic in the neighborhood (NE-E of Hornsgatan site) might have been transported along the street canyon contributing to the high load observed at Hornsgatan.

These results indicate that  $M_{LAC}$  was mainly emitted by vehicle traffic at the kerbside stations whereas the LRT contribution to the LAC concentrations was masked at these sites.  $NO_x$  was mostly emitted by traffic sources in the urban area and no agricultural wildfires influence was observed on  $NO_x$  levels at the three sites. Both  $PM_{10}$  and  $PM_{2.5}$  concentrations were strongly impacted by the LRT event at all sites.

#### 4.7 Urban versus rural contributions

Figure 7 displays the scattergram of  $NO_x$  versus  $M_{LAC}$  concentrations at Hornsgatan, Essingeleden, and Torkel. Only measurements conducted on weekdays and outside the LRT event period were averaged separately for each hour of the day and then linearly correlated. For the kerbside stations, the weekday diurnal variations of  $M_{LAC}$  and  $NO_x$  showed a high linear correlation ( $R > 0.95$ ), indicating that these compounds had common emission sources. Hornsgatan and Essingeleden sites presented similar slopes of 24.9 and 27.3, respectively. At Torkel, the linear correlation between  $M_{LAC}$  and  $NO_x$  diurnal variation was lower ( $R = 0.87$ ) than at the kerbside sites. The correlation at Torkel increased to  $R = 0.99$  when measurements were restricted to the time period 08:00–18:00 LT and the slope was  $28.5 \pm 8.6$  (not shown). The  $x$ -intercept of the regression lines in Fig. 7 represents an estimate of a “clean” regional background of  $M_{LAC}$  that is not influenced by local emissions (assuming that  $NO_x$  concentrations are zero in this “clean” background). The  $x$ -intercepts ranged from 0.1 to  $0.4 \mu\text{g m}^{-3}$ , which is consistent with mean  $M_{LAC}$  concentrations at the rural site Aspvreten. Thus,

### Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a large influence of traffic emissions on  $M_{LAC}$  and  $NO_x$  concentrations was observed on weekdays at Hornsgatan and Essingeleden, and at Torkel but restricted to the period 08:00–18:00 LT when the traffic volume is higher at Hornsgatan.

Finally, we illustrate the contribution of the urban sources to the regional background levels and discuss the potential to decrease urban concentrations by reducing local emission of atmospheric pollutants following Ketznel et al. (2004). Figure 8 displays the weekday concentrations at the different locations relative to the concentrations at the urban background site (Torkel, 100%). Rural  $NO_x$  and  $PM_{2.5}$  data correspond to Norr Malma station. For  $M_{LAC}$  and  $NO_x$ , 70–75% of the concentrations were generated in the urban area whereas only 35% of  $PM_{10}$  and 5% of  $PM_{2.5}$  corresponded to local urban sources. Regarding the kerbside stations,  $M_{LAC}$  and  $NO_x$  levels were 310–660% relatively higher than urban background levels whereas  $PM_{10}$  and  $PM_{2.5}$  concentrations were 130–230% higher relative to Torkel. Hence, the potential for reducing pollutants urban concentrations in Stockholm by controlling local emissions is much higher for  $M_{LAC}$  and  $NO_x$  compared to  $PM_{2.5}$  and  $PM_{10}$ .

## 5 Summary and conclusions

Atmospheric LAC concentrations were simultaneously sampled using custom-built PSAPs in spring 2006 in Stockholm at four locations, representing urban, urban background and rural pollution levels. These custom-built photometers were found to be a reliable alternative for the commercial Aethalometer, as indicated by the high linear correlation ( $R > 0.85$ ) observed when comparing hourly time series.

As expected, higher  $M_{LAC}$  concentrations were registered at the urban sites than at the background locations. A large hour-to-hour and day-to-day variability of  $M_{LAC}$  concentrations was observed at the kerbside locations and the variability was clearly negligible at the rural site, but for an intrusion of a polluted air mass transporting agricultural wildfire combustion products emitted in Russia and the Baltic region. Concentrations of LAC between urban sites were poorly correlated even for daily averages ( $R < 0.70$ ),

### Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



combined with highly heterogeneously distributed concentrations ( $COD > 0.30$ ) even at short spatial scales of the order of few kilometers.

Other commonly monitored atmospheric pollutants (namely  $PM_{10}$ ,  $PM_{2.5}$ , and  $NO_x$ ) were analyzed to help characterize LAC emission sources. In this study, two sources of LAC were identified in the Stockholm area: vehicle exhaust emissions and long-range transport of combustion-derived particles. This can be explained by inspection of five main indicators: (1) The high correlation ( $R > 0.75$ ) between hourly  $M_{LAC}$  and  $NO_x$  simultaneous measurements at the urban sites, when no LRT episodes were observed.  $NO_x$  is a tracer for vehicle emissions and a high correlation between  $M_{LAC}$  and  $NO_x$  suggested a common origin; (2) The statistically significantly higher  $M_{LAC}$  and  $NO_x$  concentrations at weekdays than on weekends in the early morning at Hornsgatan matching with higher traffic density of diesel vehicles; (3) The even higher correlation ( $R > 0.95$ ) between weekday average diurnal concentrations of  $M_{LAC}$  and  $NO_x$  at the kerbside stations consistent with traffic density daily cycles at both locations; (4) The same pattern of  $M_{LAC}$  and  $NO_x$  concentration roses at the kerbside stations even when the LRT episode was considered in the data analysis. When the LRT event was included in the background station calculations,  $M_{LAC}$  concentrations are highest for E and SE sectors, matching the predominant wind direction during the agricultural wild-fires episode. This suggests that local emission sources (i.e. traffic exhaust) were the main contributors to  $M_{LAC}$  and  $NO_x$  measurements at kerbside locations and the LRT contribution to the LAC fraction was masked at these sites; and (5) The average  $NO_x/M_{LAC}$  ratios in the range 20–23, characteristic of areas impacted by high traffic exhaust emissions.

This study highlights the benefit of highly resolved spatiotemporal measurements to accurately characterize the complexity of the particulate field across an urban area, and especially the carbon content fraction, to improve the intraurban assessment of human exposure to air pollution.

The potential for decreasing pollutants urban concentrations in Stockholm by reducing local emissions was analyzed by comparing the urban and urban background

## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

contributions to the air pollution levels. Abatement strategies should target reductions of vehicle exhaust emissions, after the much higher potential for  $M_{LAC}$  and  $NO_x$  to reduce pollution levels in urban areas (kerbside and urban background locations) compared to the potential of  $PM_{2.5}$  and  $PM_{10}$ .

To understand urban aerosol dynamics and long-term effects of aerosol particles on health, long continuous data sets are required. Thus, a longer field campaign should be conducted in a similar fashion in the Stockholm region. The new experiment might also allow studying the seasonal variation of the  $M_{LAC}$  spatiotemporal distribution and how the contribution of the emission sources to the LAC content varies over the year. The possible contribution of residential wood combustion to the LAC fraction should be investigated in Stockholm in the wintertime.

*Acknowledgement.* This work was supported by the Swedish Environmental Protection Agency. The authors thank Hans Karlsson, Hans Areskoug, Leif Bäcklin at Stockholm University, and Billy Sjövall at the Environment and Health Protection Administration of Stockholm for their skilled assistance during the field campaign.

## References

- Ackerman, A. S., Toon, O. B., Stevens, D. E., Heymsfield, A. J., Ramanathan, V., and Welton, E. J.: Reduction of tropical cloudiness by soot, *Science*, 288, 1042–1047, 2000.
- Areskoug, H., Johansson, C., Alesand, T., Hedberg, E., Ekengrena, T., Vesely, V., Wideqvist, U., and Hansson, H.-C.: Concentrations and sources of  $PM_{10}$  and  $PM_{2.5}$  in Sweden, ITM report 110, Stockholm University, 2004.
- Areskoug, H.: Measurements of  $PM_{10}$  - A comparison of the most common methods used in Sweden with the European reference method (Bestämning av  $PM_{10}$  – En jämförelse av de vanligaste mätmetoderna använda i Sverige och den europeiska referensmetoden, only in Swedish), ITM report 168, Stockholm University, 2007.
- Birch, M. E. and Cary, R. A.: Elemental carbon-based method for monitoring occupational exposures to particulate diesel exhaust, *Aerosol Sci. Tech.*, 25, 221–241, 1996.
- Bond, T. C. and Bergstrom, R. W.: Light absorption by carbonaceous particles: an investigative review, *Aerosol Sci. Tech.*, 39, 1–41, 2005.

## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Bond, T. C., Streets, D. G., Yarber, K. F., Nelson, S. M., Woo, J.-H., and Klimont, Z.: A technology-based global inventory of black and organic carbon emissions from combustion, *J. Geophys. Res.*, 109, D14203, doi:10.1029/2003JD003697, 2004.

Cozic, J., Verheggen, B., Weingartner, E., Crosier, J., Bower, K. N., Flynn, M., Coe, H., Henning, S., Steinbacher, M., Henne, S., Collaud Coen, M., Petzold, A., and Baltensperger, U.: Chemical composition of free tropospheric aerosol for PM<sub>1</sub> and coarse mode at the high alpine site Jungfraujoch, *Atmos. Chem. Phys.*, 8, 407–423, doi:10.5194/acp-8-407-2008, 2008.

Franco Suglia, S., Gryparis, A., Wright, R. O., Schwartz, J., and Wright, R. J.: Association of black carbon with cognition among children in a prospective birth cohort study, *Am. J. Epidemiol.*, 167, 280–286, 2007.

Gidhagen, L., Johansson, C., Langner, J., and Olivares, G.: Simulation of NO<sub>x</sub> and ultrafine particles in a street canyon in Stockholm, Sweden, *Atmos. Environ.*, 38, 2029–2044, 2004a.

Gidhagen, L., Johansson, C., Omstedt, G., Langner, J., and Olivares, G.: Model simulations of NO<sub>x</sub> and ultrafine particles close to a Swedish highway, *Environ. Sci. Technol.*, 38, 6730–6740, 2004b.

Gidhagen, L., Johansson, C., Langner, J., and Foltescu, V.: Urban scale modeling of particle number concentration in Stockholm, *Atmos. Environ.*, 39, 1711–1725, 2005.

Intergovernmental Panel on Climate Change (IPCC): Climate change 2001: the Scientific Basis, edited by: Houghton, J. T., Ding, Y., Griggs, D. J., Noguer, M., van der Linden, P. J., Dai, X., Maskell, K., and Johnson, C. A., Cambridge University Press, New York, USA, 572 pp., 2001.

Järvi, L., Junninen, H., Karppinen, A., Hillamo, R., Virkkula, A., Mäkelä, T., Pakkanen, T., and Kulmala, M.: Temporal variations in black carbon concentrations with different time scales in Helsinki during 1996–2005, *Atmos. Chem. Phys.*, 8, 1017–1027, doi:10.5194/acp-8-1017-2008, 2008.

Johansson, C., Norman, M., and Gidhagen, L.: Spatial and temporal variations of particle mass (PM<sub>10</sub>) and particle number in urban air – Implications for health impact assessment, *Environ. Monit. Assess.*, 127, 477–487, 2007a.

Johansson, C., Sjövall, B., Ferm, M., Karlsson, H., Vecely, V., Krecl, P., and Ström, J.: Relationship between PM<sub>10</sub>, PM<sub>1</sub>, and soot concentrations in Sweden (Relationerna mellan halterna av PM<sub>10</sub>, PM<sub>1</sub> och sot i Sverige, only in Swedish), ITM report 153, Stockholm University, 2007b.

## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Johansson, C., Burman, L., and Forsberg, B.: The effects of congestions tax on air quality and health, *Atmos. Environ.*, 43, 4843–4854, 2009.
- Ketzel, M., Wåhlin, P., Kristensson, A., Swietlicki, E., Berkowicz, R., Nielsen, O. J., and Palmgren, F.: Particle size distribution and particle mass measurements at urban, near-city and rural level in the Copenhagen area and Southern Sweden, *Atmos. Chem. Phys.*, 4, 281–292, doi:10.5194/acp-4-281-2004, 2004.
- Kobach, A., Johansen, B. V., Schwarze, P. E., and Namork, E.: Analytical electron microscopy of combustion particles: a comparison of vehicle exhaust and residential wood smoke, *Sci. Tot. Environ.*, 346, 231–243, 2005.
- Kondo, Y., Komazaki, Y., Miyazaki, Y., Moteki, N., Takegawa, N., Kodama, D., Deguchi, S., Nogami, M., Fukuda, M., Miyakawa, T., Morino, Y., Koike, M., Sakurai, H., and Ehara, K.: Temporal variations of elemental carbon in Tokyo, *J. Geophys. Res.*, 111, D12205, doi:10.1029/2005JD006257, 2006.
- Krecl, P., Ström, J., and Johansson, C.: Carbon content of atmospheric aerosols in a residential area during the wood combustion season in Sweden, *Atmos. Environ.*, 41, 6974–6985, 2007.
- Krecl, P., Johansson, C., and Ström, J.: Spatiotemporal variability of light-absorbing carbon concentration in a residential area impacted by woodsmoke, *J. Air Waste Manage.*, 60, 356–368, 2010.
- Kupiainen, K. and Klimont, Z.: Primary emissions of fine carbonaceous particles in Europe, *Atmos. Environ.*, 41, 2156–2170, 2007.
- Lighty, J. S., Veranth, J. M., and Sarofim, A. F.: Combustion aerosols: factors governing their size and composition and implications to human health, *J. Air Waste Manage.*, 50, 1565–1618, 2000.
- Lin, C. I., Baker, M. B., and Charlson R. J.: Absorption coefficient of the atmospheric aerosol: a method for measurement, *Appl. Opt.*, 12, 1356–1363, 1973.
- Menon, S., Hansen, J., and Nazarenko, L.: Climate effects of black carbon aerosols in China and India, *Science*, 297, 2250–2253, 2002.
- Naser, T. M., Kanda, I., Ohara, T., Sakamoto, K., Kobayashi, S., Nitta, H., and Nataami, T.: Analysis of traffic-related NO<sub>x</sub> and EC concentrations at various distances from major roads in Japan, *Atmos. Environ.*, 43, 2379–2390, 2009.
- Norman, M. and Johansson, C.: Studies of some measures to reduce road dust emissions from paved roads in Scandinavia, *Atmos. Environ.*, 40, 6154–6164, 2006.

## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Olivares, G., Johansson, C., Ström, J., and Hansson, H.-C.: The role of ambient temperature for particle number concentrations in a street canyon, *Atmos. Environ.*, 41, 2145–2155, 2007.
- Pereira, S., Wagner, F., and Silva, A. M.: Scattering properties and mass concentration of local and long-range transported aerosols over the South Western Iberia Peninsula, *Atmos. Environ.*, 42, 7623–7631, 2008.
- Putaud, J.-P., Raes, F., Van Dingenen, R., Uggemann, E., Facchini, M.-C., Decesari, S., Fuzzi, S., Gehrig, R., Ugliin, C. H., Laj, P., Lorbeer, G., Maenhaut, W., Mihalopoulos, N., Uller, K. M., Querol, X., Rodriguez, S., Schneider, J. U., Spindler, G., ten Brink, H., Tørseth, K., and Wiedensohler, A.: A European aerosol phenomenology – 2: Chemical characteristics of particulate matter at kerbside, urban, rural and background sites in Europe, *Atmos. Environ.*, 38, 2579–2595, 2004.
- Ramanathan, V. R., Crutzen, P. J., Kiehl, J. T., and Rosenfeld, D.: Aerosols, climate and the hydrological cycle, *Science*, 294, 2119–2124, 2001.
- Ramanathan, V. R., Ramana, M. V., Roberts, G., Kim, D., Corrigan, C., Chung, C., and Winker, D.: Warming trends in Asia amplified by brown cloud solar absorption, *Nature*, 448, 575–579, 2007.
- Sharma, S., Brook, J. R., Cachier, H., Chow, J., Gaudenzi, A., and Lu, G.: Light absorption and thermal measurements of black carbon in different regions of Canada, *J. Geophys. Res.*, 107(D24), 4771, doi:10.1029/2002JD002496, 2002.
- SLB: Emissions and concentrations of nitrogen oxides and nitrogen dioxide on Hornsgatan street; Evaluation of traffic measurements during autumn 2009 (Utsläpp och halter av kväveoxider och kvävedioxid på Hornsgatan analys av trafikmätningar under hösten 2009, in Swedish only), SLB report 7, 2010.
- Streets, D. G., Bond, T. C., Lee, T., and Jang, C.: On the future of carbonaceous aerosol emissions, *J. Geophys. Res.*, 109, D24212, doi:10.1029/2004JD004902, 2004.
- Targino, A. C., Krecl, P., Coraiola, G., Johansson, C.: Air quality across Sweden during transboundary biomass burning pollution episodes in spring 2006, in preparation, 2011.
- Venkatachari, P., Zhou, L., Hopke, P. K., Felton, D., Rattigan, O. V., Schwab, J. J., and Demerjian, K. L.: Spatial and temporal variability of black carbon in New York City, *J. Geophys. Res.*, 111, D10S05, doi:10.1029/2005JD006314, 2006.
- Vogt, M., Nilsson, E. D., Ahlm, L., Mårtensson, E. M., and Johansson, C.: The relationship between 0.25–2.5  $\mu\text{m}$  aerosol and  $\text{CO}_2$  emissions over a city, *Atmos. Chem. Phys. Discuss.*, 10, 21521–21545, doi:10.5194/acpd-10-21521-2010, 2010.

Wilson, J. G., Kingham, S., Pearce, J., and Sturman, A. P.: A review of intraurban variations in particulate air pollution: Implications for epidemiological research, *Atmos. Environ.*, 39, 6444–6462, 2005.

5 Wongphatarakul, V., Friedlander, S. K., and Pinto, J. P.: A comparative study of PM<sub>2.5</sub> ambient aerosol chemical databases, *Environ. Sci. Technol.*, 32, 3926–3934, 1998.

**Spatiotemporal distribution of light-absorbing carbon**

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 1.** Descriptive statistics of hourly  $M_{\text{LAC}}$  concentrations at the four sites for the period 20 April–5 July 2006.

$M_{\text{LAC}}$ [ $\mu\text{g m}^{-3}$ ]	Hornsgatan	Essingeleden	Torkel	Aspvreten
Arithmetic mean	5.39	3.23	1.13	0.36
Median	4.47	2.63	0.72	0.14
5th percentile	1.44	0.74	0.21	0.01
95th percentile	12.46	1.47	3.99	1.58
Arithmetic SD	3.60	2.27	1.15	0.50
Number of samples	1744	1791	1841	1412
Data coverage [%]	94	97	100	76

## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

**Table 2.** Linear correlation coefficients between  $M_{\text{LAC}}$  time series measured at the four sites for hourly and daily (in square brackets) concentrations in the period 20 April–5 July 2006.  $R > 0.75$  are displayed in bold.

$R$ [-]	Essingeleden	Torkel	Aspvreten
Hornsgatan	0.62 [0.70]	0.48 [0.59]	0.23 [0.40]
Essingeleden		0.50 [0.65]	0.27 [0.41]
Torkel			<b>0.78 [0.88]</b>

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

**Table 3.** COD of  $M_{LAC}$  time series measured at the four sites for hourly and daily (in square brackets) concentrations in the period 20 April–5 July 2006.

COD [–]	Essingeleden	Torkel	Aspvreten
Hornsgatan	0.37 [0.31]	0.69 [0.70]	0.88 [0.90]
Essingeleden		0.56 [0.56]	0.84 [0.86]
Torkel			0.66 [0.67]

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)




[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Table 4.** Arithmetic mean  $\text{NO}_x$ ,  $\text{PM}_{10}$ ,  $\text{PM}_{2.5}$  concentrations at the different sites for the period 20 April–5 July 2006.

Concentration [ $\mu\text{g m}^{-3}$ ]	Hornsgatan	Essingeleden	Torkel	Aspvreten	Norr Malma
$\text{NO}_x$	109.6	74.4	15.9	–	5.3
$\text{PM}_{10}$	46.1	47.4	22.0	15.2	15.0
$\text{PM}_{2.5}$	12.8 <sup>a</sup>	12.2	10.8	8.5 <sup>a</sup>	10.4

<sup>a</sup> Reduced data availability in the selected period: 36% at Hornsgatan, and 10% at Aspvreten.

## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.

**Table 5.** Linear correlation coefficients between hourly concentrations of different species measured at each site in the period 20 April–5 July 2006, but excluding the aerosol LRT event on 24 April–9 May 2006.  $n$  indicates the number of paired data for each combination.  $R > 0.75$  are displayed in bold.

$R$ [–]	$M_{\text{LAC}}^{\text{NO}_x}$	$M_{\text{LAC}}^{\text{PM}_{10}}$	$M_{\text{LAC}}^{\text{PM}_{2.5}}$	$\text{NO}_x^{\text{PM}_{2.5}}$	$\text{NO}_x^{\text{PM}_{10}}$	$\text{PM}_{2.5}^{\text{PM}_{10}}$
Hornsgatan $n$	<b>0.78</b> 1368	0.44 1382	0.64 625	0.50 664	0.51 1438	0.67 673
Essingeleden $n$	<b>0.90</b> 1443	0.45 1412	0.57 1330	0.42 1350	0.48 1433	0.38 1348
Torkel $n$	<b>0.76</b> 1389	0.36 1451	0.55 1305	0.20 1240	0.17 1386	0.52 1306
Aspvreten $n$	– 1037	0.45 1037	0.35 171	–	–	0.73 176

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Spatiotemporal  
distribution of  
light-absorbing  
carbon**

P. Krecl et al.



**Fig. 1.** Location of Stockholm City, Aspvreten and Norr Malma rural background sites.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.




**Fig. 2.** Location of the monitoring stations in Stockholm: Essingeleden (1), Hornsgatan (2), and Torkel (3). Blue indicates water surfaces, green forest/park areas, orange built-up areas, black lines important roads, and grey represent public buildings (e.g., hospitals, museums, schools).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.



**Fig. 3.** Linear correlation between hourly mean PSAP-ITM aerosol absorption coefficients and Aethalometer  $M_{LAC}$  in the period 20 April–5 July 2006. **(a)** Hornsgatan, **(b)** Torkel. The regression equation, correlation coefficient  $R$  and number of samples  $n$  are also displayed.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Spatiotemporal distribution of light-absorbing carbon**

P. Krecl et al.




**Fig. 4.** Time series of hourly mean  $M_{LAC}$  concentrations at the four sites for the period 14–20 May 2006. Date labels are at 00:00 LT. Days of the week are also displayed.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.



**Fig. 5.** Average diurnal variation of  $M_{LAC}$ ,  $PM_{10}$ ,  $PM_{2.5}$ ,  $NO_x$ , and traffic rate on weekdays and weekends for the four stations in the period 20 April–5 July 2006. Aspöreten  $PM_{2.5}$  concentrations were replaced by Norr Malma data.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion


## Spatiotemporal distribution of light-absorbing carbon

P. Krecl et al.



**Fig. 6.** Rose plots of pollutant concentrations ( $M_{LAC}$ ,  $PM_{10}$ , and  $NO_x$ ) based on hourly measurements at the four sites and expressed as  $\mu\text{g m}^{-3}$ . Top panel (a–c): whole period 20 April–7 July 2006. Bottom panel (d–f): measurements in the period 24 April–9 May 2006 were excluded (LRT event). Color code: Hornsgatan (black), Essingeleden (red), Torkel (green), and Aspvreten (blue).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




**Fig. 7.** Scatterplot of  $M_{LAC}$  vs.  $NO_x$  weekday diurnal variation concentrations for three sites (black dot: Hornsgatan, square: Essingeleden, cross: Torkel) in the period 20 April–5 July 2006, excluding the LRT event. The regression equation with the 95% confidence intervals for the slope and y-intercept, and correlation coefficient  $R$  are also displayed.

**Spatiotemporal distribution of light-absorbing carbon**

P. Krecl et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	





**Fig. 8.** Comparison of weekday  $M_{LAC}$ ,  $NO_x$ ,  $PM_{10}$  and  $PM_{2.5}$  concentrations at a rural, urban background, and kerbside stations relative to urban background concentrations. Rural  $NO_x$  and  $PM_{2.5}$  data correspond to Norr Malma station.

**Spatiotemporal distribution of light-absorbing carbon**

P. Krecl et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

