

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Fragmentation vs. functionalization: chemical aging and organic aerosol formation

H. J. Chacon-Madrid and N. M. Donahue

Center for Atmospheric Particle Studies, Carnegie Mellon Univ., Pittsburgh, PA 15213, USA

Received: 19 April 2011 – Accepted: 20 April 2011 – Published: 5 May 2011

Correspondence to: N. M. Donahue (nmd@andrew.cmu.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Fragmentation vs. functionalization

H. J. Chacon-Madrid and
N. M. Donahue

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The transformation process that a carbon backbone undergoes in the atmosphere is complex and dynamic. Understanding all these changes for all the species in detail is impossible; however, choosing different molecules that resemble progressively higher stages of oxidation or aging and studying them can give us an insight into general characteristics and mechanisms. Here we determine secondary organic aerosol (SOA) mass yields of two sequences of molecules reacting with the OH radical at high NO_x. Each sequence consists of species with similar vapor pressures but a succession of oxidation states. The first sequence consists of *n*-pentadecane, *n*-tridecanal, 2-, 7-tridecanone, and pinonaldehyde. The second sequence consists of *n*-nonadecane, *n*-heptadecanal and *cis*-pinonic acid. Oxidized molecules tend to have lower relative SOA mass yields; however, oxidation state alone was not enough to predict how efficiently a molecule forms SOA. Certain functionalities are able to fragment more easily than others, and even the position of these functionalities on a molecule can have an effect. *n*-Alkanes tend to have the highest yields, and *n*-aldehydes the lowest. *n*-Ketones have slightly higher yields when the ketone moiety is located on the side of the molecule and not in the center. In general, oxidation products remain efficient SOA sources, though fragmentation makes them less effective than comparable alkanes.

1 Introduction

Organic oxidation mechanisms in the atmosphere are very complex and dynamic. After a molecule is emitted, its transformation process (e.g. reaction with the OH radical, ozone, NO₃ radical, and photolysis, etc.) becomes progressively more complex. The difficulty of the oxidation mechanisms increases as the number of carbons of the precursor becomes larger as well as the extent of reaction (Goldstein and Galbally, 2007; Kroll and Seinfeld, 2008). Consequently, mechanisms associated with organic aerosol, which generally involve higher than average carbon numbers (Hallquist et al., 2009),

Fragmentation vs. functionalization

H. J. Chacon-Madrid and
N. M. Donahue

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



are especially rich. Smog-chamber experiments have constrained parts of some reaction mechanisms; for example the first-generation products of *n*-alkanes with the OH radical have been identified in detail (Atkinson et al., 2008; Lim and Ziemann, 2005; Atkinson and Arey, 2003), but this addresses only the first hours of reaction. A typical aerosol particle resides in the atmosphere between one to two weeks (Balkanski et al., 1993; Kanakidou et al., 2005) before removal. This time frame is significantly longer than most smog-chamber experiments. There is thus a critical need to understand later-generation chemistry of the oxygenated compounds likely to be involved in the full lifecycle of organics, and especially organic aerosol, in the atmosphere.

Studying the transformation of chemical species for weeks in smog-chamber experiments under ambient conditions is not possible; hence some studies have increased the oxidant concentration (e.g. OH radical) to simulate multiple days of oxidation (Kroll et al., 2009; Smith et al., 2009). These studies, done in flow tubes, have started to yield an insight into chemical aging. However, there is very limited research on this topic. Understanding the details of how molecules age in the atmosphere is crucial for models that predict secondary organic aerosol (SOA) formation (Kanakidou et al., 2005) as well as subsequent transformation (Rudich et al., 2007). One way to elucidate this multiple-generation chemistry is to follow it one generation at a time. The idea is to carefully select compounds that are either themselves important intermediates or that represent important classes of compounds with similar properties. Here we take this approach.

Efforts are underway to try to classify atmospherically relevant organic material in terms of properties important to the condensed phase. The most important property is volatility – it determines whether a compound is in the condensed phase at all (Donahue et al., 2006). Oxygenation has been proposed as a second important property because it progressively increases during oxidation chemistry and also appears to correlate well with water solubility, another key property (Jimenez et al., 2009). We can describe volatility in terms of the saturation concentration, C° (Donahue et al., 2006, 2011), which is operationally defined as the organic aerosol concentration (C_{OA}) at

**Fragmentation
vs. functionalization**H. J. Chacon-Madrid and
N. M. Donahue

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



which half of a given compound will be found in the condensed phase (Donahue et al., 2006). For oxygenation we employ the oxygen-to-carbon ratio O : C (Jimenez et al., 2009; Donahue et al., 2011).

It is useful to break reaction mechanisms into a succession of steps connecting one stable molecule to another. We can classify these steps as functionalization, fragmentation and oligomerization (Kroll and Seinfeld, 2008; Rudich et al., 2007) according to the effect on the product carbon number. These three processes are illustrated in Fig. 1. Functionalization usually involves the addition of oxygenated functional groups (thus an increase in O : C) to a molecule with no change in carbon number, decreasing its vapor pressure in most cases and making it more likely to partition into the aerosol phase. Fragmentation refers to carbon-carbon bond cleavage. In some cases, this path creates higher vapor pressure species, and in others, the vapor pressure decreases because the products are typically also functionalized. A good example is the ozonolysis reaction of monoterpenes (Calogirou et al., 1999). While the double bond is broken, at least two oxygens are added, often significantly decreasing the vapor pressure of the products compared to the parent alkene. Finally, oligomerization, also named accretion, refers to association reactions (typically in the condensed phase) resulting in an increase in carbon number (with no change in O : C) and a dramatic reduction in vapor pressure (Kalberer et al., 2004, 2006).

The relative effects of these three pathways on organic-aerosol levels and properties remain uncertain. Ambient organic aerosol is highly oxidized (Ng et al., 2010) (with average oxygen to carbon ratios for aerosols varying from 0.5 to 0.9, Ng et al., 2010; Aiken et al., 2008), with significant carboxylic acid functionality (Rudich et al., 2007), but this does not directly indicate how much of a role fragmentation or accretion plays.

While organic aerosol ages in the atmosphere, its total oxygen to carbon ratio (O : C) tends to increase, changing its properties with time (Ng et al., 2010). We would therefore like to examine the fragmentation susceptibility and SOA formation of individual molecules vs. their respective O : C. In order to do this, we selected two different sequences of molecules, each sequence consisting of atmospherically relevant vapor

Fragmentation vs. functionalization

H. J. Chacon-Madrid and
N. M. Donahue

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Fragmentation
vs. functionalization**H. J. Chacon-Madrid and
N. M. Donahue

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



pressures (Fig. 2). Specifically, we work with molecules that have saturation concentrations (vapor pressures in mass units) of $\sim 10^5$ and $\sim 10^3 \mu\text{g m}^{-3}$. For each sequence, we concentrate on molecules that have a different O : C, and that are atmospherically relevant. The highest O : C we have worked with so far is ~ 0.3 : atmospherically relevant molecules with higher O : C are more difficult to synthesize, separate or obtain commercially. In this manner we are systematically constructing a sequence of oxidation steps, using model compounds to represent both locations in the 2-D space as well as successive generations of oxidation. Since vapor pressure data are not available for all of the species we worked with, we used the SIMPOL model from Pankow and Asher (2008) and fits from Donahue et al. (2011) to estimate them. The major characteristic encountered is that a molecule with n carbons has a similar vapor pressure to a molecule with $n - 2$ carbons and one oxygen (with carbonyl functionality). This characteristic is used only as a guide.

Each sequence contains species with similar vapor pressures because comparison of their SOA mass yields reflects the competition between fragmentation, functionalization and possibly accretion if important. If two different chemical species with identical vapor pressures had very similar chemical oxidation paths (e.g. competition of functionalization, fragmentation and accretion) their SOA mass yields are expected to be similar as well. That is because with every generation of products formed, their overall vapor pressures increase or decrease by a similar amount. If the SOA mass yields are very different, then the different classes of reaction pathways (fragmentation, functionalization and accretion) contribute differently to the oxidation mechanism. This approach is similar to that employed by Ziemann and co-workers to explore the effects of branching, unsaturation and cyclization on hydrocarbon SOA formation (Lim and Ziemann, 2009a,b), but our focus is on the effects of oxygenated functionality as a model for later-generation chemistry.

The $10^5 \mu\text{g m}^{-3}$ sequence is *n*-pentadecane, *n*-tridecanal, pinonaldehyde, 2-tridecanone, and 7-tridecanone. *n*-Pentadecane is the “anchor” species in the sequence, defining the nominal SOA mass yields we expect from a $\sim 10^5 \mu\text{g m}^{-3}$ species. Both

n-pentadecane and *n*-tridecanal (Schauer et al., 1999a, b) are good models of important anthropogenic emissions with respect to chemical and structural characteristics (e.g. vapor pressure, oxidation state, etc.). Pinonaldehyde is a major product of the oxidation of α -pinene (Schradera et al., 2005; Glasius et al., 1997); multiple studies indicate that the molar yield formation of pinonaldehyde from both α -pinene ozonolysis and OH radical reaction can approach 50% (Hatakeyama et al., 1989, 1991), making it a good model-molecule of the first-generation products from the oxidation of monoterpenes. Finally, 2-tridecanone and 7-tridecanone allow us to explore the relationship between the position of carbonyl functionality in the molecule and its ability to fragment.

The $10^3 \mu\text{g m}^{-3}$ sequence is *n*-nonadecane, *n*-heptadecanal and *cis*-pinonic acid. Both *n*-nonadecane and *n*-heptadecanal are chosen as homologous to *n*-pentadecane and *n*-tridecanal. *cis*-Pinonic acid is similar to pinonaldehyde and it is also a first-generation product of α -pinene oxidation (Hatakeyama et al., 1991).

Our hypothesis is that there is a relationship between the oxidation state of a molecule and its ability to form SOA. More oxidized molecules can fragment more easily, as shown by Kroll et al. (2009), reducing their ability to form organic aerosol when reacting with the OH radical. Chacon-Madrid et al. (2010) showed that *n*-aldehydes fragment significantly more than *n*-alkanes with similar vapor pressures, thus forming less SOA. When examining the gas-phase chemistry of different volatile organic compounds (VOCs) with the OH radical in the presence of NO_x (Atkinson and Arey, 2003; Atkinson, 2000, 2007), it is clear the alkoxy radical is the leading intermediate that fragments molecules especially when other functionalities are already present (Atkinson, 2007; Kwok et al., 1996).

Our objectives are to a) understand the relationship between the oxidation state of a molecule and its ability to form organic aerosol, b) realize how important fragmentation vs. functionalization paths are while a molecule ages in the atmosphere and c) report SOA mass yields of atmospherically relevant species for which there is limited or no information in the literature. Our model-system does not present highly oxidized

**Fragmentation
vs. functionalization**H. J. Chacon-Madrid and
N. M. Donahue

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



precursors as seen in Fig. 2, limiting our conclusions to precursors similar to those found in the early steps of OH radical oxidation sequences. This limitation has to do with the difficulty of synthesizing, separating or obtaining commercially material that resembles low volatility-oxidized organic aerosol (LV-OOA) (Ng et al., 2010).

2 Experimental

We conducted experiments in the Carnegie Mellon University smog chamber. Many details of our experimental procedures are described elsewhere (Hildebrandt et al., 2009). The FEP Teflon (Welch Fluorocarbon) chamber has a maximum volume capacity of 12 m³. The bag is suspended inside a temperature-controlled room, which was held at 295 K for these experiments. All the experiments were conducted under high-NO_x conditions (maximum of 4 ppbC ppb⁻¹ NO_x). Table A in the Supplement lists the specific concentrations and related ratios for all the experiments. Particle-number size distributions were monitored using a scanning mobility particle sizer (SMPS, TSI 3936, 10–800 nm *D_p*). The concentrations of the different organic species were monitored with a unit mass-resolution proton transfer reaction-mass spectrometer (PTR-MS, Ionicon GmbH). For each experiment, HONO photolysis was used to create OH radicals. Additional nitric oxide (NO) was added to achieve the desired VOC:NO_x ratio. Ammonium sulfate particles (Sigma Aldrich, 99.99%) were used as inert condensation seeds, which we formed from an aqueous solution with a nebulizer (TSI 3075) then dried and neutralized. The organic species used were *n*-tridecanal (Alfa Aesar, 94%), pinonaldehyde (synthesized in Carnegie Mellon Laboratory, ~ 85%), 2-tridecanone (MP Biomedicals), 7-tridecanone (MP Biomedicals), *n*-nonadecane (Ultra, 99%), *n*-heptadecanal (synthesized in Carnegie Mellon Laboratory, ~ 85%) and *cis*-pinonic acid (98%, Aldrich). These molecules were all used without further purification. UV lights (General Electric 10526 black lights) with a $J_{\text{NO}_2} = 0.06 \text{ min}^{-1}$ were used to initiate photo-oxidation after all the components were mixed in the chamber.

Fragmentation vs. functionalization

H. J. Chacon-Madrid and
N. M. Donahue

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.1 Injection of molecules into the chamber

Molecules in the $10^5 \mu\text{g m}^{-3}$ sequence were injected directly into the chamber via a septum. Molecules in the $10^3 \mu\text{g m}^{-3}$ sequence (Fig. 2) were injected into the chamber using a resistively heated flash vaporizer. This flash vaporizer has a resistive graphite tip where the material of interest is placed, which is suspended directly in the chamber. This vaporizer heats up to about 200°C in about two minutes. A steel tube is used to allow a flow of air to dissipate the material from the tip into the chamber. Some graphite is ejected from the tip while heated, but the concentrations are significantly lower than the organic precursors. Based on PTR-MS mass spectra of the injection products, there was no evidence of precursor decomposition during flash vaporization.

2.2 SOA mass yield and wall loss calculations

The SOA mass yield from a reaction is defined as the mass of organic aerosol formed divided by the mass of precursor consumed (Odum et al., 1996).

$$Y = \frac{C_{\text{OA}}}{\Delta C_{\text{precursor}}} \quad (1)$$

Where C_{OA} is the mass of organic aerosol created, and $\Delta C_{\text{precursor}}$ is the mass of the precursor organic species consumed to form the organic aerosol (C_{OA}) and other products. We determined SOA mass yields for the different organic species with the aid of inorganic seeds, which provide surface area onto which the organic vapors can condense and allow an independent measurement of particle wall losses. We injected dried ammonium sulfate particles up to $\sim 10^4$ particles cm^{-3} or ~ 10 to $15 \mu\text{g m}^{-3}$.

Total SOA production (C_{OA}) is determined by using the ratio of suspended organic aerosol ($C_{\text{OA}}^{\text{SUS}}$) to suspended ammonium sulfate ($C_{\text{seed}}^{\text{SUS}}$) and the initial concentration of ammonium sulfate $C_{\text{seed}}^{\text{SUS}}(t = 0)$, as described by Hildebrandt et al. (2009):

Fragmentation vs. functionalization

H. J. Chacon-Madrid and
N. M. Donahue

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

$$C_{OA}(t) = \frac{C_{OA}^{SUS}(t)}{C_{seed}^{SUS}(t)} C_{seed}^{SUS}(t=0) \quad (2)$$

$C_{seed}^{SUS}(t)$ is obtained by fitting an exponential decay to the ammonium sulfate volume concentration (measured with the SMPS and compared to an AMS) over the interval between seed injection (and mixing) in the chamber and the onset of photo-oxidation.

Extrapolation of this signal after the photo-oxidation process starts defines the seed mass concentration as a function of time.

At time 0 (onset of photo-oxidation) and later, the difference between the total aerosol mass concentration in the chamber (measured with an SMPS) and the extrapolated ammonium-sulfate mass concentration is considered the $C_{OA}^{SUS}(t)$. To obtain the total $C_{OA}(t)$ from the suspended mass concentration we correct for wall losses, assuming that organic particles lost to the wall are in equilibrium with the suspended particles and vapor-phase species. This is the upper-limit estimate for SOA production. There is approximately a 10% difference in SOA mass yields for each species studied when assuming particles lost to the walls are in equilibrium with the suspended particles (upper limit) vs. not in equilibrium (lower limit). This percentage has been observed in current and past experiments (Chacon-Madrid et al., 2010).

2.3 Measurement of reactants

The concentrations of reactants were monitored with a unit mass-resolution proton transfer reaction-mass spectrometer (PTR-MS). The fragments used to track concentrations are those of the MW + 1 for *n*-tridecanal, pinonaldehyde, 2- and 7-tridecanone. The PTR-MS sensitivity at these fragment masses was previously calibrated with those species. A key assumption is that interferences from other species were minimal. An exception to this was *n*-tridecanal, explained in Chacon-Madrid et al. (2010), where a C_{n-1} dycarbonyl is formed due to isomerization.

We used *m/z* 43 and 57 to follow the concentrations and respective consumptions of *n*-heptadecanal and *n*-nonadecane. The signal corresponding to their MW + 1 was 13701

Fragmentation vs. functionalization

H. J. Chacon-Madrid and
N. M. Donahue

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



too low, thus a higher-intensity but less selective fragment was used. The specifics of using m/z 43 and 57 with a PTR-MS are described in Jobson et al. (2005) and are also applied in Presto et al. (2010). The problem with using either m/z 43 and 57 as a proxy of reactant concentration is that with time, the signal does not decay as quickly as the precursor is consumed. That is because first- and second-generation products contribute to m/z 43 and 57 as well. In some cases this interference is reduced because the products partition into the aerosol phase and do not produce a PTR-MS signal. We also use methanol as a radical tracer in all experiments. We measured it with a PTR-MS (specifically m/z 33) to determine the concentration of OH radicals during the experiments. OH concentrations and kinetic rate constants allow us to predict concentrations of the different precursors and compare with the decay of m/z 43 and 57.

2.4 Synthesis of organic species

We synthesized *n*-heptadecanal and pinonaldehyde following McMurry et al. (1987), as they are not commercially available. Products are formed via ozonolysis in solution, as shown in Fig. A in the Supplement. 1-Octadecene (Acros Organics, 90%) and α -pinene (Aldrich, 99%) are used as reactants for *n*-heptadecanal and pinonaldehyde, respectively. The synthesis consists of creating the secondary ozonide, which must be kept stable by maintaining a temperature of -78°C with a dry-ice ethanol slurry. After the ozonide is formed, dimethyl sulfide (DMS) is added to reduce the ozonide to the respective carbonyls. After adding DMS, the products are held at room temperature for about three hours, after which they are extracted with water to remove water-soluble contaminants. Finally, a simple distillation procedure vaporizes solvents and other impurities. Nuclear Magnetic Resonance (^1H NMR, HSQC and Tocsy) is used to determine products and yields from the synthesis. The spectra are shown in the Supplement in Figs. B through E.

Fragmentation vs. functionalization

H. J. Chacon-Madrid and
N. M. Donahue

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Results

SOA mass yields for the $10^5 \mu\text{g m}^{-3}$ sequence (*n*-tridecanal, pinonaldehyde, 2- and 7-tridecanone, and *n*-pentadecane) are presented in Figs. 3 and 4. All of these species are exposed to similar OH and NO_x concentrations, and none of the reagents showed significant losses to the walls before the OH-radical source was turned on, indicating that wall losses such as those reported by Matsunaga and Ziemann (2010), were not a problem. *n*-Pentadecane SOA mass yields are shown as a function reproducing data from Presto et al. (2010) and *n*-tridecanal yields are from Chacon-Madrid et al. (2010). We shall use the *n*-pentadecane mass yield curve for reference throughout this discussion. *n*-Pentadecane oxidation produces significantly more SOA than either pinonaldehyde or *n*-tridecanal, but within uncertainty, yields for those two aldehydes are identical, as shown in Fig. 3. The 2- and 7-tridecanone SOA mass yields are presented in Fig. 4. Both *n*-ketones have significantly lower SOA mass yields than the reference *n*-alkane, and the 7-tridecanone yields are identical to *n*-tridecanal yields, within error. The 2-tridecanone mass yields are larger than the *n*-tridecanal yields in the 30 to $50 \mu\text{g m}^{-3}$ range of SOA formed.

SOA mass yields for the $10^3 \mu\text{g m}^{-3}$ sequence (*n*-heptadecanal and *n*-nonadecane) are shown in Fig. 5. Because the vapor pressures of the precursors themselves are quite low, we kept the injections and subsequent SOA formation to a relatively low (and atmospherically relevant) range of $< 10 \mu\text{g m}^{-3}$. The yields for *cis*-pinonic acid are not presented in Fig. 5 because we were not confident we could accurately measure its vapor concentrations with the PTR-MS. Nonetheless, *cis*-pinonic acid oxidation produced a significant amount of SOA, with mass yields larger than *n*-heptadecanal, but lower than *n*-nonadecane.

The precision of the experiments can be seen in Fig. 3. The SOA mass yields of *n*-tridecanal are composed of two different experiments, one in the region of 0 to $\sim 25 \mu\text{g m}^{-3}$, and the other in the region of ~ 25 to $\sim 400 \mu\text{g m}^{-3}$. The two experiments form an overlapping yield curve. Pinonaldehyde has a similar mechanism to

Fragmentation vs. functionalization

H. J. Chacon-Madrid and
N. M. Donahue

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

n-tridecanal when reacting with the OH radical and a similar vapor pressure. As noticed in Fig. 3, yields of pinonaldehyde match those of *n*-tridecanal, again demonstrating the good precision of the measurements. All the experiments performed presented yields above blank experiments, including the aldehydes.

4 Discussion

4.1 *n*-Tridecanal and pinonaldehyde vs. *n*-pentadecane

The aldehydic moiety appears to dominate *n*-tridecanal and pinonaldehyde chemistry as compared with *n*-pentadecane. This seems to be responsible for the differences in SOA mass yields as seen in Fig. 3. Fragmentation overwhelms the first-generation chemical mechanism of pinonaldehyde and *n*-tridecanal as seen by Chacon-Madrid et al. (2010). The fragmentation path is important in the presence of NO_x (Atkinson and Arey, 2003). The chemical mechanism of a generic aldehyde reacting with OH radical in the presence of NO_x is presented in the Supplement, Fig. F.

It is believed that fragmentation in the first generation of aldehyde oxidation only occurs when the OH radical attacks the aldehydic moiety, which tends to be very reactive. For example, according to structure activity relationships (SAR) from Kwok and Atkinson (1995), the aldehydic moiety is attacked ~54 and ~77% of the time (relative to other sites in the molecule) for *n*-tridecanal and pinonaldehyde, respectively, when attacked by OH. It is important to mention that not every attack on the aldehydic moiety ends in fragmentation. However, the molecules that do not fragment via aldehydic attack, form peroxyacyl nitrates (PANs) (Singh et al., 1986) which in this case are too volatile to form SOA.

The fragmentation path is thought to be negligible for *n*-pentadecane, at least in the first oxidation steps (Lim and Ziemann, 2005; Atkinson et al., 2008). As shown in Fig. 3, a higher SOA mass yield is the result of adding functionality to a molecule without having significant fragmentation. Linear *n*-alkanes are especially strong SOA

Fragmentation vs. functionalization

H. J. Chacon-Madrid and
N. M. Donahue

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

precursors because branched alkanes can fragment more easily, as shown by Lim and Ziemann (2009a, b) and others (Aschmann et al., 2001). While the first-generation oxidation products from pentalane include aldehydes, they are larger and less volatile than *n*-tridecanal or pinonaldehyde. Since the first oxidation steps for *n*-pentadecane do not include fragmentation, its oxidized products will have significantly lower vapor pressures, making them more likely to partition into the aerosol phase (Pankow, 1994; Donahue et al., 2006).

4.2 *n*-Tridecanal vs. pinonaldehyde

The two important structural differences between *n*-tridecanal and pinonaldehyde are the four-member-cycle and the ketone-functionality present in pinonaldehyde but not in *n*-tridecanal. Regardless, the aldehydic moiety is the most reactive in both molecules. As mentioned above, according to SAR, the first-generation attacks by OH on the aldehydic hydrogen for *n*-tridecanal and pinonaldehyde are ~ 54 and $\sim 77\%$, respectively. Since the aldehydic moiety leads to a significant amount of fragmentation in the presence of NO_x , it is not a surprise to see low SOA mass yields.

The high fragmentation of *n*-tridecanal products was presented in Chacon-Madrid et al. (2010). In the case of pinonaldehyde, Fig. 6 offers strong evidence of fragmentation. In the figure, $t = 0$ is the start of photo-oxidation, with an initial OH radical concentration of $\sim 10^7$ molecule cm^{-3} , as determined by consumption of methanol; this concentration goes down by an order of magnitude after an hour, however. After about 1.5 h of photo-oxidation, *nor*-pinonaldehyde signal starts declining because the reaction with OH overwhelms its production. These data show that *nor*-pinonaldehyde (with 9 carbons) forms with a $\sim 50\%$ molar yield, this is much larger than the $\sim 4\%$ molar yield of *n*-dodecanal formation from *n*-tridecanal presented in Chacon-Madrid et al. (2010), showing that the C_{n-1} alkoxy radical from pinonaldehyde, with a stiff cyclic backbone, is not capable of isomerizing (Fig. F in the Supplement).

n-Tridecanal and pinonaldehyde SOA mass yields are identical within uncertainty (Fig. 3). Two factors may contribute to this similarity. First the first-generation attacks by

Fragmentation vs. functionalization

H. J. Chacon-Madrid and
N. M. Donahue

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



OH on both *n*-tridecanal and pinonaldehyde are very similar. Both result in species with one fewer carbon as their major products. Second, even though the aldehydic moiety on pinonaldehyde may be slightly more reactive than *n*-tridecanal (giving somewhat more fragmentation), the attack on the rest of the molecule (mainly the four-member-cycle) can be very efficient at adding polarity without causing the molecule to split in two different products. In the case of the linear carbon backbone in *n*-tridecanal, one fragmentation step is enough to split the molecule. Finally, the ketone functionality in pinonaldehyde might play a very minor role in the fragmentation path. The only way for the ketone functionality to play a role in the first-generation fragmentation is if the OH radical attacks its α -hydrogen (refer to Fig. 7 as a reference) on the side of the cycle, leading to an alkoxy radical.

4.3 2- vs. 7-tridecanone SOA mass yields

These two molecules are an ideal model system to explore the importance of the position of the ketone functionality in a molecule and its effect on SOA production. Broadly, if the ketone functionality promotes fragmentation, 7-tridecanone SOA yields could be very small, while 2-tridecanone SOA yields might be intermediate. A hotspot of first-generation reactivity of ketones with the OH radical is the β -hydrogens (Atkinson et al., 1982) (Fig. 7). For 7-tridecanone, the β -hydrogens are thought to account for $\sim 47\%$ of the first-generation reaction with the OH radical (following Kwok and Atkinson, 1995). In the case of 2-tridecanone, attack on β -hydrogens is only $\sim 27\%$ because there is only one β -carbon. A partial mechanism for 7-tridecanone is presented in Fig. 7, showing the attack on the β -hydrogens that can lead to fragmentation. This fragmentation path results in the formation of *n*-hexanal, which we observed by PTR-MS (Fig. G, Supplement). This particular fragmentation path might not be important for 2-tridecanone because of the low reactivity of the terminal CH_3- with the β -alkoxy radical at the isomerization stage. Attack on the α -hydrogens could also lead to fragmentation: this path can lead to an α -alkoxy radical that will preferably fragment (Kroll and Seinfeld, 2008). However, the reactivity of the α -hydrogens by OH radical is thought to be slightly

Fragmentation vs. functionalization

H. J. Chacon-Madrid and
N. M. Donahue

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



reduced because of the presence of the carbonyl (Atkinson et al., 1982), making this a minor pathway.

The consequence of having a ketone functional group in the center vs. the side can be seen in Fig. 4. Fragmenting a molecule in the center divides it into two higher vapor-pressure product molecules, which suppresses SOA formation. It is clear in Fig. 4 that 2-tridecanone is more efficient at forming SOA, as expected. However, it is somewhat surprising that the 2-tridecanone SOA yields are as low as they are. This could be due to photolysis of the oxidation products, given that Presto et al. (2005) has shown that UV lights can have an important suppression effect on SOA formation, but this requires further investigation.

4.4 *n*-Nonadecane, *n*-heptadecanal and *cis*-pinonic acid

The less volatile sequence follows a similar pattern as the more volatile sequence, as shown in Fig. 5. *n*-Nonadecane has significantly higher SOA mass yields than *n*-heptadecanal. The species with aldehydic functionality again present lower SOA mass yields due to higher fragmentation paths. This is not obvious. As an *n*-aldehyde becomes as large as *n*-heptadecane, its reactivity and SOA yields might resemble that of an *n*-alkane of comparable volatility (Chacon-Madrid et al., 2010). However, a molecule as large as *n*-heptadecane might have slightly different chemical mechanisms than those of smaller *n*-aldehydes ($< C_{11}$). For example, alkoxy radicals formed on the carbon backbone of *n*-heptadecanal are more likely to isomerize towards the aldehydic functionality, promoting fragmentation. This path was proposed by Jenkin et al. (2000), where an acyl-oxy radical formed from the ozonolysis of α -pinene attacks an internal aldehydic moiety. It is important to mention that we do not have direct evidence of attack on the aldehydic moiety by internal alkoxy radicals. Another reason for the relatively low SOA yields from *n*-heptadecanal could be higher photolysis rates for the reaction products, compared to *n*-nonadecane.

As mentioned before, we did not present SOA mass yields for *cis*-pinonic acid in Fig. 5 because of difficulty measuring the *cis*-pinonic acid concentration. Nonetheless,

Fragmentation vs. functionalization

H. J. Chacon-Madrid and
N. M. Donahue

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



based on injected quantities, yields appeared to be higher than the similar vapor pressure aldehyde but lower than the respective *n*-alkane. The probable strength of the SOA yields from *cis*-pinonic acid is related to its cyclic structure. The first-generation OH radical attack favors two of the tertiary-carbons and on the secondary-carbon placed in the cycle. Even if an intermediate alkoxy radical formed is able to create a carbon-carbon scission, only one product is formed instead of two of higher-volatility.

5 Environmental significance

Understanding how SOA formation capability evolves through the full oxidation sequence of organics in the atmosphere is critical in order to improve predictions of ambient SOA formation. Most chemical transport models (CTMs) under-predict the concentrations of organic aerosol (Appel et al., 2008; Zhang et al., 2009). This may be due to the lack of multi-generation oxidation mechanisms in CTMs. In this paper, we have systematically explored the oxidation of different molecules that are proxies for first-generation oxidation products. From *n*-pentadecane, *n*-tridecanal, 2-, 7-tridecanone and pinonaldehyde to *n*-nonadecane, *n*-heptadecanal and *cis*-pinonic acid, we noticed a higher susceptibility for more oxidized molecules to fragment more easily than less oxidized species of similar vapor pressures, based on SOA mass yields. However, neither oxidation state nor O : C is enough to predict the ability of a species to produce SOA. All oxidized organics produce less SOA than *n*-alkanes of similar vapor pressures, but there are specific moieties – such as aldehydes under high-NO_x conditions – that can fragment efficiently in the first generation of the OH radical attack. The presence of these moieties, rather than O : C in general, appears to be important to SOA formation.

Our major conclusion is that even though carbon backbones are more susceptible to fragmentation as they age, their contribution to generate more organic aerosol is very significant. For example, pinonaldehyde, a major product of α -pinene, has SOA yields very similar to those of α -pinene itself (Presto and Donahue, 2006). *cis*-Pinonic

Fragmentation vs. functionalization

H. J. Chacon-Madrid and
N. M. Donahue

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



acid also has substantial yields. It is critical to understand the contribution of SOA from aging mechanisms and implement them in CTMs in order to better predict organic aerosol levels and properties. This work represents the beginning of that process – it does not include highly oxidized species and makes no conclusions with regards to the heterogeneous oxidation of such species. We show that while more oxidized molecules are more likely to fragment, they will also contribute significantly to organic aerosol formation.

Supplementary material related to this article is available online at:
[http://www.atmos-chem-phys-discuss.net/11/13693/2011/
acpd-11-13693-2011-supplement.zip](http://www.atmos-chem-phys-discuss.net/11/13693/2011/acpd-11-13693-2011-supplement.zip).

Acknowledgement. This research was supported by the Electric Power Research Institute grant EPP25369C12290 and the EPA STAR program through the National Center for Environmental Research (NCER). This paper has not been subject to EPA's required peer and policy review, and therefore does not necessarily reflect the views of the Agency. No official endorsement should be inferred.

References

- Aiken, A. C., DeCarlo, P. F., Kroll, J. H., Worsnop, D. R., Huffman, J. A., Docherty, K. S., Ulbrich, I. M., Mohr, C., Kimmel, J. R., Sueper, D., Sun, Y., Zhang, Q., Trimborn, A., Northway, M., Ziemann, P. J., Canagaratna, M. R., Onasch, T. B., Alfarra, M. R., Prévôt, A. S. H., Dommen, J., Duplissy, J., Metzger, A., Baltensperger, U., and Jimenez, J. L.: O/C and OM/OC ratios of primary, secondary, and ambient organic aerosols with high-resolution time-of-flight aerosol mass spectrometry, *Environ. Sci. Technol.*, 42, 4478–4485, 2008.
- Appel, K. W., Bhawe, P. V., Gilliland, A. B., Sarwar, G., and Roselle, S. J.: Evaluation of the community multiscale air quality (CMAQ) model version 4.5: Sensitivities impacting model performance; Part II – Particulate matter, *Atmos. Environ.*, 42, 6057–6066, 2008.
- Aschmann, S. M., Arey, J., and Atkinson, R.: Atmospheric chemistry of three C10 alkanes, *J. Phys. Chem.*, 105, 7598–7606, 2001.

Fragmentation vs. functionalization

H. J. Chacon-Madrid and
N. M. Donahue

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fragmentation
vs. functionalization**H. J. Chacon-Madrid and
N. M. Donahue

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Atkinson, R.: Atmospheric chemistry of VOCs and NO_x, *Atmos. Environ.*, 34, 2063–2101, 2000.
- Atkinson, R.: Rate constants for the atmospheric reactions of alkoxy radicals: an updated estimation method, *Atmos. Environ.*, 41, 8468–8485, 2007.
- 5 Atkinson, R. and Arey, J.: Atmospheric degradation of volatile organic compounds, *Chem. Rev.*, 103, 4605–4638, 2003.
- Atkinson, R., Aschmann, S. M., Carter, W. P. L., and Pitts Jr., J. N.: Rate constants for the gas-phase reaction of OH radicals with a series of ketones at 299 ± 2 K, *Int. J. Chem. Kinet.*, 14, 839–847, 1982.
- 10 Atkinson, R., Arey, J., and Aschmann, S. M.: Atmospheric chemistry of alkanes: review and recent developments, *Atmos. Environ.*, 42, 5859–5871, 2008.
- Balkanski, Y. J., Jacob, D. J., Gardner, G. M., Graustein, W. C., and Turekian, K. K.: Transport and residence times of tropospheric aerosols inferred from a global three-dimensional simulation of ²¹⁰Pb, *J. Geophys. Res.*, 98, 20573–20586, 1993.
- 15 Calogirou, A., Larsen, B. R., and Kotzias, D.: Gas-phase terpene oxidation products: a review, *Atmos. Environ.*, 33, 1423–1439, 1999.
- Chacon-Madrid, H. J., Presto, A. A., and Donahue, N. M.: Functionalization vs. fragmentation: *n*-aldehyde oxidation mechanisms and secondary organic aerosol formation, *Phys. Chem. Chem. Phys.*, 12, 13975–13982, 2010.
- 20 Donahue, N. M., Robinson, A. L., Stanier, C. O., and Pandis, S. N.: Coupled partitioning, dilution, and chemical aging of semivolatile organics, *Environ. Sci. Technol.*, 40, 2635–2643, 2006.
- Donahue, N. M., Epstein, S. A., Pandis, S. N., and Robinson, A. L.: A two-dimensional volatility basis set: 1. organic-aerosol mixing thermodynamics, *Atmos. Chem. Phys.*, 11, 3303–3318, doi:10.5194/acp-11-3303-2011, 2011.
- 25 Glasius, M., Calogirou, A., Jensen, N. R., Hjorth, J., and Nielsen, C. J.: Kinetic study of gas-phase reactions of pinonaldehyde and structurally related compounds, *Int. J. Chem. Kinet.*, 29, 527–533, 1997.
- Goldstein, A. H. and Galbally, I. E.: Known and unexplored organic constituents in the Earth's atmosphere, *Environ. Sci. Technol.*, 41, 1514–1521, 2007.
- 30 Hallquist, M., Wenger, J. C., Baltensperger, U., Rudich, Y., Simpson, D., Claeys, M., Dommen, J., Donahue, N. M., George, C., Goldstein, A. H., Hamilton, J. F., Herrmann, H., Hoffmann, T., Iinuma, Y., Jang, M., Jenkin, M. E., Jimenez, J. L., Kiendler-Scharr, A., Maen-

**Fragmentation
vs. functionalization**H. J. Chacon-Madrid and
N. M. Donahue

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



haut, W., McFiggans, G., Mentel, Th. F., Monod, A., Prévôt, A. S. H., Seinfeld, J. H., Surratt, J. D., Szmigielski, R., and Wildt, J.: The formation, properties and impact of secondary organic aerosol: current and emerging issues, *Atmos. Chem. Phys.*, 9, 5155–5236, doi:10.5194/acp-9-5155-2009, 2009.

5 Hatakeyama, S., Izumi, K., Fukuyama, T., and Akimoto, H. J.: Reactions of ozone with α -pinene and β -pinene in air: yields of gaseous and particulate products, *J. Geophys. Res.*, 94, 13013–13024, 1989.

Hatakeyama, S., Izumi, K., Fukuyama, T., Akimoto, H., and Washida, N.: Reactions of OH with α -pinene and β -pinene in air: estimate of global CO production from the atmospheric oxidation of terpenes, *J. Geophys. Res.*, 96, 947–958, 1991.

10 Hildebrandt, L., Donahue, N. M., and Pandis, S. N.: High formation of secondary organic aerosol from the photo-oxidation of toluene, *Atmos. Chem. Phys.*, 9, 2973–2986, doi:10.5194/acp-9-2973-2009, 2009.

Jenkin, M. E., Shallcross, D. E., and Harvey, J. N.: Development and application of a possible mechanism for the generation of cis-pinic acid from the ozonolysis of α - and β -pinene, *Atmos. Environ.*, 34, 2837–2850, 2000.

Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prévôt, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., E, Dunlea, J., Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of organic aerosols in the atmosphere, *Science*, 326, 1525–1529, 2009.

Jobson, B. T., Alexander, M. L., Maupin, G. D., and Muntean, G. G.: On-line analysis of organic compounds in diesel exhaust using a proton transfer reaction mass spectrometer (PTR-MS), *Int. J. Mass Spectrom.*, 245, 78–89, 2005.

15 Kalberer, M., Paulsen, D., Sax, M., Steinbacher, M., Dommen, J., Prévôt, A. S. H., Fisseha, R., Weingartner, E., Frankevich, V., Zenobi, R., and Baltensperger, U.: Identification of polymers

**Fragmentation
vs. functionalization**H. J. Chacon-Madrid and
N. M. Donahue

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- as major components of atmospheric organic aerosols, *Science*, 303, 1659–1166, 2004.
- Kalberer, M., Sax, M., and Samburova, V.: Molecular size evolution of oligomers in organic aerosols collected in urban atmospheres and generated in a smog chamber, *Environ. Sci. Technol.*, 40, 5917–5922, 2006.
- 5 Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G., and Wilson, J.: Organic aerosol and global climate modelling: a review, *Atmos. Chem. Phys.*, 5, 1053–1123, doi:10.5194/acp-5-1053-2005, 2005.
- 10 Kroll, J. H. and Seinfeld, J. H.: Chemistry of secondary organic aerosol: Formation and evolution of low-volatility organics in the atmosphere, *Atmos. Environ.*, 42, 3593–3624, 2008.
- Kroll, J. H., Smith, J. D., Che, D. L., Kessler, S. H., Worsnop, D. R., and Wilson, K. R.: Measurement of fragmentation and functionalization pathways in the heterogeneous oxidation of oxidized organic aerosol, *Phys. Chem. Chem. Phys.*, 11, 8005–8014, 2009.
- 15 Kwok, E. S. C. and Atkinson, R.: Estimation of hydroxyl radical reaction rate constants for gas-phase organic compounds using a structure-reactivity relationship: an update, *Atmos. Environ.*, 29, 1685–1695, 1995.
- Kwok, E. S. C., Arey, J., and Atkinson, R.: Alkoxy radical isomerization in the OH radical-initiated reactions of C₄–C₈ *n*-alkanes, *J. Phys. Chem.*, 100, 214–219, 1996.
- 20 Lim, Y. B. and Ziemann, P. J.: Products and mechanism of secondary organic aerosol formation from reactions of *n*-alkanes with OH radicals in the presence of NO_x, *Environ. Sci. Technol.*, 39, 9229–9236, 2005.
- Lim, Y. B. and Ziemann, P. J.: Chemistry of secondary organic aerosol formation from OH radical-initiated reactions of linear, branched, and cyclic alkanes in the presence of NO_x, *Aerosol Sci. Tech.*, 43, 604–619, 2009a.
- 25 Lim, Y. B. and Ziemann, P. J.: Effects of molecular structure on aerosol yields from OH radical-initiated reactions of linear, branched, and cyclic alkanes in the presence of NO_x, *Environ. Sci. Technol.*, 43, 2328–2334, 2009b.
- Matsunaga, A. and Ziemann, P. J.: Gas-wall partitioning of organic compounds in a teflon film chamber and potential effects on reaction product and aerosol yield measurements, *Aerosol Sci. Tech.*, 44, 881–892, 2010.
- 30 McMurry, J. E. and Bosch, G. K.: Synthesis of macrocyclic terpenoid hydrocarbons by intramolecular carbonyl coupling: bicyclogermacrene, lepidozene, and casbene, *J. Org.*

**Fragmentation
vs. functionalization**H. J. Chacon-Madrid and
N. M. Donahue

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Chem., 52, 4885–4893, 1987.

Ng, N. L., Canagaratna, M. R., Zhang, Q., Jimenez, J. L., Tian, J., Ulbrich, I. M., Kroll, J. H., Docherty, K. S., Chhabra, P. S., Bahreini, R., Murphy, S. M., Seinfeld, J. H., Hildebrandt, L., Donahue, N. M., DeCarlo, P. F., Lanz, V. A., Prévôt, A. S. H., Dinar, E., Rudich, Y., and Worsnop, D. R.: Organic aerosol components observed in Northern Hemispheric datasets from Aerosol Mass Spectrometry, *Atmos. Chem. Phys.*, 10, 4625–4641, doi:10.5194/acp-10-4625-2010, 2010.

Odum, J. R., Hoffmann, T., Bowman, F., Collins, D., Flagan, R. C., and Seinfeld, J. H.: Gas/particle partitioning and secondary organic aerosol yields, *Environ. Sci. Technol.*, 30, 2580–2585, 1996.

Pankow, J. F.: An absorption model of gas/particle partitioning of organic compounds in the atmosphere, *Atmos. Environ.*, 28, 185–188, 1994.

Pankow, J. F. and Asher, W. E.: SIMPOL.1: a simple group contribution method for predicting vapor pressures and enthalpies of vaporization of multifunctional organic compounds, *Atmos. Chem. Phys.*, 8, 2773–2796, doi:10.5194/acp-8-2773-2008, 2008.

Presto, A. A. and Donahue, N. M.: Investigation of alpha-pinene + ozone secondary organic aerosol formation at low total aerosol mass., *Environ. Sci. Technol.*, 40, 3546–3543, 2006.

Presto, A. A., Huff Hartz, K. E., and Donahue, N. M.: Secondary organic aerosol production from terpene ozonolysis. 1. Effect of UV radiation, *Environ. Sci. Technol.*, 39, 7036–7045, 2005.

Presto, A. A., Miracolo, M. A., Donahue, N. M., and Robinson, A. L.: Secondary organic aerosol formation from high-NO_x photo-oxidation of low volatility precursors: *n*-alkanes, *Environ. Sci. Technol.*, 44, 2029–2034, 2010.

Rudich, Y., Donahue, N. M., and Mentel, T. F.: Aging of organic aerosol: bridging the gap between laboratory and field studies, *Annu. Rev. Phys. Chem.*, 58, 321–352, 2007.

Schauer, J. J., Kleeman, M. J., Cass, G. R., and Simoneit, B. R. T.: Measurement of emissions from air pollution sources. 1. C₁ through C₂₉ organic compounds from meat charbroiling, *Environ. Sci. Technol.*, 33, 1566–1577, 1999a.

Schauer, J. J., Kleeman, M. J., Cass, G. R., and Simoneit, B. R. T.: Measurement of emissions from air pollution sources. 2. C₁ through C₃₀ organic compounds from medium duty diesel trucks, *Environ. Sci. Technol.*, 33, 1578–1587, 1999b.

Schrader, W., Geiger, J., and Godejohann, M.: Studies of complex reactions using modern hyphenated methods: α -pinene ozonolysis as a model reaction, *J. Chromatogr. A*, 1075,

185–196, 2005.

Singh, H. B., Salas, L. J., and Viezee, W.: Global distribution of peroxyacetyl nitrate, *Nature*, 321, 588–591, 1986.

Smith, J. D., Kroll, J. H., Cappa, C. D., Che, D. L., Liu, C. L., Ahmed, M., Leone, S. R.,
5 Worsnop, D. R., and Wilson, K. R.: The heterogeneous reaction of hydroxyl radicals with
sub-micron squalane particles: a model system for understanding the oxidative aging of am-
bient aerosols, *Atmos. Chem. Phys.*, 9, 3209–3222, doi:10.5194/acp-9-3209-2009, 2009.

Zhang, Y., Vijayaraghavan, K., Wen, X.-Y., Snell, H. E., and Jacobson, M. Z.: Probing into
10 regional ozone and particulate matter pollution in the United States: 1. A 1 year CMAQ
simulation and evaluation using surface and satellite data, *J. Geophys. Res.*, 114, 1–31,
2009.

ACPD

11, 13693–13721, 2011

Fragmentation vs. functionalization

H. J. Chacon-Madrid and
N. M. Donahue

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Fragmentation vs. functionalization

H. J. Chacon-Madrid and
N. M. Donahue




Fig. 1. Major mechanisms in gas and aerosol-phase oxidation. The three major chemical mechanisms that a carbon backbone undergoes in the atmosphere are represented along with their interaction with aerosols. They are described and revised in Kroll and Seinfeld (2008) and Rudich et al. (2007). Both, functionalization and accretion tend to decrease the vapor pressure of species by either adding polarity to the molecule or by polymerizing with other molecules. Fragmentation cleaves carbon-carbon bonds of molecules, generally increasing the vapor pressure of products; however, a good exception is the ozonolysis of monoterpenes, where both fragmentation and functionalization happen simultaneously, adding polarity and decreasing the overall vapor pressure of the products. The competition of these three mechanisms define the fate of molecules in the atmosphere, but it is not clear how this competition plays out while molecules and particles age.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Fragmentation vs. functionalization

H. J. Chacon-Madrid and
N. M. Donahue




Fig. 2. Two sequences of molecules with progressively increasing oxidation state but similar vapor pressures ($\sim 10^5$ and $\sim 10^3 \mu\text{g m}^{-3}$), shown in a two-dimensional volatility-oxidation state space. The $10^5 \mu\text{g m}^{-3}$ sequence is: *n*-pentadecane, *n*-tridecanal, pinonaldehyde, and 2- and 7-tridecanone (co-located with *n*-tridecanal). The $10^3 \mu\text{g m}^{-3}$ sequence is: *n*-nonadecane, *n*-heptadecanal and *cis*-pinonic acid. The species lie in the intermediate volatile organic carbon (IVOC) range occupied by many first-generation oxidation products associated with secondary organic aerosol (SOA) formation. Pinonaldehyde and *cis*-pinonic acid are first-generation products of α -pinene oxidation. Ambient oxidized organic aerosol (OOA) lies in the ranges indicated, with SV-OOA being fresher, less oxidized and more volatile and LV-OOA being more aged and less volatile.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



Fragmentation
vs. functionalizationH. J. Chacon-Madrid and
N. M. Donahue

Fig. 3. SOA mass yields of organic species with vapor pressures of $10^5 \mu\text{g m}^{-3}$. *n*-Tridecanal, pinonaldehyde and *n*-pentadecane SOA mass yields after the OH radical reaction at high NO_x are presented here. *n*-Pentadecane yields are a fit that comes from Presto et al. (2010). The suppression of SOA yields for *n*-tridecanal and pinonaldehyde vs. *n*-pentadecane are related to the tendency of the aldehydic moiety to fragment relatively quickly as detailed in Chacon-Madrid et al. (2010). At this point, it is not clear how the ketone and cyclobutane moieties influence the SOA formation from pinonaldehyde, if at all.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Fragmentation vs. functionalization

H. J. Chacon-Madrid and
N. M. Donahue




Fig. 4. SOA mass yields of ketones with vapor pressures of $10^5 \mu\text{g m}^{-3}$. 2- and 7-tridecanone SOA mass yields are slightly different. The lower efficiency of 7-tridecanone to form SOA might be related to higher fragmentation paths when it reacts with the OH radical as compared to 2-tridecanone. Additionally, splitting the molecule in the middle might have bigger consequences on SOA production vs. splitting it on the side (e.g. 2-tridecanone). However, the differences in SOA yields are not as pronounced as the mechanisms would suggest. Photolysis might play a more important role on the oxidation products of 2-tridecanone.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Fragmentation vs. functionalization

H. J. Chacon-Madrid and
N. M. Donahue




Fig. 5. SOA mass yields of organic species with vapor pressure of $\sim 10^3 \mu\text{g m}^{-3}$. *n*-Pentadecane (from the $10^5 \mu\text{g m}^{-3}$ sequence) is shown as a reference. The differences in SOA mass yields between *n*-nonadecane and *n*-heptadecanal are not surprising. The aldehydic moiety is very reactive and causes fragmentation at high NO_x , while the *n*-nonadecane does not have relevant first-generation fragmentation paths.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




**Fragmentation
vs. functionalization**H. J. Chacon-Madrid and
N. M. Donahue

Fig. 6. Pinonaldehyde + OH radical in the presence of NO_x . Consumption of m/z 169 (molecular weight of pinonaldehyde + 1 in a PTR-MS), and the formation of m/z 155 (molecular weight of *nor*-pinonaldehyde + 1). *nor*-Pinonaldehyde molar yield from the OH radical oxidation of pinonaldehyde in the presence of NO_x is about $\sim 50\%$.




Fig. 7. A partial high- NO_x OH-radical oxidation mechanism of 7-tridecanone, emphasizing the OH radical attack of the β -hydrogens. Figure G in the Supplement shows evidence of *n*-hexanal formation from the OH radical oxidation of 7-tridecanone at high NO_x .