

**Transpacific
transport of BaP
emitted from Asia**

Y. Zhang et al.

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Transpacific transport of Benzo[a]pyrene emitted from Asia: importance of warm conveyor belt and interannual variations

Y. Zhang^{1,*}, S. Tao¹, J. Ma², and S. Simonich³

¹Laboratory for Earth Surface Processes, College of Urban and Environmental Sciences, Peking University, Beijing 100871, China

²Air Quality Research Division, Science and Technology Branch, Environment Canada, 4905 Dufferin Street, Toronto, Ontario M3H 5T4, Canada

³Department of Chemistry, Oregon State University, Corvallis, Oregon, Department of Environmental and Molecular Toxicology, Oregon State University, Corvallis, Oregon, USA

*now at: Department of Atmospheric Sciences, University of Washington, Seattle, Washington 98195, USA

Received: 17 February 2011 – Accepted: 21 June 2011 – Published: 4 July 2011

Correspondence to: S. Tao (taos@urban.pku.edu.cn)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Abstract

A global-scale three dimensional atmospheric transport and chemistry model was applied to simulate transpacific transport of Benzo[a]pyrene (BaP) emitted from Asia. The model results were compared with observations at six monitoring sites. The annual mean and seasonal variation of transport patterns and the contributions of different Asian source regions to transpacific transport flux were investigated. The episodic nature of transpacific transport was also systematically explored. Interannual variability of transpacific transport of BaP was also assessed during the period of 1948–2007. Results showed that strong enhancements of modeled BaP concentrations at 3000 m occurred in an area bounded by 70–80° E and 100–120° E. Air containing these elevated BaP concentrations was then delivered eastward by westerly winds. When approaching the West Coast of North America, the descending atmospheric motion carried BaP-laden air into the lower atmosphere. The transpacific transport flux was 1.6 times higher in the winter than in the summer. East Asian emission dominates the transpacific transport flux with a contribution of about 97 %. Near ground concentration of BaP induced by Asian sources in North America varied between 1–20 $\mu\text{g m}^{-3}$. The number of days with transpacific transport flux with a factor of 0.5, 1.0, 1.5, and 2.0 larger than the running mean were 9.4 %, 0.72 %, 0.06 % and 0.01 %, respectively, implying a mild contribution of episodic transport to the long-term mean transport flux. Significant interannual fluctuation of transpacific transport of BaP was found, including a general decreasing trend during 1948–2007, and especially after the 1970s. The transpacific transport was found to be positively correlated with the Southern Oscillation Index.

Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Polycyclic aromatic hydrocarbons (PAHs) have been included in the Convention on Long range Transboundary Air Pollution Protocol on Persistent Organic Pollutants (United Nations Economic Commission, http://www.unece.org/env/lrtap/pops_h1.htm) and linked with human health (Armstrong et al., 2004). PAHs are mainly generated from incomplete combustion of carbonaceous bio- and fossil fuels. Despite the drastic decrease of PAH emissions in developed countries due to PAH emission reduction efforts (Pacyna et al., 2003), PAH deposition on the ice sheet in Greenland showed the global trend of PAH emission was nearly constant from the beginning of the industrial period to the early 1990s (Masclat et al., 1995). One of the reasons is increasing PAH emission due to rapid economic development and population increases in Asia which compensates for the emission reduction in developed countries (Zhang et al., 2008b). According to a global emission inventory of PAHs, the total emission of sixteen PAH compounds listed in the United States Environmental Protection Agency (USEPA) priority control list from Asia was about 290 Ggy^{-1} , accounting for more than half of total global emissions (Zhang et al., 2009). Given considerable long range transport potential of many PAH compounds (Lang et al., 2008; Primbs et al., 2008; Klasmeier et al., 2006), higher emissions of PAHs from Asia might threaten ecological environments and human health on an intercontinental or even global scale.

A number of observations of transpacific transport of PAHs have been documented in the literature. Surface concentration measurements combined with backward trajectories have shown that semi-volatile organic compounds (SVOCs), including PAHs, can be transported across the Pacific Ocean in about a week under certain meteorological conditions (Primbs et al., 2007, 2008; Killin et al., 2004). For example, Kellin et al. (2004) observed significantly elevated concentrations of PAHs and other SVOCs during 15–16 March, 27–28 March and 22–23 April 2002, when backward trajectories passed over Asia, suggesting Asia as the potential source of these compounds.

Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Several modeling studies have also addressed the transpacific transport of particulate and gaseous pollutants derived from incomplete combustion of carbonaceous fuels including CO and Hg (Liu et al., 2003; Liang et al., 2004, 2005; Strode et al., 2008), and SVOCs such as PCBs (Huang et al., 2007). In these studies, the warm conveyor belt (WCB) was identified as an important pathway for transpacific transport. For instance, Liang et al. (2004) investigated transpacific transport events occurring from March 2001 to late May 2002. They found that lifting from the surface to the free troposphere ahead of surface cold fronts in WCBs was the main pathway (~70 %) of the transpacific transport events occurring in this time period; other processes, including deep convection and transport in the atmospheric boundary layer behind cold fronts, accounted for remaining ~30 %. After the lifting of pollutants from the surface to the free troposphere where the wind is stronger and fast transpacific transport is possible, single or multiple succeeding WCBs are also strongly associated with this advection process (Cooper et al., 2004). Additionally, significant seasonal variation of transpacific transport due to the frequency of cold surge events and the East Asian Monsoon system has been reported in the literature (Liu et al., 2003).

Owing to unique physical/chemical properties and emission patterns of PAHs when compared with CO – which was widely adopted as reference tracer in these studies (e.g. Liang et al., 2004; Liu et al., 2003) – the aforementioned pathways may not necessarily be applicable to PAHs. However, modeling studies of transpacific transport of PAHs have not been carried out and reported in the literature except for the study by Lang et al. (2008). In that investigation, Lang et al. applied a forward trajectory method to assess the outflow of pyrene emitted from China to different regions from 1996 to 2005, and found a mean transit time of >9 days and an outflow mass of ~0.5 tons for North America. The trajectory approach is subject to substantial uncertainty due to the large uncertainty in extensive trajectory calculations as well as a highly simplified chemistry scheme. Other PAH modeling studies either mainly focus on the regional air quality and fate (e.g. Matthias et al., 2009; Zhang et al., 2009; Sehili and Lammel, 2007) or the importance of the gas/particle partitioning scheme (e.g. Prevedouros et

al., 2008; Lammel et al., 2009).

The objectives of this study were to (1) model the transpacific transport of PAHs using a state-of-the-art model with complete atmospheric chemistry and multi-compartment fate and exchanges, (2) identify background transport patterns and episodic transport events, with a special focus on WCB transport, (3) highlight the contribution of different Asian regions to transpacific transport, (4) quantify the concentration of PAHs in North American induced by Asian sources, and (5) investigate the seasonal cycle and inter-annual variations of transpacific transport of PAHs. Due to its high emission rate and extremely strong carcinogenic effect (Zhang and Tao, 2009; Zhang et al., 2009; Nisbet and Lagoy, 1992), BaP was chosen as the representative compound for PAHs in this study.

2 Methodology

2.1 Model description

A global-scale three dimensional atmospheric transport and chemistry model Can-METOP (Canadian Model for Environmental Transport of Organochlorine Pesticides) (Ma et al., 2003; Zhang et al., 2008a, 2010) was modified and applied in this study. Other modifications to a regional scale version of this model, which was successfully used in modeling the transport of PAHs in China (Zhang et al., 2009, 2011), were also merged into this global model.

Briefly, the atmospheric transport module of this model is based on the atmospheric advection-diffusion equation, which was discretized to the model grids at $1^\circ \times 1^\circ$ latitude/longitude and vertical levels, and then solved numerically. The details of the numerical schemes are described elsewhere (Ma et al., 2003). In order to better handle the transport over high latitudes and especially over the Polar Regions, the horizontal advection was solved by a multi-dimensional flux-form semi-Lagrangian transport scheme (Lin and Rood, 1996). The meteorological data (including wind, pressure,

Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



temperature, precipitation, etc.) were obtained from the National Centers for Environmental Prediction (NCEP) reanalysis (Kalnay et al., 1996), and interpolated at each 30-minute model time step. A non-local and higher order turbulence closure scheme was used in this model to estimate the momentum and heat fluxes and to characterize the sub-grid scale process caused by the coarse meteorological data and surface heterogeneity (Ma et al., 2003). The dry deposition velocity is calculated by a resistance-in-series scheme, and the wet deposition is treated as a linear process which is proportional to the precipitation intensity, following Ma et al. (2003).

Similar to the regional-scale version of CanMETOP for PAHs (Zhang et al., 2009), a level-IV fugacity model is coupled with the atmospheric module. Fugacity is defined as the tendency of a chemical to leave a certain phase compartment (Mackay 2001), and the corresponding concentration can be calculated as the product of the fugacity and the chemical capacity. In this model, the soil/air and ocean/air exchange at the bottom of atmosphere, gas/particulate partition of semi-volatile PAH compounds, degradation by OH radical in the atmosphere, biodegradation and leaching in soils, and degradation and exchange between surface and deep ocean are considered and simulated by a fugacity approach (Zhang et al., 2009, 2011). These processes are briefly described below.

We treat the transport and fate of PAHs in soil largely following Ma et al. (2003) and Harner et al. (2001). The fugacity of PAHs in the fast exchanging layer (f_{s1} , 0.1 cm), buffer layer (f_{s2} , 1 cm), and reservoir soil layers (f_{s3} , 10 cm) are modeled by the following set of equations:

$$\begin{cases} \frac{d(f_{s1}Z_{s1})}{dt} = [f_a(D_q + D_d + D_m) + f_{s2}D_{v1,2} - f_{s1}(D_{v1,2} + D_L + D_R)] / v_{s1} \\ \frac{d(f_{s2}Z_{s2})}{dt} = [f_{s1}(D_L + D_{v1,2}) + f_{s3}D_{v2,3} - f_{s2}(D_{v1,2} + D_{v2,3} + D_L + D_R)] / v_{s2} \\ \frac{d(f_{s3}Z_{s3})}{dt} = [f_{s2}(D_L + D_{v2,3}) - f_{s3}(D_{v2,3} + D_L + D_R)] / v_{s3} \end{cases} \quad (1)$$

where f_a is the fugacity in the atmosphere above, and Z and v are the chemical capacity and volumes of the soil, respectively. The D values describe the rates of transport and transformation processes in the soil, which include vertical diffusion between layers

Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



($D_{v_{n,n+1}}$), biological reduction (D_R), leaching (D_L), dry deposition (D_q), wet deposition (D_d), and rain dissolution (D_m). The details of these D values are given by Harner et al. (2001).

The ocean is considered as a series of independent slabs without lateral advection and diffusion. These slabs have the same horizontal spatial resolution as the atmospheric grid boxes above and are divided into surface (with a depth of 30 m) and deep ocean boxes (depth 4000 m) following Wania (1995). The fugacity of PAHs in the two layers (f_{w1} and f_{w2} for the surface and deep layers, respectively) are predicted by Eq. (2):

$$\begin{cases} \frac{d(f_{w1}Z_{w1})}{dt} = [f_a(D_q + D_d + D_m) + f_{w2}D_{v1,2} - f_{w1}(D_{v1,2} + D_S + D_R)] / v_{w1} \\ \frac{d(f_{w2}Z_{w2})}{dt} = [f_{w1}(D_S + D_{v1,2}) - f_{w2}(D_{v1,2} + D_S + D_R)] / v_{w2} \end{cases} \quad (2)$$

where the D , Z and v are similar to those for soil compartment. Besides, we also consider the settling of PAHs associated with organic matter (D_S), which is calculated by Eq. (3) based on Dachs et al. (2002):

$$D_S = \text{BCF} \times F_{\text{om}} Z_{w1} \quad (3)$$

where BCF and F_{om} are the bioconcentration factor of PAHs between ocean water and phytoplankton and the settling velocity of organic matter in the surface ocean water, respectively.

The exchange fluxes ($F_{\text{soil/ocean-air}}$) of PAHs between the atmosphere and the underneath soil/water are then calculated based on the difference of fugacity across the interface between compartments:

$$F_{\text{soil/ocean-air}} = D_v(f_{\text{soil/ocean}} - f_{\text{air}}) \quad (4)$$

where the f s are the fugacity of PAHs in different compartments, while the transport coefficients (D_v) was described by Ma et al. (2003). The initial values of fugacity in the soil and ocean water are generated by an iteration approach with constant PAH emissions until a steady state is achieved.

Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A certain fraction of the PAHs is assumed to adsorb onto the surface of aerosols in the atmosphere (Pankow et al., 1987), and the fraction of particulate associated (θ) is calculated according to the surface area of aerosol (S_{tot}) and the vapor pressure of PAH compounds (P_s), as shown in Eq. (5):

$$\theta = \frac{cS_{\text{tot}}}{cS_{\text{tot}} + P_s} \quad (5)$$

where c (17.2 pa cm) is an independent constant.

The reaction of atmospheric PAHs with OH radical is considered for gaseous phases, whereas the degradation in particulate phase is neglected in this study. The reaction coefficient (k_{OH}) is temperature (T) dependent and calculated by Eq. (6):

$$k_{\text{OH}} = k_{\text{OH}}^0 \exp(a(1/T_0 - 1/T)) \quad (6)$$

where k_{OH}^0 is the preexponential factor and a is the active energy.

2.2 Emission inventory

Zhang and Tao (2009) have developed a global emission inventory based on a compiled emission factors database and emission activity data collected by the International Energy Association and other related sources. This inventory estimates the total PAH emission for each country in 2004, and also provides data on the socio-economic status of these countries including the total population, the Gross Domestic Product and the per-capita income level. Regression models were also built between the PAH emissions and these parameters. Based on these relationships, the national PAH emissions over Asian countries were interpolated into the model grid, using the population statistics of each grid cell as a weighting factor. The national population data were obtained from the LandScan™ global population dataset with a spatial resolution of 1 km × 1 km which was further re-sampled and merged into a 1° × 1° latitude/longitude grid (Oak Ridge National Laboratory, Geographic Information Science and Technology, the database can be accessed via: <http://www.ornl.gov/sci/landscan/index.shtml>).

Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



For simplicity, the Asian BaP emission west of 60° E was not taken into consideration due to the distance from North America and thus negligible influence on transpacific transport. Zhang et al. (2008) also investigated the interannual variation of PAH emissions from China from 1948–2007. The PAH emissions over other countries, the interannual variation of which are not available, were scaled by the historical total population data obtained from the Statistics Division of the United Nations (<http://unstats.un.org/unsd/demographic/sconcerns/popsizes/default.htm>).

2.3 Reference compound and year

The model was run with 2004 as the reference year and full PAH emissions over the entire Asian continent. The interannual variation of transpacific transport of PAHs was also investigated from 1948–2007 with the population-scaled emissions.

2.4 Comparison with observations

The model results were compared with the observations at six monitoring sites including Changdao, Gosan, Kanazawa, Okinawa, Cheeka Peak Observatory (CPO), and Mount Bachelor Observatory (MBO) (Feng et al., 2007; Lee et al., 2006; Tamamura et al., 2007; Primbs et al., 2007; Killin et al., 2004; Primbs et al., 2008). A map of the location of these sites as well as a comparison of their measured concentrations is shown in Fig. 1a. Changdao is an island near Shandong, China. Figure 1b shows modeled and measured seasonal mean BaP concentrations at the Changdao site during 2003–2004. The model underpredicted the concentrations in winter, partly because of underestimation of the seasonal variation of PAH emissions in this region (Zhang and Tao, 2008). At the Gosan site, modeled concentrations show a good correlation with monitoring data (Pearson correlation coefficient $r=0.67$), but the model tends to underestimate BaP concentrations in most winter cases from 2001–2004. Generally, the predictive power decreases as the distance between observation site and source areas increases, due to the weaker contribution from Asian sources as compared with

Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



local sources that were not represented in the model. Although a good agreement between modeled and measured concentrations was achieved in the western Pacific Ocean (Okinawa in the spring of 2004, $r = 0.37$, Kanazawa in 2003–2004, $r = 0.80$), the model produces larger bias in predicting the concentrations in the western seaboard of North America, as illustrated by Fig. 1f. This figure shows that the model does not entirely capture the observed data before May 2002, when the local sources may dominate. Nevertheless, the model successfully simulated the concentrations after 3 May, 2002 and satisfactorily captured the concentration peak on 12 May 2002 ($r = 0.98$ if only this time period was considered). During this period the Intercontinental Transport and Chemical Transformation (ITCT 2K2) experiment was also conducted, during which the concentrations of different pollutants were measured by an aircraft campaign over the West Coast of the United States (near Monterey, California). Results from this field campaign are discussed elsewhere (Allan et al., 2004; de Gouw et al., 2004; Goldstein et al., 2004; Nowak et al., 2004). At the MBO site, the modeled concentrations were poorly correlated with measured concentrations ($r = 0.09$), likely because of the high altitude (2700 m a.s.l.) of this site. At this site local wind patterns are strongly influenced by the local topography, which cannot be captured by course wind data in a global-scale transport model. Nevertheless, the overall agreement between the model and the observations provides a good insight into the validation of the model when it is used to quantitatively assess the transpacific transport of PAHs emitted from Asia.

3 Results and discussion

3.1 Annual mean transport pattern

Figure 2a presents the annual mean horizontal transport pattern of BaP and wind vectors at the height of 3 km across the Pacific. The vertical profile of the meridional mean concentration of BaP averaged over 25–55° N is also shown in Fig. 2b. Significant enhancements of modeled BaP concentrations occurred at the ascending motion

Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



regions between 70–80° E and 100–120° E (Fig. 2b), corresponding to West Asia and East Asia, respectively. Over East Asia, under the influence of convective injection and mid-latitude cyclones, BaP emitted from Siberia, Mongolia, and North and Northeast China were lofted to the free atmosphere and transported southeast to the western Pacific Ocean. The mechanism of this transport pattern has been extensively explored by several modeling studies (Liang et al., 2004; Lang et al. 2008; Zhang et al., 2011). Over West Asia (70–80° E), the upward transport of BaP is weaker and largely contained under 3 km, corresponding to the low concentration levels in this region, as shown in Fig. 2a. The mean ascending motion over this zone is largely forced by the Tibetan Plateau. The divergence of air parcels delivered by westerly flows produces an upward motion over the upslope of the plateau, whereas on the south of plateau the winter India monsoon and the plateau interact to form a strong descending motion in the winter (Zhang, 1991). Therefore, despite the high emissions of PAHs in South and Southeast Asia, the descending air parcels over this region (Fig. 2b) inhibit the lofting of pollutants from these regions (Wallace and Hobbs, 2005).

The BaP-laden air parcels were transported eastward by the westerly winds after they rose high in the atmosphere. Liang et al. (2004) summarized the main mechanisms of transpacific transport in two categories: advection in the free troposphere by westerly flow and direct boundary layer transport. Although under some episodic conditions transpacific transport occurs predominantly below 3 km (Jaffe et al., 1999), our model results show that the largest transport flux occurs at a height of 3–5 km each year. When approaching the western coast of North America, the descending air parcels associated with dry air streams behind cold fronts deliver BaP to a lower atmosphere.

3.2 Seasonal variation of transpacific transport

Significant seasonal variations of transpacific transport of different pollutants and tracers have been reported in the literature (Liang et al., 2004; Liu et al., 2003). Figure 3 shows the seasonal variation of the transpacific transport of BaP. Modeled mean

Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



column concentrations of BaP at North America induced by Asian source vary between 5.3×10^4 and 3.3×10^5 pg m^{-2} in different seasons. The higher transport flux in winter is partly caused by the higher BaP emission in Asia. With bio- and fossil fuel consumptions as the major sources, the emission of BaP is much higher during the cold season than the warm season in the mid- and high-latitude areas of Asia. The emission flux over the Asian continent varied between ~ 160 and ~ 320 t month^{-1} . Another important factor attributable to the lower transport flux in summer is the stronger OH radical degradation of BaP. The degradation flux was ~ 120 t month^{-1} in July, but only 70 t month^{-1} in January.

Besides the influence of emission and removal processes, significant seasonal displacements and variations of the semi-permanent pressure systems and the westerly across the North Pacific Ocean also contribute to the seasonal variation of transpacific transport (Wallace and Hobbs, 2005). Figure 3b, c present the concentrations and the corresponding wind vectors for January and July 2004, respectively. In winter, the westerly flow and winter monsoon may carry BaP emitted from northern China to the western Pacific where the westerly winds reinforced by counterclockwise air flows on the south of the Aleutian Low further deliver the pollutant eastward. Over the eastern Pacific, the eastward flow reinforced between the Aleutian Low and the Pacific High finally pushes the BaP-laden air parcels to the western seaboard of North America. However, with the onset of the East Asia summer monsoon – which features southeasterly winds extending from the western Pacific to East China (Tian et al., 2009) and the northward displacement of the semi-permanent high pressure system over the North Pacific Ocean – the outflow of BaP from North China to the western Pacific becomes almost unlikely at lower atmospheric levels. Although an eastward returning flow at the top (about 3000 m) of the East Asian summer monsoon system could transport the pollutants emitted from Asia to the western Pacific, the strength and spatial scale of this returning flow is much less than the outflow flux in winter. The northward movement of the Pacific High grows more intense in summer and extends from 15 to 60° N, which blocks the eastward transport of air masses over the North Pacific Ocean

Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

(Wallace and Hobbs, 2005).

3.3 Contribution from different Asian sources

The modeled annual mean BaP concentrations at 1200 m altitude over North America originated from different regions of Asia are shown in Fig. 4(a–f). As shown in Fig. 2, the model has predicted significant outflow flux from the Asian continent, illustrated by a plume crossing over 25° N and 45° N and extending to the western Pacific Ocean. The main contributors to this plume are emissions over East Asia (Fig. 4b). Despite of the high emission density in South Asia (Fig. 4c) and the Indochina Peninsula (Fig. 4e), the contributions from these two sources are much smaller compared with that from East Asia due to the general descending movement of air parcels in the downdraft of the Hadley cell (20–30° N). The dominant easterly winds in the subtropics further constrain the transpacific transport from this part of Asia. On the contrary, although BaP emission from West Asia can be effectively lofted to a higher atmospheric level (Fig. 2b), the contribution of BaP emissions from East Asia is less significant due to a longer travel distance (Fig. 4f). The contribution from North and Southeast Asian BaP emissions is minimal due to considerably lower emissions compared with emissions in East and South Asia (Fig. 4a, d). If the mean concentration of BaP over the western coast of North America (defined as a box between 30–50° N and 120–130° W) was taken into consideration, the East Asian emissions contributed about 97 % to BaP mixing ratios, while other regions contributed at the following percentages: South Asia at 1.3 %, West Asia at 1.1 %, North Asia at 0.68 %, the Indochina Peninsula at 0.33 %, and Southeast Asia at 0.0052 % (Fig. 4). We further divided Asia into grids of 10° × 10° latitude/longitude and estimated the contributions of BaP emissions from these grid boxes to the mean BaP concentration over North America due to transpacific transport. Results are shown in Fig. 4g. It is evident that the emission from East Asia between 20° N and 50° N made the largest contribution to the mean BaP mixing ratio over North America, primarily due to the high emissions over this region and favorable climate conditions for long-range transpacific transport. The contribution from other Asian regions

Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



becomes less important because of their lower BaP emissions and unfavorable climate and meteorological conditions.

3.4 Transport budget and near ground concentration in North America

Figure 5 shows the modeled annual mean BaP concentration at 10 m a.g.l. over North America. The concentration was enhanced by about 4 pg m^{-3} (with a range of $1 \sim 20 \text{ pg m}^{-3}$), indicating a slight enhancement of the background BaP concentration over North America (which is $\sim 100 \text{ pg m}^{-3}$, Naumova et al., 2002) due to Asian emissions. The BaP-laden air masses approaching North America spread over a wide latitudinal band between $25\text{--}70^\circ \text{ N}$. The concentrations over the western coast of North America were the highest due to the subsidence of air parcels in this region (Fig. 2b), which entrain the BaP-laden air originating from Asian sources into the boundary layer. The surface concentration was enhanced by more than 6 pg m^{-3} in the western North America, much higher than the $\sim 2 \text{ pg m}^{-3}$ in Central and Eastern America. The lowest concentration was modeled above seasonal or permanent ice surfaces (e.g. the center of Greenland) where the ground friction is minimal, and the weak eddy diffusion cannot efficiently transport the pollutants to the surface (Seinfeld and Pandis, 2000). In 2004, the total mass of BaP is about 1.4 t within a box bounded by $30\text{--}70^\circ \text{ N}$, $75\text{--}125^\circ \text{ W}$ and $0\text{--}7 \text{ km}$ altitude. Of this BaP, about 6.7 t is degraded by OH radicals and 0.6 t is deposited onto the surface. We also calculated the fraction of atmospheric BaP emitted from North American soils, and found that the reemission from North American soils enhances the background concentration by only $\sim 10^{-4} \text{ pg m}^{-3}$, and therefore is negligible even though soil is the largest reservoir for BaP (Sehili and Lammel, 2007).

3.5 Episodic transport mechanism

Many studies dedicated to investigating the episodic nature of transpacific transport indicate stronger transport of pollutants than background level under certain meteorological conditions (Lee et al., 2006; Tamamura et al., 2007; Primbs et al., 2007,

Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2008; Killin et al., 2004). On 5 May 2002, the Intercontinental Transport and Chemical Transformation (ITCT 2K2) experiment was conducted over the West Coast of North America, and air masses containing pollutants originating from Asia were measured by aircraft over a region bounded by 32–44° N, 122–125° W, and 0–8 km altitude (Allan et al., 2004). Weekly monitoring of PAH concentrations at CPO was also conducted during March 2002–May 2002 (Killin et al., 2004). A transpacific transport event was captured on 12 May 2002 (Fig. 1f). In the present study, CanMETOP also captured this transpacific transport event. As this event is closely associated with the ITCT 2K2 experiment carried out over the region about 1000 km south of CPO, which has been discussed in detail in the literature (Lee et al., 2006; Tamamura et al., 2007; Primbs et al., 2007, 2008; Killin et al, 2004), this event will be discussed as a representative case in this study.

BaP concentrations and wind vectors during this transport event at a height of 1200 m are shown in Fig. 6. In later April, a low pressure system was centered in Northeast China and the associated warm conveyor belt (WCB) helped to elevate air parcels to a relatively higher atmospheric level. This low pressure system moved eastward and headed to 170° E on 3 May and to the International Dateline on 4 May. Correspondingly, a typical comma-shaped distribution of elevated BaP can be seen on Fig. 6b (circled region). This is also a typical cold front type distribution. With further eastward movement, this low pressure system weakened considerably due to (sea) surface cooling, so the surface front associated with this low pressure system was no longer identifiable (Fig. 6c). On the same day, it can be seen that BaP concentrations extended from the central North Pacific (150° W) to the West Coast of Oregon State (near 45° N, 120° W), and some of the air parcels originating from eastern Asia even penetrated inland up to 100° W. This transport event was captured by the aircraft monitoring on 5–6 May on the western coast of North America (near 36° N, 120° W). However, the plume did not extend to the CPO monitoring site because the PAH concentration measured at this site was not enhanced. This phenomenon has been clearly modeled in this study (Fig. 6c).

Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Likewise, another low pressure system moved to the east coast of China, forming a southwesterly flow over the western North Pacific (Fig. 6c), just ahead of the cold front associated with this low pressure system. This southwesterly flow moved to the central North Pacific (about 180° E and 135° W on 7 May), resulting in another cold-front-type distribution of BaP, as shown in Fig. 6d. Because of the northward extension of a North Pacific high pressure system, this southwesterly flow was forced to turn west (Fig. 6e). Responding to the interaction of these changing meteorological conditions, BaP-laden air parcels were pushed eastward to the West Coast of Oregon (Fig. 6e and f), resulting in the enhanced BaP concentrations measured at the CPO site. On 12 May, higher concentrations were observed over the western coasts of Washington State and British Columbia (circled in Fig. 6f), and a significant peak of PAH concentrations was also observed at the CPO site (Fig. 1f).

We also conducted a systematic study to further demonstrate the episodic nature of the transpacific transport. As shown in the aforementioned case, the episodic transpacific transport occurred over a time scale of approximately one week. We calculated the eleven-day running mean of the BaP concentration in North America averaged over a box bounded by 32–44° N, 122–125° W and 0–8 km altitude between 1948 and 2007. The BaP concentration anomaly was calculated by subtracting this running mean from the daily mean concentration to filter out the low frequency variations (a.k.a. High-pass filter, e.g. Liang et al., 2005). Results reveal that the frequencies of daily concentrations at factors of 0.5, 1.0, 1.5, and 2.0 higher than the 11 day running mean concentrations were 9.4 %, 0.72 %, 0.06 % and 0.01 %, respectively. This suggests that although episodic transport can significantly enhance the concentration at a specific site or region for a short time period, its contribution to the long-term mean transpacific transport flux does not appear to be very significant.

3.6 Interannual variation of transport from East Asia

Owing to the significant contribution of trans-Pacific transport of East Asian BaP emissions to North American BaP concentrations (97 %, Fig. 4a), the interannual variability

**Transpacific
transport of BaP
emitted from Asia**

Y. Zhang et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

of such the transport merits further investigation. Figure 7a shows the interannual variability of the mean BaP concentrations averaged over North America (brown line) and corresponding total BaP emissions in East Asia (blue bar). As shown, the average BaP concentration over North America has a strong positive correlation with the East Asian emission flux ($r = 0.98$). The North American mean concentration attributable to transpacific transport has increased rapidly since the 1950s and slightly decreased during the last decade due to the declining trend in East Asian emissions. We further defined a transpacific transport index (TPI), calculated by dividing the modeled mean concentration over North America by the Asian emission flux for 1948–2007. Figure 7b illustrates the annual anomaly of the TPI as departures from its mean averaged over 1948–2007 (blue points). The decadal running mean of the TPI was also calculated and shown in this figure (pink points). Despite of the intense annual fluctuations, a decreasing trend in both annual and average TPI was observed, particularly since the 1970s, likely due to changes in interannual or interdecadal atmospheric circulations related to transpacific transport. Based on extensive studies on the outflow of pollutant tracers from East Asia to the western Pacific Ocean, the East Asian winter monsoon is a key factor influencing the outflow potential and strength (Liu et al., 2003; Zhang et al., 2011). The decreasing trend in TPI is partially caused by the continuous weakening of the East Asian winter monsoon since the late 1980s (Nakamura et al., 2002). Another important reason is the decadal weakening of the Pacific jet, which lengthens the transport time of PAHs over the North Pacific Ocean (Nakamura et al., 2002). Based on a recent review on the effect of global climate change on atmospheric transport potential, the future climate is expected to be more stagnant, with a weaker global atmospheric circulation and a decreasing frequency of mid-latitude cyclones (Jacob and Winner, 2009). Therefore, the decreasing trend in TPI would continue in the future. Although the energy consumption and PAH emissions are projected to increase in the next several decades (Zhang et al., 2009), the decreasing TPI would lower the strength of the transpacific transport of PAHs.

**Transpacific
transport of BaP
emitted from Asia**Y. Zhang et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The Pacific North American index (PNA) and the Southern Oscillation Index (SOI) are indicators for the North-South and East-West atmospheric pressure dipole patterns, respectively, and used to link interannual climate variability with changes in transpacific transport in this study. Lang et al. (2008) found a positive correlation between PNA and transpacific transport for pyrene. Liang et al. (2005) also reported a similar result. In the present study we further examined the relationship between PNA and TPI, but no significant correlation between the two indices was observed (Fig. 7c, Spearman correlation coefficient = 0.12). However, this correlation could be improved (Spearman correlation coefficient = 0.518, $p = 0.009$) if only the recent 14 years (same as the time period considered by Lang et al., 2008) were taken into account. On the other hand, a significant positive correlation exists between TPI and SOI (Fig. 7d, Spearman correlation coefficient = 0.38, $p = 0.004$). This result contradicts our first intuition because negative (positive) phase SOI is associated with El Niño (La Niña), during which zonal circulation and thus transpacific transport is stronger (weaker) (Wang, 2002). As the SOI was also closely associated with the frequency of cold episodes and intense boundary outflow behind fronts (Zhang et al., 1997), this implies that the amount of transpacific transport is largely controlled by the magnitude of the outflow flux in Asia.

4 Conclusions

The present modeling study showed strong enhancements of PAH concentration at 3000 m over the region bounded by 70–80° E and 100–120° E, and these PAH were transported eastward by the westerly winds. When the air parcels approached the western coast of North America, the general descending motion carried the PAH-laden air parcels into the lower atmosphere. The transpacific transport flux is ~1.6 times higher in the winter than in the summer. The strength of East Asian emissions dominates the transpacific transport flux, which contributed 97 % of the modeled average BaP concentrations over North America. The near ground concentration of BaP in

North America induced by Asian sources varied between 1–20 pg m⁻³.

An episodic transport event measured at the CPO site near the West Coast of North America on 12 May 2002 reveals the importance of BaP enhancement due to transpacific atmospheric transport, facilitated by the WCB and low pressure systems. The frequencies of daily BaP fluxes from transpacific transport that were a factor of 0.5, 1.0, 1.5, and 2.0 larger than the 11 day running mean were 9.4 %, 0.72 %, 0.06 % and 0.01 %, respectively, implying a mild contribution of episodic transport to the long-term mean transport flux. Significant interannual fluctuation of transpacific transport of PAHs was found. A general decreasing trend was modeled during 1948–2007, especially after the 1970s. The transpacific transport was found to be positively correlated to the Southern Oscillation Index, likely due to the higher frequency of cold episodes and intense boundary outflow within the positive phase of the SOI.

Acknowledgements. The funding for this study was provided by the National Scientific Foundation of China (40710019001, 40730737), The Ministry of Science and Technology of China (2007CB407301) and the National Institute of Health of the US (grant P42 ES016465). The authors gratefully acknowledge Nicole Wigder of University of Washington for polishing the English of the manuscript.

References

- Allan, J. D., Bower, K. N., Coe, H., Boudries, H., Jayne, J. T., Canagaratna, M. R., Millet, D. B., Goldstein, A. H., Quinn, P. K., Weber, R. J., and Worsnop, D. R.: Submicron aerosol composition at Trinidad Head, California, during ITCT 2K2: Its relationship with gas phase volatile organic carbon and assessment of instrument performance, *J. Geophys. Res.-Atmos.*, 109(D23), D23S24, doi:10.1029/2003jd004208, 2004.
- Armstrong, B., Hutchinson, E., Unwin, J., and Fletcher, T.: Lung cancer risk after exposure to polycyclic aromatic hydrocarbons: A review and meta-analysis, *Environ. Health Persp.*, 112, 970–978, 10.1289/ehp.6895, 2004.
- Cooper, O. R., Forster, C., Parrish, D., Trainer, M., Dunlea, E., Ryerson, T., Hubler, G., Fehsenfeld, F., Nicks, D., Holloway, J., de Gouw, J., Warneke, C., Roberts, J. M., Flocke, F.,

Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and Moody, J.: A case study of transpacific warm conveyor belt transport: Influence of merging airstreams on trace gas import to North America, *J. Geophys. Res.-Atmos.*, 109, doi:10.1029/2003jd003624, 2004.

Dachs, J., Lohmann, R., Ockenden, W. A., Mejanelle, L., Eisenreich, S. J., and Jones, K. C.: Oceanic biogeochemical controls on global dynamics of persistent organic pollutants, *Environ. Sci. Technol.*, 36, 4229–4237, 2002.

de Gouw, J. A., Cooper, O. R., Warneke, C., Hudson, P. K., Fehsenfeld, F. C., Holloway, J. S., Hubler, G., Nicks, D. K., Nowak, J. B., Parrish, D. D., Ryerson, T. B., Atlas, E. L., Donnelly, S. G., Schauffler, S. M., Stroud, V., Johnson, K., Carmichael, G. R., and Streets, D. G.: Chemical composition of air masses transported from Asia to the U. S. West Coast during ITCT 2K2: Fossil fuel combustion versus biomass-burning signatures, *J. Geophys. Res.-Atmos.*, 109(D23), D23S20, doi:10.1029/2003jd004202, 2004.

Feng, J. L., Guo, Z. G., Chan, C. K., and Fang, M.: Properties of organic matter in PM_{2.5} at Changdao Island, China – A rural site in the transport path of the Asian continental outflow, *Atmos. Environ.*, 41, 1924–1935, doi:10.1016/j.atmosenv.2006.10.064, 2007.

Goldstein, A. H., Millet, D. B., McKay, M., Jaegle, L., Horowitz, L., Cooper, O., Hudman, R., Jacob, D. J., Oltmans, S., and Clarke, A.: Impact of Asian emissions on observations at Trinidad Head, California, during ITCT 2K2, *J. Geophys. Res.-Atmos.*, 109(D23), D23S17, doi:10.1029/2003/JD004406, 2004.

Harner, T., Wideman, J. L., Jantunen, L. M. M., Bidleman, T. F., and Parkhurst, W. J.: Residues of organochlorine pesticides in Alabama soils, *Environ. Pollut.*, 106, 323–332, 1999.

Huang, P., Gong, S. L., Zhao, T. L., Neary, L., and Barrie, L. A.: GEM/POPs: a global 3-D dynamic model for semi-volatile persistent organic pollutants - Part 2: Global transports and budgets of PCBs, *Atmos. Chem. Phys.*, 7, 4015–4025, doi:10.5194/acp-7-4015-2007, 2007.

Jaffe, D., Anderson, T., Covert, D., Kotchenruther, R., Trost, B., Danielson, J., Simpson, W., Berntsen, T., Karlsdottir, S., Blake, D., Harris, J., Carmichael, G., and Uno, I.: Transport of Asian Air pollution to north america, *Geophys. Res. Lett.*, 26, 711–714, 1999.

Kahl, J. D. and Samson, P. J.: Uncertainty in trajectory calculation due to low resolution meteorological data, *J. Clim. Appl. Meteorol.*, 25, 1816–1831, 1986.

Kalnay, E., Kanamitsu, M., Kistley, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Leetmaa, A., Reynolds, R., Jenne, R., and Joseph, D.: The NCEP/NCAR reanalysis project, *Bull. Am. Meteorol. Soc.*, 77, 437–471, 1996.

Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Killin, R. K., Simonich, S. L., Jaffe, D. A., DeForest, C. L., and Wilson, G. R.: Transpacific and regional atmospheric transport of anthropogenic semivolatile organic compounds to Cheeka Peak Observatory during the spring of 2002, *J. Geophys. Res.-Atmos.*, 109(D23), D23S15, doi:10.1029/2003jd004386, 2004.

5 Klasmeier, J., Matthies, M., Macleod, M., Fenner, K., Scheringer, M., Stroebe, M., Le Gall, A. C., McKone, T., Van De Meent, D., and Wania, F.: Application of multimedia models for screening assessment of long-range transport potential and overall persistence, *Environ. Sci. Technol.*, 40, 53–60, 2006.

10 Lammel, G., Sehili, A. M., Bond, T. C., Feichter, J., and Grassl, H.: Gas/particle partitioning and global distribution of polycyclic aromatic hydrocarbons – A modeling approach, *Chemosphere*, 76, 98–106, 2009.

Lang, C., Tao, S., Liu, W., Zhang, Y., and Simonich, S.: Atmospheric Transport and Outflow of Polycyclic Aromatic Hydrocarbons from China, *Environ. Sci. Technol.*, 42, 5196–5201, 2008.

15 Lee, J. Y., Kim, Y. P., Kang, C. H., Ghim, Y. S., and Kaneyasu, N.: Temporal trend and long-range transport of particulate polycyclic aromatic hydrocarbons at Gosan in northeast Asia between 2001 and 2004, *J. Geophys. Res.-Atmos.*, 111(11), D11303, doi:10.1029/2005jd006537, 2006.

20 Liang, Q., Jaegle, L., Jaffe, D. A., Weiss-Penzias, P., Heckman, A., and Snow, J. A.: Long-range transport of Asian pollution to the northeast Pacific: Seasonal variations and transport pathways of carbon monoxide, *J. Geophys. Res.-Atmos.*, 109(20), D23s07, doi:10.1029/2003jd004402, 2004.

Liang, Q., Jaegle, L., and Wallace, J. M.: Meteorological indices for Asian outflow and transpacific transport on daily to interannual timescales, *J. Geophys. Res.-Atmos.*, 110(16), D18308, doi:10.1029/2005jd005788, 2005.

25 Lin, S. J. and Rood, R. B.: Multidimensional flux-form semi-Lagrangian transport schemes, *Mon. Weather Rev.*, 124, 2046–2070, 1996.

Liu, H. Y., Jacob, D. J., Bey, I., Yantosca, R. M., Duncan, B. N., and Sachse, G. W.: Transport pathways for Asian pollution outflow over the Pacific: Interannual and seasonal variations, *J. Geophys. Res.-Atmos.*, 108(D20), 8786, doi:10.1029/2002JD003102, 2003.

30 Ma, J., Daggupaty, S., Harner, T., and Li, Y.: Impacts of lindane usage in the Canadian prairies on the Great Lakes ecosystem. 1. Coupled atmospheric transport model and modeled concentrations in air and soil, *Environ. Sci. Technol.*, 37, 3774–3781, 2003.

Mackay, D.: Multimedia environmental models: The fugacity approach, Second Edition, Lewis

Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Publishers, Boca Raton Florida, 2001.

Masclat, P., V., H., Jaffrezo, J. L., and Legrand, M.: Evidence for the presence of polycyclic aromatic hydrocarbons in the polar atmosphere and in the polar ice of Greenland, *Analysis*, 23, 250–252, 1995.

5 Matthias, V., Aulinger, A., and Quante, M.: CMAQ simulations of the benzo(a)pyrene distribution over Europe for 2000 and 2001, *Atmos. Environ.*, 43, 4078–4086, 2009.

Nakamura, H. and Izumi, T.: Interannual and decadal modulations recently observed in the Pacific storm-track activity and East Asian winter monsoon. *J. Climate*, 15, 1855–1874, 2002.

10 Naumova, Y. Y., Eisenreich, S. J., Turpin, B. J., Weisel, C. P., Morandi, M. T., and Colome, S. D.: Polycyclic aromatic hydrocarbons in the indoor and outdoor air of three cities in the U.S., *Environ. Sci. Technol.*, 36, 2552–2559, 2002.

Nisbet, I. C. T. and Lagoy, P. K.: Toxic equivalency factors (tefs) for polycyclic aromatic hydrocarbons (pahs), *Regul. Toxicol. Pharm.*, 16, 290–300, 1992.

15 Nowak, J. B., Parrish, D. D., Neuman, J. A., Holloway, J. S., Cooper, O. R., Ryerson, T. B., Nicks, D. K., Flocke, F., Roberts, J. M., Atlas, E., de Gouw, J. A., Donnelly, S., Dunlea, E., Hubler, G., Huey, L. G., Schauffler, S., Tanner, D. J., Warneke, C., and Fehsenfeld, F. C.: Gas-phase chemical characteristics of Asian emission plumes observed during ITCT 2K2 over the eastern North Pacific Ocean, *J. Geophys. Res.-Atmos.*, 109(D23), D23S19, doi:10.1029/2003jd004488, 2004.

20 Pacyna, J. M., Breivik, K., Műch, J., and Fudala, J.: European atmospheric emissions of selected persistent organic pollutants, 1970–1995, *Atmos. Environ.*, 37, 119–131, 2003.

Pankow, J. F.: Review and comparative analysis of the theories on partitioning between the gas and aerosol particulate phases in the atmosphere, *Atmos. Environ.*, 21, 2275–2283, 1987.

25 Prevedouros, K., Palm-Cousins, A., Gustafsson, O., and Cousins, I. T.: Development of a black carbon-inclusive multi-media model: Application for PAHs in Stockholm, *Chemosphere*, 70, 607–615, 2008.

Primbs, T., Simonich, S., Schmedding, D., Wilson, G., Jaffe, D., Takami, A., Kato, S., Hatakeyama, S., and Kajii, Y.: Atmospheric outflow of anthropogenic semivolatile organic compounds from East Asia in spring 2004, *Environ. Sci. Technol.*, 41, 3551–3558, doi:10.1021/es062256w, 2007.

30 Primbs, T., Wilson, G., Schmedding, D., Higginbotham, C., and Simonich, S. M.: Influence of Asian and Western United States Agricultural Areas and Fires on the Atmospheric Transport of Pesticides in the Western United States, *Environ. Sci. Technol.*, 42, 6519–6525, 2008.

**Transpacific
transport of BaP
emitted from Asia**

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Sehili, A. M. and Lammel, G.: Global fate and distribution of polycyclic aromatic hydrocarbons emitted from Europe and Russia, *Atmos. Environ.*, 41, 8301–8315, doi:10.1016/j.atmosenv.2007.06.050, 2007.

Shrestha, A. B., Wake, C. P., Dibb, J. E., Mayewski, P. A., Whitlow, S. I., Carmichael, G. R., and Ferm, M.: Seasonal variations in aerosol concentrations and compositions in the Nepal Himalaya, *Atmos. Environ.*, 34, 3349–3363, 2000.

Strode, S. A., Jaegle, L., Jaffe, D. A., Swartzendruber, P. C., Selin, N. E., Holmes, C., and Yantosca, R. M.: Trans-Pacific transport of mercury, *J. Geophys. Res.-Atmos.*, 113(12), D15305, doi:10.1029/2007jd009428, 2008.

Tamamura, S., Sato, T., Ota, Y., Wang, X., Tang, N., and Hayakawa, K.: Long-range transport of polycyclic aromatic hydrocarbons (PAHs) from the eastern Asian continent to Kanazawa, Japan with Asian dust, *Atmos. Environ.*, 41, 2580–2593, doi:10.1016/j.atmosenv.2006.11.021, 2007.

Tian, C., Ma, J., Liu, L., Jia, H., Xu, D., and Li, Y.: A modeling assessment of association between East Asian summer monsoon and fate/outflow of a-HCH in Northeast Asia, *Atmos. Environ.* 43, 3891–3901, 2009.

Wallace, M. J. and Hobbs, P. V.: *Atmospheric science: an introductory survey*, Elsevier, 2005.

Wania, F. and Mackay, D.: *A Global Distribution Model for Persistent Organic-Chemicals*, *Sci. Total Environ.*, 161, 211–232, 1995.

Wang, C.: Atmospheric circulation cells associated with the El Niño-southern oscillation, *J. Climate*, 15, 399–419, 2002.

Zellner, R., Seinfeld, J. H., and Pandis, S. N.: *Atmospheric Chemistry and Physics, from Air Pollution to Climate Change*, *J. Atmos. Chem.*, 37, 212–214, doi:10.1023/a:1006483708571, 2000.

Zhang, J. C.: *China Climate*, China Meteorological Press, Beijing, 1991.

Zhang, L., Ma, J., Venkatesh, S., Li, Y. F., and Cheung, P.: Modeling evidence of episodic inter-continental long-range transport of lindane, *Environ. Sci. Technol.*, 42, 8791–8797, 2008a.

Zhang, Y. and Tao, S.: Seasonal variation of polycyclic aromatic hydrocarbons (PAHs) emissions in China, *Environ. Pollut.*, 156, 657–663, 2008.

Zhang, Y. and Tao, S.: Global atmospheric emission inventory of polycyclic aromatic hydrocarbons (PAHs) for 2004, *Atmos. Environ.*, 43, 812–819, 2009.

Zhang, L., Ma, J., Tian, C., Li, Y., and Hung, H.: Atmospheric transport of persistent semi-volatile organic chemicals to the Arctic and cold condensation in the mid-troposphere - Part

**Transpacific
transport of BaP
emitted from Asia**

Y. Zhang et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2: 3-D modeling of episodic atmospheric transport, *Atmos. Chem. Phys.*, 10, 7315–7324, doi:10.5194/acp-10-7315-2010, 2010.

Zhang, Y., Sperber, K. R., and Boyle, J. S.: Climatology and interannual variation of the East Asian winter monsoon: Results from the 1979–95 NCEP/NCAR reanalysis, *Mon. Weather Rev.*, 125, 2605–2619, 1997.

Zhang, Y., Dou, H., Chang, B., Wei, Z., Qiu, W., Liu, S., Liu, W., and Tao, S.: Emission of Polycyclic Aromatic Hydrocarbons from Indoor Straw Burning and Emission Inventory Updating in China, *Annals of the New York Academy of Sciences*, 1140, 218–227, doi:10.1196/annals.1454.006, 2008b.

Zhang, Y., Tao, S., Shen, H., and Ma, J.: Inhalation exposure to ambient polycyclic aromatic hydrocarbons and lung cancer risk of Chinese population, *Proc. Nat. Acad. Sci.*, 106, 21063–21067, 2009.

Zhang, Y., Shen, H., Tao, S., and Ma, J.: Modeling the atmospheric transport and outflow of polycyclic aromatic hydrocarbons emitted from China, *Atmos. Environ.*, 45, 2820–2827, 2011.

Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 1. Comparison of the modeled BaP concentrations with observations at different sites: (a) a map of the locations of these sites; (b) seasonally averaged total BaP concentration at Changdao, China during 2003–2004; (c) comparison of weekly averaged particulate phase total PAH concentration at Gosan, Korea with modeled total BaP concentration during 2001–2004; (d) comparison of total PAH concentration at Okinawa, Japan with modeled total BaP concentration during May, 2004; (e) comparison of total BaP concentration at Kanazawa, Japan with modeled total BaP concentration during 2003–2004; (f) comparison of the total PAH concentration at CPO, US with modeled total BaP concentration during March–May, 2002; (g) comparison of total PAH concentrations over MBO, US with modeled BaP concentration during 2004–2006.

Transpacific transport of BaP emitted from Asia

Y. Zhang et al.




Fig. 2. Annual mean total BaP concentration (pg m^{-3}) and associated wind vectors over the northern Pacific Ocean in 2004: **(a)** horizontal distribution of total BaP and wind concentration at 1200 m; **(b)** a cross section showing the meridional mean concentration and wind between 25–55° N, the vertical winds are magnified by 1000.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 3. Seasonal variation of transpacific transport of BaP in 2004: **(a)** monthly mean column BaP concentration over North America (blue curve), which was chosen as an indicator of transpacific transport of BaP; the monthly emission flux over Asia (brown curve) is also shown; **(b), (c)** monthly mean total BaP and associated wind vectors concentration in January and July, 2004 at a height of 1200 m over the northern Pacific Ocean; the color bar for concentration and the scale for wind vectors are the same as that in Fig. 2.

Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Transpacific transport of BaP emitted from Asia

Y. Zhang et al.




Fig. 4. Annual mean concentration of BaP induced by emissions from different Asian sources: (a)–(f) North Asia, East Asia, South Asia, Southeast Asia, the Indochina Peninsula, and West Asia, respectively; (g) contribution of the Asian BaP emission sources to the average BaP concentration over North America in 2004; each box represents a 10 degree by 10 degree area.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

**Transpacific
transport of BaP
emitted from Asia**

Y. Zhang et al.




Fig. 5. Annual mean BaP concentration (pg m^{-3}) at 10 m above ground level induced by Asian emission sources in 2004.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Transpacific transport of BaP emitted from Asia

Y. Zhang et al.




Fig. 6. BaP concentration (background color) and wind field (arrows) at 1200m during the transport event on 12 May 2002; **(a)–(f)** show 30 April, 3 May, 5 May, 7 May, 10 May and 12 May, respectively. The black triangle in the panel c marks the location of CPO.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 7. Interannual variability of transpacific transport of BaP during 1948–2007. **(a)** interannual variation of BaP emission over East Asia and mean BaP column concentration over the region of North America within the boundaries of 30–70° N, 75–125° W, and 0–7 km altitude; **(b)** interannual variation of transpacific transport index (TPI) anomaly, and its decadal running mean; **(c)** the relationship between TPI anomaly and the anomaly of the Pacific North America Index (PNA); **(d)** the relationship between TPI anomaly and the anomaly of the Southern Oscillation Index (SOI).

Transpacific transport of BaP emitted from Asia

Y. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

