

Received: 9 October 2010 – Accepted: 1 December 2010 – Published: 20 January 2011

Correspondence to: L. X. Zhou (zhoulx@cams.cma.gov.cn)

Published by Copernicus Publications on behalf of the European Geosciences Union.

ACPD

11, 1939–1969, 2011

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

High frequency measurements of carbon monoxide (CO) recorded over three years at Mount Waliguan (WLG), a global background station in remote western China, were examined using back trajectory analysis. Corrections for the drift in reference gases were also included in the data revision. Between July 2004 and June 2007, a time series of CO exhibited large fluctuations and the 5%, 50% and 95%-percentiles of relevant CO mixing ratios were 102 ppb, 126 ppb and 194 ppb. Approximately 50% of all observed data have been selected as CO background data using a mathematical procedure of robust local regression with the remainder affected by regional-scale pollution. The monthly mean background CO mixing ratios showed a minimum in summer and a maximum in late winter, although all seasons were effected by short-term enhancements that exceeded background levels two or more times. The CO data were compared to the values observed at the high alpine research station Jungfrauoch, Switzerland. Smaller seasonal amplitudes were observed at WLG compared to the Jungfrauoch due to lower winter and spring CO levels, however, episodic enhancements of polluted air were much greater at the site in China. The air parcels arriving at WLG came predominately from the West, except in summer when advection from the East and Southeast prevailed. Transport from the East typically brought polluted air to the site, having passed over populated urban areas upwind. A large number of elevated CO mixing ratios could also be associated with advection from the Northwest of WLG via the central Xinjiang Uygur Autonomous Region (XUAR) and the Ge'ermu urban area where growing industrial activities as well as crops residue burning provide large sources of CO. These background conditions were observed most frequently when air masses originated from remote Tibet west of WLG. The probability that air parcels pass over regions of clean or polluted regions was further identified using potential source contribution function (PSCF) analysis.

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Carbon monoxide (CO) effects the tropospheric oxidizing capacity as it is the main sink of the hydroxyl radical (OH) (Logan et al., 1981; Thompson, 1992). Through its role in removing OH, CO has been identified as an important indirect greenhouse gas (Daniel and Solomon, 1998). With an atmospheric lifetime ranging from 10 days in summer over continental regions to more than a year over polar regions in winter (Novelli et al., 1992), the uneven distribution of sources causes large spatial and temporal CO variations. The main sources of CO are associated with primary emissions from combustion processes (transport, heating, industrial activities and biomass burning) and the oxidation of methane and non-methane hydrocarbons (e.g., Logan et al., 1981; Holloway et al., 2000).

The gas chromatographic technique combined with a mercuric oxide detector (GC-HgO) has been widely used in quasi-continuous measurements of CO mixing ratios at many measurement stations over the globe because of its low detection limit and good precision (Novelli et al., 1992, 1998). However, the nonlinearity of the technique requires careful calibration. The drift of CO mixing ratios in standard gases over time may further affect the measurement accuracy (Novelli et al., 2003; Zellweger et al., 2009). As a result, the observed discrepancies between different labs during inter-comparison activities have often exceeded the limits stated in the quality objectives of the World Meteorological Organization/Global Atmosphere Watch (WMO/GAW) (WMO, 2009) and have, to a certain extent, limited the accuracy of CO regional source and sink estimation from their spatial and temporal distributions.

Although industrial growth in Asia has the potential to emit large amounts of CO, there are few monitoring stations on the continent. The location of WLG is influenced both by background and polluted air, from which changes in regional CO abundance and sources can be examined. For the proper use of the WLG data it is essential to first evaluate the quality of the CO measurements and the calibration history. In this paper, we establish the link between the WLG measurements and the WMO2000 CO scale.

ACPD

11, 1939–1969, 2011

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



We then characterize the CO time series in terms of background vs. polluted air parcels and compare the results with those observed at Junfraujoch, Switzerland. Lastly, using two trajectory-based statistical approaches, we examine how the meteorological conditions and potential source regions influence WLG observation.

2 Site description and experimental method

2.1 Description of the site

Waliguan Baseline Observatory (WLG) (36.28° N, 100.09° E, 3816 m a.s.l.), situated in remote western China (Fig. 1), is one of the 27 WMO/GAW global baseline stations. It is located at the edge of the northeastern part of the Tibetan Plateau and run by the China Meteorological Administration (CMA). The area surrounding the station is pristine with sparse vegetation, mostly arid and semiarid grassland. Small farms are located in the lower valley area, with yak and sheep grazing during the summer months. Xining city lies approximately 90 km northeast of WLG with a total population of 2.2 million. The predominant wind directions are southwest in winter and northeast/southeast in summer (Zhou et al., 2003). Measurements from WLG could provide essential information on sources and sinks within the Eurasian continent because of its unique location (Zhou et al., 2004, 2005, 2006).

2.2 Experimental methods

2.2.1 Instrumentation and analysis

CO mixing ratios at WLG were measured using an RGA-3 gas chromatograph (GC) equipped with an HgO reduction detector (Trace Analytical Inc.). The automated sampling module was designed to sample from separate gas streams (standard tanks and ambient air) supplied to the instrument. The ambient air was delivered to the GC at approximately 5 L min⁻¹ by a KNF Neuberger N2202 vacuum pump via a dedicated 80 m

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



0.95 cm i.d. Dekabon tube sampling from the intake on an 89 m steel triangular tower located approximately 15 m away from the laboratory. The flow rate through the sample line was about 1 L min⁻¹. After the pumping, a valve bleed reduced the flow to approximately 0.23 L min⁻¹. The air was then filtered via a 7 μm inline filter and cryogenically dried to a dew point of -65 °C. The ambient air was then distributed to the GC system. The flow through the 3 mL sampling loop was stopped a few seconds before injection to equilibrate the pressure. The flow rate of the carrier gas nitrogen (N₂, 99.999%) was 0.03 L min⁻¹. The ambient air first passed through a pre-column (Unibeads 1S 60/80) and thereafter into an analytical column (Molecular Sieve 5A 60/80), maintained at 90 °C. The sample then passed into a reaction chamber (265 °C) where CO was oxidized to CO₂ and HgO reduced to Hg; Hg was then detected by UV absorption. A HP3396 integrator acquired and processed the detector analog information. Peak height was used for CO quantification. Sixteen injections were made every hour including two reference gases. The hourly, daily and monthly averages were aggregated from 14 ambient air samples per hour (see Sect. 2.2.2). The overall precision, defined as the standard deviation of all determined results by continuously measuring a series of reference tanks with CO mixing ratios of approximately 60–300 ppb (30 times for each tank), was 1.0–2.5 ppb.

In some observed periods, no data were collected when new analytical columns and HgO beds were exchanged for the old ones. Other missing data resulted from periods of instrument calibration or, more commonly, instrument malfunction (e.g. problems with stream selection valves, which sometimes shifted to a wrong position). Those data have been flagged and are not included. During the period of study acceptable data were collected approximately 75% of the time.

2.2.2 Standards and calibrations

Both the absolute response and the instrument linearity can change over time using the GC-HgO technique (Novelli et al., 1992). Therefore a calibration sequence using working low (WL, ~100 ppb) and working high (WH, ~290 ppb) reference gases was applied

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(WH followed by 7 air samples, WL followed by another 7 air samples). Ambient CO mixing ratios were determined by the updated linear curve of the WH and WL in each hour. Target tanks with assigned CO mixing ratios of 203.9 ppb (July 2004–December 2004) and 254.9 ppb (January 2005–June 2007) were injected every six hours to track and evaluate the performance of the system (Fig. 2). 92% of the measurements of the target tanks were within ± 5.0 ppb of their assigned value, indicating good long-term system stability.

To account for non-linear instrument response, a weekly calibration sequence with working gases followed by a set of six primary gases (S1–S6, 29.5 L aluminum cylinders purchased from Scott Marrin, Inc., with CO ranging from ~ 50 to 300 ppb) at WLG was used. The primary gases were calibrated in 1997 at Environment Canada (EC, formally the Meteorological Service of Canada, MSC) against standards referenced to WMO1988 CO scale (Novelli et al., 1994a, b). The assigned values for S1–S6 are listed in Table 1. Assigned CO mixing ratios for the working standards were updated every week using the quadratic fitting function derived from the six primary gases at WLG.

2.3 Calibration scale conversions

The CO reference scale maintained at National Oceanic and Atmospheric Administration/Earth System Research Laboratory (NOAA/ESRL) was designated by WMO as the reference scale for the GAW program. As the Central Calibration Laboratory (CCL) for GAW, NOAA provides calibrated standards to other laboratories. The certified values from the CCL were revised in 2002 to account for drift in the original WMO1988 scale and are denoted as WMO2000 (WMO, 2010).

The six primary gases (S1–S6) at WLG were calibrated at EC against the original WMO1988 scale. After the EC standards were revised by the CCL in response to a WMO scale correction (Novelli et al., 2003), EC provided WLG with a conversion algorithm to update the WLG primary standards to WMO2000. The

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



revised WLG CO mixing ratios were calculated using a quadratic conversion curve ($y=0.0002\cdot x^2-0.1036\cdot x+13.99$, $R^2=0.9856$). In the function, “x” refers to CO mixing ratios in WLG working scale; “y” refers to the difference of calibrated results (WLG working scale minus WMO2000 scale). The results are listed in Group 1 of Table 1.

We also used results from the WMO World Calibration Centre (WCC)-Empa CO Audits in 2000 and 2004 at WLG to revise the WLG standards (Zellweger et al. 2000, 2004). In 2000, four cylinders from WCC were calibrated against working standards at WLG. Based on differences in the WLG and WCC results, a quadratic fit of $y=0.0002\cdot x^2-0.1233\cdot x+17.896$ ($R^2=1$) was applied to update the S1–S6 values. Here “x” refers to CO mixing ratios in WLG working scale, “y” is the estimated bias (WLG working scale minus WMO2000). The results are shown in Table 1, Group 2. Because the CO values of WCC cylinders covered the range from 58.6 to 207.1 ppb, the function is more accurate and robust for calculating S1–S4; the results for S5 and S6 contain larger uncertainties and were not used in the revision.

We further tied the WLG working standards to the WMO2000 scale using the results of the WCC audit at WLG in 2004. Six WCC CO traveling standards (referenced to WMO2000 scale, covering the range 73.4 to 304.5 ppb) were used to re-calibrate the S1–S6 primary standards; in this case, the CO mixing ratios of the S1–S6 cylinders were regarded as unknown. We use the assigned values of the six WCC cylinders to calculate CO values of S1–S6. The results of these on-site calibrations are shown in Table 1, Group 3.

We averaged only results (marked with an asterisk in Table 1) that fell within the range of CO mixing ratios in these transfer gases – as the GC non-linear response may not be accurately characterized above and below this range. Then the ambient CO mixing ratios were updated and recalculated using the newly assigned CO mixing ratios.

In summary, the CO mixing ratios for the six WLG primary standards were revised based upon results from the re-analysis by EC and the two WCC audits. The WLG working standards were recalculated based on the new primary assignments and all

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



CO measurements were revised based on the corrected working standards. All the results are reported on the WMO2000 scale.

3 Results and discussions

3.1 Revised CO data

5 New calibration functions were generated using the revised CO mixing ratios of the S1–S6 references. These functions were then used to re-process ambient CO mixing ratios. Figure 3 displays an overlay of the original and the revised daily averages of the ambient CO mixing ratios. As can be seen in Table 1, the largest correction to the original standard CO assignments occurred at the lower end of the scale. Therefore, as
10 the levels of ambient CO varied, the magnitude of the correction to the ambient samples changed accordingly. Changes in CO were typically greatest in samples with a lower mixing ratio (7–8 ppb below ~100 ppb) and decreased to 2 ppb at 200 ppb (Fig. 4).

3.2 Time series and monthly variations

15 Figure 5 shows the hourly data of CO at WLG from July 2004 to June 2007. Enhanced CO mixing ratios were observed with some episodes exceeding 200 ppb, and back trajectories for the specific periods of high CO have traced these episodes to regions with large sources (see next section). To define CO mixing ratios representing background air, we used a statistical approach for extracting background concentrations from measurements. This method is based on robust local regression called REBS
20 (robust extraction of background signal). It is a modified version of a technique of robust baseline estimation (RBE) and developed for baseline removal from chemical analytical spectra (Ruckstuhl et al., 2001, 2010). Approximately 50% of all observed data were thus classified as CO background data. The distribution of CO mixing ratios is characterized by value of 102 ppb, 126 ppb and 194 ppb for the 5%, 50% and

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



95%-percentile. Data classified as background (Fig. 5) exhibited low mixing ratios and smooth changes; data classified as pollution influenced exhibit large fluctuations compared to background values.

To investigate regional influences on mixing ratios of CO observed at WLG, we compared both background and pollution values with data from another baseline station (Jungfrauoch, Switzerland, 46.55° N, 7.99° E, 3580 m a.s.l. of <http://gaw.empa.ch/gawsis/>) in the Northern Hemisphere at a similar altitude (Fig. 6). Jungfrauoch (JFJ), situated in Central Europe, is influenced by both clean air masses and polluted boundary layer air (Zellweger et al., 2003) and observations at JFJ has been used to assess locations of emissions from the European continent (Reimann et al., 2008). The CO observations at JFJ are performed by Empa as part of the Swiss National Air Pollution Monitoring Network and as a contribution to the GAW programme. The data selection algorithm used to define background CO at WLG was also applied to the JFJ data.

The background CO monthly means at WLG exhibited a seasonal cycle typical of the Northern Hemisphere where in the late winter/early spring CO reaches its annual maximum before a rapid decrease to its summer minimum resulting from reaction with the OH radical. Similarly, the general characteristics of background CO at WLG are consistent with JFJ. However, while the summer CO background mixing ratios at WLG and JFJ are similar, we find larger differences in background CO in winter when JFJ is 20–40 ppb higher, possibly reflecting different surface sources and greater wintertime OH at WLG.

Pollution episodes were characterized by significantly enhanced mixing ratios and large and erratic variations. This apparently reflects significant influence of regional emission sources on WLG. The greatest enhancements above background occur in the winter and spring months due to transport of pollution from the northwest of the site (as further discussed in Sects. 3.3 and 3.4). The relatively lower enhancements in summer may reflect CO loss during transport.

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.3 Long-distance transport impact on ambient CO

3.3.1 Trajectories analysis of selected observation episodes

Statistical analysis of air mass back trajectories combined with long-term ambient air pollution measurements can be used for analyzing large-scale air pollutant transport and source identification at a receptor site (Stohl, 1996; Rousseau et al., 2004). Trajectory position error of the method is typically considered to be about 20% of the traversed distance (Stohl, 1998), but the statistical uncertainty can be reduced with large sets of trajectories. In our study, the CO data from July 2004 to June 2007 were analyzed using trajectory statistics. Five day (120 h) back trajectories originating at 00:00, 06:00, 12:00 and 18:00 UTC were calculated by the Hybrid Single-Particle Lagrangian Integrated Trajectory model (HYSPPLIT) (Draxler and Hess, 1998) using NCEP(National Centers for Environmental Prediction) reanalysis data. The arrival height of the trajectories was 500 m above ground level which has been shown to be the most representative for air sampled at WLG (Zhou et al., 2003, 2004). Figure 7 shows hourly data of ambient CO for 2006. Two pollution episodes (20–23 January 2006 and 2–5 July 2006) and one baseline period (26–29 October 2006) were selected as examples for evaluating source regions. Figure 8 shows the trajectories for the selected days. The period in October represents background conditions when corresponding trajectories track to the Tibet and south Xinjiang Uygur Autonomous Region (XUAR). The episode of high CO in July is associated with the trajectories from the heavily populated regions east or southeast of WLG suggesting a large source area due to human activities. During the event on 20–23 January 2006, the highest CO values were observed when the trajectories indicated transport of air parcels arrive from the area northwest of WLG and via the Ge'ermu urban area (the second largest city of Qinghai), a major source region due to rapid industrial development caused by abundant petroleum oil and natural gas resources, additional emissions are expected from biofuel burning in the region.

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.3.2 Cluster analysis

Cluster analysis, which combines the trajectories into representative spatial groups using multivariate statistics, was employed to identify distinct transport patterns (Fig. 9). Angle distance was chosen as the clustering criterion (Sirois and Bottenheim, 1995).

5 We used CO hourly averages corresponding to trajectory arrival times only at 00:00, 06:00, 12:00 and 18:00 UTC in the analysis. Each trajectory in a cluster was identified and then used to obtain cluster statistics (Table 2, polluted CO criteria in each season was determined by the maximum background CO: 132.4 ppb for spring, 121.6 ppb for summer, 109.6 ppb for autumn and 124.9 ppb for winter). Table 3 lists regional characteristics for each cluster by season.

10 Within each cluster, air parcels transported to WLG are characterized as either background or polluted using the background subtraction method described previously. In spring, cluster 3 contains air parcels mainly from the area southeast of the WLG, 85%, which are characterized as polluted (Table 2 and Fig. 9). The air parcels from cluster 2, arriving from the Northeast and representing 24% of the transport, had also elevated CO levels, reflecting potential sources probably due to the biomass burning in Northwest Gansu province, the largest agricultural region in Northwest China. Cluster 1 (with 71% of the trajectories) corresponds to the air parcels coming from the central region of XUAR and north of the Qinghai province. The relatively remote area defined in cluster 1 is still a potential CO source region mainly due to agricultural biomass burning

20 (Yan et al., 2006) and a growing economy (from fossil fuel combustion in the region). In summer, the most elevated CO mixing ratios are associated with cluster 3 which passed through the urbanized area southeast of WLG (e.g. Lanzhou city, the central region and southeast of Gansu province). The clusters 1 and 2 that represent advection from the North or Northwest (central region of XUAR and region northwest of Qinghai) with relatively low CO mixing ratios compared to those of observed in cluster 3, but they still contain a large number of trajectories with enhanced CO originating in more polluted areas. As in spring, fossil fuel combustion and biomass burning add CO to

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



this region. However, in summer, unlike other times of the year, WLG is significantly influenced by the air masses from the populated southeastern regions. In autumn and winter the high CO mixing ratios were shown in which the airflow originated in the East or Southeast. All year round, air parcels originating from this direction exhibit the highest CO levels.

As seen in spring and summer, during autumn and winter polluted air arrives from the West or Northwest. This further indicates that the XUAR region and the North Qinghai province is a potential CO source region, which is also consistent with model estimations (GEOS-CHEM) using biofuel and fossil fuel emissions determined by Yevich and Logan (2003) and Bey et al. (2001), as reported in Liang et al. (2004). These showed a band of “hot spots” in the region, in agreement with studies having found strong CO emissions from biomass burning especially during cold winters (Yan et al., 2006). The large number of baseline data in which the air masses originated from the West or a southwestern area (via sparsely inhabited Tibet) suggest a relatively clean environment.

3.3.3 PSCF analysis

We also used a potential source contribution function analysis approach (PSCF) for further investigation. PSCF can compute probability density functions reflecting the residence times of an air parcel over a given geographical area prior to its arrival at the measurement site using back trajectories (Ashbaugh et al., 1985; Begum et al., 2005). To calculate the PSCF, the geographic region covered by the trajectories is divided into an array of 0.3×0.3 degree grid cells (with a height of up to 500 m from ground). The PSCF is a location function defined by cell indices i and j while the number of trajectories that pass through the ij -th cell is denoted by n_{ij} . The number of trajectories resulting in CO values that exceed the background value is defined by m_{ij} . The probability that any given cell can be related to enhance CO is defined following (Polissar et al., 1999):

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



$$\text{PSCF}_{ij} = m_{ij}/n_{ij}$$

This method tends to give good angular but poor radial resolution because the trajectories converge as they approach the receptor (Vasconcelos et al., 1996). Hence, because grid cells always have the same PSCF value regardless of whether CO mixing ratios are slightly higher or much higher than the threshold, making it difficult to distinguish between moderate and strong CO sources. To reduce this effect, PSCF values were weighted with an arbitrary coefficient W_{ij} (Polissar et al., 1999), which reduced them when the total number of the endpoints in a particular cell was less than about three times the average value of the endpoints in each cell. Here W_{ij} was defined as:

$$W_{ij} = \begin{cases} 1.00 & n_{ij} > 80 \\ 0.70 & 20 < n_{ij} \leq 80 \\ 0.42 & 10 < n_{ij} \leq 20 \\ 0.05 & n_{ij} \leq 10 \end{cases}$$

Figure 10 shows the potential source regions resulting from trajectory statistics of CO in the four seasons. Regions in black and white respectively indicate strong and weak potential CO sources. In general, the CO source locations identified by PSCF are very similar to the results obtained by the clustering method. In summer, the black and dark gray colored regions (areas more likely to be CO source areas) showed large spatial extensions to the northeast and southeast of WLG, areas with dense populations and industry. Otherwise, in the other three seasons, the PSCF are very similar and the source regions were mainly situated northwest of WLG, influences which may be increasing especially under the strategy of China's western region development. This policy could significantly change the CO source distributions in the future. The largest increases in regional emissions throughout China will have an impact on global CO.

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



4 Conclusions

In this paper, we report a revision to the quasi-continuous observations of CO at WLG to the WMO 2000 scale. The three year record of CO mixing ratios exhibited large fluctuations showing periodic transport of polluted air to the site. We have used a “robust local regression” statistical method to distinguish the background data from polluted air and find that approximately 50% of all observations can be identified as background air. The seasonal variation of background CO shows a minimum in summer and a maximum in winter, which is consistent with that observed in the Northern Hemisphere CO surface distribution and the high altitude measurements at JFJ. Larger differences in background CO in winter between the two sites (when JFJ is 20–40 ppb higher) reflect different surface sources and possibly greater wintertime OH at WLG. Compared to the JFJ, air masses identified at WLG as polluted contained more CO relative to the background values and displayed large and irregular fluctuations suggesting greater influence from regional emission sources. By combining the CO data with trajectory-based statistics and cluster analysis, we have demonstrated that air masses originating from the East to the Southeast of WLG have the strongest impact on CO, however, are only significant in the summer transport from this direction. In the other seasons, air parcels arriving via the Northwest are more common. These exhibit both background and polluted characteristics. Air from the central XUAR and the Ge’ermu urban area have shown enhanced CO levels due to the growing economy in West China as well as biomass burning in the region. Air parcels coming from sparsely populated Tibet contribute least to the enhanced CO values observed at WLG.

Acknowledgement. This work was supported by National Natural Science Foundation of China (40775078), Non-profit Research Project (Grant No. GYHY200806026) International S&T Co-operation Program of MOST (Grant No. 2007DFA20650) and National Basic Research Program of China “973” (Grant No. 2010CB950601). The station personnel have extensively supported the operation at WLG station. Yaqiang Wang is acknowledged for his help with the calculation of trajectories and clustering/PSCF.

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

- Ashbaugh, L. L., Malm, W. C., and Sadeh, W. Z.: A residence time probability analysis of sulfur concentrations at Grand Canyon National Park, *Atmos. Environ.*, 19, 1263–1270, 1985.
- Begum, B. A., Kim, E., Jeong, C. H., Lee, D. W., and Hopke, P. K.: Evaluation of the potential source contribution function using the 2002 Quebec forest fire episode, *Atmos. Environ.*, 39, 3719–3724, 2005.
- Bey, I., Jacob, D. J., Yantosca, R. M., Logan, J. A., Field, B. D., Fiore, A. M., Li, Q., Liu, H. Y., Mickley, L. J., and Schultz, M. G.: Global modeling of tropospheric chemistry with assimilated meteorology: model description and evaluation, *J. Geophys. Res.*, 106, 23073–23096, 2001.
- Daniel, J. S. and Solomon, S.: On the climate forcing of carbon monoxide, *J. Geophys. Res.*, 103, 13249–13260, 1988.
- Draxler, R. R. and Hess, G. D.: An overview of the HYSPLIT_4 modeling system for trajectories, dispersion, and deposition, *Aust. Meteorol. Mag.*, 47, 295–308, 1998.
- Holloway, T., Levy II, H., and Kasibhatla, P.: Global distribution of carbon monoxide, *J. Geophys. Res.*, 105, 12123–12147, 2000.
- Liang, Q., Jaeglé, L., Jaffe, D. A., Weiss-Penzias, P., Heckman, A., and Snow, J. A.: Long-range transport of Asian pollution to the Northeast Pacific: seasonal variations and transport pathways of carbon monoxide, *J. Geophys. Res.*, 109, D23S07, doi:10.1029/2003JD004402, 2004.
- Logan, J. A., Prather, M. J., Wofsy, S. C., and McElroy, M. B.: Tropospheric chemistry: a global perspective, *J. Geophys. Res.*, 86, 7210–7254, 1981.
- Novelli, P. C., Masarie, K. A., Lang, P. M., Hall, B. D., Myers, R. C., and Elkins, J. W.: Re-analysis of tropospheric CO trends: Effects of the 1997, 1998 wildfires, *J. Geophys. Res.*, 108(D15), 4464–4477, doi:10.1029/2002JD003031, 2003.
- Novelli, P. C., Steele, L. P., and Tans, P. P.: Mixing ratios of carbon monoxide in the troposphere, *J. Geophys. Res.*, 102, 12855–12861, 1992.
- Novelli, P. C., Collins Jr., J. E., Myers, R. C., Sachse, G. W., and Scheel, H. E.: Re-evaluation of NOAA/CMDL carbon monoxide reference scale and comparisons with CO reference gases at NASA-Langley and the Fraunhofer Institute, *J. Geophys. Res.*, 99, 12833–12839, 1994a.
- Novelli, P. C., Masarie, K. A., Tans, P. P., and Lang, P. M.: Recent changes in atmospheric carbon monoxide, *Science*, 263, 1587–1590, 1994b.
- Polissar, A. V., Hopke, P. K., Paatero, P., Kaufmann, Y. J., Hall, D. K., Bodhaine, B. A., Dut-

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ton, E. G., and Harris, J. M.: The aerosol at Barrow, Alaska: long-term trends and source locations, *Atmos. Environ.*, 33, 2441–2458, 1999.

Reimann, S., Vollmer, M. K., Folini, D., Steinbacher, M., Hill, M., Buchmann, B., Zander, R., and Mahieu, E.: Observations of long-lived anthropogenic halocarbons at the high-Alpine site of Junfraujoch (Switzerland) for assessment of trends and European sources, *Sci. Total Environ.*, 391, 224–231, doi:10.1016/j.scitotenv.2007.10.022, 2008.

Rousseau, D. D., Duzer, D., Etienne, J. L., Cambon, G., Jolly, D., Ferrier, J., and Schevin, P.: Pollen record of rapidly changing air trajectories to the North Pole, *J. Geophys. Res.*, 109, D06116, doi:10.1029/2003JD003985, 2004.

Ruckstuhl, A. F., Matthew, P. J., Robert, W. F., and James, A. D.: Baseline subtraction using robust local regression estimation, *J. Quant. Spectrosc. Ra.*, 68, 179–193, 2001.

Ruckstuhl, A. F., Henne, S., Reimann, S., Steinbacher, M., Buchmann, B., and Hueglin, C.: Robust extraction of baseline signal of atmospheric trace species using local regression, *Atmos. Meas. Tech. Discuss.*, 3, 5589–5612, 2010, doi:10.5194/amtd-3-5589-2010, 2010.

Sirois, A. and Bottenheim, J. W.: Use of backward trajectories to interpret the 5-year record of PAN and O₃ ambient air concentrations at Kejimikujik National Park, Nova Scotia, *J. Geophys. Res.*, 100, 2867–2881, 1995.

Stohl, A.: Trajectory statistics – a new method to establish source-receptor relationships of air pollutants and its application to the transport of particulate sulfate in Europe, *Atmos. Environ.*, 30, 579–587, 1996.

Stohl, A.: Computation, accuracy and applications of trajectories – a review and bibliography, *Atmos. Environ.*, 32, 947–966, 1998.

Thompson, A. M.: The oxidizing capacity of the Earth's atmosphere: probable past and future changes, *Science*, 256, 1157–1168, 1992.

Vasconcelos, L. A. P., Kahl, J. D. W., Liu, D., Macias, E. S., and White, W. H.: Spatial resolution of a transport inversion technique, *J. Geophys. Res.*, 101, 19337–19342, 1996.

WMO: Guidelines for the Measurement of Atmospheric Carbon Monoxide, GAW Report No. 192, World Meteorological Organization, Geneva, Switzerland, 2010.

Yan, X. Y., Ohara, T., and Akimoto, H.: Bottom-up estimate of biomass burning in mainland China, *Atmos. Environ.*, 40, 5262–5273, 2006.

Yevich, R. and Logan, J. A.: An assessment of biofuel use and burning of agricultural waste in the developing world, *Global Biogeochem. Cy.*, 17, 1095, doi:10.1029/2002GB001952, 2003.

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Zellweger, C., Hofer, P., and Buchmann, B.: System and Performance Audit of Surface Ozone and Carbon Monoxide at the China GAW Baseline Observatory Waliguan Mountain, WCC-Empa Report 00/3, [gaw.empa.ch/audits/WLG_2000.pdf](http://www.empa.ch/plugin/template/empa/*/7571), available at: http://www.empa.ch/plugin/template/empa/*/7571, last access: September 2000, 2000.
- 5 Zellweger, C., Forrer, J., Hofer, P., Nyeki, S., Schwarzenbach, B., Weingartner, E., Ammann, M., and Baltensperger, U.: Partitioning of reactive nitrogen (NO_y) and dependence on meteorological conditions in the lower free troposphere, *Atmos. Chem. Phys.*, 3, 779–796, doi:10.5194/acp-3-779-2003, 2003.
- Zellweger, C., Klausen, J., and Buchmann, B.: System and Performance Audit of Surface Ozone Carbon Monoxide and Methane at the Global GAW Station Mt. Waliguan, China, October 2004, WCC-Empa Report 04/3, [gaw.empa.ch/audits/WLG_2004.pdf](http://www.empa.ch/audits/WLG_2004.pdf), available at: http://www.empa.ch/plugin/template/empa/*/7571, last access: October 2004, 2004.
- 10 Zellweger, C., Hüglin, C., Klausen, J., Steinbacher, M., Vollmer, M., and Buchmann, B.: Inter-comparison of four different carbon monoxide measurement techniques and evaluation of the long-term carbon monoxide time series of Jungfrauoch, *Atmos. Chem. Phys.*, 9, 3491–3503, doi:10.5194/acp-9-3491-2009, 2009.
- Zhou, L., White, J. W. C., Conway, T. J., Mukai, H., MacClune, K., Zhang, X., Wen, Y., and Li, J.: Long-term record of atmospheric CO_2 and stable isotopic ratios at Waliguan Observatory: seasonally averaged 1991–2002 source/sink signals, and a comparison of 1998–2002 record to the 11 selected sites in the Northern Hemisphere, *Global Biogeochem. Cy.* 20, GB2001, doi:10.1029/2004GB002431, 2006.
- 20 Zhou, L. X., Tang, J., Wen, Y. P., Li, J. L., Yan, P., and Zhang, X. C.: The impact of local winds and long-range transport on the continuous carbon dioxide record at Mount Waliguan, China, *Tellus B*, 55, 145–158, 2003.
- Zhou, L. X., Worthy, D. E., Lang, J. P. M., Ernst, M. K., Zhang, X. C., Wen, Y. P., and Li, J. L.: Ten years of atmospheric methane observations at a high elevation site in Western China, *Atmos. Environ.*, 38, 7041–7054, 2004.
- 25 Zhou, L. X., Conway, T. J., White, J. W. C., Mukai, H., Zhang, X., Wen, Y., Li, J., and MacClune, K.: Long-term record of atmospheric CO_2 and stable isotopic ratios at Waliguan Observatory: background features and possible drivers, 1991–2002, *Global Biogeochem. Cy.*, 19, GB3021, doi:10.1029/2004GB002430, 2005.
- 30

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. S1–S6 primary standards cylinders revision for CO mixing ratios.

Cylinder#	CA01488	CA01500	CA01441	CA01457	CA01459	CA01449
Used as	S1	S2	S3	S4	S5	S6
Assigned (ppb) ^a	49.4	95.2	150.2	202	254.9	297.4
Group1 ^b	58.8*	101.1*	153.1*	203.2*	255.5*	298.3*
Group2 ^c	61.7*	103.2*	154.1*	203.2*	254.4	296.3
Group3 ^d	59.0	103.9*	155.7*	205.5*	259.1*	298.5*
Mean (see text)	60.2	102.7	154.3	204.0	257.3	298.4
S.D.	2.1	1.4	1.3	1.3	2.6	0.2
Expanded uncertainty	4.2	2.8	2.6	2.6	5.2	0.4

^a WLG working scale, in ppb, assigned by MSC in 1997 against the WMO88 scale.

^b WMO2000, in ppb, calculated based on the curves from EC.

^c WMO2000, in ppb, calculated based on the data of WCC 2000 CO audit at WLG (CO span of WCC cylinders covering from 58.6 to 207.1 ppb).

^d WMO2000, in ppb, calculated based on the data of WCC 2004 CO audit at WLG (Range of the 2004 audit gases from 73.4 to 304.5 ppb).

* Results used to assigned mixing ratios to the WLG standards. These are referenced to the WMO2000 scale.

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Table 2. Statistical CO results of clusters in different seasons at WLG.

	Cluster	Background conditions			Polluted conditions		
		Number of Trajectories (–)	Mean (CO) (ppb)	s.d. (CO) (ppb)	Number of Trajectories (–)	Mean (CO) (ppb)	s.d. (CO) (ppb)
Spring	1	202	121.8	6.2	169	154.5	26.8
	2	26	125	6.1	100	177.8	33.2
	3	4	122.9	4.9	23	183.3	38.2
Summer	1	107	107.4	7.1	61	141.3	18
	2	21	111.3	4.8	53	150.3	25.3
	3	16	115.9	5.4	153	178.8	41.3
	4	40	112.9	5.3	120	154.7	28
Autumn	1	281	99.2	6.4	176	141.4	42.5
	3	10	97.2	3.4	16	149.2	31.1
	2	2	94.8	4.3	38	181.1	67.8
Winter	1	240	112.8	6.4	148	183.3	75.8
	2	103	113.7	5.9	78	181.4	68.6
	3	0	NA	NA	12	206.4	65.5

The “cluster number” indicates different air parcel origin/routes (see Fig. 9). The statistical results for background and polluted CO mixing ratios in different seasons are listed here: “Number of Trajectories(–)” refers to quantity of trajectories in each corresponding cluster; “Mean(CO) (ppb)” refers to averaged CO mixing ratios; s.d.(CO) (ppb) refers to standard deviation of the background CO mixing ratios.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. Characteristics for each cluster in all seasons.

Season	Cluster	Region	Characteristic
Spring	1	Central region of XUAR and north of Qinghai province	Fossil fuel resources and agricultural
	2	northwest of Gansu province	Rural
	3	southeast of Gansu province	Populated
Summer	1	Central region of XUAR and northwest of Qinghai province	Fossil fuel resources and agricultural
	2	northwest of Gansu province	Rural
	3	southeast of Gansu province	Populated
	4	central region of Gansu province	Rural
Autumn	1	Northwest of Qinghai and some region of WSoutheast XUAR	Fossil fuel resources and agricultural
	2	northwest of Gansu province	Rural
	3	southeast of Gansu province	Populated
Winter	1	Central region of XUAR and northwest of Qinghai	Fossil fuel resources and agricultural
	2	northwest of Tibet and Qinghai province (only in this season)	Undeveloped
	3	centrally and southeast of Gansu province	Populated




Fig. 1. Map of Waliguan (black triangle) and location of main cities (black dots).

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 2. Deviation of CO mixing ratios for target gases from their assigned value.

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 3. Overlay of the revised (black dots) and original (white circles) daily averages of CO; the gray line shows the bias of the original data which indicates a bias of the WLG working scale relative to the WMO2000 scale.

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 4. Ambient CO: WMO2000 minus WLG working scale.

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.




Fig. 5. Ambient CO mixing ratios obtained from in situ measurement (hourly averages, black lines represent background conditions, and gray lines represent data influenced by polluted air masses) at WLG (July 2004–June 2007).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 6. Monthly mean mixing ratios of ambient CO and its standard deviations at Waliguan and Jungfrauoch for both air identified as background and as polluted.

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.

**Fig. 7.** Hourly data of ambient CO for the whole year of 2006.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.




Fig. 8. Trajectories for selected days (5d back trajectories at 00:00, 06:00, 12:00 and 18:00 UTC), black cross is the location of Geer'mu city.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Atmospheric carbon monoxide at Mount Waliguan, China

F. Zhang et al.




Fig. 9. Isobaric trajectory clusters (+500 m) in spring, summer, autumn and winter at WLG, July 2004–June 2007. Numbers beside each cluster indicate cluster sequence; the percentages present the trajectory counts of the total.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Atmospheric carbon
monoxide at Mount
Waliguan, China

F. Zhang et al.




Fig. 10. Potential source regions calculated from trajectory statistics during July 2004–June 2007. WPSCF values have an arbitrary unit, higher values (e.g. WPSCF > 0.6) indicating a higher probability for a grid cell of CO source.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)