

**Aerosol mass
spectrometer data in
Las Vegas, with EPA
PMF**

S. G. Brown et al.

Receptor modeling of near-roadway aerosol mass spectrometer data in Las Vegas, Nevada, with EPA PMF

**S. G. Brown^{1,2}, T. Lee², G. A. Norris³, P. T. Roberts¹, J. L. Collett Jr.²,
P. Paatero⁴, and D. R. Worsnop⁵**

¹Sonoma Technology, Inc., Petaluma, California, USA

²Colorado State University, Atmospheric Science Department, Fort Collins, Colorado, USA

³US EPA, Office of Research and Development, Research Triangle Park, North Carolina, USA

⁴University of Helsinki, Dept. of Physics, Helsinki, Finland

⁵Aerodyne Research, Billerica, Massachusetts, USA

Received: 1 June 2011 – Accepted: 27 June 2011 – Published: 15 August 2011

Correspondence to: S. G. Brown (sbrown@sonomatech.com)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

Ambient non-refractory PM₁ aerosol particles were measured with an Aerodyne High Resolution Time-of-Flight Aerosol Mass Spectrometer (HR-AMS) at an elementary school 20 m from the US 95 freeway in Las Vegas, Nevada, during January 2008. Additional collocated continuous measurements of black carbon (BC), carbon monoxide (CO), nitrogen oxides (NO_x), and meteorological data were collected. The US Environmental Protection Agency's (EPA) positive matrix factorization (PMF) data analysis tool was used to apportion organic matter (OM) as measured by HR-AMS, and rotational tools in EPA PMF were used to better characterize the solution space and pull resolved factors toward known source profiles. Three- to six-factor solutions were resolved. The four-factor solution was the most interpretable, with the typical AMS PMF factors of hydrocarbon-like organic aerosol (HOA), low-volatility oxygenated organic aerosol (LV-OOA), biomass burning organic aerosol (BBOA), and semi-volatile oxygenated organic aerosol (SV-OOA). When the measurement site was downwind of the freeway, HOA composed about half the OM, with SV-OOA and LV-OOA accounting for the rest. Attempts to pull the PMF factor profiles toward source profiles were successful but did not qualitatively change the results, indicating that these factors are very stable. Oblique edges were present in G-space plots, suggesting that the obtained rotation may not be the most plausible one. Since solutions found by pulling the profiles or using F_{peak} retained these oblique edges, there appears to be little rotational freedom in the base solution. On average, HOA made up 26 % of the OM, and it made up nearly half of the OM when the monitoring site was downwind of US 95 during morning rush hour. LV-OOA was highest in the afternoon and accounted for 26 % of the OM. BBOA occurred in the evening hours, was predominantly from the residential area to the north, and on average constituted 12 % of the OM; SV-OOA accounted for the remaining third of the OM. Use of the pulling techniques available in EPA PMF and ME-2 suggested that the four-factor solution was very stable.

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

A number of studies have demonstrated the significant health impacts of air pollution such as particulate matter of aerodynamic diameter less than 2.5 microns ($PM_{2.5}$), including increased asthma rates, detrimental fetal development during pregnancy, and decreased lung capacity (Brunekreef et al., 1997; McDonald et al., 2004; Dockery et al., 1993; Dockery and Stone, 2007). In particular, recent literature has demonstrated that adverse health effects are evident in subjects exposed to ambient air near major roadways (Edwards et al., 1994; Nitta et al., 1993; Kunzli et al., 2000; Hoek et al., 2002; Finkelstein et al., 2004; Kim et al., 2004b). A wide body of literature of previous near-roadway air pollution studies (Zhu et al., 2002; K. M. Zhang et al., 2004; Phuleria et al., 2007; Ntziachristos et al., 2007; Fruin et al., 2008) has demonstrated that, for a number of pollutants such as black carbon (BC), carbon monoxide (CO), and ultrafine particles, concentrations are very high next to the freeway but rapidly decrease to background urban levels after a few hundred meters under persistent winds. In pre-sunrise hours and low wind conditions, the influence of freeway emissions can reach even further (Hu et al., 2009).

Organic matter (OM) is a large and important part of $PM_{2.5}$, particularly in near-roadway environments, and is often the largest component of PM in urban areas in the western United States (Phuleria et al., 2007; Riddle et al., 2008; Minguillon et al., 2008). OM is a complicated mixture of thousands of individual molecules and is a combination of both primary particulate emissions and secondary aerosol formed from gaseous precursors. It is a major component of vehicular exhaust emissions and, in addition to being a large part of $PM_{2.5}$ mass, also includes polycyclic aromatic hydrocarbons (PAHs), which are carcinogenic (Larsen and Baker, 2003; Lobscheid and McKone, 2004; Adonis et al., 2003; Flowers et al., 2002).

The composition of OM has been analyzed under a number of different methods, including molecular marker analysis of aerosol filter samples and real-time analysis with instruments such as the Aerodyne Aerosol Mass Spectrometer (AMS) and its more

ACPD

11, 22909–22950, 2011

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper



recent version, the High Resolution Time-of-Flight Aerosol Mass Spectrometer (HR-AMS) (Drewnick et al., 2005; Jimenez et al., 2003; Zhang et al., 2005a; Allan et al., 2003a, b, 2004; DeCarlo et al., 2006). With the AMS, individual molecular marker compounds are not typically quantified, but with this loss of molecular specificity we gain high time resolution and a more complete representation of the full organic fraction. Rather than individual molecules, specific groups of mass-to-charge ratio fragments (m/z) can be used to identify differences between less oxidized, hydrocarbon-like organic aerosol (HOA) and oxygenated organic aerosol (OOA). Some examples include data from Pittsburgh, Pennsylvania (Q. Zhang et al., 2004, 2005a, b), Riverside, California (Docherty et al., 2008), Zurich (Lanz et al., 2007, 2008a), Mexico City (Dzepina et al., 2007; Johnson et al., 2008; Aiken et al., 2008), and Fraser Valley, British Columbia (Alfarra et al., 2004, 2007).

Laboratory and field studies have found that m/z 44 (i.e. CO_2^+) is a major fragment when long range transport is important and during periods of active photochemistry and that m/z 44 is representative of OOA. In prior field studies in Pittsburgh, British Columbia, and elsewhere (Q. Zhang et al., 2004, 2005a, b; Alfarra et al., 2004, 2007), saturated hydrocarbon fragments such as m/z 57 (i.e. C_4H_9^+) were found to be typical of HOA.

AMS data have, in recent years, been further evaluated with positive matrix factorization (PMF) or other mathematical methods (Q. Zhang et al., 2004, 2005a, b) to decompose the mass spectra and quantify the amount of OOA and HOA impacting a monitoring site. Source studies of primary emissions – such as diesel exhaust in a vehicle-chasing experiment (Canagaratna et al., 2004), source profile acquisition (Mohr et al., 2009), and reaction chamber secondary OA (SOA) (Alfarra et al., 2006) – show similarities to spectra from ambient aerosol. These two types of primary OA (POA) and SOA profiles resemble those observed in the ambient air under conditions more conducive to primary (high m/z 57 concentrations) and secondary (high m/z 44 concentrations) influences, respectively. In addition, two types of OOA spectra have been observed in Riverside and elsewhere (Docherty et al., 2008; Huffman et al., 2009)

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

(Jimenez et al., 2009), one with more “low volatility” fragments (LV-OOA) including m/z 44, but also one with significant mass from other, less oxidized fragments. This spectrum has also been observed in chamber experiments with diesel exhaust and resembles semi-volatile OA with some oxidized OA, termed SV-OOA.

2 Methods

2.1 Aerosol and gaseous measurements

Measurements were made next to a classroom and play yard at Fyfe Elementary School, directly adjacent to and 18 m from the US 95 soundwall (Fig. 1). In 2007, the annual average daily traffic (AADT) was between 189 000 and 201 000 vehicles on the stretch of US 95 near Fyfe Elementary School. The wind sector from 90 to 250 degrees encompasses the adjacent freeway. CO, NO_x, Aethalometer BC, wind speed, and wind direction were measured continuously, yielding 5-min averages. A two-channel (370 nm and 880 nm) Magee Scientific Aethalometer was used to measure BC in 5-min intervals. Aerosol was collected on a glass fiber tape, passing through a Harvard impactor with a size-cut of 2.5 microns. Raw data were post-processed with the Washington University Air Quality Lab AethDataMasher Version 6.0e to format date-time stamps and perform data validation. A Thermo Scientific 42i NO/NO₂/NO_x analyzer and a Thermo Scientific 48i CO analyzer provided 5-min NO_x and CO data. Zero-checks and span-checks were performed nightly; data were zero-corrected if the zero-check was greater than 5 % of expected. Continuous gaseous instruments were calibrated at setup, take down and quarterly, plus on an as needed basis via remote control over the internet. 5-min data were visually reviewed daily and after the study for additional QC. Wind speed and direction were measured with an RM Young AQ 5305-L at 1-min intervals, and were used to calculate vector-averaged 5-min average data.

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2 HR-AMS description and data processing

The operation of the Aerodyne HR-AMS has been described in detail elsewhere (DeCarlo et al., 2006; Drewnick et al., 2005; Jayne et al., 2000; Jimenez et al., 2003, 2009). Ambient air is drawn through a URG cyclone ($D_{50} = 2.5 \mu\text{m}$, 3 lpm) and is sampled with
5 2-min time resolution through a critical orifice into an aerodynamic lens, creating a narrow particle beam, with a 50 % efficiency of $1 \mu\text{m}$ particles, so that essentially PM_{10} is measured (Sun et al., 2009; Canagaratna et al., 2007). The particles are accelerated in the supersonic expansion of gas molecules into a vacuum at the end of the lens. Particles are collected by inertial impaction and non-refractory species such as nitrate,
10 sulfate, ammonium, chloride, potassium and OM are thermally vaporized. Vaporized gases undergo electron impact ionization and the charged fragments enter the ToF-MS region, where they are separated by mass-to-charge ratio (m/z). After correction for ambient gases such as N_2 , mass spectra are analyzed for each 2-min averaged sample, and the sum of organic aerosol peaks is used to calculate total OM. AMS data
15 were processed and analyzed using the standard AMS analysis software, Squirrel version 1.48, implemented with Wavemetric's Igor Pro (version 6.12). Concentration and uncertainty data for PMF were regenerated using Squirrel v 1.51. Uncertainty estimates are based on counting statistics methods and generated from Squirrel (Allan et al., 2003b). For a given AMS signal, the error estimate is:

$$20 \quad S(j) \equiv \alpha \sqrt{\frac{I(j_o) + I(j_b)}{t(s)}} \quad (1)$$

where the ion signal is j , the signal when the beam is open is ($I(j_o)$), and the signal when it is blocked is ($I(j_b)$) over the sampling time $t(s)$, with a distribution factor α .

2.3 Positive Matrix Factorization

PMF is a multivariate factor analysis tool (Paatero and Hopke 2009; Paatero, 1997; Paatero and Tapper, 1994, 1993) that has been applied to a wide range of data,
25

22914

ACPD

11, 22909–22950, 2011

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



be allowed. These pulls are implemented in ME-2 by the generation of a moreparams file, which is generated by EPA PMF when a user specifies a pull, or can be generated as a text file by a user outside of EPA PMF and ME-2. When the moreparams file is present, ME-2 generates a solution where the base solution is pulled as indicated in the moreparams file. Additional details are available in Paatero 2004 and Norris et al. (2009).

2.4 AMS data for EPA PMF application

A matrix of 7455 2-min HR-AMS V-mode observations of 198 unit mass resolution (UMR) fragments made every 4 min during January 2008 at Fyfe was used in EPA PMF analysis. Fragments predominantly from inorganic species such as nitrate and sulfate were not retained for EPA PMF analysis. Fragments m/z 15, 18, 19, 20, 29, and 30 were excluded from EPA PMF analysis because of potential interference with nitrogen and oxygen. While data up to m/z 700 are available, many fragments above m/z 200 had low signal-to-noise (S/N) ratios (i.e. less than 6), made a minimal contribution to total OM, and were collinear with a number of other fragments. Fragments up to m/z 240 were retained for EPA PMF, a total of 198 fragments. Fragments with low S/N may bias the results (Norris et al., 2008), so the uncertainties of a given fragment were multiplied by 3 if the S/N for the fragment was less than 6; this reduces the fragment's influence on the solution. A global 10% uncertainty was also applied to account for additional modeling uncertainty (Norris et al., 2008). EPA PMF was run in the robust mode, which reduces the influence of outliers.

Each observation was also classified as downwind (wind speed greater than 2 m s^{-1} and wind direction between 90 and 270 degrees; $N = 1360$); upwind (wind speed greater than 2 m s^{-1} and wind direction between 310 and 60 degrees; $N = 949$); other (wind speed greater than 2 m s^{-1} and wind direction between 60 and 90 degrees or between 270 and 310 degrees; $N = 461$); or calm (wind speed less than 2 m s^{-1} ; $N = 4907$). As an additional set of runs, EPA PMF was also applied to downwind-only data to examine whether factor profiles change, and how factor contributions change.

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Since downwind-only data are more heavily influenced by the freeway, we may expect that a downwind-only HOA factor would be more similar to vehicle exhaust source profiles than when all data are used.

The AMS PMF results were averaged up to 20-min intervals to allow for matching with the collocated 5-min data, which were also averaged up to 20-min intervals. The 20-min averaged AMS data ($N = 1491$) were also re-analyzed with EPA PMF to evaluate the impact of high-mass transient events; profiles and contributions were then compared to the results using 2-min data. Factor profiles were compared to PMF factor profiles from earlier ambient studies (Lanz et al., 2007, 2008a; Ulbrich et al., 2009a) as well as to source profiles (Mohr et al., 2009; Sage et al., 2008; Weimer et al., 2008) originating from a publicly available online database of reference spectra hosted by University of Colorado (Ulbrich et al., 2009a, b). Specifically, source profiles of Honda gasoline exhaust and diesel exhaust (Mohr et al., 2009), PMF-resolved OOA and HOA factor profiles from Pittsburgh (Zhang et al., 2005a), BBOA factor profiles from Switzerland (Lanz et al., 2007, 2008b), aged diesel exhaust profile from chamber experiments (Sage et al., 2008), and burning and smoldering oak and chestnut wood source profiles (Weimer et al., 2008) were used.

3 Results

3.1 Ambient aerosol variability and composition

OM averaged $3.3 \mu\text{g m}^{-3}$ during the January intensive campaign and was typically highest during the evening hours (i.e. 19:00 through 21:00 LST) with a secondary peak in the morning (i.e. 06:00–09:00 LST, during rush hour commute) as shown in Fig. 2. AMS fragments associated with HOA, such as m/z 57 and m/z 43, showed a similar diurnal pattern. AMS fragments associated with OOA, such as m/z 44 (COO^+), showed only minor fluctuations throughout the day, while those used as tracers of biomass burning, such as m/z 60 (Lanz et al., 2007; Lanz et al., 2008b), were evident only during the evening and overnight hours.

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



environment, predominantly originates from emissions along the roadway. Like BC, OM was higher during stagnant conditions (those with winds less than 0.5 m s^{-1}), as a shallow boundary layer and minimal dispersion quickly lead to the buildup of pollutants.

Temporal patterns for both sulfate and nitrate differed from the temporal pattern for OM. Sulfate concentrations were extremely low throughout the study, with a median concentration of $0.16 \mu\text{g m}^{-3}$, and a maximum 20-min average of $0.88 \mu\text{g m}^{-3}$. Nitrate concentrations were episodic but were also typically low, with a median concentration of $0.54 \mu\text{g m}^{-3}$. In contrast, BC had a median concentration of $1.24 \mu\text{g m}^{-3}$, more than twice as high as nitrate, an order of magnitude higher than sulfate, and a third of OM. BC is more than twice as high at this site compared to a site 2 km away in the urban center but away from freeways (Hancock Elementary School), where BC was on average $0.5 \mu\text{g m}^{-3}$. These sulfate levels are quite low compared to most other areas in the U.S., though typical of the western US, as there is minimal transport of sulfate into Las Vegas and no major sources of SO_2 or sulfate upwind of the urban area. In many prior studies elsewhere, sulfate is higher and is correlated with LV-OOA; however, as the sulfate levels are so low in Las Vegas, it is unlikely that LV-OOA will correlate with sulfate concentrations.

Nitrate had little relationship with OM or other measured pollutants. In a one-day episode, nitrate peaked above $8 \mu\text{g m}^{-3}$ for three hours, after which it decreased to approximately $2 \mu\text{g m}^{-3}$ for the next two days. Nitrate was greater than $1 \mu\text{g m}^{-3}$ on a few other days, but these periods typically lasted only a few hours. These higher concentrations occurred during the day and night, with no distinct, consistent diurnal pattern, unlike OM, BC, CO and other pollutants which peaked in the evening and early morning hours. As Las Vegas is in an arid desert environment with little agriculture in the area, ammonia emissions are relatively low (e.g. <http://nadp.sws.uiuc.edu/nh3net/>), so ammonium nitrate formation may be limited by the availability of ammonia. Nitrate has been observed to often correlate with SV-OOA, as both may be associated with similar transport and formation mechanisms. Since nitrate concentrations did not show any pattern consistent with OM, and were often higher when OM was relatively low,

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

there may be little correlation between SV-OOA and nitrate here.

3.2 EPA PMF application to AMS data summary

Three- to six-factor solutions were explored with EPA PMF. Initially, 50 runs from a random seed were performed for each number of factors. Random starting seeds were used to increase the likelihood of finding a global minimum of the goodness-of-fit parameter, Q . The stability of Q over these runs, the ratio of Q to expected (theoretical) Q , scaled residuals, the Q/Q_{expected} by fragment and sample, and factor independence (G-space plots) were examined. If these parameters are not stable for a given number of factors, it indicates a global minimum

was not consistently achieved, and that a solution may not be stable (Norris et al., 2008, 2009). In all solutions, the total OM was well apportioned (i.e. slope equal to 1.0 ± 0.10) and r^2 was greater than 0.95 between apportioned and total OM.

PMF factors were classified by their temporal pattern and the comparison of their profile with source profiles and profiles from previous studies. The typical AMS PMF factors of HOA, LV-OOA, and BBOA were resolved in every solution with three or more factors, and the identification of each factor profile was based on its similarity to profiles available in the literature, the abundance of key fragments in each profile, and each factor's temporal pattern. For example, the LV-OOA factors displayed a significant amount of m/z 44 and were similar to the OOA factor identified in Pittsburgh and elsewhere. BBOA factors had typical tracer fragments of m/z 60 and 73, which are produced during AMS analysis of levoglucosan and related anhydrosugars produced during biomass combustion (Lanz et al., 2008b; Alfarra et al., 2004; Lee et al., 2010). The HOA factors were similar to the HOA factor from Pittsburgh and a diesel exhaust source profile.

With more than three factors, semi-volatile OOA (SV-OOA) was also resolved. This factor's profile was similar to that of aged diesel exhaust and was more episodic than HOA or LV-OOA profiles. With five and six factors, additional factors that occurred during the nighttime were resolved. The additional nighttime factors occurred nearly

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



every night coincident with BBOA between 17:00 and 02:00 LST, and contributions were higher with low wind speed conditions and with winds from the north (upwind). With peaks of m/z 41, 43, 55, and 91, it is unclear what these “night OA” factors may represent. The change in the BBOA between the four- and five-factor solutions and its temporal variability suggest the additional fifth factor may be related to biomass burning, but the lack of known tracer fragments and correlation with source profiles makes this link difficult to prove. We focused the remainder of the analyses on the four-factor solution. Figure 5 summarizes the amount of OM apportioned by factor. Table 1 summarizes the correlation of factor profiles with selected source, aged source, and PMF factor profiles from other studies.

3.3 Four-factor solution

The factors in the four-factor solution were HOA, LV-OOA and BBOA factors, plus a semi-volatile OOA (SV-OOA) factor. The HOA and LV-OOA factors were better resolved than in the three-factor solution. Profiles of each factor, the average factor concentration plus other species’ concentrations by hour, and a time series of concentrations are provided in Figs. 6 through 8. Figure 9 provides scatter plots of factor contributions with selected collocated measurements: HOA with BC, HOA with CO, SV-OOA with nitrate, and LV-OOA with sulfate. Bootstrapping, in which many runs are used to gauge the uncertainty of the base solution (in this case 300 runs with an r^2 of 0.60), showed good reproducibility of the factors. All factors were reproduced at least 98 % of the time, demonstrating that these factors are stable and characterize the solution space well.

The LV-OOA factor displayed the typical high amount of m/z 44 but with a lower amount of m/z 43 than in the three-factor solution, and it showed a high correlation with the Pittsburgh OOA factor profile (r^2 of 0.99; Table 1). LV-OOA accounted for 26 % of the OM with four factors and showed only small diurnal variability in its concentration, though it was a much greater percentage of the OM during the daytime than at other times. Similar to other factors and total OM, LV-OOA concentrations were lower at

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

higher wind speeds, though it was a higher percentage of OM at higher wind speeds. LV-OOA showed little correlation with other pollutants; ozone, which often showed a moderate correlation with LV-OOA in other studies, was not measured here. Sulfate is also often correlated with LV-OOA, but during this study sulfate levels were extremely low, with a median of $0.16 \mu\text{g m}^{-3}$. LV-OOA is heavily oxidized and likely part of a background OM, and it may be transported into Las Vegas over multiple days. In addition, there are very few SO_2 sources upwind of Las Vegas, so there is very little sulfate transported into the area. Thus, we may not expect LV-OOA concentrations to be correlated with sulfate here.

The BBOA factor accounted for 12% of OM, on average, and was similar to the BBOA factor found in the three-factor solution. This factor had more than 60% of the m/z 60 fragment, which is associated with levoglucosan and related anhydrosugars (Lanz et al., 2008b; Alfara et al., 2004); the contribution from BBOA was well correlated with m/z 60 ($r^2 = 0.86$). This factor is most likely from residential wood burning in the evenings rather than wildfire emissions since there was little regional wildfire activity in the winter; furthermore, the contribution was nearly zero during the daytime, with a sharp rise in concentrations nearly every evening after 17:00 LST that peaked around 21:00 LST. While the BBOA factor concentration began to decrease after around 21:00 LST, its relative contribution to OM remained above 15% until after midnight. The factor was highest in terms of both concentration and relative contribution to OM under low wind speed conditions (i.e. less than 2 m s^{-1}). This profile is similar to a smoldering Chestnut profile ($r^2 = 0.80$) and the levoglucosan combustion profile ($r^2 = 0.80$) (Schneider et al., 2006). This BBOA factor is mostly associated with winds from the north and west – the direction of a large residential neighborhood upwind of the freeway. When using five factors, the BBOA factor profile has even better correlations with both of these profiles (r^2 of 0.89 and 0.91), plus an r^2 of 0.95 with an oak flaming profile. However, since the fifth factor is unidentified, we have focused on the four-factor solution.

of concentrations greater than $4 \mu\text{g m}^{-3}$ and a few hours during which concentrations intermittently exceeded $2 \mu\text{g m}^{-3}$. In contrast, SV-OOA was present during nearly the entire study, and had a modest diurnal pattern similar to other factors where it was highest in evening hours.

To help confirm factor identification and understand the OM composition difference between upwind and downwind conditions, we examined the factor contributions during downwind ($N = 1360$) and upwind ($N = 949$) conditions and compared them to the average over the study period. We also examined the OM composition under downwind conditions ($N = 195$) from 05:00 to 09:00 LST, when the impact from emissions on the freeway is expected to be highest. Results are summarized in Fig. 10. As expected, HOA contributions are higher under downwind conditions; during morning downwind conditions, HOA accounts for 49 % of the OM. SV-OOA was on average similar during upwind, downwind and stagnant conditions, except during morning downwind situations when it was only 23 % of the OM. The minimal difference with different wind directions suggests that SV-OOA is a slightly aged factor that is not characteristic of direct, primary emissions. BBOA contributions were low on average under downwind conditions and higher (16 % on average) during upwind conditions. LV-OOA contributions were relatively lower during morning downwind conditions and relatively highest during midday periods, regardless of wind direction.

3.4 Further analysis using ME-2 rotational tools in EPA PMF

F_{peak} is a parameter available in PMF to rotate the entire solution, a process that can help indicate if there is rotational freedom in the solution. G-space plots of the base solution show distinct edges, indicating some factor interdependence. To ascertain whether the solution changes or whether these edges can be rotated to the y- and x-axes, rotation using F_{peak} was performed. PMF runs were conducted with F_{peak} values at increments of 2 between 8 and -8 for a total of eight runs. In general, minimal change was seen in the factor profile, contributions, and G-space plots; Q increased by

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

less than 0.1 % for the runs with the highest F_{peak} value and by even less under other F_{peak} values. Since there is little change in the solution with F_{peak} -induced rotations, the base solution appears to be rotationally unique. The oblique edges in the G-space plots may be due to co-dependence among factors, or modeling errors, such as variation in true source profiles during the monitoring campaign.

The four-factor solution was further explored with ME-2 rotational tools available in EPA PMF. In each scenario, fragment ratios in factor profiles were pulled toward source profile ratios. In one scenario, the ratio of m/z 43/44 in SV-OOA factor (6.75) was pulled toward the m/z 43/44 ratio in the 5-h aged diesel exhaust profile (1.34). In another scenario, the m/z 57/55 and 41/43 ratios in HOA (0.78 and 0.80) were pulled to the ratios in the diesel exhaust profile (1.03 and 0.69, respectively). The BBOA factor ratio of m/z 60/91 was pulled to the chestnut smolder profile ratio (1.96 to 7.20). Lastly, 157 points along an apparent edge in the SV-OOA versus LV-OOA G-space plot were pulled in an attempt to force additional independence between these factors. Seeing how the solutions change, in terms of factor profiles, dQ , and G-space plots, can help us understand the stability of the original solutions. Different maximum dQ values were allowed for each combination; 1 % and 3 % of $Q(\text{robust})$ were calculated, and these values were used for the maximum dQ allowed for a given pull. In the G-space pull, each point was allowed a dQ of 0.2 %, for a total dQ of 13 %.

The results from these pulls are summarized in Table 2 and Fig. 11. Since results were similar with dQ values of 1 % and 3 %, only the dQ of 1 % are shown, except for the BBOA pull, where both are provided as an example. In general, most pulls resulted in only minor changes in the HOA and LV-OOA factor profiles and contributions. For example, when the HOA factor was pulled toward the diesel exhaust profile, the correlation between the two increased by only 0.02. In all pulls, the correlation between BBOA and the chestnut smolder profile improved (e.g. correlation improved from 0.79 to 0.88 with the BBOA pull). However this was often at the expense of the SV-OOA profile, where the amount of m/z 44 was much lower compared to the base solution, and in some pulls, was actually zero. While LV-OOA and HOA did not

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

5 vary much between these pulled solutions, the changes in SV-OOA and BBOA profiles and contributions suggest some rotational freedom in these two factors. LV-OOA and HOA factors are similar across many studies, but the SV-OOA and BBOA factors vary among studies, and within this study under different pulling scenarios, because these factor profiles represent semi-fresh factors that are likely changing minute-to-minute in the atmosphere. Overall, the pulling results indicate that the base solution is likely at a global Q minima, and that there is little rotational freedom in the unpulled, base solution, in particular regarding the contributions of SV-OOA and BBOA.

10 With the G-space pull, 157 points along an edge in the LV-OOA/SV-OOA scatter plot were pulled to reduce LV-OOA to zero. LV-OOA concentrations on these points were reduced towards zero, so that the edge was less well defined. This resulted in a dQ of 9%, but an improvement in the comparison of the BBOA profile to the chestnut smolder profile (r^2 from 0.79 to 0.93) and in the SV-OOA to aged diesel profile comparison (r^2 from 0.69 to 0.72). The SV-OOA m/z 43/44 ratio also decreased from 4.4 to 2.9, which is similar to the ratio of other studies summarized in Ng, et al. (2010). HOA and BBOA contributions increased, while LV-OOA and SV-OOA contributions decreased. Even though Q increased by 9%, these results are useful to show that contribution pulls based on the G-space plots helped improve the factor profiles, and to show what may be the bounds in the base solution results.

20 4 Discussion

Evaluating solutions with different numbers of factors, comparing profiles to source profiles, examining temporal trends, and exploring rotational ambiguity with the rotational tools available in ME-2 can lead to a greater understanding of the AMS data set. In general, all the factors were consistent under multiple scenarios, suggesting high confidence in their apportionment. The addition of factors after four factors helped to better characterize the solution space, but the additional “night OA” factors are not easily attributable to known sources. While G-space plots suggested there was rotational

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ambiguity in all solutions, we were unable to rotate the solutions to reduce this ambiguity, and pulling edge points to axes resulted in an increase in Q but minimal change in the factors.

The Q/Q_{exp} ratios for most fragments were around unity (i.e. between 0.8 and 1.3), indicating that the obtained Q values were approximately equal to the expected values (see Fig. 1 in Supplement). Some fragments had Q/Q_{exp} ratios below 0.5, indicating that computed Q values were significantly smaller than the expected Q values. This discrepancy is most likely due to the global uncertainty (s_{ij} increase of 10%); since the Q/Q_{exp} ratios for some m/z were fairly low, the 10% value may be too high for these m/z but appropriate for many other fragments. The Q/Q_{exp} ratios were between 1.3 and 2 for several m/z , indicating that the average residuals are between 13 and 20% of x_{ij} for these m/z . The Q contributions drop sharply beginning at $m/z = 198$, as most of these fragments had low S/N ratios and were downweighted. The large residuals may indicate that the PMF solution does not fully characterize these fragments¹. Inaccurate subtraction of the inorganic component from x_{ij} before PMF analysis is a possibility, though many of the fragments with high Q/Q_{exp} do not have an inorganic component. More likely, the large residuals are due to the variation of factor profiles with time and/or to the presence of an occasional, spurious, or localized source(s).

While factors in the four-factor solution are similar to those observed elsewhere, the apportionment of mass among them is different from that seen in other studies. This difference is expected because a number of the previous studies occurred in the summer and/or in environments with a higher amount of oxidized aerosol than the present study. Previous studies in Los Angeles, Pittsburgh, rural British Columbia, and elsewhere have typically found that at least one-third of the OM was attributable to LV-OOA, originally termed OOA I (Allan et al., 2003a; Alfarra et al., 2004; Dzepina et al., 2007; Sun et al., 2009). The lower amount of LV-OOA observed during the wintertime Las Vegas study could be due to less transported/aged aerosol, lower biogenic emissions,

¹Fragments with high Q/Q_{exp} include $m/z = [44, 60, 73, 85, 86, 111, 112, 113, 114, 123, 124, 125, 126, 137, 138, 140, 141, 154, 155, 156]$.

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and/or less overall atmospheric oxidation compared to summertime. In Zurich, a wintertime study found 52 % to 57 % of the OM to be LV-OOA (Lanz et al., 2008a), 69 % of which originated from non-fossil sources such as wood burning. In Las Vegas, there is a much lower concentration of BBOA than in Zurich, leading to a smaller concentration of LV-OOA from non-fossil sources and a smaller concentration of LV-OOA overall.

5 Conclusions

EPA PMF v4.0, with its new rotational tools, was successfully applied to a near-road, high time resolution AMS data set. HOA was a quarter of the OM (24 %), and higher under downwind conditions (about 40 %). In addition to this local, primary OA, there was a highly oxidized background of OA (LV-OOA) that on average constituted 29 % of the OM, and a less oxidized, semi-volatile fraction that accounted for 34 % of the OM. During the evening hours, biomass burning (BBOA) was also seen, likely from the surrounding residential area. Rotational tools allowed for additional analysis of the PMF solution space, increasing our confidence in the results.

Supplementary material related to this article is available online at:
<http://www.atmos-chem-phys-discuss.net/11/22909/2011/acpd-11-22909-2011-supplement.pdf>.

Acknowledgements. The authors thank the Nevada Department of Transportation, US EPA Office of Research and Development, and Sonoma Technology, Inc., for supporting this work. We also thank Jennifer DeWinter (STI) and Ram Vedantham (EPA ORD) for data analysis support and analysis of the collocated data at Fyfe Elementary, Mary Jo Teplitz (STI) for editorial support, and Bryan Penfold (STI) for the site map. We greatly appreciate the development of the data processing and analysis methods done by the AMS community, particularly by Jose Jimenez and Ingrid Ulbrich (University of Colorado). The acquisition of the CSU Aerosol Mass Spectrometer (AMS) was supported by the NSF-MRI program under grant number ATM-0521643. The AMS was operated in the CSU Mobile Laboratory, which is supported by the National Park Service (NPS).

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Disclaimer

The United States Environmental Protection Agency through its Office of Research and Development partially funded the research described here under PO 720002129 to Sonoma Technology, Inc. through Lockheed Martin. It has been subjected to Agency review and approved for publication.

References

Adonis, M. I., Riquelme, R. M., Martinez, V. D., Gil, L., Rios, C., Rodriguez, L., and Rodriguez, E. M.: PAHs and mutagenicity of inhalable and respirable diesel particulate matter in Santiago, Chile, *Polycyc. Aromatic Compounds* 23, 495–514, 2003.

Aiken, A. C., DeCarlo, P., Kroll, J. H., Worsnop, D. R., Huffman, J. A., Docherty, K. S., Ulbrich, I. M., Mohr, C., Kimmel, J. R., Sueper, D., Sun, Y., Zhang, Q., Trimborn, A., Northway, M., Ziemann, P. J., Canagaratna, M. R., Onasch, T. B., Alfarra, M. R., Prevot, A. S. H., Dommen, J., Duplissy, J., Metzger, A., Baltensperger, U., and Jimenez, J. L.: O/C and OM/OC ratios of primary, secondary, and ambient organic aerosols with high-resolution time-of-flight aerosol mass spectrometry, *Environ. Sci. Technol.*, 42(12), 4478–4485, doi:10.1021/es703009q, 2008.

Alfarra, M. R., Coe, H., Allan, J. D., Bower, K. N., Boudries, H., Canagaratna, M. R., Jimenez, J. L., Jayne, J. T., Garforth, A. A., Li, S.-M., and Worsnop D. R.: Characterization of urban and rural organic particulate in the Lower Fraser Valley using two Aerodyne aerosol mass spectrometers, *Atmos. Environ.*, 38, 5745–5758, doi:10.1016/j.atmosenv.2004.01.054, 2004.

Alfarra, M. R., Paulsen, D., Gysel, M., Garforth, A. A., Dommen, J., Prévôt, A. S. H., Worsnop, D. R., Baltensperger, U., and Coe, H.: A mass spectrometric study of secondary organic aerosols formed from the photooxidation of anthropogenic and biogenic precursors in a reaction chamber, *Atmos. Chem. Phys.*, 6, 5279–5293, doi:10.5194/acp-6-5279-2006, 2006.

Alfarra, M. R., Prevot, A. S. H., Szidat, S., Sandradewi, J., Weimer, S., Lanz, V. A., Schreiber, D., Mohr, M., and Baltensperger, U.: Identification of the mass spectral signature of organic aerosols from wood burning emissions, *Environ. Sci. Technol.*, 41, 5770–5777, doi:10.1021/es062289b, 2007.

Allan, J. D., Alfarra, M. R., Bower, K. N., Williams, P. I., Gallagher, M. W., Jimenez, J. L.,

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

McDonald, A. G., Nemitz, E., Canagaratna, M. R., Jayne, J. T., Coe, H., and Worsnop, D. R.: Quantitative sampling using an Aerodyne aerosol mass spectrometer – 2. Measurements of fine particulate chemical composition in two UK cities, *J. Geophys. Res.-Atmos.*, 108(D3), 4091, doi:10.1029/2002JD002359, 2003a.

5 Allan, J. D., Jimenez, J. L., Williams, P. I., Alfarra, M. R., Bower, K. N., Jayne, J. T., Coe, H., and Worsnop, D. R.: Quantitative sampling using an Aerodyne aerosol mass spectrometer – 1. Techniques of data interpretation and error analysis, *J. Geophys. Res.-Atmos.*, 108(D3), 4090, doi:10.1029/2002JD002358, 2003b.

Allan, J. D., Delia, A. E., Coe, H., Bower, K. N., Alfarra, M. R., Jimenez, J. L., Middlebrook, A. M., Drewnick, F., Onasch, T. B., Canagaratna, M. R., Jayne, J. T., and Worsnop, D. R.: A generalised method for the extraction of chemically resolved mass spectra from Aerodyne aerosol mass spectrometer data, *J. Aerosol Sci.*, 35(7), 909–922, 2004.

Brown, S. G. and Hafner, H. R.: Source apportionment of VOCs in the Houston, Texas, area. Presented at the NARSTO Workshop on Innovative Methods for Emission-Inventory Development and Evaluation, Austin, TX, 14–16 October (STI-2356), 2003.

15 Brown, S. G., DeWinter, J. L., McCarthy, M. C., Vaughn, D. L., and Roberts, P. T.: Contributions of mobile sources to near-roadway concentrations of air toxics in Las Vegas, Nevada, (STI-3889), in preparation, 2011.

Brunekreef, B., Janssen, N. A., de Hartog, J., Harssema, H., Knape, M., and van Vliet, P.: Air pollution from truck traffic and lung function in children living near motorways, *Epidemiology*, 8(3), 298–303, 1997.

20 Canagaratna, M. R., Jayne, J. T., Ghertner, D. A., Herndon, S., Shi, Q., Jimenez, J. L., Silva, P. J., Williams, P., Lanni, T., Drewnick, F., Demerjian, K. L., Kolb, C. E., and Worsnop, D. R.: Chase studies of particulate emissions from in-use New York City vehicles, *Aerosol Sci. Technol.*, 38(6), 555–573, 2004.

25 Canagaratna, M. R., Jayne, J. T., Jimenez, J. L., Allan, J. D., Alfarra, M. R., Zhang, Q., Onasch, T. B., Drewnick, F., Coe, H., Middlebrook, A., Delia, A., Williams, L. R., Trimborn, A. M., Northway, M. J., DeCarlo, P. F., Kolb, C. E., Davidovits, P., and Worsnop, D. R.: Chemical and microphysical characterization of ambient aerosols with the aerodyne aerosol mass spectrometer, *Mass Spectrom. Rev.*, 26(2), 185–222, doi:10.1002/mas.20115, 2007.

30 DeCarlo, P., Kimmel, J. R., Trimborn, A., Northway, M., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T., Docherty, K. S., Worsnop, D. R., and Jimenez, J. L.: Field-deployable, high-resolution, time-of-flight aerosol mass spectrometer, *Anal. Chem. (Wash.)*,

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

78(24), 8281–8289, doi:10.1021/ac061249n, 2006.

Docherty, K. S., Stone, E. A., Ulbrich, I. M., DeCarlo, P., Snyder, D. C., Schauer, J. J., Peltier, R. E., Weber, R. J., Murphy, S. M., Seinfeld, J. H., Grover, B. D., Eatough, D. J., and Jimenez, J. L.: Apportionment of primary and secondary organic aerosols in Southern California during the 2005 Study of Organic Aerosols in Riverside (SOAR-1), *Environ. Sci. Technol.*, 42(20), 7655–7662, doi:10.1021/es8008166, 2008.

Dockery, D. W. and Stone, P. H.: Cardiovascular risks from fine particulate air pollution, *New Engl. J. Med.*, 356, 511–513, 2007.

Dockery, D. W., Pope, C. A., Xu, X. P., Spengler, J. D., Ware, J. H., Fay, M. E., Ferris, B. G., and Speizer, F. E.: An association between air pollution and mortality in six U.S. cities, *New Engl. J. Med.*, 329, 1753–1759, 1993.

Drewnick, F., Hings, S. S., DeCarlo, P., Jayne, J. T., Gonin, M., Fuhrer, K., Weimer, S., Jimenez, J. L., Demerjian, K. L., Borrmann, S., and Worsnop, D. R.: A new time-of-flight aerosol mass spectrometer (TOF-AMS) – instrument description and first field deployment, *Adv. Sci. Technol.* 39, 637–658, doi:10.1080/02786820500182040, 2005.

Dzepina, K., Arey, J., Marr, L. C., Worsnop, D. R., Salcedo, D., Zhang, Q., Onasch, T. B., Molina, L. T., Molina, M. J., and Jimenez, J. L.: Detection of particle-phase polycyclic aromatic hydrocarbons in Mexico City using an aerosol mass spectrometer, *Int. J. Mass Spectrom.*, 263, 152–170, doi:10.1016/j.ijms.2007.01.010, 2007.

Edwards, J., Walters, S., and Griffiths, R. K.: Hospital admissions for asthma in preschool children: relationship to major roads in Birmingham, United Kingdom, *Arch. Environ. Health*, 49, 223–227, 1994.

Finkelstein, M. M., Jerrett, M., and Sears, M. R.: Traffic air pollution and mortality rate advancement periods, *Am. J. Epidemiol.*, 160(2), 173–177, 2004.

Flowers, L., Rieth, S. H., Coglianò, V. J., Foureman, G. L., Hertzberg, R., Hofmann, E. L., Murphy, D. L., Nesnow, S., and Schoeny, R. S.: Health assessment of polycyclic aromatic hydrocarbon mixtures: current practices and future directions, *Polycyc. Aromatic Compounds*, 22, 811–821, 2002.

Fruin, S., Westerdahl, D., Sax, T., Sioutas, C., and Fine, P. M.: Measurements and predictors of on-road ultrafine particle concentrations and associated pollutants in Los Angeles, *Atmos. Environ.*, 42, 207–219, doi:10.1016/j.atmosenv.2007.09.057, 2008.

Hoek, G., Brunekreef, B., Goldbohm, S., Fischer, P., and van den Brandt, P. A.: Association between mortality and indicators of traffic-related air pollution in the Netherlands: a cohort

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

study, *Lancet*, 360, 1203–1209, 2002.

Hopke, P. K.: A guide to positive matrix factorization. Prepared for the Positive Matrix Factorization Program, Potsdam, NY, by the Department of Chemistry, Clarkson University, Potsdam, NY, 2003.

5 Hu, S., Fruin, S., Kozawa, K., Mara, S., Paulson, S. E., and Winer, A. M.: A wide area of air pollutant impact downwind of a freeway during pre-sunrise hours, *Atmos. Environ.*, 43, 2541–2549, doi:10.1016/j.atmosenv.2009.02.033, 2009.

Huffman, J. A., Docherty, K. S., Mohr, C., Cubison, M., Ulbrich, I. M., Ziemann, P. J., Onasch, T. B., and Jimenez, J. L.: Chemically-resolved volatility measurements of organic aerosol from
10 different sources, *Environ. Sci. Technol.*, 43, 5351–5357, doi:10.1021/es803539d, 2009.

Jayne, J. T., Leard, D. C., Zhang, X. F., Davidovits, P., Smith, K. A., Kolb, C. E., and Worsnop, D. R.: Development of an aerosol mass spectrometer for size and composition analysis of submicron particles, *Adv. Sci. Technol.*, 33(1–2), 49–70, 2000.

15 Jimenez, J. L., Jayne, J. T., Shi, Q., Kolb, C. E., Worsnop, D. R., Yourshaw, I., Seinfeld, J. H., Flagan, R. C., Zhang, X. F., Smith, K. A., Morris, J. W., and Davidovits, P.: Ambient aerosol sampling using the Aerodyne Aerosol Mass Spectrometer, *J. Geophys. Res.-Atmos.*, 108(D7), 8425, doi:10.1029/2001JD001213, 2003.

20 Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dunlea, E. J., Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A.,
25 Miyoshi, T., Hatakeyama, S., Shimojo, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of organic aerosols in the atmosphere, *Science*, 326, 1525–1529, doi:10.1126/science.1180353, 2009.

30 Johnson, K. S., Laskin, A., Jimenez, J. L., Shutthanandan, V., Molina, L. T., Salcedo, D., Dzepina, K., and Molina, M. J.: Comparative analysis of urban atmospheric aerosol by Particle-Induced X-ray Emission (PIXE), Proton Elastic Scattering Analysis (PESA), and Aerosol Mass Spectrometry (AMS), *Environ. Sci. Technol.*, 42(17), 6619–6624, doi:10.1021/es800393e, 2008.

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Kim, E. and Hopke, P. K.: Improving source identification of fine particles in a rural northeastern U.S. area utilizing temperature-resolved carbon fractions, *J. Geophys. Res.*, 109(D9), D09204, doi:09210.01029/02003JD004199, 2004.

Kim, E., Hopke, P. K., and Edgerton, E. S.: Source identification of Atlanta aerosol by positive matrix factorization, *J. Air Waste Manag. Assoc.*, 53, 731–739, 2003.

Kim, E., Hopke, P. K., and Edgerton, E. S.: Improving source identification of Atlanta aerosol using temperature resolved carbon fractions in positive matrix factorization, *Atmos. Environ.*, 38, 3349–3362, 2004a.

Kim, J. J., Smorodinsky, S., Lipsett, M., Singer, B. C., Hodgson, A. T., and Ostro, B.: Traffic-related air pollution near busy roads: the East Bay Children's Respiratory Health Study, *Am. J. Respir. Crit. Care Med.*, 170(5), 520–526, 2004b.

Kunzli, N., Kaiser, R., Medina, S., Studnicka, M., Chanel, O., and Filliger, P.: Public-health impact of outdoor and traffic-related air pollution: a European assessment, *Lancet*, 356, 795–801, 2000.

Lanz, V. A., Alfarra, M. R., Baltensperger, U., Buchmann, B., Hueglin, C., and Prévôt, A. S. H.: Source apportionment of submicron organic aerosols at an urban site by factor analytical modelling of aerosol mass spectra, *Atmos. Chem. Phys.*, 7, 1503–1522, doi:10.5194/acp-7-1503-2007, 2007.

Lanz, V. A., Alfarra, M. R., Baltensperger, U., Buchmann, B., Hueglin, C., Szidat, S., Wehrli, M. N., Wacker, L., Weimer, S., Caseiro, A., Puxbaum, H., and Prevot, A. S. H.: Source attribution of submicron organic aerosols during wintertime inversions by advanced factor analysis of aerosol mass spectra, *Environ. Sci. Technol.*, 42(1), 214–220, doi:10.1021/es0707207, 2008a.

Lanz, V. A., Alfarra, M. R., Baltensperger, U., Buchmann, B., Hueglin, C., Szidat, S., Wehrli, M. N., Wacker, L., Weimer, S., Caseiro, A., Puxbaum, H., and Prevot, A. S. H.: Source attribution of submicron organic aerosols during wintertime inversions by advanced factor analysis of aerosol mass spectra, *Environ. Sci. Technol.*, 42, 214–220, doi:10.1021/es0707207, 2008b.

Lanz, V. A., Prévôt, A. S. H., Alfarra, M. R., Weimer, S., Mohr, C., DeCarlo, P. F., Gianini, M. F. D., Hueglin, C., Schneider, J., Favez, O., D'Anna, B., George, C., and Baltensperger, U.: Characterization of aerosol chemical composition with aerosol mass spectrometry in Central Europe: an overview, *Atmos. Chem. Phys.*, 10, 10453–10471, doi:10.5194/acp-10-10453-2010, 2010.

Larsen, R. K. and Baker, J. E.: Source apportionment of polycyclic aromatic hydrocarbons in

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)




[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

the urban atmosphere: a comparison of three methods, *Environ. Sci. Technol.*, 37(9), 1873–1881, 2003.

Lee, T., Sullivan, A. P., Mack, L., Jimenez, J. L., Kreidenweis, S. M., Onasch, T. B., Worsnop, D. R., Malm, W., Wold, C. E., Hao, W. M., and Jeffrey, L. C. J.: Chemical smoke marker emissions during flaming and smoldering phases of laboratory open burning of wildland fuels, *Adv. Sci. Technol.*, 44(9), i–v, doi:10.1080/02786826.2010.499884, 2010.

Lobscheid, A. B. and McKone, T. E.: Constraining uncertainties about the sources and magnitude of polycyclic aromatic hydrocarbon (PAH) levels in ambient air: the state of Minnesota as a case study, *Atmos. Environ.*, 38, 5501–5515, 2004.

McCarthy, M. C., Brown, S. G., Landreneau, J. V., Vaughn, D. L., Roberts, P. T., and Ludwig, J. F.: Filtration efficiency of HVAC systems at near-roadway schools in Las Vegas, Nevada, (STI-906034-3629), in preparation, 2011.

McDonald, J. D., Harrod, K. S., Seagrave, J., Seilkop, S. K., and Mauderly, J. L.: Effects of low sulfur fuel and a catalyzed particle trap on the composition and toxicity of diesel emissions, *Environ. Health Perspect.*, 112(13), 1307–1312, 2004.

Minguillon, M. C., Arhami, M., Schauer, J. J., and Sioutas, C.: Seasonal and spatial variations of sources of fine and quasi-ultrafine particulate matter in neighborhoods near the Los Angeles-Long Beach harbor, *Atmos. Environ.*, 42, 7317–7328, 2008.

Mohr, C., Huffman, J. A., Cubison, M., Aiken, A. C., Docherty, K. S., Kimmel, J. R., Ulbrich, I. M., Hannigan, M., and Jimenez, J. L.: Characterization of primary organic aerosol emissions from meat cooking, trash burning, and motor vehicles with high-resolution aerosol mass spectrometry and comparison with ambient and chamber observations, *Environ. Sci. Technol.*, 43(7), 2443–2449, doi:10.1021/es8011518, 2009.

Ng, N. L., Canagaratna, M. R., Zhang, Q., Jimenez, J. L., Tian, J., Ulbrich, I. M., Kroll, J. H., Docherty, K. S., Chhabra, P. S., Bahreini, R., Murphy, S. M., Seinfeld, J. H., Hildebrandt, L., Donahue, N. M., DeCarlo, P. F., Lanz, V. A., Prévôt, A. S. H., Dinar, E., Rudich, Y., and Worsnop, D. R.: Organic aerosol components observed in Northern Hemispheric datasets from Aerosol Mass Spectrometry, *Atmos. Chem. Phys.*, 10, 4625–4641, doi:10.5194/acp-10-4625-2010, 2010.

Nitta, H., Sato, T., Nakai, S., Maeda, K., Aoki, S., and Ono, M.: Respiratory health associated with exposure to automobile exhaust. 1. Results of cross-sectional studies in 1979, 1982, and 1983, *Arch. Environ. Health*, 48(1), 53–58, 1993.

Norris, G., Vedantham, R., Wade, K. S., Brown, S. G., Prouty, J. D., and Foley, C.: EPA positive

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

matrix factorization (PMF) 3.0 fundamentals and user guide. Prepared for the U.S. Environmental Protection Agency, Washington, DC, by the National Exposure Research Laboratory, Research Triangle Park; Sonoma Technology, Inc., Petaluma, CA; and Lockheed Martin Systems Engineering Center, Arlington, VA, EP-D-05-004; STI-907045.05-3347-UG, October, 2008.

Norris, G., Vedantham, R., Wade, K., Zahn, P., Brown, S., Paatero, P., Eberly, S., and Foley, C.: Guidance document for PMF applications with the Multilinear Engine. Prepared for the U.S. Environmental Protection Agency, Research Triangle Park, NC, by the National Exposure Research Laboratory, Research Triangle Park, NC; Sonoma Technology, Inc., Petaluma, CA; University of Helsinki, Helsinki, Finland; Geometric Tools, Inc., Phoenix, AZ; and Lockheed Martin, Systems Engineering Center, Arlington, VA, EPA 600/R-09/032, April, 2009.

Ntziachristos, L., Ning, Z., Geller, M. D., and Sioutas, C.: Particle concentration and characteristics near a major freeway with heavy-duty diesel traffic, *Environ. Sci. Technol.*, 41(7), 2223–2230, doi:10.1021/es062590s, 2007.

Paatero, P.: Least squares formulation of robust non-negative factor analysis, *Chemometrics Intellig. Lab. Syst.*, 37, 23–35, 1997.

Paatero, P.: User's guide for the multilinear engine program "ME2" for fitting multilinear and quasi-multilinear models, February, 2000.

Paatero, P.: End user's guide to multilinear engine applications, August, 2004.

Paatero, P. and Hopke, P.: Rotational tools for factor analytic models, *Journal of Chemometrics*, 23, 91–100, 2009.

Paatero P. and Tapper U.: Analysis of different modes of factor analysis as least squares fit problems, *Chemometrics Intellig. Lab. Syst.*, 18, 183–194, 1993.

Paatero, P. and Tapper, U.: Positive matrix factorization: a non-negative factor model with optimal utilization of error estimates of data values, *Environmetrics* 5, 111–126, 1994.

Phuleria, H. C., Sheesley, R. J., Schauer, J. J., Fine, P. M., and Sioutas, C.: Road-side measurements of size-segregated particulate organic compounds near gasoline and diesel-dominated freeways in Los Angeles, CA, *Atmos. Environ.*, 41, 4653–4671, doi:10.1016/j.atmosenv.2007.03.031, 2007.

Poirot, R. L., Wishinski, P. R., Hopke, P. K., and Polissar, A. V.: Comparative application of multiple receptor methods to identify aerosol sources in northern Vermont, *Environ. Sci. Technol.*, 35(23), 4622–4636, 2001.

Polissar, A. V., Hopke, P. K., and Poirot, R. L.: Atmospheric aerosol over Vermont: chemical

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- composition and sources, *Environ. Sci. Technol.*, 35(23), 4604–4621, 2001.
- Riddle, S. G., Robert, M. A., Jakober, C. A., Hannigan, M. P., and Kleeman, M. J.: Size-resolved source apportionment of airborne particle mass in a roadside environment, *Environ. Sci. Technol.*, 42(17), 6580–6586, doi:10.1021/es702827h, 2008.
- 5 Sage, A. M., Weitkamp, E. A., Robinson, A. L., and Donahue, N. M.: Evolving mass spectra of the oxidized component of organic aerosol: results from aerosol mass spectrometer analyses of aged diesel emissions, *Atmos. Chem. Phys.*, 8, 1139–1152, doi:10.5194/acp-8-1139-2008, 2008.
- Schneider, J., Weimer, S., Drewnick, F., Borrmann, S., Helas, G., Gwaze, P., Schmid, O.,
10 Andreae, M. O., and Kirchner, U.: Mass spectrometric analysis and aerodynamic properties of various types of combustion-related aerosol particles, *Int. J. Mass Spec.*, 258(1–3), 37–49, doi:10.1016/j.ijms.2006.07.008, 2006.
- Sun, Y., Zhang, Q., Macdonald, A. M., Hayden, K., Li, S. M., Liggio, J., Liu, P. S. K., Anlauf, K. G., Leaitch, W. R., Steffen, A., Cubison, M., Worsnop, D. R., van Donkelaar, A., and
15 Martin, R. V.: Size-resolved aerosol chemistry on Whistler Mountain, Canada with a high-resolution aerosol mass spectrometer during INTEX-B, *Atmos. Chem. Phys.*, 9, 3095–3111, doi:10.5194/acp-9-3095-2009, 2009.
- Ulbrich, I. M., Canagaratna, M. R., Zhang, Q., Worsnop, D. R., and Jimenez, J. L.: Interpretation of organic components from Positive Matrix Factorization of aerosol mass spectrometric data, *Atmos. Chem. Phys.*, 9, 2891–2918, doi:10.5194/acp-9-2891-2009, 2009a.
- 20 Ulbrich, I. M., Lechner, M., and Jimenez, J. L.: AMS spectral database (unit mass resolution). Jimenez Group, University of Colorado, Boulder, CO, available at: <http://cires.colorado.edu/jimenez-group/AMSSd/>, 2009b.
- Weimer, S., Alfarra, M. R., Schreiber, D., Mohr, M., Prevot, A. S. H., and Baltensperger, U.: Organic aerosol mass spectral signatures from wood-burning emissions: influence of burning conditions and wood type, *J. Geophys. Res.-Atmos.*, 113, D10304, doi:10.1029/2007JD009309, 2008.
- 25 Zhang, K. M., Wexler, A. S., Zhu, Y. F., Hinds, W. C., and Sioutas, C.: Evolution of particle number distribution near roadways. Part II: The “road-to-ambient” process, *Atmos. Environ.*, 38, 6655–6665, 2004.
- 30 Zhang, Q., Stanier, C. O., Canagaratna, M. R., Jayne, J. T., Worsnop, D. R., Pandis, S. N., and Jimenez, J. L.: Insights into the chemistry of new particle formation and growth events in Pittsburgh based on aerosol mass spectrometry, *Environ. Sci. Technol.*, 38(18), 4797–4809,

2004.

Zhang, Q., Alfarra, M. R., Worsnop, D. R., Allan, J. D., Coe, H., Canagaratna, M. R., and Jimenez, J. L.: Deconvolution and quantification of hydrocarbon-like and oxygenated organic aerosols based on aerosol mass spectrometry, *Environ. Sci. Technol.*, 39, 4938–4952, doi:10.1021/es048568l, 2005a.

Zhang, Q., Worsnop, D. R., Canagaratna, M. R., and Jimenez, J. L.: Hydrocarbon-like and oxygenated organic aerosols in Pittsburgh: insights into sources and processes of organic aerosols, *Atmos. Chem. Phys.*, 5, 3289–3311, doi:10.5194/acp-5-3289-2005, 2005.

Zhu, Y. F., Hinds, W. C., Kim, S., and Sioutas, C.: Concentration and size distribution of ultrafine particles near a major highway, *J. Air Waste Manage. Assoc.*, 52(9), 1032–1042, 2002.

ACPD

11, 22909–22950, 2011

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

Table 1. Correlation (r^2) of PMF factor profiles with Pittsburgh (Pittsburgh profiles from Zhang et al., 2005a) OOA PMF profile, Pittsburgh HOA PMF profile, diesel exhaust source profile (Mohr et al., 2009), aged diesel exhaust profile (Sage et al., 2008), charbroil (Lanz et al., 2007) source profile, oak-flame (Wood burning profiles from Wiemer et al., 2008) source profile, oak-smolder source profile, chestnut-flame source profile, chestnut-smolder source profile, and levoglucosan (Schneider et al., 2006) profile. Correlations from 0.80 to 0.90 are denoted in italics, and those greater than 0.90 are denoted in bold.

N	Factor	Pitt HOA	gasoline	Diesel	Pitt OOA	Aged diesel	charbroil	Oak smolder	Oak flame	levo-glucosan	Chestnut flame	Chestnut smolder
3	LV-OOA	0.59	0.58	0.58	0.98	0.95	0.26	<i>0.88</i>	0.91	0.74	<i>0.83</i>	<i>0.89</i>
	HOA	0.97	0.92	0.95	0.46	0.75	0.57	0.29	0.71	0.49	0.50	0.62
4	BBOA	<i>0.80</i>	0.73	0.76	0.67	<i>0.82</i>	0.40	0.44	<i>0.89</i>	0.77	0.77	<i>0.85</i>
	LV-OOA	0.38	0.39	0.38	0.99	<i>0.87</i>	0.13	0.93	<i>0.83</i>	0.70	0.80	<i>0.84</i>
	HOA	0.99	0.96	0.98	0.48	0.77	0.60	0.32	0.69	0.48	0.50	0.62
	BBOA	0.47	0.41	0.44	0.59	0.64	0.20	0.39	0.75	<i>0.80</i>	0.77	<i>0.80</i>
5	SV-OOA	0.91	<i>0.84</i>	<i>0.87</i>	0.57	0.79	0.49	0.37	0.80	0.58	0.60	0.71
	LV-OOA	0.46	0.46	0.46	0.99	0.90	0.18	0.91	<i>0.88</i>	0.72	<i>0.82</i>	<i>0.87</i>
	HOA	0.99	0.97	0.98	0.47	0.78	0.61	0.32	0.68	0.49	0.50	0.62
	BBOA	0.71	0.69	0.69	<i>0.86</i>	0.91	0.37	0.69	0.95	<i>0.89</i>	0.91	0.97
	SV-OOA	<i>0.90</i>	<i>0.82</i>	<i>0.85</i>	0.56	0.78	0.47	0.37	0.79	0.56	0.58	0.69
6	Night OA I	0.60	0.51	0.59	0.42	0.61	0.27	0.26	0.57	0.30	0.35	0.45
	LV-OOA	0.13	0.16	0.15	0.59	0.46	0.05	0.54	0.47	0.44	0.51	0.51
	HOA	0.12	0.17	0.14	0.55	0.38	0.11	0.60	0.42	0.43	0.49	0.48
	BBOA	0.24	0.26	0.24	0.72	0.57	0.13	0.66	0.59	0.56	0.63	0.64
	SV-OOA	0.24	0.28	0.25	0.61	0.48	0.12	0.60	0.54	0.54	0.59	0.61
	Night OA I	0.33	0.36	0.34	0.54	0.48	0.16	0.47	0.53	0.55	0.56	0.61
Night OA II	0.14	0.16	0.15	0.49	0.35	0.03	0.47	0.42	0.42	0.44	0.47	

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

Table 2. Summary of pulls on the four-factor solutions, all with a maximum allowed dQ of 1 %, except for one iteration with BBOA to Chestnut smolder ($dQ = 3 %$), and except for the edge points pull, which had a total allowed dQ of 31 %, or 0.2 % per point with a total of 157 points.

Pull	SV-OOA to Aged Diesel	HOA to Diesel	BBOA to Chestnut Smolder ($dQ=1 %$)	BBOA to Chestnut Smolder ($dQ=3 %$)	Edge points on SV-OOA vs. LV-OOA
Target	Ratio m/z 43/44 from 6.75 to 1.34	Ratio m/z 57/55 from 0.78 to 1.03 and ratio m/z 41/13 from 0.80 to 0.69	Ratio m/z 60/91 from 1.96 to 7.20	Ratio m/z 60/91 from 1.96 to 7.20	157 points pulled to axis
Change in target values?	Yes	Yes	Yes	Yes	Yes
dQ	0.1 %	0.5 %	0.4 %	1.3 %	8.8 %
Improved G-space plot?					
LV-OOA vs. Pittsburgh OOA	Declined r^2 from 0.99 to 0.93	No change	No change	No change	No change
HOA vs. Diesel	No change	Slightly better	No change	No change	No change
BBOA vs. Chestnut smolder	Slightly better; excl. m/z 44	Improved r^2 from 0.79 to 0.87	Improved r^2 from 0.79 to 0.88	Improved r^2 from 0.79 to 0.86	Improved r^2 from 0.79 to 0.93
SV-OOA vs. Aged Diesel	Improved r^2 from 0.62 to 0.69;	Slightly worse r^2 from 0.62 to 0.57	Worse r^2 0.62 to 0.56, due to m/z 44 = 0	Worse r^2 .62 to 0.56, due to m/z 44 = 0	Improved r^2 from 0.69 to 0.72;
Other Changes of Note	0 m/z 44 in BBOA factor; unreasonable result	Large decrease in m/z 44 in SV-OOA, resulting in 43/44 ratio of 21, higher than HOA ratio of 10	In SV-OOA m/z 44 = 0; unreasonable result	In SV-OOA m/z 44 = 0; in BBOA m/z 55 = 0; unreasonable result	SV-OOA m/z 43/44 ratio changed from 4.4 to 2.9



Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)**Fig. 1.** Location of Fyfe Elementary School in Las Vegas, Nevada.

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.




Fig. 2. (a) Box plot of AMS OM ($\mu\text{g m}^{-3}$) by hour, and (b) average concentrations of selected m/z (43, 44, 57, and 60) by hour.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.




Fig. 3. Time series of wind speed (m s^{-1}), AMS OM ($\mu\text{g m}^{-3}$), AMS nitrate, AMS sulfate, AethalometerTM BC ($\mu\text{g m}^{-3}$), CO (ppm), NO (ppb) and NO₂ (ppb). Major tick marks indicate midnight for each day.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.




Fig. 4. Box plot of OM concentrations ($\mu\text{g m}^{-3}$) during downwind, upwind and stagnant (wind speed less than 0.5 m s^{-1}) conditions, grouped by time of day.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.

**Fig. 5.** Summary of OM apportioned by factor number.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.




Fig. 6. PMF factor profiles through m/z 200 for the four factor solution.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.




Fig. 7. PMF factor contributions, OM, BC, CO and wind speed averaged by hour.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.




Fig. 8. Time series of PMF factor contributions, BC, sulfate, nitrate and OM ($\mu\text{g m}^{-3}$).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.




Fig. 9. Scatter plot comparisons of PMF factor contributions for: **(a)** HOA and BC; **(b)** HOA and CO; **(c)** SV-OOA and nitrate; and **d)** LV-OOA and sulfate.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.




Fig. 10. Attribution of OM by factor in the four-factor solution over all data, during downwind conditions only, during downwind conditions between 05:00 and 09:00 LST only, during upwind conditions only, during stagnant conditions, and during 23:00–05:00 LST only.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Aerosol mass spectrometer data in Las Vegas, with EPA PMF

S. G. Brown et al.




Fig. 11. Attribution of OM with four factors for base run, with ratio of m/z 43/44 in the SV-OOA profile pulled toward the 5-h aged diesel profile; ratio of m/z 41/43 in the HOA profile pulled toward the diesel exhaust profile; ratio of m/z 60/91 in the BBOA profile pulled toward the oak-flame source profile; m/z 44 in LV-OOA profile pulled up maximally; and edge points on the HOA/LV-OOA G-space plot pulled down to the y- and x-axes.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion