Atmos. Chem. Phys. Discuss., 11, 22987–23027, 2011 www.atmos-chem-phys-discuss.net/11/22987/2011/ doi:10.5194/acpd-11-22987-2011 © Author(s) 2011. CC Attribution 3.0 License.



This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

# The comparison of MODIS-Aqua (C5) and CALIOP (V2&V3) aerosol optical depth

J. Redemann<sup>1</sup>, M. A. Vaughan<sup>2</sup>, Q. Zhang<sup>1</sup>, Y. Shinozuka<sup>1</sup>, P. B. Russell<sup>3</sup>, J. M. Livingston<sup>4</sup>, M. Kacenelenbogen<sup>5</sup>, and L. A. Remer<sup>6</sup>

<sup>1</sup>BAER Institute/NASA Ames, 4742 Suffolk Ct., Ventura, CA 93003, USA
 <sup>2</sup>NASA Langley Research Center, Hampton, VA 23681, USA
 <sup>3</sup> NASA Ames Research Center, MS 245-5, Bldg. 245, P.O. Box 1, Moffett Field, CA 94035, USA
 <sup>4</sup>SRI, International, 333 Ravenswood Ave., Menlo Park, CA 94025, USA
 <sup>5</sup>ORAU/NASA Ames Research Center, MS 245-5, Moffett Field, CA 94035, USA

<sup>6</sup>NASA Goddard Space Flight Center, code 613.2, Bldg. 33, Greenbelt, MD 20771, USA

Received: 11 July 2011 - Accepted: 2 August 2011 - Published: 16 August 2011

Correspondence to: J. Redemann (jens.redemann-1@nasa.gov)

Published by Copernicus Publications on behalf of the European Geosciences Union.



#### Abstract

We assess the consistency between instantaneously collocated level-2 aerosol optical depth (AOD) retrievals from MODIS-Aqua (C5) and CALIOP (Version 2&3), comparing the standard MODIS AOD (MYD04\_L2) data to the AOD calculated from CALIOP
aerosol extinction profiles for both the previous release (V2) and the latest release (V3) of CALIOP data. Based on data collected in January 2007, we investigate the most useful criteria for screening the MODIS and CALIOP retrievals to achieve the best agreement between the two data sets. Applying these criteria to eight months of data (January, April, July, October 2007 and 2009), we find an order of magnitude increase for the CALIOP V3 data density (by comparison to V2), that is generally accompanied by equal or better agreement with MODIS AOD. Differences in global, monthly mean, over-ocean AOD (532 nm) between CALIOP and MODIS range between 0.03 and 0.04 for CALIOP V3, with CALIOP generally biased low, when all available data from both sensors are considered. Root-mean-squares (RMS) differences in instan-

- <sup>15</sup> taneously collocated AOD retrievals by the two instruments are reduced from values ranging between 0.14 and 0.19 using the unscreened V3 data to values ranging from 0.09 to 0.1 for the screened data. A restriction to scenes with cloud fractions less than 1 % (as defined in the MODIS aerosol retrievals) generally results in improved correlation ( $R^2 > 0.5$ ), except for the month of July when correlations remain relatively lower.
- <sup>20</sup> Regional assessments show hot spots in disagreement between the two sensors in Asian outflow during April and off the coast of South Africa in July.

### 1 Introduction

25

Aerosols introduce a major uncertainty in predictions of possible future changes to the Earth system in general, and its climate in particular, owing to the incomplete knowledge of aerosol physicochemical properties and their spatial distribution. The IPCC-2007 estimates of the direct aerosol radiative forcing (DARF) of climate are largely



based on climate model simulations. Observationally-based estimates are sparse, and their mean result for DARF is about a factor of two larger than the mean of the modelbased estimates (IPCC, 2007). A recent study by Myhre (2009) suggests that the differences between model- and observation-based estimates is largely due to a change

- in aerosol optical properties attributable to anthropogenic activity (i.e., a stronger increase in soot aerosols by comparison to purely scattering aerosols since pre-industrial times) not accounted for in the observation-based estimates. Myhre (2009) concludes that "remaining uncertainty (in DARF) is probably related to the vertical profiles of the aerosols and their location in relation to clouds".
- <sup>10</sup> Designated aerosol satellite sensors such as MODIS, MISR, CALIOP and OMI (Remer et al., 2005; Kaufman et al., 2002; Zhang et al., 2005; Torres et al., 2007; Winker et al., 2010) provide important contributions to our understanding of the effects of aerosols on climate. MODIS and MISR provide reliable retrievals of spectral AOD with reasonably well known uncertainties under most conditions (e.g., Kahn et al., 2010; Rede-
- <sup>15</sup> mann et al., 2006, Levy et al., 2010; Kleidman et al., 2011), with particular strengths over oceans where surface conditions are fairly well known. OMI has begun to investigate aerosol absorption, although extensive validation of the absorption optical depth product is still to be undertaken (Torres et al., 2007; Satheesh et al., 2009). CALIOP provides vertical profiles of aerosol properties over a narrow swath (~90 m)
- <sup>20</sup> along the satellite ground track (Winker et al., 2007). It should be noted that we use the term aerosol "retrieval" in this paper to denote a data product that was inverted from an observation which is not a direct measurement of the aerosol property, but instead requires assumptions about specific aerosol properties such as the aerosol phase function or particle size distribution.
- To provide the full complement of aerosol properties required to improve the accuracy of aerosol radiative effect calculations, these observations have to be combined effectively with each other and with suborbital aerosol measurements. For example, studies of aerosol direct radiative effects (DRE) depend critically on the vertical profile of radiative properties, yet CALIOP, which provides aerosol extinction profile retrievals,



does so at only two wavelengths, and the maturity of these products is still low. Estimates of the semi-direct effect of aerosols depend on reliable measurements of aerosol absorption near the clouds whose developments they are affecting, yet OMI has a large foot-print and its observations become less reliable in cloudy environments (Satheesh

- et al., 2009; Livingston et al., 2009). Finally, estimates of aerosol indirect effects require detailed knowledge of aerosol activation processes in areas difficult to access with satellite sensors, e.g., below opaque clouds. Hence, it is paramount for the scientific community to develop techniques to combine the A-Train aerosol observations with each other and with auxiliary suborbital observations to improve the estimates of
   the specific aerosol properties relevant for the various processes governing aerosol-
- climate interactions.

Although a few studies combining MODIS, OMI and CALIOP observations exist (e.g., Jeong and Hsu, 2008), little is known about the degree of consistency between these data sets. The study presented here investigates the consistency between instanta-

- <sup>15</sup> neously collocated MODIS and CALIOP aerosol retrievals for globally distributed data in eight months (January, April, July, October 2007 and 2009). Investigating intraannual differences allows us to study the impact of different regions and aerosol types on global mean AOD estimates, while studying the inter-annual differences gives insight into potential sensor calibration or performance changes. The most easily com-
- 20 parable observation between the two sensors, that is also relevant for aerosol radiative effects, is AOD, which MODIS provides at seven wavelengths over-ocean (one of which is extrapolated) and three wavelengths over-land (of which two hold no independent information), and which can be calculated from CALIOP profiles of aerosol extinction at 532 nm and 1064 nm. We focus our discussions in Sect. 3 on the quantitative com-
- <sup>25</sup> parison of collocated AOD retrievals over-ocean, because the uncertainty and variability of MODIS over-ocean AOD retrievals is considerably less than over-land and well documented in the literature (e.g., Redemann et al., 2005, 2006, 2009a, c; Russell et al., 2007; Remer et al., 2005, 2008). Therefore, the over-ocean comparisons are more useful in identifying potential shortcomings in the retrievals from either sensor.



Hence, we consider the MODIS over-ocean retrievals a standard that the CALIOP retrievals can be measured by and should agree with within uncertainties. We note that the CALIOP-MODIS AOD comparisons presented here are an imperfect tool to assess CALIOP data quality, because the MODIS retrievals themselves are subject to uncertainties that vary with cloud, surface and aerosol conditions, and because the daytime CALIOP measurements required are adversely affected by the high-intensity solar background signal. However, as previously noted by many investigators (e.g., Schuster et al., 2007), comparing CALIOP to AERONET retrievals is limited to a few hundred opportunities per year if reasonable collocation criteria are applied. Hence, the MODIS-CALIOP comparisons presented here, while imperfect, provide an oppor-

- the MODIS-CALIOP comparisons presented here, while imperfect, provide an opportunity for a statistically relevant assessment of the CALIOP-derived AOD against a data product with peer-reviewed accuracy and limitations. Over-land comparisons are provided in our study only to assess whether or not there is consistency with the overocean retrievals as far as the geographical distribution of AOD differences between the
- two sensors is concerned. Kittaka et al. (2011) provide a similar analysis to the one presented in this paper. However, their study is restricted to CALIOP V2 retrievals, they do not distinguish between MODIS ocean and land retrievals, and their analysis is considerably more qualitative than the analysis described here. Oo and Holz (2011) analyzed the same data in 2007 as we did here, and found a greater bias difference between MODIS and CALIOP AOD. However, their comparisons were screened very
- 20 between MODIS and CALIOP AOD. However, their comparisons were screened ver differently from our data set, making a quantitative comparison of results difficult.

#### 2 Data sets

### 2.1 MODIS-Aqua MYD04\_L2 AOD

MODIS-Aqua (Moderate Resolution Imaging Spectroradiometer) is one of six Earthobserving instruments aboard the Aqua satellite. The MODIS aerosol product (MYD04\_L2) is derived from radiance measurements in channels 1–7 and 20 of the



36 MODIS bands. Over the ocean the measured radiances are inverted into the aerosol optical depths at 470, 550, 659, 865, 1240, 1640 and 2130 nm and volume distribution (in the range of  $0.08-5 \,\mu m$  radius). In the inversion, it is assumed that the aerosol size distribution is bi-modal log-normal (Remer et al., 2005). Aerosol retrievals are produced at a resolution of nominally  $10 \times 10 \text{ km}^2$  at nadir. The MODIS aerosol algorithm 5 uses the standard deviation of 0.55 µm reflectances in groups of 3 by 3 pixels within a box (Martins et al., 2002) to define its cloud mask. If any group of 9 contiguous pixels has a standard deviation greater than 0.0025, the center pixel is labeled as "cloudy" and discarded. This test separates aerosol from most cloud types, but may fail at the centers of large, thick clouds and with cirrus, both of which can be spatially homoge-10 neous. It may also erroneously identify inhomogeneous aerosol fields (e.g., dust) as clouds. In an effort to avoid both scenarios, additional spectral dependence filters are applied. There is an extensive body of literature documenting the validation of MODIS aerosol properties since the beginning of data collection by the MODIS-Terra instrument (e.g., Redemann et al., 2005, 2006, 2009a, c; Russell et al., 2007, Remer et al., 15 2005, 2008). Over oceans, MODIS AOD uncertainties have been shown to be about

 $\pm$ (0.03 + 0.05 AOD); over land retrieval uncertainties are generally  $\pm$ (0.05 + 0.15 AOD) (Remer et al., 2005; Levy et al., 2010; Kleidman et al., 2011).

### 2.2 CALIOP AOD derived from aerosol extinction profiles

- Within the A-Train, the CALIPSO satellite carries an elastic backscatter lidar (CALIOP), a three-channel imaging infrared radiometer (IIR) and a wide-field camera (WFC). The key instrument for this work is CALIOP, a two-wavelength (532 and 1064 nm), polarization-sensitive lidar that makes continuous profile measurements in the plane-tary boundary layer (PBL) and free troposphere at single-shot vertical and horizontal
   resolutions of 30 m and 333 m at 532 nm, respectively. While boundary layer clouds are readily detected in the full resolution date the single shot single chart aircraft (CND) of
- readily detected in the full resolution data, the single-shot signal-to-noise ratio (SNR) of the CALIOP profile data is too low to reliably retrieve aerosol spatial and optical properties (Hunt et al., 2009; Winker et al., 2009). Version 2 of the CALIOP level-2 data



products reported aerosol spatial properties (in the layer product files) at a horizontal resolution of 5 km, and range-resolved aerosol optical properties (in the profile product files) at a horizontal resolution of 40 km. The new version 3 data products now report aerosol optical properties at the same 5-km horizontal resolution used for the spa-

- tial properties. However, the same optical properties retrieval strategy is used in both versions 2 and 3 of the CALIOP data products (Young and Vaughan, 2009). This technique does not provide a direct measurement of AOD, but instead derives estimates of AOD by first assessing the geo-spatial location and optical characteristics of aerosol layers, and then selecting a "best-match" extinction-to-backscatter ratio (also known as the lidar ratio) from a look-up table based on a compilation of in-situ observations and
- the lidar ratio) from a look-up table based on a compilation of in-situ observations an a cluster analysis of AERONET data (Omar et al., 2009).

It should be noted that the CALIOP instrument was not primarily designed to provide AOD, but instead vertical profiles of aerosol backscatter, depolarization and extinction. As such, the derivation of AOD from integration of extinction profiles is subject to sev-

eral limitations and uncertainties. Kacenelenbogen et al. (2011) provide a description of the most important potential uncertainties in CALIOP AOD on the basis of a detailed multi-sensor case study. These are:

20

- i. CALIOP's low signal-to-noise ratio (SNR), which can lead to the misclassification and/or lack of aerosol layer identification, especially for daytime measurements and/or in optically thick layers which rapidly attenuate the backscatter signal;
- ii. possible cloud contamination of CALIOP aerosol backscatter and extinction profiles;
- iii. potentially erroneous assumptions of the aerosol extinction-to-backscatter ratio  $(S_a)$  used in CALIOP's extinction retrievals; and
- <sup>25</sup> iv. calibration coefficient biases in the CALIOP attenuated backscatter coefficient profiles.



In going from CALIOP V2 to V3, major code and algorithm modifications were implemented. They were intended to result in more accurate V3 estimates of layer spatial and optical properties and can be summarized as follows:

- 1. improved daytime calibration procedures, as described by Powell et al. (2010);
- refinements to the layer detection algorithm including the elimination of a bug in the cloud clearing code and an alternative retrieval philosophy for low-lying aerosol layers – for certain conditions, the aerosol layer base estimate is revised to a new, generally lower altitude near the surface (Vaughan et al., 2010);
  - development of a new five-dimensional PDF-based cloud and aerosol discrimination algorithm, especially leading to a better classification of dense aerosol layers (Liu et al., 2010);
  - 4. implementation of an entirely new algorithm for determining cloud thermodynamic phase (Hu et al., 2009).
- In addition, the V3 products also contain new diagnostics, quality assurance parame ters and uncertainty estimates that were not available in prior releases (e.g., extinction QC flags, cloud and aerosol feature fraction, atmospheric volume description, column optical depths, ice water content, particulate depolarization ratio profile and associated uncertainties). These algorithm improvements influenced the CALIPSO project's decision to upgrade the maturity level of the optical depth data from a "beta" product in V2
   to a "provisional" product in V3, the former not considered suitable for use in scientific publication. Nonetheless, the V2 data were used extensively in several published stud-
- ies (e.g., Peyridieu et al., 2010, Jones and Christopher, 2010), partly motivating the quantitative comparisons presented in this paper.

### 2.3 Data collocation

10

As pointed out previously, the CALIOP V2 retrievals are based on CALIOP extinction profiles at a horizontal resolution of 40 km. CALIOP V3 extinction profiles (and by 22994



extension our AOD retrievals) have a horizontal resolution of 5 km. Figure 1 shows the basic strategy for collocating MODIS AOD retrievals and CALIOP V2 and V3 derived AOD. For the comparison of CALIOP V2 AOD to MODIS AOD retrievals, we averaged all available, valid MODIS retrievals in a 40 × 40 km<sup>2</sup> grid box centered at the CALIOP  $_{5}$  V2 retrieval. As an example, this is illustrated by the two  $40 \times 40 \text{ km}^2$  boxes plotted in Fig. 1. MODIS AOD retrievals are shown by filled parallelograms, roughly measuring  $10 \times 10$  km<sup>2</sup>, as the CALIPSO ground track is near the nadir point of the MODIS scans. Along the CALIOP track,  $40 \times 40 \text{ km}^2$  boxes centered around the CALIOP V2 AOD retrieval center points are shown in red. Light green dots denote the center points of the MODIS retrieval boxes. The MODIS retrievals that were averaged for 10 comparison to the two CALIOP V2 retrievals are denoted by red dots near the light green center dots. Black squares in Fig. 1 denote 12 × 12 km<sup>2</sup> boxes centered around the CALIOP V3 AOD retrieval center points. The 12 × 12 km<sup>2</sup> boxes were chosen because they present an optimum sampling of the MODIS retrievals. Smaller boxes (e.g.,  $10 \times 10 \text{ km}^2$ ) often did not contain any center points of MODIS retrieval boxes; larger 15 boxes (e.g.,  $15 \times 15$  km<sup>2</sup>) would have resulted in undue spatial averaging of several MODIS retrieval boxes, as opposed to using just one closely collocated MODIS re-

- trieval box. As an example, for 1 October 2007, using the  $12 \times 12 \text{ km}^2$  boxes resulted in an 84% probability of having exactly one MODIS retrieval in the box (16% had two retrievals); using  $13 \times 13 \text{ km}^2$  boxes resulted in a 74% probability of having exactly one MODIS retrieval in the box (24% had two retrievals, 2% had four retrievals); using  $14 \times 14 \text{ km}^2$  boxes resulted in a less than 64% probability of having exactly one MODIS retrieval in the box (32% had two retrievals, more than 4% had four retrievals). At the same time, the average daily AOD changed by less than 0.001 (i.e., less than 1%)
- <sup>25</sup> between any one sampling option and the next, while RMS differences increased with box size. Hence, we conclude that 12 × 12 km<sup>2</sup> boxes centered at CALIOP V3 AOD retrievals provide an optimum sampling of MODIS retrievals for comparison of the two data sets.



3 Results

# 3.1 One month comparison (January 2007) between MODIS AOD and CALIOP (V2&V3) AOD over ocean

To illustrate the technical details of our comparison between MODIS and CALIOP AOD,
this section discusses the comparison of MODIS and CALIOP AOD at 532 nm for January 2007. As pointed out above, to calculate the CALIOP AOD at 532 nm, we integrate the aerosol extinction profile contained in the scientific data segment (sds) "Extinction\_Coefficient\_532" with respect to altitude after replacing all fill values with a numerical value of zero. To facilitate the comparison with MODIS, we use a quadratic fit of log(AOD) versus log(wavelength) for the seven over-ocean MODIS spectral AOD (sds "Effective\_Optical\_Depth\_Average\_Ocean") and interpolate to a wavelength of 532 nm. As described in Sect. 2.3, we use different spatial matching between MODIS and CALIOP V2 on one hand, and between MODIS and CALIOP V3 on the other, because the CALIOP extinction profile data is presented at the higher horizontal resolution of 5km in V3 along the CALIPSO ground track. This results in a comparison of CALIOP V2 data are com-

pared to spatially more closely collocated MODIS retrievals at the 12 × 12 km<sup>2</sup> scale. In a heterogeneous aerosol environment, the coarser spatial sampling for CALIOP V2 is likely to introduce some variability in the comparisons. Spatial variability of aerosol properties is discussed in more detail in the discussion section of our paper.

Figure 2 shows scatter plot comparisons between MODIS and CALIOP V2 in the first row (Fig. 2a–c) and between MODIS and CALIOP V3 (Fig. 2d–f) in the second row. The symbol color represents the MODIS derived fine mode fraction (FMF), i.e., the fraction of AOD that is due to the fine particle mode in the bi-modal fine-coarse mode <sup>25</sup> retrieval. The solid red lines show the linear fits to the data using a model-2 regression method, i.e., the least-squares bisector method. In general, model-2 regressions are more appropriate when both data sets have measurement errors and when such errors are deemed to be of similar order of magnitude (e.g., Sokal and Rohlf, 1995). The fit



parameters of this line are determined by calculating the slope of the line that bisects the minor angle between the regression of Y-on-X and X-on-Y. The fit line parameters along with monthly mean AOD from the two sensors, the mean and standard deviation of the difference between the two sensors, the number of available data points, the

- <sup>5</sup> square of the linear correlation coefficient and the rms difference between the two data sets are given in Table 2. Figure 2a and d shows comparisons for all available collocated MODIS and CALIOP AOD retrievals in the month of January, i.e., 36859 for CALIOP V2 and 247976 for CALIOP V3. These numbers essentially reflect the 8-fold increase in data density in going from CALIOP V2 to V3, although not all valid
- <sup>10</sup> V2 40 km retrievals necessarily contain eight valid V3 5 km retrievals and not all valid V3 5km retrieval are collocated with valid V2 40 km retrievals. In addition, changes in the retrieval algorithm, most notably with respect to cloud clearing, would result in differences between the two data sets. Comparing the correlations of CALIOP V2 and V3 with the MODIS data, we note a significantly better correlation for V3 ( $r^2$  of 0.25 as opposed to 0.04 for V2) and smaller rms differences (0.19 as opposed to 0.33 for V2).
- Mean differences from MODIS AOD, however, are smaller for V2 than for V3.

To investigate if an application of the various quality criteria published in the MODIS and CALIOP data sets result in better agreement between the two AOD data sets, we carried out several tests, which successively increased the complexity of the data screening process. Table 1 provides the details of the various comparisons. Starting

- with the comparisons of all data (comparison #1), we added the following restrictions, requiring:
  - MODIS AOD data to have an aerosol quality flag (Quality\_Assurance\_Ocean) of 1 (marginal), 2 (good), or 3 (very good) (comparisons # 2–6 in Table 1);
- CALIOP extinction retrievals to fall within the "valid range" (identified in the CALIPSO data products catalog as 0 to 1.25 km<sup>-1</sup>) (comparisons # 2–4 in Table 1);
  - 3. CALIOP extinction retrievals to have quality flags of 0, 1 or 2 and to eliminate 22997

	ACPD								
	11, 22987–23027, 2011								
	The comparison of MODIS-Aqua and CALIOP AOD J. Redemann et al.								
	Title	Page							
D r	Abstract	Introduction							
_	Conclusions	References							
	Tables	Figures							
	14	۶I							
5	•	•							
-	Back	Close							
	Full Scre	en / Esc							
<u>.</u>	Printer-frier	Idly Version							
	Interactive	Discussion							
5	C	•							

profiles for which any extinction retrievals do not have said quality flags (comparisons # 3–5 in Table 1);

- CALIOP extinction profiles to have collocated cloud optical depth retrievals equal to zero (comparisons # 3–6 in Table 1);
- 5. MODIS cloud fractions (as determined by the aerosol algorithm) to be below 1 % (comparison # 4 in Table 1);
  - CALIOP extinction retrievals to have uncertainties less than 200 % when extinction is below negative 0.2 km<sup>-1</sup>, or less than 500 % when extinction is greater than 2.5 km<sup>-1</sup> and to eliminate profiles for which any extinction retrievals do not have said extinction coefficients and uncertainty limits (as described in comparison # 5 in Table 1);

10

7. CALIOP relative AOD uncertainty calculated from the extinction uncertainties to be below 100% (comparison # 6 in Table 1).

Figure 2b and e shows the same comparisons as Fig. 2a and d, after the application of the data quality criteria summarized as comparison #3 in Table 1. First, we restrict the comparison to MODIS data with an aerosol guality flag (Quality\_Assurance\_Ocean) of 1 (marginal), 2 (good), or 3 (very good), all designated as appropriate for quantitative use. Second, we only use extinction retrievals that were in the documented valid range of 0 to 1.25 km<sup>-1</sup>. Third, we only use comparisons for which the CALIOP data indicates a cloud optical depth of zero. Fourth, we only use extinction profiles for which 20 all extinction retrievals have a quality flag of 0, 1 or 2. This combination of quality criteria is summarized as comparison # 3 in Table 1. V3 aerosol profile data files contain an sds entitled "Column\_Optical\_Depth\_Cloud\_532", which we used for the second step described above. The CALIOP V2 aerosol profile data, however, contained no analogous data field. We therefore used the V2 cloud layer data (at 5 km resolution) 25 and only allowed data that showed no cloud layers as indicated by the sds "Number\_Layers\_Found". Because the V2 profile data is reported at a horizontal resolution



of 40 km, yet the cloud layer products are reported at a resolution of 5 km, we only used V2 40 km aerosol extinction profile data for which all eight 5 km sub-segments showed no cloud layers present. Effectively, this represents a very stringent cloud screening, as no V2 extinction data were used in our analysis if even a single 5 km sub-segment was

reported to contain a cloud. Consequently, the application of the three quality criteria (MODISQA = 1–3, CALIOPcloudOD = 0, extQCflag = 0,1,2) has a much more dramatic impact on the V2 comparisons than on the V3 comparisons to MODIS. For V2, only about 15% of the original data shown in Fig. 2a are still present in Fig. 2b. For V3, more than 72% of the data survive the application of these quality criteria. There is
 significant improvement after application of the quality criteria for both CALIOP V2 and

V3, especially with respect to rms differences.

As an additional restriction to even more cloud-free conditions, Fig. 2c and f show the comparisons between MODIS and CALIOP for cloud fractions of less than 1 % as defined by the MODIS aerosol product (Cloud\_Fraction\_Ocean). The statistical param-

- eters for these comparisons are given as comparison # 4 in Table 2. Both V2 and V3 show larger correlation coefficients (both  $R^2$  greater than 0.5). It should be noted, however, that there is 20 times more data in CALIOP V3 (N = 37740) than in CALIOP V2 (N = 1883) that remains after application of the four quality criteria and restriction to cloud fractions less than 1%. In terms of absolute numbers, the remaining V2 re-
- trievals represent about 5 % of all valid V2 retrievals, while the remaining V3 retrievals represent more than 15 % of all valid V3 retrievals.

We tested several other methodologies for combining the various restrictions described in the previous paragraphs. Two such combinations are described as comparison # 5 and # 6 in Table 1. Comparison # 5 tests the usefulness of the CALIOP

extinction uncertainties, and comparison # 6 tests the usefulness of the published uncertainty in the CALIOP column AOD. In comparison # 5, we restrict our AOD comparison to use only CALIOP extinction retrievals whose uncertainties are below 200% at the low (negative) end of the extinction spectrum and below 500% at the high end of the extinction spectrum. We allowed negative values in the extinction, because some



range of small negative values is plausible based on the retrieval technique and simple statistical scatter of solutions around the correct retrieval. Table 2 shows that an application of comparison # 5 and # 6 to the January 2007 data results in less good correlation between the two data sets. Similar tests with different ranges of allowable 5 CALIOP extinction values (not shown here) also provided no improvement in correlation when compared to the methodology of comparison # 3. Therefore, we summarize

- the best methodology for comparing MODIS and CALIOP V3 AOD to consist of the following 4 steps:
  - 1. use MODIS aerosol quality flag (Quality\_Assurance\_Ocean) of 1-3;

15

- use only CALIOP extinction retrievals that fall within the "valid range" of 0 to 1.25 km<sup>-1</sup>;
  - 3. require CALIOP extinction profiles to have collocated cloud Optical depth retrievals equal to zero; and
  - 4. eliminate CALIOP profiles for which any extinction retrievals have quality flags different from 0, 1 or 2.

As we will see in Sect. 3.2, applying these quality criteria eliminates only 20-30% of all collocated cases and results in the best correlation between the two AOD data sets.

# 3.2 Eight months of comparisons (January, April, July, October 2007 and 2009) between MODIS AOD and CALIOP (V2&V3) AOD over land and ocean

In this section we extend the results shown in Sect. 3.1 to eight months of data: January, April, July and October of 2007 and 2009. Considering several months within one year will allow us to study whether certain aerosol types in certain geographical regions contribute differently to the global MODIS-CALIOP AOD differences. By considering two years of data (2007 and 2009) we are able to assess potential sensor calibration or performance changes.



Figure 3a shows the comparison of MODIS to CALIOP V2 and V3 AOD (532 nm) for four months in 2007 in terms of three parameters: (1)  $\Delta$ AOD (M-C), the global monthly mean difference in AOD (MODIS-CALIOP), (2) the square of the linear correlation coefficient ( $r^2$ ), and (3) the RMS difference between MODIS and the respective CALIOP

- <sup>5</sup> retrievals. The left hand side of Fig. 3a shows these parameters for all retrievals in CALIOP V2 and V3, while the right hand side only shows the comparison using the quality criteria summarized as comparison # 3 in Table 1. Hereafter, we will refer to these comparisons as V2QA and V3QA, respectively. Figure 3b shows the same data for four months in 2009.
- In going from V2 to V3 for all data (left hand side in Fig. 3a and b, as well as comparison # 1 in Tables 2–9), the analysis is consistent between different months and even between 2007 and 2009. The mean optical depth differences between MODIS and CALIOP increase in going from V2 to V3. However, the correlation coefficients (dashed lines) increase from V2 to V3, and RMS differences (dotted lines) decrease from values greater than 0.3 to values between 0.15 and 0.2.

The comparisons for the quality assured data sets (right hand sides of Fig. 3a and b as well as comparison # 3 in Tables 2–9) are less straight forward. In 2007 (Fig. 3a), the mean differences from MODIS increase between V2QA and V3QA, while in 2009 the differences to MODIS for three of the four months essentially stay the same. The

- <sup>20</sup> correlation with MODIS data increases between V2QA and V3QA for January 2007, April 2007, October 2007 and July 2009. For January and October 2009 it is essentially unchanged between V2QA and V3QA, and for July 2007 and April 2009, the correlation with MODIS data decreases between V2QA and V3QA. However, as discussed in Sect. 3.1, the quality-assured V2QA data only represents a small fraction
- of the original V2 data and the four quality criteria effectively represent a strong initial cloud filter, making the comparison between V2QA and V3QA somewhat misleading. Comparison 4 in Tables 2–9 illustrates that after application of the criterion that MODIS cloud fractions as determined by the aerosol algorithm have to be below 1 %, the difference between V2QA and V3QA in terms of correlation with MODIS data is negligible,



with most RMS differences near 0.1.

Because of the relatively better performance of the data screening methodology summarized as comparison # 3 in Table 1 to all other methodologies, starting with Fig. 4, all MODIS versus CALIOP V3 AOD comparisons were screened according to that methodology. Figure 4 shows the difference between MODIS and V3QA data as histograms. The blue curves represent over-ocean data. Comparing all eight months in 2007 and 2009, we observe a consistent difference between MODIS and CALIOP V3QA of the order of 0.03–0.04. The AOD difference is essentially normally distributed. In addition to the over-ocean data, we present the over-land comparisons (using MODIS sds "Corrected\_Optical\_Depth\_Land") for each month as red distribution curves. We note the much broader distributions with mode values near –0.01. The broader distributions are attributable to the larger uncertainties in MODIS over-land AOD retrievals, as well as differences in the aerosol type identified by the CALIOP aerosol classification scheme.

## **3.3** Geographical distribution of AOD differences between MODIS and CALIOP V3

In this section we present results regarding the geographical distribution of the AOD differences between MODIS and CALIOP V3QA. Figure 5 shows the latitudinally averaged AOD differences for over-ocean retrievals (blue lines) and over-land retrievals

- (red lines). The thin lines represent 1-degree averaged data, while the thick lines represent 5-degree averages. Figure 5a–d shows the data for 2007, while Fig. 5e–h shows the data for 2009. As the thin lines in Fig. 5 indicate, the 1-degree data is considerably noisier over-land than over-ocean. This is due to both the decreased quality of the over-land MODIS retrievals, as well as the data sparseness in some latitude regions.
- Figure 6 shows the same comparisons but restricted to data with MODIS cloud fractions less than 1 % (Cloud\_Fraction\_Ocean/Land < 0.01, comparison # 4 in Table 1). Focusing on the over-ocean comparisons, there are some remarkably similar features between 2007 and 2009. For example, there is a consistent difference in April of near



0.1 in the Northern Hemisphere. In July, centered near 5° S, there is a similar, though more localized, increase in the difference between MODIS and CALIOP V3QA. Interestingly, both of these features are also present in the cloud-screened data in Fig. 6, making cloud contamination rather unlikely as a possible cause of the discrepancy.

- <sup>5</sup> Figures 7 and 8 show the global distribution of the number of instantaneously collocated comparisons between MODIS and CALIOP V3QA in a 5 × 5 degree box for each month in the left hand plots (a–d of each figure) and the AOD difference in the right hand plots. Several interesting features are observable. The aforementioned AOD difference in April in the Northern hemisphere seems to be due mostly to outflow from
- Asia. The AOD difference feature in the Southern hemisphere in July seems to be due almost exclusively to transport of biomass burning aerosols off of the African subcontinent into the Atlantic Basin. On the other hand, these plots side by side also illustrate that most emphasis should be placed on locations where the vast majority of instantaneously collocated observations are made. This area is generally confined to the
- <sup>15</sup> region 40 degrees to the North and South of the equator, with the notable exception of the region between 10° N and 40° N in July, where the CALIOP retrievals are inside the MODIS glint region and no coincident, collocated observations can be made by the two sensors.

We note further that the collocated over-land comparisons, while much less frequent, generally provide a consistent picture with the adjacent regional over-ocean comparisons. For example, the large positive difference between MODIS and CALIOP V3QA along the West-African coast near 5° S in July 2009 is also found in the inland retrievals.

#### 4 Discussion

In this paper, we present a general consistency analysis between MODIS and CALIOP

V2 and V3 AOD. We use only instantaneously collocated retrievals by the two sensors. A quantitative comparison of the agreement between MODIS and CALIOP V2 AOD on one hand and MODIS and CALIOP V3 AOD on the other is difficult, because of differing

iscussion Pa	AC 11, 22987–2	<b>PD</b> 23027, 2011
per   Discussion	The comp MODIS-A CALIO J. Redem	oarison of Aqua and P AOD ann et al.
Pape	Title	Page
er.	Abstract	Introduction
	Conclusions	References
iscussi	Tables	Figures
on P	14	►I
aper	•	•
_	Back	Close
Discu	Full Scre	een / Esc
ssion	Printer-frier	ndly Version
Pap	Interactive	Discussion
)er	6	•

spatial sampling of the MODIS data that are compared to the CALIOP data. Our analysis of the most suitable comparison scale showed that using  $12 \times 12 \text{ km}^2$  boxes around the CALIOP V3 retrievals resulted in optimal number of collocations with minimal impact of spatial variability in aerosol properties on the data comparison. Specifically, we

- found a steady increase in the rms difference between MODIS and CALIOP V3 AOD when going from 11 × 11 km<sup>2</sup> boxes to 15 × 15 km<sup>2</sup> boxes. However, there was a significant drop in the number of collocated retrievals when we went from 12 × 12 km<sup>2</sup> boxes to 11 × 11 km<sup>2</sup> boxes, hence our choice of 12 × 12 km<sup>2</sup> boxes. The steady increase in rms difference between the two sensors' AOD leads us to conclude that there is a finite
   rms difference caused by the spatial variability of aerosol properties, as is the case for any comparison of AOD retrievals based on data from sensors with different spatial
  - and temporal sampling (Shinozuka and Redemann, 2011).

In general, we find increased data density and equal or better agreement of CALIOP V3 than CALIOP V2 when compared to MODIS AOD over-ocean. A restriction to data with MODIO exceeded evaluation for (Overline Accuracy) of 1.0. CALIOP evidence

- <sup>15</sup> with MODIS aerosol quality flag (Quality\_Assurance\_Ocean) of 1–3, CALIOP extinction retrievals that fall within the "valid range" of 0 to 1.25 km<sup>-1</sup>, eliminating CALIOP profiles for which any extinction retrievals have quality flags different from 0, 1 or 2, and CALIOP extinction profiles that have collocated cloud optical depths equal to zero, yields the strongest correlation between the MODIS and CALIOP derived AOD (with-
- out restricting the comparisons to severely cloud-free cases). In absolute terms, the comparison between the two data sets still yields rms differences of 0.1, which may be too large for many applications that attempt to combine these data sets quantitatively. It should be noted that some fraction of this difference is due to the fact that CALIOP only estimates aerosol properties in those regions where it "detects" an aerosol layer.
- In regions where aerosol layers are more tenuous, i.e., distributed over greater geometric depth, CALIOP likely fails to detect such aerosol layers, contributing to the rms difference found here. This is corroborated by independent aircraft validation exercises in the ARCTAS experiment (Redemann et al., 2009b). The geographical distribution of over-ocean AOD difference shows latitudinally averaged differences of generally less



than 0.05, although some latitude bands with 0.1 AOD differences occur. These bands can often be traced to regional AOD difference hot spots, for example in Asian outflow during April and African outflow into the Atlantic Ocean basin during July. We find no statistically significant difference in the consistency between MODIS and CALIOP

- <sup>5</sup> V2/V3 between 2007 and 2009, leading to the conclusion that sensor calibration and performance did not change significantly between 2007 and 2009. As stated in the introduction, the study presented here is worthwhile because the MODIS over-ocean AOD retrieval uncertainty is reasonably well documented in the literature, and because few other opportunities exist to assess CALIOP AOD retrieval quality. However, our
- study is hampered by the necessary use of CALIOP daytime data and by the imperfect collocation of sampling between the two sensors. The latter is a limitation that is inherent to all data comparisons with different spatial sampling, but that should only matter if spatial variability of AOD is large. Future studies to assess CALIOP data quality using specifically designed suborbital observations and CALIOP nighttime retrievals should be encouraged.

*Acknowledgements.* This study was partially supported by CALIPSO ST funding under NASA grant NNX10AN60G. We would like to thank the MODIS and CALIOP algorithm development teams for their efforts in providing and discussing these data sets.

#### References

<sup>20</sup> Hu, Y., Winker, D., Vaughan, M., Lin, B., Omar, A., Trepte, C., Flittner, D., Yang, P., Nasiri, S., Baum, B., Holz, R., Sun, W., Liu, Z., Wang, Z., Young, S., Stamnes, K., Huang, J., and Kuehn, R.: CALIPSO/CALIOP Cloud Phase Discrimination Algorithm, J. Atmos. Oceanic Technol., 26, 2293–2309, doi:10.1175/2009JTECHA1280.1, 2009.

Hunt, W. H., Winker, D. M., Vaughan, M. A., Powell, K. A., Lucker, P. L., Weimer, C.: CALIPSO

Lidar Description and Performance Assessment, J. Atmos. Oceanic Technol., 26, 1214– 1228, doi:10.1175/2009JTECHA1223.1, 2009.

IPCC: Contribution of Working Groups I, II and III to the Fourth Assessment Report of the



23006

Intergovernmental Panel on Climate Change, Core Writing Team: Pachauri, R. K. and Reisinger, A., IPCC, Geneva, Switzerland, 2007.

- Jeong, M.-J. and Hsu, N. C.: Retrievals of aerosol single-scattering albedo and effective aerosol layer height for biomass-burning smoke: Synergy derived from "A-Train" sensors, Geophys.
- <sup>5</sup> Res. Lett., 35, L24801, doi:10.1029/2008GL036279, 2008.

20

25

30

Jones, T. A. and Christopher, S. A.: Statistical properties of aerosol-cloud-precipitation interactions in South America, Atmos. Chem. Phys., 10, 2287–2305, doi:10.5194/acp-10-2287-2010, 2010.

Kacenelenbogen, M., Vaughan, M. A., Redemann, J., Hoff, R. M., Rogers, R. R., Ferrare, R.

A., Russell, P. B., Hostetler, C. A., Hair, J. W., and Holben, B. N.: An accuracy assessment of the CALIOP/CALIPSO version 2/version 3 daytime aerosol extinction product based on a detailed multi-sensor, multi-platform case study, Atmos. Chem. Phys., 11, 3981–4000, doi:10.5194/acp-11-3981-2011, 2011.

Kahn, R. A., Gaitley, B. J., Martonchik, J. V., Diner, D. J., Crean, K. A., and Holben, B.: Mul-

- tiangle Imaging Spectroradiometer (MISR) global aerosol optical depth validation based on 2 years of coincident Aerosol Robotic Network (AERONET) observations, J. Geophys. Res., 110, D10S04, doi:10.1029/2004JD004706, 2005.
  - Kahn, R. A., Gaitley, B. J., Garay, M. J., Diner, D. J., Eck, T., Smirnov, A., and Holben, B. N.: MISR global aerosol product assessment by comparison with Aerosol Robotic Network, J. Geophys. Res., 115, D23209, doi:10.1029/2010JD014601, 2010.
  - Kaufman, Y., Tanré, D., and Boucher, O.: A satellite view of aerosols in the climate system, Nature, 419, 215–223, 2002.
  - Kittaka, C., Winker, D. M., Vaughan, M. A., Omar, A., and Remer, L. A.: Intercomparison of column aerosol optical depths from CALIPSO and MODIS-Aqua, Atmos. Meas. Tech., 4, 131-141, doi:10.5194/amt-4-131-2011, 2011.
  - Kleidman, R. G., Smirnov, A., Levy, R. C., Mattoo, S., and Tanre, D.: Evaluation and wind speed dependence of MODIS aerosol retrievals over open ocean, IEEE TGARS May, accepted, 2011.

Levy, R. C., Remer, L. A., Kleidman, R. G., Mattoo, S., Ichoku, C., Kahn, R., and Eck, T. F.: Global evaluation of the Collection 5 MODIS dark-target aerosol products over land, Atmos. Chem. Phys., 10, 10399–10420, doi:10.5194/acp-10-10399-2010, 2010.

Liu, Z., Kuehn, R., Vaughan, M., Winker, D., Omar, A., Powell, K., Trepte, C., Hu, Y., and Hostetler, C.: The CALIPSO Cloud and Aerosol Discrimination: Version 3 Algorithm and



Test Results, 25th International Laser Radar Conference (ILRC), St. Petersburg, Russia, ISBN 978-5-94458-109-9, 2010.

- Livingston, J. M., Redemann, J., Russell, P. B., Torres, O., Veihelmann, B., Veefkind, P., Braak, R., Smirnov, A., Remer, L., Bergstrom, R. W., Coddington, O., Schmidt, K. S., Pilewskie,
- P., Johnson, R., and Zhang, Q.: Comparison of aerosol optical depths from the Ozone Monitoring Instrument (OMI) on Aura with results from airborne sunphotometry, other space and ground measurements during MILAGRO/INTEX-B, Atmos. Chem. Phys., 9, 6743–6765, doi:10.5194/acp-9-6743-2009, 2009.
- Martins, J. V., Tanré, D., Remer, L. A., Kaufman, Y. J., Mattoo, S., and Levy, R.: MODIS Cloud screening for remote sensing of aerosol over oceans using spatial variability. Geophys. Res.
  - Lett., 29(12), 8009, doi:10.1029/2001GL013252, 2002. Myhre, G.: Consistency between satellite-derived and modeled estimates of the direct aerosol

20

- effect, Science, 325(5937), 187–190, doi:10.1126/science.1174461, 2009.
- Omar, A. H., Won, J., Winker, D. M., Yoon, S., Dubovik, O., and McCormick, M. P.: Development of global aerosol models using cluster analysis of Aerosol Robotic Network (AERONET) measurements, J. Geophys. Res., 110, D10S14, doi:10.1029/2004JD004874, 2005.
  - Omar, A. H., Winker, D. M., Vaughan, M. A., Hu, Y., Trepte, C. R., Ferrare, R. A., Lee, K.-P., Hostetler, C. A., Kittaka, C., Rogers, R. R., Kuehn, R. E., and Liu, Z.: The CALIPSO Automated Aerosol Classification and Lidar Ratio Selection Algorithm. J. Atmos. Oceanic Technol., 26, 1994–2014. doi:10.1175/2009JTECHA1231.1, 2009.
  - Oo, M. and Holz, R.: Improving the CALIOP aerosol optical depth using combined MODIS-CALIOP observations and CALIOP integrated attenuated total color ratio, J. Geophys. Res., 116, D14201, doi:10.1029/2010JD014894, 2011.

Peyridieu, S., Chédin, A., Tanré, D., Capelle, V., Pierangelo, C., Lamquin, N., and Armante,

- R.: Saharan dust infrared optical depth and altitude retrieved from AIRS: a focus over North Atlantic – comparison to MODIS and CALIPSO, Atmos. Chem. Phys., 10, 1953–1967, doi:10.5194/acp-10-1953-2010, 2010.
  - Powell, K. A., Vaughan, M. A., Rogers, R. R., Kuehn, R. E., Hunt, W. H., Lee, K.-P., and Murray, T. D.: The CALIOP 532-nm Channel Daytime Calibration: Version 3 Algorithm, Proceedings
- <sup>30</sup> of the 25th International Laser Radar Conference, 1367–1370, ISBN 978-5-94458-109-9, 2010.
  - Redemann, J., Schmid, B., Eilers, J. A., Kahn, R. A., Levy, R. C., Russell, P. B., Livingston, J. M., Hobbs, P. V., Smith Jr., W. L., and Holben, B. N.: Suborbital measurements of spectral



aerosol optical depth and its variability at sub-satellite grid scales in support of CLAMS, 2001, J. Atmos. Sci. , 62(4), 993–1007, doi:10.1175/JAS3387.1, 2005.

- Redemann, J., Zhang, Q., Schmid, B., Russell, P. B., Livingston, J. M., Jonsson, H., and Remer, L. A.: Assessment of MODIS-derived visible and near-IR aerosol optical properties
- and their spatial variability in the presence of mineral dust, Geophys. Res. Lett., 33, L18814, doi:10.1029/2006GL026626, 2006.
  - Redemann, J., Pilewskie, P., Russell, P. B., Livingston, J. M., Howard, S., Schmid, B., Pommier, J., Gore, W., Eilers, J., and Wendisch, M.: Airborne measurements of spectral direct aerosol radiative forcing in the Intercontinental chemical Transport Experiment/Intercontinental
- <sup>10</sup> Transport and Chemical Transformation of anthropogenic pollution, 2004, J. Geophys. Res., 111, D14210, doi:10.1029/2005JD006812, 2006.
  - Redemann, J., Zhang, Q., Russell, P. B., Livingston, J. M., and Remer, L. A.: Case studies of aerosol remote sensing in the vicinity of clouds, J. Geophys. Res., 114, D06209, doi:10.1029/2008JD010774, 2009a.
- Redemann, J., Livingston, J., Shinozuka, Y., Russell, P., Zhang, Q., Johnson, R., Ramachandran, S., Clarke, A., McNaughton, C., and Veheilmann, B.: Airborne sunphotometer (AATS-14) measurements in ARCTAS – first insights into their combined use with satellite observations to study Arctic aerosol radiative effects, ARCTAS Science Team Meeting, Virginia Beach, VA, January 27, 2009b.
- Redemann, J., Zhang, Q., Livingston, J., Russell, P., Shinozuka, Y., Clarke, A., Johnson, R., and Levy, R.: Testing aerosol properties in MODIS Collection 4 and 5 using airborne sunphotometer observations in INTEX-B/MILAGRO, Atmos. Chem. Phys., 9, 8159-8172, doi:10.5194/acp-9-8159-2009, 2009c.

Remer, L. A., Kaufman, Y. J., Tanré, D., Mattoo, S., Chu, D. A., Martins, J. V., Li, R.-R., Ichoku,

- C., Levy, R. C., Kleidman, R. G., Eck, T. F., Vermote, E., and Holben, B. N.: The MODIS aerosol algorithm, products and validation. J. Atmos. Sci., 62(4), 947–973, 2005.
  - Remer, L. A., Kleidman, R. G., Levy, R. C., Kaufman, Y. J., Tanre, D., Mattoo, S., Martins, J. V., Ichoku, C., Koren, I., Yu, H. B., and Holben, B. N.: Global aerosol climatology from the MODIS satellite sensors, J. Geophys. Res., 113(D14), D14S07, doi:10.1029/2007JD009661, 2008.

30

Russell, P. B., Livingston, J. M., Redemann, J., Schmid, B., Ramirez, S. A., Eilers, J., Kahn, R., Chu, A., Remer, L., Quinn, P. K., Rood, M. J., and Wang, W.: Multi-grid-cell validation of satellite aerosol property retrievals in INTEX/ITCT/ICARTT 2004, J. Geophys. Res., 112,



D12S09, doi:10.1029/2006JD007606, 2007.

- Satheesh, S. K., Torres, O., Remer, L. A., Babu, S. S., Vinoj, V., Eck, T. F., Kleidman, R. G., and Holben, B. N.: Improved assessment of aerosol absorption using OMI-MODIS joint retrieval, J. Geophys. Res., 114, D05209, doi:10.1029/2008JD011024, 2009.
- Shinozuka, Y. and Redemann, J.: Horizontal variability of aerosol optical depth observed during the ARCTAS airborne experiment, Atmos. Chem. Phys. Discuss., 11, 16245-16264, doi:10.5194/acpd-11-16245-2011, 2011.
  - Schuster, G., MacDonnell, D., and Trepte, C.: Can AERONET ground sites be used to validate CALIPSO aerosol products, CALIPSO ST meeting, 2007.
- Sokal, R. R. and Rohlf, F. J.: Biometry, 3rd Edn., W.H. Freeman and Co., New York, 1995. Torres, O., Tanskanen, A., Veihelmann, B., Ahn, C., Braak, R., Bhartia, P.K., Veefkind, P., and Levelt, P.: Aerosols and surface UV products from Ozone Monitoring Instrument observations: An overview, J. Geophys. Res., 112, D24S47, doi:10.1029/2007JD008809, 2007.
  - Vaughan, M., Kuehn, R., Tackett, J., Rogers, R., Liu, Z., Omar, A., Getzewich, B., Powell, K.,
- Hu, Y., Young, S., Avery, M., Winker, D., and Trepte, C.: Strategies for Improved CALIPSO Aerosol Optical Depth Estimates, Proceedings of the 25th International Laser Radar Conference, 1340–1343, ISBN 978-5-94458-109-9, 2010.
  - Winker, D. M., Hunt, W. H., and McGill, M. J.: Initial performance assessment of CALIOP, Geophys. Res. Lett., 34, L19803, doi:10.1029/2007GL030135, 2007.
- Winker, D. M., Vaughan, M. A., Omar, A. H., Hu, Y., Powell, K. A., Liu, Z., Hunt, W. H., and Young, S. A.: Overview of the CALIPSO Mission and CALIOP Data Processing Algorithms, J. Atmos. Oceanic Technol., 26, 2310–2323, doi:10.1175/2009JTECHA1281.1, 2009.
  - Winker, D. M., Pelon, J., Coakley Jr., J. A., Ackerman, S. A., Charlson, R. J., Colarco, P. R., Flamant, P., Fu, Q., Hoff, R. M., Kittaka, C., Kubar, T. L., Le Treut, H., McCormick, M.
- P., Mégie, G., Poole, L., Powell, K., Trepte, C., Vaughan, M. A., and Wielick, B. A.: The CALIPSO Mission: A Global 3D View Of Aerosols And Clouds, Bull. Am. Meteorol. Soc., 91, 1211–1229, doi:10.1175/2010BAMS3009.1, 2010.
  - Young, S. A. and Vaughan, M. A.: The retrieval of profiles of particulate extinction from Cloud Aerosol Lidar Infrared Pathfinder Satellite Observations (CALIPSO) data: Algorithm descrip-
- tion, J. Atmos. Oceanic Technol., 26, 1105–1119, doi:10.1175/2008JTECHA1221.1, 2009. Zhang, J., Christopher, S. A., Remer, L. A., and Kaufman, Y. J.: Shortwave Aerosol Cloud-Free Radiative Forcing from Terra, II : Global and Seasonal Distributions, J. Geophys. Res., 110, D10S24, doi:10.1029/2004jd005009, 2005.



**Discussion** Paper **ACPD** 11, 22987-23027, 2011 The comparison of **MODIS-Aqua and CALIOP AOD Discussion** Paper J. Redemann et al. Title Page Introduction Abstract Conclusions References **Discussion** Paper Tables Figures 14 ١ ► ◀ Back Close **Discussion Paper** Full Screen / Esc **Printer-friendly Version** Interactive Discussion  $(\mathbf{\hat{H}})$ 

(cc)

Table 1. Data attributes for different AOD comparisons.

Version → Comparison #↓	CALIOP V2 vs MODIS	CALIOP V3 vs MODIS
1	CALIOP V2: all valid ext532 retrievals* (*: from 40 km profile product) MODIS: all valid AOD retrievals	CALIOP V3: all valid ext532 retrievals‡ (‡: from 5 km profile product) MODIS: all valid AOD retrievals
2	CALIOP V2: 0 <ext532<1.25* (*: from 40 km profile product) MODIS: MODISQA=1,2,3</ext532<1.25* 	CALIOP V3: 0 <ext532<1.25‡ (‡: from 5 km profile product) MODIS: MODISQA=1,2,3</ext532<1.25‡ 
3	CALIOP V2: 0 <ext532<1.25* &<br="">cloudOD=0** &amp; extQCflag=0,1,2** (*: from 40 km profile product **: from 5 km layer product) MODIS: MODISQA=1,2,3</ext532<1.25*>	CALIOP V3: $0 < ext532 < 1.25 \ddagger \&$ cloudOD=0 $\ddagger \& extQCflag=0,1,2\ddagger$ Delete profile if any extQCflag $\neq 0,1,2$ ( $\ddagger$ : from 5 km profile product) MODIS: MODISQA=1,2,3
4	CALIOP V2: 0 <ext532<1.25* &<br="">cloudOD=0** &amp; extQCflag=0,1,2** (*: from 40 km profile product **: from 5 km layer product) MODIS: MODISQA=1,2,3 &amp; FOC&lt;0.01</ext532<1.25*>	CALIOP V3: $0 < ext532 < 1.25 \ddagger \&$ cloudOD=0 $\ddagger \& extQCflag=0,1,2 \ddagger$ Delete profile if any extQCflag $\neq 0,1,2$ ( $\ddagger$ : from 5 km profile product) MODIS: MODISQA=1,2,3 & FOC<0.01

Table. 1. Continued.

			P		
Version → Comparison #↓	CALIOP V2 vs MODIS	CALIOP V3 vs MODIS	aper	The comp	parison o
5		CALIOP V3: all valid ext532 retrievals‡, where the atmo- spheric volume descriptor indi-	Dia	MODIS-A CALIO	Aqua and P AOD
		cates "aerosol" or "clear air", except delete profiles if for any	scussi	J. Redem	ann et al.
		ext532: ext532 < -0.2 km <sup>-1</sup> &&	on F		
		$ unc_{ext}532  > 0.4 \text{ km}^{-1}$	ape	Title	Page
		or ext532 > 2.5 km <sup>-1</sup> && km/ma_art522 > 500 %	br	Abstract	Introductio
		cloudOD=0‡	D	Conclusions	Reference
		(‡: from 5 km profile product) MODIS: MODISQA=1,2,3	SCUS	Tables	Figures
6		CALIOP V3: all valid col_AOD retrievals <sup>±</sup> .	sion	14	۲I
		except delete profiles if for any	Pap		
		ext532: rel.err>100% where	)er		•
		rel_err = unc_col_AOD/col_AOD	—	Back	Close
		(‡: from 5 km profile product) cloudOD=0‡	Die	Full Scre	een / Esc
		(‡: from 5 km profile product) MODIS: MODISQA=1,2,3	CUSS	Printer-frier	ndly Version
			on		
			Dap	Interactive	Discussion
			er	6	•

**ACPD** 

11, 22987-23027, 2011

Discussion



AC	ACPD						
11, 22987–	11, 22987–23027, 2011						
The comparison of MODIS-Aqua and CALIOP AOD							
J. Redem	nann et al.						
Title	Page						
Abstract	Introduction						
Conclusions	References						
Tables	Figures						
I	►I						
•	•						
Back	Close						
Full Scr	een / Esc						
Printer-friendly Version							
Interactive	Discussion						
œ	BY						

**Discussion** Paper

**Discussion** Paper

**Discussion** Paper

**Discussion** Paper

 Table 2. MODIS vs. CALIOP AOD532, January 2007, over-ocean.

	MOD mean	CAL mean	MeanDiff./ StdDiff.	Num.	R <sup>2</sup>	Slope/ offset	RMS Abs/rel
CALIOP Version 2							
Comparison #1 Comparison #2 Comparison #3 Comparison #4	0.180 0.180 0.131 0.147	0.180 0.132 0.137 0.120	0.0003/0.330 0.048/0.180 -0.007/0.124 0.027/0.082	36859 36758 5750 1883	0.043 0.147 0.342 0.655	1.300/-0.054 0.923/-0.034 1.065/-0.002 0.917/-0.015	0.330/183.2 % 0.186/103.6 % 0.124/94.6 % 0.087/58.9 %
CALIOP Version 3							
Comparison #1 Comparison #2 Comparison #3 Comparison #4 Comparison #5 Comparison #6	0.170 0.170 0.147 0.147 0.147 0.152	0.123 0.111 0.105 0.092 0.112 0.109	0.047/0.189 0.059/0.127 0.042/0.093 0.055/0.085 0.036/0.130 0.043/0.124	247976 247951 177741 37740 177743 176378	0.247 0.390 0.474 0.635 0.382 0.375	1.258/-0.091 0.859/-0.035 1.080/-0.054 0.875/-0.037 1.408/-0.096 1.091/-0.057	0.194/114.0% 0.140/82.3% 0.102/69.6% 0.101/68.9% 0.135/91.9% 0.131/86.4%

Discussion Pa	AC 11, 22987–2	<b>PD</b> 23027, 2011
Iper   Discussion	The comp MODIS-A CALIO J. Redem	parison of Iqua and P AOD ann et al.
Pap	Title	Page
er	Abstract	Introduction
_	Conclusions	References
iscussi	Tables	Figures
on P	I	►I.
aper	•	Þ
_	Back	Close
Discus	Full Scre	en / Esc
sion	Printer-frien	dly Version
Paper		

Table 3. MODIS vs. CALIOP AOD532, April 2007, over-ocean.

	MOD mean	CAL mean	MeanDiff./ StdDiff.	Num.	R <sup>2</sup>	Slope/ offset	RMS Abs/rel
CALIOP Version 2							
Comparison #1 Comparison #2 Comparison #3 Comparison #4	0.169 0.169 0.126 0.141	0.164 0.123 0.124 0.112	0.005/0.308 0.046/0.160 0.003/0.121 0.029/0.079	34024 33943 3426 805	0.032 0.095 0.214 0.632	1.356/-0.065 1.002/-0.046 1.164/-0.023 1.018/-0.032	0.308/182.3 % 0.166/98.5 % 0.121/95.7 % 0.084/59.7 %
CALIOP Version 3							
Comparison #1 Comparison #2 Comparison #3 Comparison #4	0.156 0.156 0.140 0.144	0.112 0.104 0.101 0.099	0.044/0.164 0.052/0.113 0.039/0.089 0.046/0.083	221680 221650 158534 32047	0.187 0.295 0.424 0.586	1.322/-0.094 0.964/-0.046 1.090/-0.051 1.010/-0.047	0.170/109.1 % 0.124/79.5 % 0.097/69.6 % 0.095/65.9 %

AC	ACPD							
11, 22987–	11, 22987–23027, 2011							
The com MODIS-/ CALIC	The comparison of MODIS-Aqua and CALIOP AOD							
J. Reden	nann et al.							
Title	Page							
Abstract	Introduction							
Conclusions	References							
Tables	Figures							
14	۶I							
•	•							
Back	Close							
Full Scr	Full Screen / Esc							
Printer-frie	Drinter friendly Version							
	Printer-friendly Version							
Interactive	DISCUSSION							
	<b>O</b> BY							

**Discussion** Paper

**Discussion Paper** 

**Discussion** Paper

**Discussion** Paper

Table 4. MODIS vs. CALIOP AOD532, July 2007, over-ocean.

	MOD mean	CAL mean	MeanDiff./ StdDiff.	Num.	R <sup>2</sup>	Slope/ offset	RMS Abs/rel
CALIOP Version 2							
Comparison #1 Comparison #2 Comparison #3 Comparison #4 CALIOP Version 3	0.149 0.149 0.105 0.100	0.162 0.115 0.118 0.106	-0.013/0.320 0.034/0.144 -0.013/0.111 -0.006/0.077	23121 23051 3245 1035	0.024 0.148 0.299 0.571	1.356/-0.040 1.015/-0.036 1.064/0.006 1.123/-0.006	0.320/215.4 % 0.147/99.2 % 0.111/106.1 % 0.077/77.4 %
Comparison #1 Comparison #2 Comparison #3 Comparison #4	0.131 0.131 0.117 0.100	0.099 0.091 0.087 0.069	0.032/0.157 0.040/0.100 0.030/0.084 0.031/0.082	156614 156592 121252 20392	0.091 0.221 0.248 0.316	1.272/-0.067 0.938/-0.032 0.997/-0.030 0.829/-0.014	0.160/122.4 % 0.108/82.2 % 0.089/75.9 % 0.088/87.5 %

Discussion Pa	<b>ACPD</b> 11, 22987–23027, 2011						
per   Di	The comparison of MODIS-Aqua and CALIOP AOD						
scussion	J. Redem	ann et al.					
Pap	Title Page						
	Abstract	Introduction					
	Conclusions	References					
iscuss	Tables	Figures					
ion P	14	►I					
aper	•	•					
	Back	Close					
Discu	Full Scre	en / Esc					
ISSIO	Printer-frier	dly Version					
n Pap	Interactive	Discussion					
ber	$\odot$	BY					

 Table 5. MODIS vs. CALIOP AOD532, October 2007, over-ocean.

	MOD mean	CAL mean	MeanDiff./ StdDiff.	Num.	R <sup>2</sup>	Slope/ offset	RMS Abs/rel
CALIOP Version 2							
Comparison #1 Comparison #2 Comparison #3 Comparison #4 CALIOP Version 3	0.149 0.149 0.115 0.123	0.161 0.114 0.113 0.094	-0.013/0.309 0.035/0.141 0.002/0.103 0.029/0.064	35345 35275 5814 2129	0.016 0.058 0.231 0.577	1.387/-0.045 1.109/-0.051 1.211/-0.026 1.000/-0.029	0.309/208.0 % 0.145/97.6 % 0.103/89.8 % 0.070/57.2 %
Comparison #1 Comparison #2 Comparison #3 Comparison #4	0.141 0.141 0.129 0.124	0.102 0.094 0.092 0.079	0.038/0.154 0.047/0.098 0.037/0.080 0.046/0.068	243572 243539 183311 39198	0.112 0.212 0.307 0.487	1.391/-0.093 1.029/-0.051 1.142/-0.056 0.987/-0.044	0.159/112.7 % 0.109/77.1 % 0.088/68.2 % 0.082/65.8 %

	MOD mean	CAL mean	MeanDiff./ StdDiff.	Num.	R <sup>2</sup>	Slope/ offset	RMS Abs/rel
CALIOP Version 2							
Comparison #1	0.153	0.149	0.004/0.349	37824	0.008	1.257/-0.043	0.349/228.3%
Comparison #2	0.153	0.092	0.061/0.130	37715	0.102	0.931/-0.051	0.144/94.2%
Comparison #3	0.122	0.087	0.035/0.091	4689	0.357	0.834/-0.015	0.098/80.1 %
Comparison #4	0.128	0.082	0.046/0.074	1632	0.564	0.725/-0.011	0.087/68.3%
CALIOP Version 3							
Comparison #1	0.144	0.108	0.036/0.155	244148	0.170	1.365/-0.088	0.160/110.6 %
Comparison #2	0.144	0.099	0.045/0.104	244136	0.278	0.994/-0.044	0.113/78.3%
Comparison #3	0.130	0.097	0.032/0.091	180852	0.332	1.093/-0.045	0.096/74.1 %
Comparison #4	0.122	0.077	0.044/0.074	35324	0.484	0.895/-0.032	0.087/71.1%



Discussion Pa	<b>ACPD</b> 11, 22987–23027, 2011						
Iper   Discussion	The comp MODIS-A CALIO J. Redem	The comparison of MODIS-Aqua and CALIOP AOD J. Redemann et al.					
Pap	Title	Page					
er	Abstract	Introduction					
_	Conclusions	References					
iscussi	Tables	Figures					
on P	I	►I.					
aper	•	Þ					
_	Back	Close					
Discus	Full Scre	en / Esc					
sion	Printer-frien	aly version					
Paper							

Table 7. MODIS vs. CALIOP AOD532, April 2009, over-ocean.

	MOD mean	CAL mean	MeanDiff./ StdDiff.	Num.	R <sup>2</sup>	Slope/ offset	RMS Abs/rel
CALIOP Version 2							
Comparison #1 Comparison #2 Comparison #3 Comparison #4	0.165 0.165 0.141 0.156	0.153 0.100 0.111 0.118	0.012/0.343 0.065/0.145 0.030/0.094 0.038/0.083	38617 38543 4835 2169	0.015 0.129 0.465 0.575	1.265/-0.056 0.895/-0.047 1.025/-0.034 1.079/-0.050	0.343/207.8 % 0.159/96.4 % 0.098/69.9 % 0.092/58.6 %
CALIOP Version 3							
Comparison #1 Comparison #2 Comparison #3 Comparison #4	0.154 0.154 0.138 0.147	0.113 0.105 0.104 0.107	0.041/0.163 0.050/0.116 0.034/0.096 0.040/0.092	246057 246049 179128 36121	0.189 0.288 0.393 0.568	1.322/-0.091 1.008/-0.051 1.173/-0.058 1.161/-0.063	0.168/109.1 % 0.126/81.6 % 0.102/73.7 % 0.100/67.9 %

ACPD									
11, 22987–23027, 2011									
The comparison of MODIS-Aqua and CALIOP AOD									
Title	Page								
Abstract	Abstract Introduction								
Conclusions	References								
Tables	Tables Figures								
I	►I								
•	•								
Back	Back Close								
Full Scr	Full Screen / Esc								
Printer-friendly Version									
Interactive	Interactive Discussion								

**Discussion** Paper

**Discussion** Paper

**Discussion** Paper

**Discussion** Paper

Table 8. MODIS vs. CALIOP AOD532, July 2009, over-ocean.

	MOD mean	CAL mean	MeanDiff./ StdDiff.	Num.	R <sup>2</sup>	Slope/ offset	RMS Abs/rel
CALIOP Version 2							
Comparison #1 Comparison #2 Comparison #3 Comparison #4 CALIOP Version 3	0.168 0.168 0.124 0.116	0.159 0.110 0.096 0.078	0.008/0.351 0.057/0.179 0.028/0.183 0.038/0.098	27077 27026 3959 1219	0.016 0.082 0.062 0.368	1.211/-0.044 0.903/-0.041 0.819/-0.006 0.876/-0.024	0.351/209.6 % 0.188/112.1 % 0.185/149.7 % 0.105/91.0 %
Comparison #1 Comparison #2 Comparison #3 Comparison #4	0.142 0.142 0.127 0.117	0.103 0.097 0.093 0.077	0.039/0.159 0.045/0.116 0.034/0.097 0.040/0.110	167115 167105 131583 22652	0.124 0.224 0.234 0.296	1.157/-0.061 0.873/-0.027 0.973/-0.031 0.862/-0.024	0.164/115.5 % 0.125/88.0 % 0.103/80.8 % 0.117/99.9 %

	MOD mean	CAL mean	MeanDiff./ StdDiff.	Num.	R <sup>2</sup>	Slope/ offset	RMS Abs/rel
CALIOP Version 2							
Comparison #1 Comparison #2 Comparison #3 Comparison #4	0.154 0.154 0.126 0.142	0.156 0.110 0.123 0.121	-0.002/0.316 0.044/0.136 0.003/0.103 0.021/0.077	40742 40650 6645 2344	0.022 0.111 0.411 0.691	1.440/-0.065 1.100/-0.060 1.290/-0.040 1.255/-0.057	0.316/205.0 % 0.143/93.0 % 0.103/81.5 % 0.079/55.8 %
CALIOP Version 3							
Comparison #1 Comparison #2 Comparison #3 Comparison #4	0.142 0.143 0.131 0.133	0.104 0.099 0.098 0.097	0.038/0.136 0.044/0.095 0.033/0.080 0.036/0.072	267050 258462 206608 42350	0.185 0.313 0.425 0.625	1.320/-0.084 1.022/-0.048 1.148/-0.052 1.088/-0.048	0.142/ 99.6 % 0.105/73.6 % 0.087/66.2 % 0.081/60.9 %

Table 9. MODIS vs. CALIOP AOD532, October 2009, over-ocean.



**Discussion** Paper

**Discussion** Paper



**Fig. 1.** Geometry for comparing MODIS AOD retrievals (solid boxes) to CALIOP V2 (40 kmaverage) and V3 (5 km-average) AOD retrievals. Black boxes around V3 retrieval locations represent the optimal sampling of the MODIS AOD retrievals (see text).





**Fig. 2.** (a) Comparison of all instantaneously collocated MODIS AOD retrievals for January of 2007 to AOD calculated from CALIOP V2 aerosol extinction profiles. (b) Same as (a), but comparison screened with the methodology summarized as comparison # 3 in Table 1. (c) same as a), but comparison screened with the methodology summarized as comparison # 4 in Table 1. (d), (e) and (f) are the same as (a), (b) and (c), respectively, but for CALIOP V3.







**Fig. 3. (a)** Left – comparison of agreement between MODIS AOD and CALIOP V2 and V3 AOD, respectively, in terms of difference in mean AOD, square of the linear correlation coefficient, and rms difference, for four months in 2007. Right – same as left, but for data after application of comparison # 3 in Table 1. (b) Same as (a), but for four months in 2009.











**Fig. 5.** Latitudinal distribution of the mean difference between MODIS and CALIOP V3 AOD (using comparison # 3 screening from Table 1) over-ocean (blue lines) and over-lean (red lines), with thin lines representing 1-degree averages and thick lines indicating 5-degree averages, for all 8 months considered in this study.





**Fig. 6.** Same as Fig. 5, but for the more severely cloud-screened data, as described by comparison # 4 in Table 1.

23025



**Fig. 7.** (a) Global distribution of the number of instantaneously collocated MODIS versus CALIOP V3 AOD comparisons in 5 × 5 degree boxes for January 2007. (e) Global distribution of the mean difference between instantaneously collocated MODIS and CALIOP V3 AOD retrievals for January 2007. (b–d) same as (a), but for April, July, and October of 2007. (f–h) same as (e), but for April, July, and October of 2007.





Fig. 8. Same as Fig. 7, but for data collected in 2009. 23027

