

**Pyridine-containing  
clusters reacting with  
NH<sub>3</sub>**

M. J. Ryding et al.

**Reactions of H<sup>+</sup>(pyridine)<sub>m</sub>(H<sub>2</sub>O)<sub>n</sub> and  
H<sup>+</sup>(NH<sub>3</sub>)<sub>1</sub>(pyridine)<sub>1</sub>(H<sub>2</sub>O)<sub>n</sub> with NH<sub>3</sub>:  
experiments and kinetic modelling under  
tropospheric conditions**

M. J. Ryding<sup>1</sup>, Å. M. Jonsson<sup>2</sup>, A. S. Zatula<sup>3</sup>, P. U. Andersson<sup>1</sup>, and E. Uggerud<sup>3</sup>

<sup>1</sup>Department of Chemistry, Atmospheric Science, University of Gothenburg, 412 96 Göteborg, Sweden

<sup>2</sup>IVL Swedish Environmental Research Institute Ltd., Box 5302, 400 14 Göteborg, Sweden

<sup>3</sup>Mass Spectrometry Laboratory and Centre for Theoretical and Computational Chemistry, Department of Chemistry, University of Oslo, P.O. Box 1033 Blindern, 0315 Oslo, Norway

Received: 23 June 2011 – Accepted: 22 August 2011 – Published: 1 September 2011

Correspondence to: P. U. Andersson (pan@chem.gu.se)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Reactions between pyridine containing water cluster ions,  $\text{H}^+(\text{pyridine})_1(\text{H}_2\text{O})_n$ ,  $\text{H}^+(\text{pyridine})_2(\text{H}_2\text{O})_n$  and  $\text{H}^+(\text{NH}_3)_1(\text{pyridine})_1(\text{H}_2\text{O})_n$  ( $n$  up to 15) with  $\text{NH}_3$  have been studied experimentally using a quadrupole time-of-flight mass spectrometer. The product ions in the reaction between  $\text{H}^+(\text{pyridine})_m(\text{H}_2\text{O})_n$  and  $\text{NH}_3$  have been determined for the first time. It is found that the reaction mainly leads to cluster ions of the form  $\text{H}^+(\text{NH}_3)_1(\text{pyridine})_1(\text{H}_2\text{O})_{n-x}$ , with  $x = 1$  or  $2$  depending on the initial size of the reacting cluster ion. For a given number of water molecules (from 5 to 15) in the cluster ion, rate coefficients are found to be slightly lower than those for protonated pure water clusters reacting with ammonia. The rate coefficients obtained from this study are used in a kinetic cluster ion model under tropospheric conditions. The results from the model suggest that cluster ions containing ammonia and more than one pyridine, picoline or lutidine molecule should dominate at ground level under typical conditions.

## 1 Introduction

Atmospheric aerosols can either be of primary origin, i.e. emitted to the atmosphere from anthropogenic or natural sources, or they can be of secondary origin, i.e. grown from molecules pre-existing in the atmosphere. The first steps of the secondary formation mechanism are unclear; suggestions include binary or ternary homogeneous nucleation as well as ion-induced nucleation followed by ion-ion recombination (Kulmala et al., 2004). Atmospheric ions are initially formed by solar radiation, galactic cosmic rays and radioactive decay. The ions are found in the entire atmosphere, although the formation mechanisms vary with altitude, region and time of day. The main products of the ionisation of air are  $\text{O}_2^+$ ,  $\text{NO}^+$  and free electrons (Wayne, 2000). Attachment of neutral polar molecules to the ions leads to charged molecular clusters. Subsequent to cluster formation and growth, recombination of positively and negatively charged cluster ions is possible. For small clusters, this is believed to be associated

## Pyridine-containing clusters reacting with $\text{NH}_3$

M. J. Ryding et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Pyridine-containing clusters reacting with $\text{NH}_3$

M. J. Ryding et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with extensive fragmentation; for large clusters coalescence may occur. The neutral clusters formed by the recombination are considered large enough to continue to grow spontaneously by condensation (Yu and Turco, 2000; Yu, 2003). The charged clusters formed by ion-induced nucleation are more stable than the corresponding neutral clusters; hence this process is more effective than the homogeneous nucleation. Even so, it is estimated that the major part of the total atmospheric particle formation has its origin in homogeneous nucleation, since the neutral clusters are thought to be 10 to 100 times as numerous as the ionic clusters (Kulmala et al., 2007).

Several air ion mobility measurements have identified cluster ions in the troposphere. However, the chemical nature of these cluster ions is often difficult to identify (Horrak et al., 2000; Vana et al., 2008). By contrast, there have been measurements and identification of molecule ions in the troposphere during the last two decades. A large fraction of the molecule ions observed in these studies has likely originated from cluster ions that fragment before mass analysis (Eisele, 1983, 1986, 1988; Eisele and McDaniel, 1986; Eisele and Tanner, 1990; Schulte and Arnold, 1990). The first ground based measurement of atmospheric ion composition was performed by Perkins and Eisele in 1983. In this study, several unidentified positive ions were observed (Perkins and Eisele, 1984). Improved measurements conducted a few years later revealed the unidentified ions that had a mass-to-charge ratio of 80, 94 and 108 to be protonated pyridine ( $\text{C}_5\text{H}_5\text{NH}^+$ ), protonated picoline (methyl-pyridine) and protonated lutidine (dimethyl-pyridine), respectively (Eisele, 1988). Several other ions have been identified in the troposphere in addition to these, although pyridinium is often found to dominate the mass spectrum. For example, Schulte and Arnold (1990) identified pyridinium as the dominating ion in air-plane based measurements in the free troposphere over Europe. Recently, Junninen et al. (2010) measured day-time air ions at an urban site (the SMEAR III station in Helsinki), using an Atmospheric Pressure Interface Time-of-Flight instrument. They identified protonated poly(alkyl) pyridines as one of the main positive compound types. Ehn et al. (2010) measured day and night-time air ions at a remote site (the SMEAR II station in Hyytiälä) using the same instrument. They

observed pyridine ions and alkyl substituted pyridine ions in both the day- and night-time ion spectra, with approximately a factor two higher concentration during night-time.

Sources of atmospheric pyridine and pyridine derivatives are supposed to be biomass burning, automobile exhaust, coal tars and tobacco smoke (Clemons, 1973; Saintjalm and Moreetesta, 1980; Beig, 2008). The main atmospheric sink is considered to be reaction with OH radicals (Eisele, 1986, 1988; Atkinson et al., 1987; Yeung and Elrod, 2003). Yeung and Elrod (2003) calculated atmospheric lifetimes based on experimentally determined reaction rate coefficients for pyridine and for various substituted pyridine compounds to be 44 days and around 1 to 10 days, respectively. Other suggested atmospheric sinks of significance are reaction with HNO<sub>3</sub> in polluted environments (Atkinson et al., 1987) and reaction with atomic chlorine (Zhao et al., 2007). Due to the localised and sometimes irregular nature of the sources – as well as the relatively short atmospheric lifetimes – the concentration of pyridine is expected to be highly variable with time and location (Beig and Brasseur, 2000; Yeung and Elrod, 2003). Few measurements of pyridine concentrations in the atmosphere exist. Among these, Tanner and Eisel (1991) measured a concentration of about 2.5 ppt ± 50 % ( $6.2 \times 10^7 \text{ cm}^{-3}$ ) of molecular pyridine at Mauna Loa Observatory, Hawaii. In the measurements by Junninen et al. (2010) about  $1 \text{ cm}^{-3}$  of protonated pyridine was observed at the urban SMEAR III station in Helsinki. However, in contrast to most previous measurements they found up to six times higher concentrations for ionic alkyl substituted pyridine compounds  $\text{H}^+\text{C}_5\text{H}_5\text{N}(\text{CH}_2)_n$ ,  $1 \leq n \leq 6$ , including picoline and lutidine. The reason for this is unknown but interesting and the findings show that there is a need to better understand the atmospheric chemistry of these compounds. Ehn et al. (2010) reported average concentrations of pyridinium and alkyl substituted pyridine ions from the SMEAR II station in Hyytiälä during 4 days in early May 2009: pyridinium  $36.4 \text{ cm}^{-3}$ , picolinium  $57.3 \text{ cm}^{-3}$ , lutidinium  $33.5 \text{ cm}^{-3}$ . Also in this case, alkyl substituted pyridine ions are higher in concentration than the pyridine ion.

A kinetic cluster ion model by Beig and Brasseur (2000) indicate that pyridine containing clusters may be the dominating positive ions in the lower free troposphere (from

## Pyridine-containing clusters reacting with NH<sub>3</sub>

M. J. Ryding et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version


Interactive Discussion



1 to 6 km above ground). More specifically, the pyridinated cluster ions in the model were  $H^+(X)_1(H_2O)_n$  and  $H^+(NH_3)_m(X)_1(H_2O)_n$ , where X = pyridine, picoline or lutidine. Beig and Brasseur proposed two reaction pathways for the formation of these pyridinated cluster ions from  $H^+(H_2O)_n$  clusters. The first reaction pathway starts with addition of  $NH_3$  to a protonated water cluster. The formed cluster can thereafter react with a pyridine type molecule X:



In the second reaction pathway, a pyridine molecule reacts with a protonated water cluster. The pyridine is thereafter ejected when an ammonia molecule attaches to the cluster in a second step:



The second step can thereafter be followed by Reaction (R1b) above to form a cluster containing both ammonia and pyridine. The driving force behind these reactions – forming cluster ions containing ammonia and pyridine derivatives – appears connected to the high basicities of ammonia and the pyridine derivatives. Note that loss of water is likely to occur also in the first step of the first mechanism (Reaction R1a) and in the second step of the second mechanism (Reaction R2b); however, this was not included in the notation by Beig and Brasseur.

While the reaction rate coefficients are known for the first three reactions in the case of X = pyridine, it is unknown for Reaction (R2b) (Viggiano et al., 1988a, b). Beig and Brasseur assumed  $10^{-11} \text{ cm}^3 \text{ s}^{-1}$  as an upper limit for the rate coefficient of Reaction (R2b) for all pyridine derivatives in their study, which is two orders of magnitude lower than the rate coefficient for Reaction (R1b) at 298 K.

## Pyridine-containing clusters reacting with $NH_3$

M. J. Ryding et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

In the present work we investigate the reactions of two types of cluster ions with  $\text{NH}_3$  in a cluster beam experiment; the clusters being  $\text{H}^+(\text{pyridine})_m(\text{H}_2\text{O})_n$  ( $m = 1$  and  $2$ ) and  $\text{H}^+(\text{NH}_3)_1(\text{pyridine})_1(\text{H}_2\text{O})_n$ . The results from the experiments are thereafter used to improve the present kinetic model by Beig and Brasseur for atmospheric positive ions. The atmospheric implications of the experimental results and the results from the kinetic modelling are discussed.

## 2 Methods

### 2.1 Experimental

The experiments were performed using a modified QTOF 2 mass spectrometer (Micro-mass/Waters, Manchester UK). The instrument and experimental procedure has been described in detail previously (Andersson et al., 2008; Ryding et al., 2011), and therefore only a brief overview is presented here. The cluster ions were produced from aqueous solutions at atmospheric pressure by electrospray ionization (ESI) and thereafter entered into the high vacuum part of the instrument. Two different solutions were used: 2.5 mM pyridine(aq) for production of  $\text{H}^+(\text{pyridine})_m(\text{H}_2\text{O})_n$  ions and a mixture with 2.5 mM pyridine(aq) and 30 mM  $\text{NH}_3$ (aq) for production of  $\text{H}^+(\text{NH}_3)_1(\text{pyridine})_1(\text{H}_2\text{O})_n$  ions. A quadrupole mass filter (with better than unit resolution) allowed for selection of single sized clusters based on their mass-to-charge ratio,  $m/z$ . The selected clusters then entered the collision cell, where they were brought to collide with gaseous ammonia at a centre-of-mass (COM) collision energy  $E_{\text{COM}} = 8 \text{ kJmol}^{-1}$  (0.085 eV). The resulting products were analyzed by a reflectron time-of-flight (TOF) mass analyzer set at a mass resolution,  $m/\Delta m$ , of about 5000 (full-width-half-height). The ammonia was introduced into the collision cell through an Ultra-High-Vacuum (UHV) leak valve giving a pressure of about  $10^{-5}$  mbar. This pressure of  $\text{NH}_3$  was chosen to keep the number of ion-neutral collisions below 10% to ensure single-collision conditions for the entire range of cluster sizes. Reference measurements were collected by

## Pyridine-containing clusters reacting with $\text{NH}_3$

M. J. Ryding et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



measuring the cluster ion  $\text{H}^+(\text{pyridine})_1(\text{H}_2\text{O})_{11}$  at regular intervals during the experiment to make sure that the collision gas pressure was constant. For each measurement, a corresponding background measurement was collected with an empty collision cell. For the cluster ions  $\text{H}^+(\text{pyridine})_1(\text{H}_2\text{O})_{10}$  and  $\text{H}^+(\text{NH}_3)_1(\text{pyridine})_1(\text{H}_2\text{O})_{10}$  measurements were also performed at various collision energies between  $1.45 \text{ kJ mol}^{-1}$  and  $120 \text{ kJ mol}^{-1}$  (COM). Reagents used in the experiments:  $\text{H}_2\text{O}$  (no. 95270 for HPLC, Fluka), pyridine (99.5 %, BDH Chemicals Ltd.),  $\text{NH}_3$  (99.96 %, AGA), 25 %  $\text{NH}_3$  (aq) (Pro analysi, Merck).

## 2.2 Positive ion model description

The positive cluster ion reaction schemes in this paper are based on the aforementioned model by Beig and Brasseur (2000). In their model, protonated water clusters,  $\text{H}^+(\text{H}_2\text{O})_n$  are continuously produced in a series of reactions starting from  $\text{O}_2^+$  and  $\text{N}_2^+$ . Subsequent to this, the protonated water clusters may then react with ammonia, pyridine, picoline, lutidine, acetone and acetonitrile. The production of cluster ions is balanced by the loss of cluster ions through ion-ion recombination and attachment to aerosol particles. In our first model (Model A), we have made two modifications to the model by Beig and Brasseur. Firstly, we have included the experimental results from this study. That is, in the reaction between  $\text{H}^+(\text{pyridine})_{1,2}(\text{H}_2\text{O})_n$  and  $\text{NH}_3$  virtually no exchange of pyridine for ammonia will take place. Instead, ammonia is incorporated into the cluster ion with subsequent loss of one or two water molecules, i.e.  $\text{H}^+(\text{pyridine})_1(\text{H}_2\text{O})_n + \text{NH}_3 \rightarrow \text{H}^+(\text{NH}_3)_1(\text{pyridine})_1(\text{H}_2\text{O})_{n-x} + x\text{H}_2\text{O}$ . Hence, the former reaction was excluded from the model and the latter was added (the corresponding modifications were made for the reactions with clusters containing picoline and lutidine). For simplicity, the rate coefficient for  $\text{H}^+(\text{pyridine})_1(\text{H}_2\text{O})_4 + \text{NH}_3$  (obtained in this study) was used for all pyridine/water clusters. Secondly, we have included reactions leading to clusters containing two amines (pyridine, picoline and lutidine). The reactions included in Model A are shown in Fig. 1 and the corresponding rate coefficients

## Pyridine-containing clusters reacting with $\text{NH}_3$

M. J. Ryding et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



are given in Table 1. In our second model, Model B, we have omitted the amines picoline and lutidine since pyridine containing cluster ions are found to be the dominating cluster ions in Model A. We also allowed for up to five pyridine molecules in each cluster. The reactions included in this model are shown in Fig. 2 and the corresponding rate coefficients are given in Table 1. In the absence of detailed data for evaporation of pyridine from protonated water clusters and since we do not observe evaporation of pyridine under our experimental conditions (see Sect. 3.1), we did not include evaporation in our models presented in this study. The differential rate equations were solved in the program FACSIMILE (2007). All model calculations in this paper were performed under ground level conditions at a temperature of 298 K. Steady-state concentrations were reached after approximately 500 s.

### 3 Results

#### 3.1 Experimental results

Figure 3a shows the TOF mass spectrum obtained for mass selected (by the quadrupole)  $\text{H}^+(\text{pyridine})_1(\text{H}_2\text{O})_{11}$  ions passing an empty collision cell. The intensity of the parent ion dominates but minor peaks corresponding to loss of 1 to 5 water molecules are also seen. The loss of water is caused by evaporation from the parent cluster during the flight time of about  $160 \mu\text{s}$  from the quadrupole to the TOF unit. Evaporation is inevitable for the reactions studied here, applying this experimental setup including the setting of the collision energy. In agreement with a previous study by Ryding et al. (2011), evaporation of pyridine does not occur. Figure 3b shows the TOF mass spectrum obtained after reaction between the cluster ion  $\text{H}^+(\text{pyridine})_1(\text{H}_2\text{O})_{11}$  and  $\text{NH}_3$  at a collision energy of  $8 \text{ kJ mol}^{-1}$  (COM). Evaporation of water molecules during the passage through the QTOF 2 is found in approximately equal amounts as in the background measurement. That is, the area of the peaks at  $m/z - 18$ ,  $-36$  and  $-54$  relative the parent ion is basically unaffected by the presence of  $\text{NH}_3$  in the collision cell. For the reaction with  $\text{NH}_3$ , the main product formed is  $\text{H}^+(\text{NH}_3)_1(\text{pyridine})_1(\text{H}_2\text{O})_9$

## Pyridine-containing clusters reacting with $\text{NH}_3$

M. J. Ryding et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Pyridine-containing clusters reacting with NH<sub>3</sub>

M. J. Ryding et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



at  $-19u$  relative the parent ion mass. This means that NH<sub>3</sub> is incorporated into the cluster and typically two H<sub>2</sub>O molecules evaporate from the short-lived collision complex formed. Product clusters corresponding to loss of one, three, four and five H<sub>2</sub>O molecules are also seen; however, these products are significantly less abundant.

5 Clusters H<sup>+</sup>(NH<sub>3</sub>)<sub>2</sub>(pyridine)<sub>1</sub>(H<sub>2</sub>O)<sub>*n*</sub> are produced in small amounts due to multiple collisions with NH<sub>3</sub>. Essentially no pyridine is lost from the cluster ions upon reaction, as seen by the absence of peaks at the indicated locations in Fig. 3b.

Due to the above mentioned evaporation of H<sub>2</sub>O molecules from the parent ion, some of the detected products will have originated from reactions of pre-  
10 formed evaporation products. In order to estimate the contribution of these reactions, we devised the following simple model. The  $m/z - 1$  peak represents the reaction  $A^+(H_2O)_n + NH_3 \rightarrow A^+(NH_3)_1(H_2O)_{n-1} + H_2O$  (with  $A^+ = H^+$ ,  $H^+(\text{pyridine})_1$ ,  $H^+(\text{pyridine})_2$  or  $H^+(NH_3)_1(\text{pyridine})_1$ ). We assume that all of the evaporation products  $A^+(H_2O)_{n-x}$ ,  $x = 1, 2, 3\dots$  in a mass spectrum would form the products  
15  $A^+(NH_3)_1(H_2O)_{n-x-1}$ ,  $x = 1, 2, 3\dots$  to a degree that corresponds to the relative intensity of the  $m/z - 1$  peak in the mass spectrum of the correspondingly sized parent ion. This gives a small contribution from the  $A^+(H_2O)_{n-1}$  cluster to the  $A^+(NH_3)_1(H_2O)_{n-2}$  peak, located at  $m/z - 19$ . The remaining intensity in this peak originate from the parent ion reaction  $A^+(H_2O)_n + NH_3 \rightarrow A^+(NH_3)_1(H_2O)_{n-2} + 2H_2O$ . We calculated the magnitude  
20 of this reaction for all parent ion cluster sizes, and applied it to the correspondingly sized evaporation products as well. We were then able to estimate the magnitude of the third parent ion reaction:  $A^+(H_2O)_n + NH_3 \rightarrow A^+(NH_3)_1(H_2O)_{n-3} + 3H_2O$  in the same way.

Figure 4 shows the branching ratios of different products from the  
25  $H^+(\text{pyridine})_m(H_2O)_n + NH_3$  reaction ( $m = 0$  to  $2$ ) at  $E_{COM} = 8 \text{ kJmol}^{-1}$ . For  $H^+(H_2O)_n + NH_3$  the main product peaks represent incorporation of the NH<sub>3</sub> molecule and loss of two or three water molecules. The simple model described above attributes the intensity of these peaks to reactions of the parent ion (by 100% and  $\geq 90\%$ , respectively). Peaks corresponding to addition of the ammonia molecule and loss of

## Pyridine-containing clusters reacting with NH<sub>3</sub>

M. J. Ryding et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



four and five H<sub>2</sub>O from the parent cluster can be seen; they tend to be higher for the larger clusters. A significant part of the abundance of these products is likely resulting from reactions of evaporation products; they have been included for comparison. Loss of a single water molecule after reaction with NH<sub>3</sub> is effectively not observed for this cluster type.

The branching ratios of the H<sup>+</sup>(pyridine)<sub>1</sub>(H<sub>2</sub>O)<sub>*n*</sub> and H<sup>+</sup>(pyridine)<sub>2</sub>(H<sub>2</sub>O)<sub>*n*</sub> clusters are rather similar, and are dominated by loss of two water molecules after addition of the ammonia molecule. For smaller clusters, loss of a single water molecule occurs with a frequency similar to the loss of two water molecules. However, the –H<sub>2</sub>O curve drops off with size, giving about an order of magnitude lower abundance compared to the –2H<sub>2</sub>O products as *n* approaches 15. Both of these peaks are due to parent ion reactions, while the peaks corresponding to loss of three or four water molecules after addition of ammonia is likely to contain large contributions from reactions of evaporation clusters. Again, we notice that virtually no pyridine leaves the cluster ions after reaction with ammonia, neither for H<sup>+</sup>(pyridine)<sub>1</sub>(H<sub>2</sub>O)<sub>*n*</sub> nor for H<sup>+</sup>(pyridine)<sub>2</sub>(H<sub>2</sub>O)<sub>*n*</sub>. This is in contrast to the assumption made by Beig and Brasseur (2000).

Branching ratios for the H<sup>+</sup>(NH<sub>3</sub>)<sub>1</sub>(pyridine)<sub>1</sub>(H<sub>2</sub>O)<sub>*n*</sub> + NH<sub>3</sub> reaction are shown in Fig. 5 for *n* = 1 to 15. As seen, the products are dominated by incorporation of the reactant NH<sub>3</sub> and loss of one or two water molecules in more or less equal amounts, except for *n* = 2. Again, these reactions can be attributed to the parent ion by 100 % and ≥90 %, respectively. Products with three or four water molecules fewer than the parent ion are also observed and are likely formed from evaporation product reactions. Common for all the curves in Fig. 5 is that they show less size dependence than the clusters in Fig. 4.

The reaction rate coefficients for the clusters H<sup>+</sup>(H<sub>2</sub>O)<sub>*n*</sub>, H<sup>+</sup>(pyridine)<sub>1</sub>(H<sub>2</sub>O)<sub>*n*</sub>, H<sup>+</sup>(pyridine)<sub>2</sub>(H<sub>2</sub>O)<sub>*n*</sub> and H<sup>+</sup>(NH<sub>3</sub>)<sub>1</sub>(pyridine)<sub>1</sub>(H<sub>2</sub>O)<sub>*n*</sub> reacting with NH<sub>3</sub> at 8 kJmol<sup>-1</sup> (COM) are found in Fig. 6 as a function of the number of water molecules in the cluster, for the range *n* = 1–15 (for the pure water clusters *n* = 4–15). The reaction rate coefficients are expressed relative to the rate coefficient of the H<sup>+</sup>(H<sub>2</sub>O)<sub>4</sub> cluster, in the

absence of an exactly calibrated  $\text{NH}_3$  pressure in the collision cell. It should, however, be mentioned that reference measurements show the pressure to be constant during the course of the experiments. The thermal rate coefficients for the reference cluster  $\text{H}^+(\text{H}_2\text{O})_4$  is  $1.91 \times 10^{-9} (300/T)^{0.39} \text{ cm}^3 \text{ s}^{-1}$ , (Viggiano et al., 1988a). For the same cluster reacting with  $\text{ND}_3$  at  $E_{\text{COM}} = 8 \text{ kJmol}^{-1}$  (0.085 eV) Honma et al. (1992) reported a reaction cross section of approximately  $1.5 \times 10^{-14} \text{ cm}^2$ . For the present case, given a velocity of  $1090 \text{ m s}^{-1}$  through the collision cell, this corresponds to a reaction rate coefficient of  $1.6 \times 10^{-9} \text{ cm}^3 \text{ s}^{-1}$  for the reference cluster. As can be seen in Fig. 6, pure water clusters exhibit a somewhat higher reaction rate in the size range  $n = 5\text{--}15$  compared to the pyridine containing clusters.

In order to estimate the rate coefficient for exchanging a pyridine molecule in a cluster with an ammonia molecule i.e. Reaction (R2b) above, we calculated the total abundance of the peaks corresponding to incorporation of ammonia and loss of pyridine and 0 to 2 water molecules. We found that for  $\text{H}^+(\text{pyridine})_m(\text{H}_2\text{O})_n$  clusters with  $m = 1\text{--}2$  and  $n = 1\text{--}15$ , the maximum relative rate coefficient for loss of pyridine was  $8.0 \times 10^{-4}$  (the standard deviation due to signal statistics is  $2.4 \times 10^{-4}$ ) relative the total rate coefficient for  $\text{H}^+(\text{H}_2\text{O})_4 + \text{NH}_3$ . Using the value by Viggiano et al. (1988a) for the latter, this gives a rate coefficient of  $1.5 \times 10^{-12} \text{ cm}^3 \text{ s}^{-1}$  (at 298 K) for the reaction  $\text{H}^+(\text{pyridine})_1(\text{H}_2\text{O})_n + \text{NH}_3 \rightarrow \text{H}^+(\text{NH}_3)_1(\text{H}_2\text{O})_{n-x} + x\text{H}_2\text{O} + \text{pyridine}$ . This is a factor of 7 lower than  $1 \times 10^{-11} \text{ cm}^3 \text{ s}^{-1}$  as assumed by Beig and Brasseur. However, for the majority of the clusters in our study the reaction rate coefficient is even lower. Typical values of the rate coefficient for the above reaction is in the range of  $1 \times 10^{-4}$  to  $4 \times 10^{-4}$  relative  $\text{H}^+(\text{H}_2\text{O})_4$ .

### 3.2 Modelling results

We have modelled the reaction kinetics using three different models. First we performed a calculation using the model of Beig and Brasseur applying their set of parameters, and successfully reproduced their results. We thereafter used Model A at

## Pyridine-containing clusters reacting with $\text{NH}_3$

M. J. Ryding et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

different pyridine concentrations. The concentrations used for all molecules in the model are shown in Table 2. The results are shown in Fig. 7 on a linear scale (in the Supplement the results are shown on a logarithmic scale to include also ions with low concentrations). At pyridine concentrations below  $10^4 \text{ cm}^{-3}$ , clusters of the type  $\text{H}^+(\text{NH}_3)_m(\text{H}_2\text{O})_n$  dominate completely but already at pyridine concentrations of  $2.8 \times 10^6 \text{ cm}^{-3}$ , 50 % of the clusters contain both pyridine and ammonia. At concentrations above  $5 \times 10^6 \text{ cm}^{-3}$ , clusters with ammonia, water and two pyridine molecules dominate. Since the lutidine and picoline concentration is 10 times lower than the pyridine concentration, clusters containing both pyridine and lutidine or picoline are found in concentrations about 10 times lower than the concentration of  $\text{H}^+(\text{NH}_3)_1(\text{pyridine})_2(\text{H}_2\text{O})_n$ . In order to evaluate the importance of including the reaction  $\text{H}^+(\text{pyridine})_1(\text{H}_2\text{O})_n + \text{NH}_3 \rightarrow \text{H}^+(\text{NH}_3)_1(\text{pyridine})_1(\text{H}_2\text{O})_{n-x} + x\text{H}_2\text{O}$ , we also compared the results using Model A with the results obtained by substituting this reaction with  $\text{H}^+(\text{pyridine})_1(\text{H}_2\text{O})_n + \text{NH}_3 \rightarrow \text{H}^+(\text{NH}_3)_1(\text{H}_2\text{O})_{n-x} + x\text{H}_2\text{O} + \text{pyridine}$ . Essentially no difference in the ion concentrations from the two models were observed, neither at an ammonia concentration of  $2.46 \times 10^{10} \text{ cm}^{-3}$  nor at an ammonia concentration 50 times lower.

A typical pyridine concentration in the troposphere is around 4 ppt ( $9.9 \times 10^7 \text{ cm}^{-3}$  at 298 K) (Eisele, 1988; Tanner and Eisele, 1991). From Fig. 7 it is clear that more than one pyridine molecule may be present in the cluster ions in the atmosphere. In order to study this in more detail we constructed Model B, where up to five pyridine molecules are allowed to be incorporated in each cluster. In Fig. 8, we show the ion concentration as a function of pyridine concentration at a typical ammonia concentration (1.0 ppb, or  $2.46 \times 10^{10} \text{ cm}^{-3}$ ). Assuming no evaporation, a pyridine concentration of 4 ppt ( $9.9 \times 10^7 \text{ cm}^{-3}$ ) would give more than four pyridine molecules in most clusters. Neglecting evaporation of pyridine may not be realistic, but this model shows that clusters with a multiple number of pyridine molecules may be present in the atmosphere. In Fig. 8b, the concentrations from Fig. 8a are shown on a logarithmic scale. The concentration of  $\text{H}^+(\text{CH}_3\text{COCH}_3)_1(\text{H}_2\text{O})_n$ ,  $\text{H}^+(\text{H}_2\text{O})_n$ , and  $\text{H}^+(\text{CH}_3\text{CN})_1(\text{H}_2\text{O})_n$  are all below

## Pyridine-containing clusters reacting with $\text{NH}_3$

M. J. Ryding et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

0.1 cm<sup>-3</sup> at all pyridine concentrations used. The concentration of H<sup>+</sup>(pyridine)<sub>m</sub>(H<sub>2</sub>O)<sub>n</sub> 1 ≤ m ≤ 5 are also well below 0.1 cm<sup>-3</sup> at pyridine concentrations below 3 × 10<sup>10</sup> cm<sup>-3</sup> and an ammonia concentration of 2.46 × 10<sup>10</sup> cm<sup>-3</sup>.

We also studied the influence of the ammonia concentration on the number of pyridine molecules in the clusters using Model B. Decreasing the ammonia concentration 50 times (shown in Fig. 8c–d) gave no significant change in the concentration of H<sup>+</sup>(pyridine)<sub>m</sub>(H<sub>2</sub>O)<sub>n</sub> and H<sup>+</sup>(NH<sub>3</sub>)<sub>m</sub>(pyridine)<sub>1</sub>(H<sub>2</sub>O)<sub>n</sub>. However, the concentration of H<sup>+</sup>(CH<sub>3</sub>COCH<sub>3</sub>)<sub>1</sub>(H<sub>2</sub>O)<sub>n</sub> is increased to about 2 cm<sup>-3</sup> up to a pyridine concentration of about 10<sup>9</sup> cm<sup>-3</sup>. The concentrations of H<sup>+</sup>(pyridine)<sub>m</sub>(H<sub>2</sub>O)<sub>n</sub> 1 ≤ m ≤ 5 are below 0.1 cm<sup>-3</sup> except for H<sup>+</sup>(pyridine)<sub>5</sub>(H<sub>2</sub>O)<sub>n</sub> at pyridine concentrations above 3 × 10<sup>9</sup> cm<sup>-3</sup>.

#### 4 Discussion and atmospheric implications

Honma et al. (1992) studied the reaction of H<sup>+</sup>(H<sub>2</sub>O)<sub>4</sub> with ND<sub>3</sub> and for all products where the ammonia-d3 molecule entered the cluster all three deuterium atoms remained there after evaporation of H<sub>2</sub>O. Effective H/D exchange between molecules in the reaction complex requires a mobile proton (Yamaguchi et al., 2003; Honma and Armentrout, 2004; Andersson et al., 2008; Ryding et al., 2011). This observation must be the consequence of strong proton binding most likely due to an adamant NH<sub>4</sub><sup>+</sup> core ion, corresponding to its comparably high pK<sub>a</sub> value. It has been shown that this absence of H/D exchange extends to larger sizes (Andersson et al., 2008). Thus we expect protonation exclusively on the ammonia molecule for all clusters formed in the reaction H<sup>+</sup>(H<sub>2</sub>O)<sub>n</sub> + NH<sub>3</sub> → H<sup>+</sup>(NH<sub>3</sub>)<sub>1</sub>(H<sub>2</sub>O)<sub>n-x</sub> + xH<sub>2</sub>O. Since the pyridine molecule of a H<sup>+</sup>(pyridine)<sub>1</sub>(H<sub>2</sub>O)<sub>n</sub> or H<sup>+</sup>(NH<sub>3</sub>)<sub>1</sub>(pyridine)<sub>1</sub>(H<sub>2</sub>O)<sub>n</sub> cluster does not leave after addition of NH<sub>3</sub>, the product will contain two or three basic molecules. This raises a number of interesting questions regarding the cluster structure, dynamics and protonation site. In the case of pure pyridine water clusters, one pyridine molecule in a

## Pyridine-containing clusters reacting with NH<sub>3</sub>

M. J. Ryding et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



protonated water cluster leads to locking of the proton (i.e. the proton is bound to the pyridine molecule), while with two or three pyridine molecules in the cluster the proton becomes mobile (Ryding et al., 2011). The situation is more complicated for the present mixed clusters, since we are now dealing with two different types of nitrogen bases. The gas phase proton affinity of pyridine is greater than that of ammonia; they are  $930 \text{ kJmol}^{-1}$  and  $853.6 \text{ kJmol}^{-1}$ , respectively (Lide, 2006c). However, in bulk water, ammonia is the stronger base as indicated by its higher acid dissociation constant ( $pK_a = 9.25$  for ammonia,  $pK_a = 5.23$  for pyridine, Lide, 2006a, b). Since molecular clusters represent a bridge between the domains of gas phase chemistry and bulk, it is difficult to determine the effective proton affinities, and therefore to which degree the proton will be mobile within the reaction complex. Further studies of clusters containing both ammonia and pyridine using deuterated reactants – for instance  $\text{D}_2\text{O}$  – should provide additional clues.

The current setup of the TOF unit does not allow for accurate quantification of product ions below  $m/z = 50$ . For the  $\text{H}^+(\text{H}_2\text{O})_4$  cluster we were only able to measure the reaction products  $\text{H}^+(\text{NH}_3)_1(\text{H}_2\text{O})_3$  and  $\text{H}^+(\text{NH}_3)_1(\text{H}_2\text{O})_2$  at  $m/z = 72$  and  $m/z = 54$ , respectively. Consequently, the  $-3\text{H}_2\text{O}$  peak is missing for this cluster in Fig. 4a. The experiments by Honma et al. (1992) indicate that the reaction leading to formation of  $\text{H}^+(\text{NH}_3)_1(\text{H}_2\text{O})_1$  should represent no more than a few percent of the total abundance; the main product is in fact  $\text{H}^+(\text{NH}_3)_1(\text{H}_2\text{O})_2$  ( $-2\text{H}_2\text{O}$  in Fig. 4a) which constitutes almost the entire reaction cross section, which is in good agreement with the results in Fig. 4a.

For the experimental setup described in Sect. 2.1, applying collision energy in the lab frame below approximately  $0.3 \text{ eV}$  was observed to result in inefficient ion transmission and major loss of signal. The collision energy used in the measurements,  $8 \text{ kJmol}^{-1}$  (COM), was chosen since it allowed for a satisfactory beam intensity to be obtained for all cluster ions studied. This energy is a factor of two higher than typical tropospheric collision energies ( $\sim 4 \text{ kJmol}^{-1}$  at room temperature). However, the heating of the formed reaction complex due to addition of  $\text{NH}_3$  (i.e. the dissociation energy)

## Pyridine-containing clusters reacting with $\text{NH}_3$

M. J. Ryding et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



is larger still, with values in the range of 65 to 120 kJmol<sup>-1</sup> depending on cluster size (based on calculations for clusters with  $n = 4$  to 7 using data from Meotner, 1984 and Lide, 2006c). The use of higher collision energies would therefore be expected to have only minor effects on the results, as long as the collision energy is sufficiently small compared to the dissociation energy of NH<sub>3</sub> for the cluster in question.

The above mentioned experiment by Honma et al. (1992) show that the total cross section of the H<sup>+</sup>(H<sub>2</sub>O)<sub>4</sub> cluster reacting with ND<sub>3</sub> remains virtually unchanged up to collision energies of 20 kJmol<sup>-1</sup> (COM). For all measured collision energies in this range, the cross section is made up almost entirely by the reaction forming the product H<sup>+</sup>(NH<sub>3</sub>)<sub>1</sub>(H<sub>2</sub>O)<sub>2</sub>, indicating that the energy released into the cluster upon addition of an ammonia molecule corresponds to evaporation of two H<sub>2</sub>O. Based on values calculated using literature thermochemical data (Meotner, 1984; Lide, 2006c), the energy released by introducing NH<sub>3</sub> into the H<sup>+</sup>(H<sub>2</sub>O)<sub>4</sub> cluster is 120 kJmol<sup>-1</sup>. The dissociation energies for losing first one H<sub>2</sub>O and then a second H<sub>2</sub>O from the intermediate cluster H<sup>+</sup>(NH<sub>3</sub>)<sub>1</sub>(H<sub>2</sub>O)<sub>4</sub> are 44 kJmol<sup>-1</sup> and 52 kJmol<sup>-1</sup>, respectively. For larger protonated water clusters reacting with NH<sub>3</sub> at 8 kJmol<sup>-1</sup> (COM), we see approximately equal amounts of evaporation of two and three H<sub>2</sub>O from the reaction complex (Fig. 4a), implying a somewhat different balance between the dissociation energies of NH<sub>3</sub> and H<sub>2</sub>O. As cluster size increases, the energy released into the cluster when NH<sub>3</sub> enters the reaction complex becomes smaller compared to the energy cost of evaporating first one and then a second water molecule. For instance, for H<sup>+</sup>(H<sub>2</sub>O)<sub>7</sub>, the former is 65 kJmol<sup>-1</sup> while the latter two are 35 kJmol<sup>-1</sup> and 38 kJmol<sup>-1</sup>, respectively. This is contrary to the experimental findings in Fig. 4a. The discrepancy could be due to the fact that the experimental branching ratios do not necessarily represent a sequential loss of H<sub>2</sub>O molecules since there is also a possibility of loss of water as a dimer or a trimer (dissociation energies of the dimer and trimer are 20.7 kJmol<sup>-1</sup> and 21.7 kJmol<sup>-1</sup>, respectively, Santra et al., 2007). Furthermore, the temperature is higher for larger clusters in the beam, which might be part of the explanation. For

## Pyridine-containing clusters reacting with NH<sub>3</sub>

M. J. Ryding et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the pyridine containing clusters in Fig. 4b–c, the tendency is loss of – on average – fewer water molecules post reaction. Even fewer water molecules are lost from the  $\text{H}^+(\text{NH}_3)_1(\text{pyridine})_1(\text{H}_2\text{O})_n$  cluster, with the  $+\text{NH}_3 -\text{H}_2\text{O}$  and  $+\text{NH}_3 -2\text{H}_2\text{O}$  peaks being of equal size.

Separate measurements of  $\text{H}^+(\text{pyridine})_1(\text{H}_2\text{O})_{10}$  and  $\text{H}^+(\text{NH}_3)_1(\text{pyridine})_1(\text{H}_2\text{O})_{10}$  reacting with  $\text{NH}_3$  at different collision energies give further insights into the impact of  $E_{\text{COM}}$  on the respective branching ratios. In the case of the former, the branching ratios are essentially unchanged below  $8 \text{ kJmol}^{-1}$ , with the branching ratio of  $-2\text{H}_2\text{O}$  changing from 0.64 at  $8 \text{ kJmol}^{-1}$  to 0.60 at  $3 \text{ kJmol}^{-1}$ . The corresponding change for  $-\text{H}_2\text{O}$  is from 0.23 to 0.31. For the latter cluster, the changes are somewhat more significant, with both curves having a branching ratio of 0.43 at  $E_{\text{COM}} = 8 \text{ kJmol}^{-1}$ . A change in  $E_{\text{COM}}$  to  $3 \text{ kJmol}^{-1}$  results in the  $-2\text{H}_2\text{O}$  curve dropping to 0.35 and the  $-\text{H}_2\text{O}$  curve increasing to 0.56. It would seem that for this cluster, the collision energy of choice ( $8 \text{ kJmol}^{-1}$ ) also happens to be the point where the two curves representing the collision energy dependence of the  $+\text{NH}_3 -\text{H}_2\text{O}$  product and the  $+\text{NH}_3 -2\text{H}_2\text{O}$  product cross each other. Consequently, for typical tropospheric conditions the  $+\text{NH}_3 -\text{H}_2\text{O}$  product would have a higher abundance than the  $+\text{NH}_3 -2\text{H}_2\text{O}$  product for this cluster.

The pure water clusters in Fig. 6 have a somewhat higher reaction rate coefficient for collision with  $\text{NH}_3$  compared to the pyridine containing cluster for a large part of the size range. This is very similar to the behaviour exhibited by these three cluster types when reacting with  $\text{D}_2\text{O}$ , as reported in a previous study (Ryding et al., 2011).

Model A gave results that were in excellent agreement with those of Beig and Brasseur (2000) with regards to the concentration of pyridinated cluster ions. Our model is also in agreement with Beig and Brasseur in that the pyridinated cluster ions dominates the tropospheric ion spectrum. Our experiments show that the reaction between  $\text{H}^+(\text{pyridine})_1(\text{H}_2\text{O})_n$  and  $\text{NH}_3$  does not give loss of pyridine and have a higher rate coefficient than assumed by Beig and Brasseur. However, this did not have any significant effect on the modelled cluster distribution. Model A allows for two pyridine

## Pyridine-containing clusters reacting with $\text{NH}_3$

M. J. Ryding et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



derivatives in each cluster ion, and the model calculations show that it should be possible for these clusters to form in the troposphere. In Model B we allowed for up to five pyridines in each cluster ion. The model calculations showed that already at a pyridine concentration of  $10^7 \text{ cm}^{-3}$  the cluster ion distribution is dominated by clusters containing ammonia, water and five pyridines. The model is likely to overestimate the number of pyridines in each cluster since evaporation or other removal mechanisms are neglected. However, in our experiments evaporation of pyridine from cluster ions containing one, two, or three pyridine has not been observed, even after reaction with  $\text{NH}_3$  (this work) or  $\text{D}_2\text{O}$  (Ryding et al., 2011). In addition, our models indicate that it is also possible to have clusters containing both ammonia and pyridine at tropospheric pyridine concentrations significantly lower than typical.

Measurements by Junninen et al. (2010) and Ehn et al. (2010) suggest that the ions of alkyl substituted pyridine compounds may be more abundant than ordinary protonated pyridine at ground level in urban and boreal environments. This may seem contradictory when considering the atmospheric concentrations of the compounds in question, as well as the lifetime calculations (for instance by, Yeung and Elrod, 2003), both of which indicate pyridine as the more common neutral species. However, as pointed out by Junninen et al., the transition from neutral molecule to cation takes place by addition of a proton, which will lead to compounds with higher proton affinity being relatively more abundant in the tropospheric ion spectrum. The proton affinities in question are  $930 \text{ kJmol}^{-1}$  for pyridine,  $943.4$  to  $949.1 \text{ kJmol}^{-1}$  (depending on isomer) for picoline and  $955.4$  to  $963.0 \text{ kJmol}^{-1}$  (depending on the isomer) for lutidine (Lide, 2006c). For cluster ions, the type of pyridine or pyridine derivative that enters the clusters (Reactions R1b and R2a) is likely more dependent on concentration than proton affinity. In case of large water cluster ions with more than one pyridine type molecule the actual protonation site becomes a matter of basicity.  $pK_a$  values are as follows: 5.23 for pyridine, 5.70 to 6.00 for picoline (depending on isomer), 6.15 to 6.99 depending on isomer for lutidine (Lide, 2006b). Since the transition from atmospheric pressure to high vacuum probably leads to larger tendency for fragmentation and/or

## Pyridine-containing clusters reacting with $\text{NH}_3$

M. J. Ryding et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

evaporation of a cluster ion, actual measurements of clusters containing both pyridine and an alkyl substituted variant are likely to be detected as the latter, since both the proton affinity and the acid dissociation constants are higher. If also ammonia is present – as indicated by our calculations – the proton may also be situated on the ammonia molecule ( $pK_a = 9.25$ ). The location of the proton is probably also dependent on the cluster size. The fragmentation and evaporation upon sampling could be the reason why clusters containing at least two amines are not observed in the studies by Junninen et al. (2010) and Ehn et al. (2010). In order to better understand the fragmentation and evaporation processes of these ions during atmospheric measurements, we suggest experimental studies on collision induced dissociation of water cluster ions containing two or more amines. We also suggest detailed studies on evaporation of amines from charged water containing clusters under tropospheric conditions.

**Supplementary material related to this article is available online at:**  
**[http://www.atmos-chem-phys-discuss.net/11/24535/2011/  
acpd-11-24535-2011-supplement.pdf](http://www.atmos-chem-phys-discuss.net/11/24535/2011/acpd-11-24535-2011-supplement.pdf)**

*Acknowledgements.* This work was supported by the Swedish Research Council, the Norwegian Research Council by the Grant No. 179568/V30 to the Centre for Theoretical and Computational Chemistry through their Centre of Excellence program, and the Nanoparticle in Interactive Environments platform at the Faculty of Science at University of Gothenburg. MJR is grateful for travel grants awarded by the University of Gothenburg through Jubileumsfonden and Filosofiska fakultetens gemensamma donationsnämnd.

ACPD

11, 24535–24566, 2011

## Pyridine-containing clusters reacting with $\text{NH}_3$

M. J. Ryding et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## References

- Andersson, P. U., Ryding, M. J., Sekiguchi, O. and Uggerud, E.: Isotope exchange and structural rearrangements in reactions between size-selected ionic water clusters,  $\text{H}_3\text{O}^+(\text{H}_2\text{O})_{(n)}$  and  $\text{NH}_4^+(\text{H}_2\text{O})_{(n)}$ , and  $\text{D}_2\text{O}$ , *Phys. Chem. Chem. Phys.*, 10, 6127–6134, 2008.
- 5 Arijis, E. and Brasseur, G.: Acetonitrile in the Stratosphere and Implications for Positive-Ion Composition, *J. Geophys. Res. Atmos.*, 91, 4003–4016, 1986.
- Atkinson, R., Tuazon, E. C., Wallington, T. J., Aschmann, S. M., Arey, J., Winer, A. M., and Pitts, J. N.: Atmospheric Chemistry of Aniline, N,N-Dimethylaniline, Pyridine, 1,3,5-Triazine, and Nitrobenzene, *Environ. Sci. Technol.*, 21, 64–72, 1987.
- 10 Beig, G.: Global change induced trends in ion composition of the troposphere to the lower thermosphere, *Ann. Geophys.*, 26, 1181–1187, doi:10.5194/angeo-26-1181-2008, 2008.
- Beig, G. and Brasseur, G. P.: Model of tropospheric ion composition: A first attempt, *J. Geophys. Res. Atmos.*, 105, 22671–22684, 2000.
- Beig, G., Walters, S., and Brasseur, G.: A 2-Dimensional Model of Ion Composition in the Stratosphere .1. Positive-Ions, *J. Geophys. Res. Atmos.*, 98, 12767–12773, 1993.
- 15 Clemo, G. R.: Some Aromatic Basic Constituents of Coal Soot, *Tetrahedron*, 29, 3987–3990, 1973.
- Ehn, M., Junninen, H., Petäjä, T., Kurtzn, T., Kerminen, V.-M., Schobesberger, S., Manninen, H. E., Ortega, I. K., Vehkamäki, H., Kulmala, M., and Worsnop, D. R.: Composition and temporal behavior of ambient ions in the boreal forest, *Atmos. Chem. Phys.*, 10, 8513–8530, doi:10.5194/acp-10-8513-2010, 2010.
- 20 Eisele, F. L.: Direct Tropospheric Ion Sampling and Mass Identification, *Int. J. Mass Spectrom.*, 54, 119–126, 1983.
- Eisele, F. L.: Identification of Tropospheric Ions, *J. Geophys. Res. Atmos.*, 91, 7897–7906, 1986.
- 25 Eisele, F. L.: First tandem mass spectrometric measurement of tropospheric ions, *J. Geophys. Res. Atmos.*, 93, 716–24, 1988.
- Eisele, F. L. and McDaniel, E. W.: Mass-Spectrometric Study of Tropospheric Ions in The Northeastern and Southwestern United-States, *J. Geophys. Res. Atmos.*, 91, 5183–5188, 1986.
- 30 Eisele, F. L. and Tanner, D. J.: Identification of ions in continental air, *J. Geophys. Res. Atmos.*, 95, 20539–20550, 1990.

## Pyridine-containing clusters reacting with $\text{NH}_3$

M. J. Ryding et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



FACSIMILE v4.0 User Guide. MCPA Software Ltd, Oxford, UK, 2007.

Hauck, G. and Arnold, F.: Improved Positive-Ion Composition Measurements in the Upper Troposphere and Lower Stratosphere and the Detection of Acetone, *Nature*, 311, 547–550, 1984.

5 Honma, K. and Armentrout, P. B.: The mechanism of proton exchange: Guided ion beam studies of the reactions,  $\text{H}(\text{H}_2\text{O})_n^+ (n = 1-4) + \text{D}_2\text{O}$  and  $\text{D}(\text{D}_2\text{O})_n^+ (n = 1-4) + \text{H}_2\text{O}$ , *J. Chem. Phys.*, 121, 8307–8320, 2004.

Honma, K., Sunderlin, L. S., and Armentrout, P. B.: Reactions of Protonated Water Clusters with Deuterated Ammonia –  $\text{H}(\text{H}_2\text{O})_n^+ (n = 1-4) + \text{ND}_3$ , *Int. J. Mass Spectrom.*, 117, 237–259, 1992.

10 Horrak, U., Salm, J., and Tammet, H.: Statistical characterization of air ion mobility spectra at Tahkuse Observatory: Classification of air ions, *J. Geophys. Res.-Atmos.*, 105, 9291–9302, 2000.

15 Junninen, H., Ehn, M., Petäjä, T., Luosujrvi, L., Kotiaho, T., Kostianen, R., Rohner, U., Gonin, M., Fuhrer, K., Kulmala, M., and Worsnop, D. R.: A high-resolution mass spectrometer to measure atmospheric ion composition, *Atmos. Meas. Tech.*, 3, 1039–1053, doi:10.5194/amt-3-1039-2010, 2010.

Kulmala, M., Vehkamäki, H., Petaja, T., Dal Maso, M., Lauri, A., Kerminen, V. M., Birmili, W. and McMurry, P. H.: Formation and growth rates of ultrafine atmospheric particles: a review of observations, *J. Aerosol. Sci.*, 35, 143–176, 2004.

20 Kulmala, M., Riipinen, I., Sipila, M., Manninen, H. E., Petaja, T., Junninen, H., Dal Maso, M., Mordas, G., Mirme, A., Vana, M., Hirsikko, A., Laakso, L., Harrison, R. M., Hanson, I., Leung, C., Lehtinen, K. E. J., and Kerminen, V. M.: Toward direct measurement of atmospheric nucleation, *Science*, 318, 89–92, 2007.

25 Lide, D. R.: *Dissociation Constants of Inorganic Acids and Bases*, available at: [CRCHandbookofChemistryandphysics,internetversion2006](http://CRCHandbookofChemistryandphysics,internetversion2006), D. R. Lide. Boca Raton, FL, USA, Taylor and Francis, 2006a.

Lide, D. R.: *Dissociation Constants of Organic Acids and Bases*, available at: [CRCHandbookofChemistryandphysics,internetversion2006](http://CRCHandbookofChemistryandphysics,internetversion2006), D. R. Lide. Boca Raton, FL, USA, Taylor and Francis, 2006b.

30 Lide, D. R.: *Proton Affinities*, [CRCHandbookofChemistryandphysics,internetversion2006](http://CRCHandbookofChemistryandphysics,internetversion2006), D. R. Lide. Boca Raton, FL, USA, Taylor and Francis, 2006c.

Meotner, M.: *The Ionic Hydrogen-Bond and Ion Solvation. 2. Solvation of Onium Ions by One*

**Pyridine-containing clusters reacting with  $\text{NH}_3$**

M. J. Ryding et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Pyridine-containing  
clusters reacting with  
NH<sub>3</sub>**

M. J. Ryding et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- to 7 H<sub>2</sub>O Molecules – Relations between Monomolecular, Specific, and Bulk Hydration, *J. Am. Chem. Soc.*, 106, 1265–1272, 1984.
- Perkins, M. D. and Eisele, F. L.: First mass-spectrometric measurements of atmospheric ions at ground level, *J. Geophys. Res. Atmos.*, 89, 9649–9657, 1984.
- 5 Ryding, M. J., Zatula, A. S., Andersson, P. U., and Uggerud, E.: Isotope exchange in reactions between D<sub>2</sub>O and size-selected ionic water clusters containing pyridine, H<sup>+</sup>(pyridine)<sub>m</sub>(H<sub>2</sub>O)<sub>n</sub>, *Phys. Chem. Chem. Phys.*, 13, 1356–1367, 2011.
- Saintjalm, Y. and Moreetesta, P.: Study of Nitrogen-Containing Compounds in Cigarette-Smoke by Gas Chromatography-Mass Spectrometry, *J. Chromatography*, 198, 188–192, 1980.
- 10 Santra, B., Michaelides, A. and Scheffler, M.: On the accuracy of density-functional theory exchange-correlation functionals for H bonds in small water clusters: Benchmarks approaching the complete basis set limit, *J. Chem. Phys.*, 127–135, 2007.
- Schlager, H., Fabian, R., and Arnold, F.: A new cluster ion source/ion drift cell apparatus for atmospheric ion studies – First mobility and reaction rate coefficient measurements. 3rd Int. Swarm Seminar, Innsbruck, Austria, 1983.
- 15 Schulte, P. and Arnold, F.: Pyridinium Ions and Pyridine in the Free Troposphere, *Geophys. Res. Lett.*, 17, 1077–1080, 1990.
- Tanner, D. J. and Eisele, F. L.: Ions in Oceanic and Continental Air Masses, *J. Geophys. Res. Atmos.*, 96, 1023–1031, 1991.
- 20 Vana, M., Ehn, M., Petaja, T., Vuollekoski, H., Aalto, P., de Leeuw, G., Ceburnis, D., O'Dowd, C. D. and Kulmala, M.: Characteristic features of air ions at Mace Head on the west coast of Ireland, *Atmos. Res.*, 90, 278–286, 2008.
- Wayne, R. P.: *Chemistry of Atmospheres*, Oxford, Oxford University Press, 2000.
- Viggiano, A. A., Dale, F. and Paulson, J. F.: Proton transfer reactions of aquated hydrogen ions (H<sup>+</sup>(H<sub>2</sub>O)<sub>n</sub>, n = 2–11) with methanol, ammonia, pyridine, acetonitrile, and acetone, *J. Chem. Phys.*, 88, 2469–2477, 1988a.
- 25 Viggiano, A. A., Morris, R. A., Dale, F. and Paulson, J. F.: Tropospheric Reactions of H<sup>+</sup>(NH<sub>3</sub>)<sub>m</sub>(H<sub>2</sub>O)<sub>n</sub> with Pyridine and Picoline, *J. Geophys. Res. Atmos.*, 93, 9534–9538, 1988b.
- 30 Yamaguchi, S., Kudoh, S., Okada, Y., Orii, T., Takeuchi, K., Ichikawa, T. and Nakai, H.: Size-dependent reaction cross section of protonated water clusters H<sup>+</sup>(H<sub>2</sub>O)<sub>(n)</sub> (n = 2–11) with D<sub>2</sub>O, *J. Phys. Chem. A*, 107, 10904–10910, 2003.
- Yeung, L. Y. and Elrod, M. J.: Experimental and computational study of the kinetics of OH plus

pyridine and its methyl- and ethyl-substituted derivatives, J. Phys. Chem. A, 107, 4470–4477, 2003.

5 Yu, F. Q.: Nucleation rate of particles in the lower atmosphere: Estimated time needed to reach pseudo-steady state and sensitivity to H<sub>2</sub>SO<sub>4</sub> gas concentration, Geophys. Res. Lett., 30, 1526, doi:10.1029/2003GL017084, 2003.

Yu, F. Q. and Turco, R. P.: Ultrafine aerosol formation via ion-mediated nucleation, Geophys. Res. Lett., 27, 883–886, 2000.

10 Zhao, Z., Huskey, D. T., Olsen, K. J., Nicovich, J. M., McKee, M. L., and Wine, P. H.: Kinetics, mechanism, and thermochemistry of the gas-phase reaction of atomic chlorine with pyridine, Phys. Chem. Chem. Phys., 9, 4383–4394, 2007.

ACPD

11, 24535–24566, 2011

## Pyridine-containing clusters reacting with NH<sub>3</sub>

M. J. Ryding et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 1.** Rate coefficients for the reactions used in Model A and Model B.

Reaction	Rate coefficient, $\text{cm}^3 \text{s}^{-1}$	Reference
Formation of $\text{H}^+(\text{H}_2\text{O})_n$	$\text{Rate}_1 = 2$ (unit: $\text{cm}^{-3} \text{s}^{-1}$ )	Beig and Brasseur (2000) <sup>a</sup>
$\text{H}^+(\text{H}_2\text{O})_n + \text{CH}_3\text{CN} \rightarrow \text{H}^+(\text{CH}_3\text{CN})_m(\text{H}_2\text{O})_n$	$k_1 = 3.06 \times 10^{-9}(300/T)$	Viggiano et al. (1988a)
$\text{H}^+(\text{H}_2\text{O})_n + \text{NH}_3 \rightarrow \text{H}^+(\text{NH}_3)_m(\text{H}_2\text{O})_n$	$k_2 = 1.91 \times 10^{-9}(300/T)^{0.39}$	Viggiano et al. (1988a)
$\text{H}^+(\text{H}_2\text{O})_n + \text{CH}_3\text{COCH}_3 \rightarrow \text{H}^+(\text{CH}_3\text{COCH}_3)_1(\text{H}_2\text{O})_n$	$k_3 = 2.04 \times 10^{-9}(300/T)^{0.59}$	Viggiano et al. (1988a)
$\text{H}^+(\text{CH}_3\text{COCH}_3)_1(\text{H}_2\text{O})_n + \text{NH}_3 \rightarrow \text{H}^+(\text{NH}_3)_m(\text{H}_2\text{O})_n$	$k_4 = 2 \times 10^{-9}$	Hauck and Arnold (1984)
$\text{H}^+(\text{CH}_3\text{CN})_m(\text{H}_2\text{O})_n + \text{CH}_3\text{COCH}_3 \rightarrow \text{H}^+(\text{CH}_3\text{COCH}_3)_1(\text{H}_2\text{O})_n$	$k_5 = 1.8 \times 10^{-9}$	Hauck and Arnold (1984)
$\text{H}^+(\text{CH}_3\text{CN})_m(\text{H}_2\text{O})_n + \text{NH}_3 \rightarrow \text{H}^+(\text{NH}_3)_m(\text{H}_2\text{O})_n$	$k_6 = 1.8 \times 10^{-9}$	Schlager et al. (1983)
$\text{H}^+(\text{NH}_3)_m(\text{H}_2\text{O})_n + \text{picoline} \rightarrow \text{H}^+(\text{NH}_3)_x(\text{picoline})_1(\text{H}_2\text{O})_n$	$k_7 = 2.6 \times 10^{-9}(300/T)^{0.7}$	Viggiano et al. (1988b)
$\text{H}^+(\text{NH}_3)_m(\text{H}_2\text{O})_n + \text{lutidine} \rightarrow \text{H}^+(\text{NH}_3)_x(\text{lutidine})_1(\text{H}_2\text{O})_n$	$k_8 = 2 \times 10^{-9}$	assumed <sup>b</sup>
$\text{H}^+(\text{NH}_3)_m(\text{H}_2\text{O})_n + \text{pyridine} \rightarrow \text{H}^+(\text{NH}_3)_x(\text{pyridine})_1(\text{H}_2\text{O})_n$	$k_9 = 2.1 \times 10^{-9}(300/T)^{0.7}$	Viggiano et al. (1988b)
$\text{H}^+(\text{H}_2\text{O})_n + \text{picoline} \rightarrow \text{H}^+(\text{picoline})_1(\text{H}_2\text{O})_n$	$k_{10} = 2 \times 10^{-9}$	assumed <sup>b</sup>
$\text{H}^+(\text{H}_2\text{O})_n + \text{lutidine} \rightarrow \text{H}^+(\text{lutidine})_1(\text{H}_2\text{O})_n$	$k_{11} = 2 \times 10^{-9}$	assumed <sup>b</sup>
$\text{H}^+(\text{H}_2\text{O})_n + \text{pyridine} \rightarrow \text{H}^+(\text{pyridine})_1(\text{H}_2\text{O})_n$	$k_{12} = 2.08 \times 10^{-9}(300/T)^{0.89}$	Viggiano et al. (1988a)
$\text{H}^+(\text{X})_1(\text{H}_2\text{O})_n + \text{NH}_3 \rightarrow \text{H}^+(\text{NH}_3)_m(\text{H}_2\text{O})_n + \text{X}$	$k_{13} = k_{14} = k_{15} = 0^c$	assumed
cluster + aerosol $\rightarrow$ aerosol ion	$k_{16} = 5 \times 10^{-6}$	Beig and Brasseur (2000)
cluster + cluster recombination	$k_{17} = 6 \times 10^{-8}(300/T)^{0.5}$ $+ 1.25 \times 10^{-25}[\text{M}](300/T)^{4d}$	Arijs and Brasseur (1986); Beig et al. (1993)
$\text{H}^+(\text{X})_1(\text{H}_2\text{O})_n + \text{NH}_3 \rightarrow \text{H}^+(\text{NH}_3)_m(\text{X})_1(\text{H}_2\text{O})_n$	$k_{18} = 0.907 \times k_2$	this paper
$\text{H}^+(\text{X})_1(\text{H}_2\text{O})_n + \text{X} \rightarrow \text{H}^+(\text{X})_2(\text{H}_2\text{O})_n$	$k_{19} = k_{12}$	assumed
$\text{H}^+(\text{NH}_3)_m(\text{X})_1(\text{H}_2\text{O})_n + \text{pyridine} \rightarrow \text{H}^+(\text{NH}_3)_m(\text{X})_2(\text{H}_2\text{O})_n$	$k_{20} = k_9$	assumed
$\text{H}^+(\text{NH}_3)_m(\text{X})_1(\text{H}_2\text{O})_n + \text{picoline} \rightarrow \text{H}^+(\text{NH}_3)_m(\text{X})_2(\text{H}_2\text{O})_n$	$k_{21} = k_7$	assumed
$\text{H}^+(\text{NH}_3)_m(\text{X})_1(\text{H}_2\text{O})_n + \text{lutidine} \rightarrow \text{H}^+(\text{NH}_3)_m(\text{X})_2(\text{H}_2\text{O})_n$	$k_{22} = k_8$	assumed
$\text{H}^+(\text{X})_2(\text{H}_2\text{O})_n + \text{NH}_3 \rightarrow \text{H}^+(\text{NH}_3)_m(\text{X})_2(\text{H}_2\text{O})_n$	$k_{23} = k_{18}$	assumed
$\text{H}^+(\text{NH}_3)_m(\text{X})_2(\text{H}_2\text{O})_n + \text{pyridine} \rightarrow \text{H}^+(\text{NH}_3)_m(\text{X})_3(\text{H}_2\text{O})_n$	$k_{24} = k_9$	assumed
$\text{H}^+(\text{NH}_3)_m(\text{X})_3(\text{H}_2\text{O})_n + \text{pyridine} \rightarrow \text{H}^+(\text{NH}_3)_m(\text{X})_4(\text{H}_2\text{O})_n$	$k_{25} = k_9$	assumed
$\text{H}^+(\text{NH}_3)_m(\text{X})_4(\text{H}_2\text{O})_n + \text{pyridine} \rightarrow \text{H}^+(\text{NH}_3)_m(\text{X})_5(\text{H}_2\text{O})_n$	$k_{26} = k_9$	assumed

<sup>a</sup> The formation of  $\text{H}^+(\text{H}_2\text{O})_n$  was set as to give the same rate as used by Beig and Brasseur (2000).

<sup>b</sup> The value is the same as used by Beig and Brasseur (2000).

<sup>c</sup> A value of  $1 \times 10^{-11} \text{ cm}^{-3} \text{ s}^{-1}$  was used by Beig and Brasseur (2000).

<sup>d</sup>  $[\text{M}]$  is the neutral number density in  $\text{cm}^{-3}$ .

## Pyridine-containing clusters reacting with $\text{NH}_3$

M. J. Ryding et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Pyridine-containing clusters reacting with NH<sub>3</sub>

M. J. Ryding et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




**Table 2.** Initial concentrations in cm<sup>-3</sup> for the simulations presented in Figs. 7 and 8.

Figure	Model	NH <sub>3</sub>	H <sub>2</sub> O	Pyridine	Picoline	Lutidine	Acetone	CH <sub>3</sub> CN	Aerosol
7	A	$2.46 \times 10^{10}$	$4.61 \times 10^{17}$	variable	[pyridine]/10	[pyridine]/10	$3.69 \times 10^{10}$	$4.92 \times 10^8$	$1.0 \times 10^3$
8a–b	B	$2.46 \times 10^{10}$	$4.61 \times 10^{17}$	variable	–	–	$3.69 \times 10^{10}$	$4.92 \times 10^8$	$1.0 \times 10^3$
8c–d	B	$4.92 \times 10^8$	$4.61 \times 10^{17}$	variable	–	–	$3.69 \times 10^{10}$	$4.92 \times 10^8$	$1.0 \times 10^3$



## Pyridine-containing clusters reacting with NH<sub>3</sub>

M. J. Ryding et al.



**Fig. 1.** Tropospheric cluster ion reaction scheme used in Model A. Dotted lines represent reactions removed from the model employed by Beig and Brasseur (2000). Dashed-dotted lines represent reactions not included in the model by Beig and Brasseur (2000).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Pyridine-containing clusters reacting with NH<sub>3</sub>**

M. J. Ryding et al.




**Fig. 2.** Tropospheric cluster ion reaction scheme used in Model B.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



## Pyridine-containing clusters reacting with $\text{NH}_3$

M. J. Ryding et al.



**Fig. 3.** Mass spectra of  $\text{H}^+(\text{pyridine})_1(\text{H}_2\text{O})_{11}$  with peak intensities expressed relative to the parent ion (PI) mass and height. **(a)** Background measurement. **(b)** The cluster ion reacting with  $\text{NH}_3$  at  $8 \text{ kJ mol}^{-1}$ . The product  $\text{H}^+(\text{NH}_3)_1(\text{pyridine})_1(\text{H}_2\text{O})_{11-y}$  peaks are designated by open blue triangles. Red circles designate peaks arising from evaporation of  $\text{H}_2\text{O}$  from the parent ion. Green vertical lines designate the  $m/z$  where a possible  $\text{NH}_3/\text{pyridine}$  exchange product,  $\text{H}^+(\text{NH}_3)_1(\text{H}_2\text{O})_{10-z}$ , would appear.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Pyridine-containing clusters reacting with NH<sub>3</sub>


M. J. Ryding et al.



**Fig. 4.** Branching ratios for the reactions of  $\text{H}^+(\text{pyridine})_m(\text{H}_2\text{O})_n$  ( $m = 0$  to  $2$ ) with  $\text{NH}_3$  at  $E_{\text{COM}} = 8 \text{ kJ mol}^{-1}$ . The number of water molecules,  $n$ , is indicated for some of the data points to improve readability. Dashed lines indicate products likely resulting from clusters that have lost water molecules prior to reaction.

## Pyridine-containing clusters reacting with NH<sub>3</sub>

M. J. Ryding et al.



**Fig. 5.** Branching ratios for  $H^+(NH_3)_1(pyridine)_1(H_2O)_n$  reacting with  $NH_3$  at  $E_{COM} = 8 \text{ kJmol}^{-1}$ . The curves represent peaks corresponding to the parent ion incorporating  $NH_3$  and losing one to four  $H_2O$ . Dashed lines indicate products likely resulting from clusters that have lost water molecules prior to reaction. The numbers next to the curves indicate the number of water molecules, *n*.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Pyridine-containing clusters reacting with NH<sub>3</sub>

M. J. Ryding et al.




**Fig. 6.** Relative reaction rate coefficients for cluster ions  $\text{H}^+(\text{pyridine})_m(\text{H}_2\text{O})_n$  with  $m = 0$  to 2 and  $\text{H}^+(\text{NH}_3)_1(\text{pyridine})_1(\text{H}_2\text{O})_n$ , reacting with  $\text{NH}_3$  at  $E_{COM} = 8 \text{ kJ mol}^{-1}$ . The rate coefficients are normalized with the rate coefficient for  $\text{H}^+(\text{H}_2\text{O})_4$  reacting with  $\text{NH}_3$ . The numbers 1, 5, 10, 15 indicate the number of water molecules to improve readability.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


## Pyridine-containing clusters reacting with NH<sub>3</sub>

M. J. Ryding et al.



**Fig. 7.** Concentration of cluster ions as a function of pyridine concentration. The values were calculated using Model A and the initial concentrations are given in Table 2. The total concentration of pyridinated cluster ions in the model by Beig and Brasseur (2000) is also indicated. Py, Pic and Lu indicate pyridine, picoline and lutidine, respectively.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Pyridine-containing clusters reacting with NH<sub>3</sub>**

M. J. Ryding et al.



**Fig. 8.** Concentration of cluster ions as a function of pyridine concentration. The values were calculated using Model B and the initial concentrations are given in Table 2. Panel (a) and (b) shows the results using an ammonia concentration of  $2.46 \times 10^{10} \text{ cm}^{-3}$  on a linear and logarithmic scale, respectively. Panel c) and d) shows the results for an ammonia concentration of  $4.92 \times 10^8 \text{ cm}^{-3}$ . Py, Pic and Lu indicate pyridine, picoline and lutidine, respectively.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

