

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

# Soil-air exchange controls on background atmospheric concentrations of organochlorine pesticides

A. Cabrerizo<sup>1</sup>, J. Dachs<sup>1</sup>, K. C. Jones<sup>2</sup>, and D. Barceló<sup>1</sup>

<sup>1</sup>Department of Environmental Chemistry, IDAEA-CSIC, Jordi Girona 18-26, Barcelona, 08034 Catalunya, Spain

<sup>2</sup>Lancaster Environment Centre, Lancaster University, Lancaster, LA1 4YQ, UK

Received: 7 July 2011 – Accepted: 13 September 2011 – Published: 19 September 2011

Correspondence to: J. Dachs (jordi.dachs@idaea.csic.es)

Published by Copernicus Publications on behalf of the European Geosciences Union.

ACPD

11, 25937–25965, 2011

## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Soils are the major terrestrial reservoir of persistent organic pollutants, and thus net volatilization from soil, when it happens, may exert a control on the atmospheric occurrence and variability of organic pollutants. Here, we report and discuss the concentrations of legacy organochlorine pesticides (OCPs) such as hexachlorobenzene (HCB), hexachlorocyclohexanes (HCH) and dichlorodiphenyltrichloroethane (DDT) in the atmosphere and in soils, their measured fugacities in soil, the soil-air partition coefficients ( $K_{SA}$ ) and soil-air fugacity ratios ( $f_s/f_a$ ) in rural background areas of N-NE Spain and N-NW England. Four sampling campaigns were carried out in Spain and UK to assess seasonal variability and differences between sampling sites.  $K_{SA}$  values were significantly dependent on soil temperature and soil organic matter quantity, and to a minor extent on organic matter type. HCH isomers and DDT metabolites in soil are close to equilibrium with the atmosphere at rural background areas of Spain with a tendency to volatilize and deposit during warm and cold periods, respectively. The mixture of HCH and DDT found in the atmosphere is clearly strongly influenced by the mixture of HCH and DDT which escapes from soil, with significant correlations between them ( $r^2$  ranging between 0.74–0.76 and p-level < 0.001 for the Ebro sampling sites), thus suggesting a close coupling of air and soil concentrations demonstrating that net volatilization from soil control the atmospheric levels of OCPs in the Northern Spain background atmosphere. Conversely, soils at rural UK sites were usually a sink for atmospheric DDT and HCH, but not for HCB. The negative statistically significant relationship found between log  $K_{SA}$  and the log ( $f_s/f_a$ ) ratio, suggests that high latitude regions, due to the high soil organic matter content and lower temperatures, will act as larger traps and accumulate more atmospheric OCPs. Thus, the extent to which soils are secondary sources to the atmosphere is currently dependent on the reservoir potential of soils for OCPs and shows a marked seasonality in their strength.

ACPD

11, 25937–25965, 2011

## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 1 Introduction

Organochlorine pesticides (OCPs) were used heavily in the past, such as hexachloro-cyclohexanes (HCH) for pest control, dichlorodiphenyltrichloroethane (DDT) for malaria control, or for industrial and agricultural purposes such as hexachlorobenzene (HCB).

5 Most have been banned or restricted in industrialized countries and are now being eliminated under the United Nations Economic Commission for Europe (UN-ECE) and the Stockholm Convention on persistent organic pollutant (POPs). However, due to their long environmental lifetimes, semi-volatility, and hydrophobicity, residues still persist in the local and global environment (Blais et al., 1998; Gregor and Gummer, 1989; 10 Grimalt et al., 2001; Ribes et al., 2002). In background regions with no historical direct inputs, the occurrence of OCPs in the atmosphere and soils is influenced by their volatilization from soils and atmospheric deposition events, with atmospheric transport as the main vector for their spatial re-distribution from secondary and potential primary regional sources. Several studies over the past years (Zhang et al., 2007, 2011) 15 have recognized the importance of the amount and composition of soil organic matter (SOM) to explain the sorption and the availability for biological degradation, vertical transport or volatilization of OCPs from the soil surface to the atmosphere. Soils are the major environmental reservoir of POPs (Dalla Valle et al., 2005), and volatilization of OCPs from soil can be a continuous and diffuse source of atmospheric contamination, which must be assessed to determine the relative contribution of “legacy” residues 20 (secondary sources) and ongoing usage (primary sources). Sampling air and soil at the same sites may allow statements on the atmosphere-surface exchange of OCPs and the determination of in-situ soil and ambient air fugacities will provide quantitative evidence of the direction of soil-air exchange of OCPs. In this study, we investigate in 25 detail the factors controlling OCPs gradients between soils and the atmosphere and the soils controls on atmospheric concentrations in selected temperate background areas. This is achieved by means of determining the OCPs concentrations in the atmosphere, in soils, their fugacities in surface soils, and fugacity ratios between soil and

ACPD

11, 25937–25965, 2011

### Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the atmosphere by using the soil fugacity sampler (Meijer et al., 2003a; Cabrerizo et al., 2009). Therefore, the main objective is to study the influence of soils as a control of the atmospheric levels of OCPs by means of (i) studying the effect that temperature and soil organic matter quantity and quality have on soil OCPs fugacity and concentrations in soil and the soil-air partition coefficients ( $K_{SA}$ ), (ii) quantifying soil/air fugacity ratios to determine the direction of soil-air exchange and (iii) exploring the factors affecting the atmospheric concentrations of OCPs. This study has been performed in background rural and semi-rural areas of the Ebro river basin (Spain) and northwest England and is a companion assessment of the recently reported studies on polycyclic aromatic hydrocarbons (PAHs) (Cabrerizo et al., 2011a) and polychlorinated biphenyls (PCBs) (Cabrerizo et al., 2011b).

## 2 Materials and methods

### 2.1 Soil and air sampling

Nine background sites in Spain and UK were selected. All of them were non-agricultural rural or semirural sites (see Fig. S1 and Table S1 in the Supplement (SI) for sampling sites locations and information). There were six rural sites: Borau, Alfaro, Nájera, Lasieso, Uruñuela located along the Ebro river basin (N-NE Spain) and Langden (near Lancaster, UK); and three semi-rural: Tudela, Sabiñanigo (Ebro river basin-Spain) and Hazelrigg (near Lancaster, UK). Sampling sites in Alfaro and Tudela were close to land used for fruit crops but these and all other sites are non agricultural sites. Three sampling campaigns were carried out in June 2006, November 2006 and September 2007 in the Ebro river basin. Sampling in the UK sites was performed during August–September 2008.

Ambient air samples and air equilibrated in-situ with the soil surface samples were collected using a low volume sampler proposed by Cabrerizo et al. (2009). As in commercial high volume samplers, the air passes through a glass fiber filter to remove dust

## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



particles and a polyurethane foam (PUF) plug in which the compounds from the gas phase will be retained. Filters and PUF used in this sampler are quartz fiber filters (GF/F) of 47 mm of diameter and 10 × 2 cm PUF. The main advantage of this sampler is that allows for accurately determining the OCPs soil fugacity by sampling the air which has been equilibrated for enough time with the soil, thus OCPs soil fugacities were determined. Unlike a commercial high volume sampler, the flow rate is much lower (8–10 l min<sup>-1</sup>), thus allowing for air that passes below the sampler to equilibrate in terms of OCPs concentrations with the soil surface (Cabrerizo et al., 2009). In parallel, another sampler located at 1.5 m is operating at the same condition in order to determine ambient air fugacities. Each sample is an integration of 24 h of sampling with a total air volume of 10–14 m<sup>3</sup>.

Soil samples were taken after soil fugacity sampling, by gently collecting the soil surface layer (approximately top 0.5–1 cm). Concerning this study, a total of 43 soil fugacity measurements, 41 ambient air concentrations, and 20 top surface soils were sampled and analyzed. After sampling, air and soil samples were stored in freezers at –20 °C until analysis.

## 2.2 Analytical methods

Soil and air samples were Soxhlet extracted for 24 h in dichloromethane:methanol (2:1) and acetone:hexane (3:1), respectively. Extracts were cleaned and fractionated using alumina chromatography. Soil samples were also treated with activated copper to avoid chromatogram interferences by sulfur. Samples taken in the Ebro river basin (Spain) were analyzed by gas chromatography coupled with electron capture detection (GC/ECD) and those taken in the UK by gas chromatography coupled to mass spectrometry (GC/MS) with an EI+ source operating in selected ion mode (SIM). Details of the temperature programs and monitored ions are given elsewhere (Thomas et al., 1998; Gouin et al., 2002; Ockenden et al., 1998; Cabrerizo et al., 2011b). The following OCPs were analyzed: HCB, HCH isomers ( $\alpha$ -HCH,  $\beta$ -HCH,  $\gamma$ -HCH,  $\delta$ -HCH) and DDT and its metabolites (p,p'-DDT, o,p'-DDT, p,p'-DDD, o,p'-DDD, p,p'-DDE, o,p'-DDE).

## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Unfortunately, o,p'-DDD was omitted in this study due to a co-elution with PCB82 when analyzed by GC/ECD. In the same way, HCB did show a co-elution in some air samples taken only in the first sampling campaign (June 2006), so these samples were not considered. No breakthrough of OCPs is observed with the sampling conditions, due to the small sampling volumes (10–14 m<sup>3</sup>) and the PUFs size which are 10 cm long, ×2 cm of diameter. The breakthrough for OCPs has been estimated to be around 80 m<sup>3</sup> for the more volatile OCI (HCB) during the summer. Reference to the sum of isomers groups of HCH (ΣHCH) and DDT (ΣDDT) considers the aforementioned isomers.

### 2.2.1 Quality assurance/quality control

All analytical procedures were monitored using strict quality assurance and control measures. Laboratory blanks and field blanks constituted 21 % of the total number of samples processed. δ-HCH, p,p'-DDE and p,p'-DDT were detected in blanks and field blanks at low concentrations, ranging between 2 and 10 % of levels found in samples, indicating minimal contamination during storage, sampling, transport and processing. Therefore, samples were not blank corrected. Method quantification limits were derived from the lowest standard in the calibration curve, ranged from 0.35 pg m<sup>-3</sup> and 0.5 pg g<sup>-1</sup> for air and soil samples processed with GC/ECD and between 3.5 pg m<sup>-3</sup> and 10 pg g<sup>-1</sup> for air and soil samples processed with GC/MS. Recoveries were routinely monitored using PCB65 and PCB200 and <sup>13</sup>C<sub>12</sub>-PCBs (28, 52, 101, 138, 153, 180) and they ranged from 53 ± 17 to 76 ± 14 and from 58 ± 3 to 101 ± 26 for soil and gas phase respectively, so samples were not corrected by recoveries.

### 2.2.2 Meteorological data

Concurrent meteorological data were monitored where possible using a wireless weather station or were provided by the nearest weather station to the sampling site by the Spanish Meteorological Agency (AEMET). The following physico-chemical characteristics of samples were measured by standard methods described elsewhere

## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(Cabrerizo et al., 2011a, b): soil temperature (soil  $T$ ), ambient air temperature (air  $T$ ), soil water content (SWC) and precipitation, soil organic carbon (TOC) or organic carbon fraction ( $f_{OC}$ ), surface soil pH and redox potential, fraction of nitrogen ( $f_N$ ) and altitude. The four last parameters were used as indirect “proxy parameters” for the organic matter type/quality.

### 3 Results and discussion

#### 3.1 Occurrence of organochlorine pesticides in ambient air

All the measured ambient air concentrations, air  $T$  and precipitation in the Ebro river basin and UK sites are listed in the SI (Tables S2a–d). Average concentrations of individual organochlorine pesticides  $\Sigma HCH$ ,  $\Sigma DDT$  and HCB are also reported. The range in the  $\Sigma HCH$ ,  $\Sigma DDT$  and HCB concentration in ambient air for Ebro sites (Spain) were 0.034–0.352 ng m<sup>-3</sup>, 0.006–0.083 ng m<sup>-3</sup> and 0.018–0.066 ng m<sup>-3</sup>, respectively. Overall, the highest concentrations of HCH isomers and DDT metabolites were observed in warm periods campaigns (June 2006 and September 2007), suggesting a seasonal variability. However, the air concentrations of legacy pesticides do not change dramatically throughout the year, as other POPs, such as PCBs do (Cabrerizo et al., 2011b), which is in agreement with previous studies (Hayward et al., 2003; Yao et al., 2006).  $\Sigma HCH$ ,  $\Sigma DDT$  and HCB concentrations in ambient air for UK sites (Hazelrigg and Langden) were in the ranges 0.023–0.078 ng m<sup>-3</sup>, 0.012–0.016 ng m<sup>-3</sup>, and 0.034–0.037 ng m<sup>-3</sup> respectively. These atmospheric levels are in the same order of magnitude to those reported in other studies at background areas (Halse et al., 2011; Růžicková et al., 2007; Barber et al., 2005; Gioia et al., 2007). Overall, the concentration of OCPs in ambient air increased, when air  $T$  also increased (p-level < 0.05) (Fig. 1), suggesting that atmospheric concentrations of OCPs vary seasonally, with concentrations reaching a maximum in warmer periods of ambient air concentration. The 24 h back trajectories were calculated using the US National Atmospheric and

## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Oceanic Administration (NOOA) H-Split model for individual OCPs in those samples for which there was a significant difference between maximum and minimum at the same temperature. Examples of back trajectories for selected OCPs were calculated at 10 m and 100 m above ground level in order to identify where the air masses were originated.

5 Figures S2a–c shows, that the back trajectories for maximum and minimum concentrations at the same ambient air temperature do not correspond to a predominant direction of air masses, which suggest that re-emissions of OCPs from local sources in these background environments, consistent with volatilization from soil, may be playing an important role in controlling their ambient air concentration. Ambient air concentra-  
10 tions for selected sites in the Ebro river basin are shown in Fig. S3 and presented as fugacities (or partial pressures in Pa) by

$$f_a = 10^{-9} C_A RT / MW \quad (1)$$

where  $C_A$  is the measured air concentration in  $\text{ng m}^{-3}$ ,  $R$  is the gas constant ( $8.314 \text{ Pa m}^3 \text{ mol}^{-1} \text{ K}^{-1}$ ),  $MW$  is the chemical molecular weight ( $\text{g mol}^{-1}$ ), and  $T$  is the  
15 temperature (K).

### 3.2 Occurrence of pesticides in surface soils

Soil concentrations for individual OCPs,  $\Sigma\text{HCH}$  and  $\Sigma\text{DDT}$  in  $\text{ng g dw}^{-1}$  ( $\text{ng g soil dry weight (dw)}$ ) at each site are shown in the SI (Tables S3a, b). Concentrations of HCB,  $\Sigma\text{HCH}$  and  $\Sigma\text{DDT}$  in soils are in the range between  $0.004\text{--}0.49 \text{ ng g dw}^{-1}$  with a mean value of  $0.11 \text{ ng g dw}^{-1}$ ; between detection limit and  $0.52 \text{ ng g dw}^{-1}$  with a mean value of  $0.06 \text{ ng g dw}^{-1}$ ; and between  $0.02$  and  $4.73 \text{ ng g dw}^{-1}$  with a mean value of  $0.79 \text{ ng g dw}^{-1}$ , respectively. The concentrations in soil reported here are similar to those reported in previous studies for non-agricultural and background soils (Grimalt et al., 2004; Borgini et al., 2005; Meijer et al., 2003b). Overall, p,p'-DDE was the main contributor (up to 60 %) to the  $\Sigma\text{DDT}$  for most soils. This is consistent with the fact that DDT in soil is subject to microbial degradation to more stable and toxic metabo-  
25 lites such as p,p'-DDE. The ratios p,p'-DDE/p,p'-DDT have been successfully used

## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





to distinguish the source of DDT (Harner et al., 2004; Liu et al., 2009; Wang et al., 2009). Therefore, if the ratio of  $p,p'$ -DDE/ $p,p'$ -DDT is  $>1$ , it indicates aged DDT, while ratios  $<1$  indicate fresh inputs. From the data set reported here ( $N=20$ ), almost all soil samples had ratios of  $p,p'$ -DDE/ $p,p'$ -DDT  $>1$ , indicating aged DDT. Only 3 samples collected in the Ebro basin in different sampling campaigns had values  $<1$  (Sabiñánigo (0.53), Lasieso (0.74) and (Nájera (0.03)) which could suggest a short recent use of this OCPs in the area. Regarding HCH isomers,  $\gamma$ -HCH and  $\delta$ -HCH were found to be the dominant isomers in these soils. In order to study the influence of physico-chemical properties on the occurrence of OCPs in soils, a regression analysis was performed. Regressions were calculated between the soil OCPs concentrations and the following variables: soil  $T$ , air  $T$ , SOM quantity and proxies for soil type. HCB,  $\Sigma$ DDT and  $\Sigma$ HCH concentrations in soils ( $\text{ng g dw}^{-1}$ ) were significantly correlated ( $p$ -level  $< 0.05$ ) with soil organic carbon (Fig. 2) suggesting that SOM is clearly an important parameter influencing the concentration of OCPs in background soils, consistent with previous studies (Meijer et al., 2003b). The major contributors among HCH isomers and DDT metabolites to  $\Sigma$ HCH and  $\Sigma$ DDT correlated against soil organic carbon are presented in Fig. S4. A strong relationship (steep slope, less scatter) was observed for HCB ( $r^2 = 0.73$  and  $p$ -level  $< 0.001$ ). This implies that the reservoir of OCPs is larger for soils with larger SOM, and that processes like atmospheric transport and re-distribution through air-soil exchange are important in influencing the soil OCPs concentration, since no direct sources are known for the sampling sites. A statistically significant correlation ( $p$ -level  $< 0.05$ ) was observed between soil concentration normalized to soil organic carbon and SOM quality ( $f_{\text{OC}}/f_{\text{N}}$ ) for selected DDT metabolites ( $p,p'$ -DDE and  $o,p'$ -DDT), with  $r^2$  ranging between 0.23–0.24, while no significant correlation were found for HCH isomers, thus suggesting a variable but modest influence of the soil OCPs reservoir with respect of the SOM type. Concerning temporal trends, there were no statistically significant differences in soil concentrations of OCPs with respect to soil or ambient air temperature for the three sampling periods in the Ebro, consistent with the fact that the inventory of POPs in soil is generally believed

## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to be considerably larger than the annual fluxes in and out of the soil (Kurt-Karakus et al., 2006). No significant correlation was observed with soil pH or soil redox. There were significant ( $p$ -level  $< 0.001$  and  $p$ -level  $< 0.05$ ) correlations between the different POPs families' concentrations (Table S4) for HCB,  $\Sigma$ HCH,  $\Sigma$ DDT,  $\Sigma$ PCBs and  $\Sigma$ PAHs across the whole set of soils, which is clear evidence of the common role of organic matter quantity as a descriptor of the burden of hydrophobic pollutants. This was also observed by Nam et al. (2008) in other European background soils.

### 3.3 OCPs fugacity in soils

Only those OCPs present in the exchangeable fraction of soil will be able to escape from the soil surface to the atmosphere. The recent development of a new soil fugacity sampler (Cabrerizo et al., 2009) allows for direct determination of the pollutants fugacity in soils. In this study, OCPs fugacity in soil was determined directly under field controlled conditions by:

$$f_s = 10^{-9} C_{SA} RT / MW \quad (2)$$

where  $C_{SA}$  ( $\text{ng m}^{-3}$ ) is the gas phase concentration that has been equilibrated with the soil surface as measured using the soil fugacity sampler. To establish the role of temperature and SOM in the magnitude of the emission of OCPs in soils, a least squares regressions of  $\log f_s = a + b(1/T)$  and  $\log f_s = a + b(\log f_{OC})$  were applied for individual HCH isomers, DDT metabolites and HCB which were detected in at least 15 of the samples ( $N \geq 15$ ). Results show that lower soil fugacities correspond to low temperatures (Fig. S5) with  $r^2$  ranging between 0.25–0.38 ( $p$ -level  $< 0.05$ ) and to higher amounts of SOM, with  $r^2$  ranging between 0.15 and 0.35 ( $p$ -level  $< 0.05$ ) for  $\alpha$ -HCH,  $\gamma$ -HCH and  $o,p'$ -DDT and  $p,p'$ -DDT (Table S5a). Soil fugacities of those DDT metabolites which were not affected by the SOM quantity (ex.  $o,p'$ -DDE,  $p,p'$ -DDE,  $p,p'$ -DDD) were regressed against SOM quality ( $f_{OC}/f_N$ ) and soil pH (see Tables S5b–c). Results show that lower soil fugacities correspond to high  $f_{OC}/f_N$  ratios with  $r^2$  ranging between 0.26–0.42 ( $p$ -level  $< 0.05$ ) for  $o,p'$ -DDE,  $p,p'$ -DDE,  $p,p'$ -DDD, and that

## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the soil fugacity decreases with decreasing pH with  $r^2$  ranging between 0.47–0.67, thus suggesting that both SOM quantity and quality (especially for DDT metabolites) may be important parameters retarding the emission of OCPs to the atmosphere and therefore soils with high concentration of SOM and lower pH are favored in the accumulation of POPs as described in previous studies (Ockenden et al., 2003; Cabrerizo et al., 2011b). There was much variability for HCB soil fugacities and no statistically significant trend could be observed.

### 3.4 Factors affecting the soil-air partitioning of organochlorine pesticides

As there are currently no fresh applications of the studied OCPs in Spain and UK, their air concentration variability, with higher concentrations during warm periods, is presumably related to their volatilization from soils. As shown above, the soil fugacity depends on temperature, SOM quantity and type, thus the soil-air exchange and partitioning will be dominated by several physical and chemical properties of the soil and atmosphere, among which the vapor pressure, the enthalpy of vaporization of the chemicals ( $\Delta H$ ) (Mackay et al., 1986; Bidleman et al., 1999) and SOM quantity and type are the most important.  $\Delta H$  has been calculated from the slope of the regression resulting when the soil-air partitioning coefficient ( $K_{SA}$ ) is plotted against  $1/T$  (Fig. 3).  $K_{SA}$  was calculated as:

$$K_{SA} = C_S / C_{SA} \quad (3)$$

Characteristic examples of the influence of temperature on  $K_{SA}$  for selected HCH isomers and DDT metabolites are shown in Fig. 3. Log  $K_{SA}$  was significantly positively correlated with  $1/T$ , in agreement with previous studies (Su et al., 2006; Haugen et al., 1998; Cortes et al., 1998). For those HCH isomers and DDT metabolites which show a significant dependence of temperature (p-level < 0.05), it is possible to calculate the enthalpy of volatilization ( $\Delta H$ , KJ mol<sup>-1</sup>) from the slope of the regression by:

$$\Delta H = \frac{\text{slope}}{2.3} R \quad (4)$$

## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where  $R$  is the gas constant and 2.3 the conversion factor from decimal to natural log. The values of  $\Delta H$  for  $\alpha$ -HCH,  $\gamma$ -HCHs and  $\delta$ -HCH are  $40 \pm 12$ ,  $80 \pm 13$  and  $64 \pm 12$  KJ mol<sup>-1</sup> respectively. The difference in temperature dependence between  $\alpha$ -HCH and  $\gamma$ -HCH isomers has also been observed in other studies (Haugen et al., 1998; Cortes et al., 1998; Poissant and Kropivnjak, 1996). Calculated values of  $\Delta H$  for p,p'-DDE, o,p'-DDT and p,p'-DDT are  $49 \pm 14$ ,  $29 \pm 16$  and  $38 \pm 12$  KJ mol<sup>-1</sup>. These air-soil enthalpies are consistent with previously reported values from other studies. Grimalt et al. (2004) reported  $\Delta H$  values of 88 kJ mol<sup>-1</sup> for  $\alpha$ -HCH and 43 for  $\gamma$ -HCH; Buehler et al. (2003) reported  $\Delta H$  between 50–65 kJ mol<sup>-1</sup> for  $\gamma$ -HCH, between 21–27 kJ mol<sup>-1</sup> for  $\alpha$ -HCH and between 47–61 kJ mol<sup>-1</sup> for  $\Sigma$ DDT. Aulagnier and Poissant (2005) also reported values in the range 24–36 kJ mol<sup>-1</sup> for DDE and between 14–92 kJ mol<sup>-1</sup> for DDT. Air-soil phase exchange enthalpies reported in this study for OCPs and in other studies for other POPs are consistent, but in general slightly lower than the enthalpies associated with the phase change of a chemical from octanol to air ( $\Delta H_{OA}$ ) which are 61.9 and 65.4 for  $\alpha$ -HCH,  $\gamma$ -HCHs (Hinckley et al., 1999) respectively and 97.8 and 88 for p,p'-DDE and p,p'-DDT respectively (Shoeib and Harner, 2002), suggesting that OCPs sorption to SOM is weaker than sorption to the octanol phase.

However, no significant correlation with temperature was observed for HCB. A breakthrough of HCB has been considered in the literature (Lee et al., 1999) as a potential artifact to explain the lack of temperature dependence, due to the high vapor pressure of HCB in comparison to the rest of OCPs. However its vapor pressure is in the same range of those for phenanthrene for which temperature dependence was observed in a previous reported study at the same sites (Cabrerizo et al., 2011a). This together with the low sampling air volumes (10–14 m<sup>3</sup>) and the estimated breakthrough to be on 80 m<sup>3</sup> for this study, suggest that this artifact may not be important in our results. The dependence of HCB partitioning with temperature is not clear in the literature. While some studies (Cortes et al., 1998; Lee et al., 1999; Kaupp et al., 1996) show temperature dependence with relatively low  $\Delta H$  values, such as 5 kJ mol<sup>-1</sup> at Eagle Harbour, 1.9 and 17 kJ mol<sup>-1</sup> in the UK, other studies show a lack of dependence. Su

## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al. (2006) reported that the monthly variations of HCB in the Arctic were less than a factor of 2, while the temperature varied up to 20 °C. Liu et al. (2009), reported a spatial trend for HCB in winter and spring but not in summer and autumn while Aulagnier and Poissant (2005) reported a negative correlation of HCB with temperature and so apparent negative enthalpies of volatilization. Then, it is possible that the enthalpy of volatilization from the pure compound is similar to the enthalpy of solubilization to SOM in the case of HCB. These suggest that HCB is weakly temperature dependent and other processes are strongly affecting the partitioning behavior, consistent with the lack of temperature dependence of its  $K_{SA}$  in this study and other reported observations.

Several studies over the last years (Finizio et al., 1998; Bidleman and Leone, 1998, 2004) have provided evidence that biodegradation of OCPs is occurring in field soils by looking at chiral signatures of selected OCPs such as  $\alpha$ -HCH, o,p'-DDT and o,p'-DDD and studying the preferential breakdown in field samples in order to assess fresh inputs or aged OCPs re-emissions. In this study, soil fugacity measurements were done under equilibrium conditions, thus degradation may not be affecting the  $K_{SA}$ . If degradation is occurring in these soils, the degradation products of OCPs will potentially be more bioavailable than the parent compound and so will be more prone to undergo in soil-air exchange.

To elucidate differences in soil retention capacity for the selected OCPs in the different sampling sites, the soil organic carbon-air partition coefficient ( $K_{OCA}$ ) was calculated from the soil-air partition coefficient ( $K_{SA}$ ), where  $K_{OCA}$  is given by:

$$K_{OCA} = K_{SA} / f_{OC} \quad (5)$$

$K_{OCA}$  were regressed against the octanol-air partition coefficient ( $K_{OA}$ ). Figure S6 shows the log  $K_{OCA}$  versus log  $K_{OA}$  for the sampling campaigns, in different seasons, in the rural and semi-rural Ebro sites (Spain) and for the 2 sampling sites (Hazelrigg and Langden) selected in the UK.  $K_{OA}$  values for individual HCH isomers, DDT metabolites and HCB were estimated using the values proposed by Shoeib and Harner (2002) and corrected by the in-situ measured temperature at each sampling site. The plots of log

$K_{OCA}$  versus  $\log K_{OA}$  suggest that  $K_{OA}$  is an excellent descriptor of soil-air partitioning, with  $r^2$  ranging from 0.27–0.74 and p-level < 0.001 when all pesticides and all sampling sites are considered. As shown in Fig. S6,  $\log K_{OCA}$  values obtained at different sampling sites lie in a range of 2 log units for HCH isomers and HCB (left part of the figure) and 5 log units for DDT metabolites (right part of the figure), which indicates differences in the soil retention capacity depending on the site. Sampling sites in the UK show, in general, similar or slightly higher values of  $\log K_{OCA}$  for low molecular weight OC pesticides (HCH isomers and HCB) at the same  $\log K_{OA}$  value than sampling sites in the Ebro River's watershed. However, higher values of  $\log K_{OCA}$  were observed for DDT isomers in UK sites than in the Ebro River's watershed, highlighting the Langden site. In order to obtain compound specific values for the fitting parameters, explaining the potential role of temperature and organic matter type, compound specific  $K_{OCA}$  values were correlated with the various variables (temperature, altitude,  $f_{OC}/f_N$ , pH....). Single regressions were performed using the equation  $\log K_{OCA} = a + b(X_n)$ , with  $a$  and  $b$  being the fitting parameters and  $X_n$  the parameters considered. Statistical analyses were only done for those OCPs having at least  $n = 15$  pairs of soil-air measurements (Fig. S7 and Tables S6a–c). Altitude, Soil  $T$  and pH were statistically correlated ( $p < 0.05$ ) when regressed against HCH isomers and DDT metabolites  $K_{OCA}$  in single regressions, while  $f_{OC}/f_N$ , an indirect parameter of the organic matter quality, was only statistically positively correlated when regressed against  $K_{OCA}$  for DDT metabolites but not statistically significant correlations were observed for HCH, thus suggesting that the differences in the soil retention capacity may be attributed to differences in the SOM type as well. Altitude was not correlated to soil  $T$  for these data, so altitude may be also related to differences in SOM due to different vegetation types which can contribute to the SOM quality.

# Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3.5 Secondary sources and soil-air exchange controls on atmospheric concentrations of OCPs

Measurements of the air concentration equilibrated with the soil and the atmospheric concentration, both transformed into fugacities of OCPs in soil and ambient air, were compared to infer the direction of air-soil exchange. When fugacity in soil is higher than in ambient air there is a net volatilization of chemical. In contrast, if the fugacity in ambient air is higher than in soil, there is a net deposition of the chemical. Figure 4 shows soil/air fugacity ratios ( $f_s/f_a$ ), in logarithmic scale, for the three sampling campaigns, at different seasons, in the rural and semi-rural Ebro sites and for UK sampling. Bars represent the different sampling sites. Since there is an analytical error of 20 % when analyzing gas phase concentrations with the low volume of sampled air and the soil fugacity sampler is averaging the signal for 24 h, there is an uncertainty associated to the soil-air fugacity ratios, estimated to be lower than 70 % (Cabrerizo et al., 2011a, b). With this in mind, equilibrium would be represented by  $\log f_s/f_a$  in the range of +0.23 and -0.52, being the real uncertainty, lower than this range for many sampling sites.  $f_s/f_a$ , in the Ebro river campaigns, indicates near equilibrium conditions for selected HCH isomers ( $\alpha$ -HCH,  $\gamma$ -HCH) and DDT metabolites (p,p'-DDE, p,p'-DDT), although a trend for soils to move from equilibrium towards becoming a net source to the atmosphere was observed during warmer periods (sampling campaigns carried out in early and late summer, June 2006 and September 2007). Conversely, during colder sampling periods (November 2006) the soils tended to move from equilibrium towards a net sink of atmospheric OCPs. Soils appear to move from equilibrium (p,p'-DDT) to net atmospheric sinks ( $\gamma$ -HCH and p,p'-DDE) even in the late summer in the NW England environment. The fact that UK soils behave as sinks of atmospheric OCPs may be due to higher SOM content or lower temperatures. In fact, Fig. 5 shows  $f_s/f_a$  ratios regressed against  $1/T$  for individual HCH isomers and DDT metabolites. Higher  $f_s/f_a$  ratios were observed when increasing the temperature, suggesting that the tendency of the soil to act as a secondary source of OCPs to the atmosphere is temperature

## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





dependent. The behavior of HCB is interesting since all soils seem to be moving from equilibrium to act as a clear net source of HCB to the atmosphere, even in Ebro's sites in winter and in the UK. Unfortunately the behavior of Ebro soils in summer could not be assessed due to the chromatographic co-elutions as previously was pointed.

In those sampling periods when there is net volatilization, and to a lesser extent when air and soil are close to equilibrium, it is possible that soils act as significant sources controlling the atmospheric occurrence of OCPs. Indeed, the left part of Fig. 6 shows significant correlations between fugacities in soil and in ambient air when all HCH isomers ( $\alpha$ -HCH,  $\beta$ -HCH,  $\gamma$ -HCH, and  $\delta$ -HCH), and all DDT metabolites (p,p'-DDE, o,p'-DDD, o,p'-DDT and p,p'-DDT) for the Ebro sampling sites are considered. Therefore, in the Ebro River's water shed atmosphere, in which there is a seasonal net source of OCPs to the atmosphere with a net volatilization of HCH and DDT during warm periods, the soils exert a clear control on the atmospheric occurrence of OCPs. Indeed, the levels of HCH and DDT found in the atmosphere are significantly correlated with HCH and DDT which escape from soil (fugacity in soil) with  $r^2$  ranging between 0.74 and 0.76 ( $p$ -level  $< 0.001$ ) independent on the season. Similarly, the soil fugacity of individual HCH isomers was statistically correlated ( $p$ -level  $< 0.001$ ) with their ambient air fugacity, with  $r^2$  ranging between 0.69–0.84 (Fig. S8a). This trend is in agreement with results for PCBs reported elsewhere (Cabrerizo et al., 2011b), and indicates that atmospheric levels of banned HCH, DDT, and PCB in the Ebro River basin atmosphere are mostly controlled by re-emissions from soils. This trend is not as clear at UK sampling sites (right part of Fig. 6), as we have observed the soil is still a net sink for most OCPs. Therefore, the comparisons between regions show a spatial pattern. These findings suggest that secondary sources at mid latitudes, as is the case of background soils from NE Spain are playing an important role of secondary sources, and OCPs can be exchanging between the soil and the atmosphere. In contrast, regions where  $\log K_{SA}$  values are elevated, due to higher SOM content or lower temperatures, as is the case at the UK sampling sites, soils will act as larger traps and accumulate more OCPs, since soils are still under-saturated with respect to the

## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





atmospheric OCPs levels. Therefore, these soils will need longer times, during which primary sources of OCPs decrease, to become a secondary source to the atmosphere. The extent of secondary sources may be related to  $K_{SA}$ , which depends on  $T$ , SOM and to a lesser extent soil type. Soils with higher retention potential (higher  $K_{SA}$ ) may still be net sinks of legacy POPs. If this is the correct scenario, then the log  $K_{SA}$  should be inversely correlated with the log  $f_s/f_a$  ratio. Figure 7 shows the negative correlation between log  $K_{SA}$  and log  $f_s/f_a$  for the data set reported here, for selected HCH isomers, DDT metabolites and HCB. This is consistent with the results reported in a previous paper (Cabrerizo et al., 2011b) for other POPs, PCBs, for which we observed the same trend only for some of the more volatile PCB congeners, those having similar physico-chemical properties than OCPs, while no statistically significant correlation was observed for most chlorinated congeners. Figure 7 clearly shows that at high  $K_{SA}$  values, soils act as a sink of atmospheric OCPs ( $f_s/f_a < 1$ ), while in regions and periods with low  $K_{SA}$  (low SOM content and high  $T$ ), soils are a source of OCPs to the atmosphere ( $f_s/f_a > 1$ ). This finding is important since after decades of restriction in the use of OCPs, soils at higher latitudes and in cold environments are still important sinks of atmospheric POPs. This is in agreement with the studies of Schuster et al. (2010, 2011), in European background regions of UK and Norway. In these regions with soils with high SOM content and low temperatures, they found evidence that diffuse primary sources of POPs are still dominating the atmospheric burden of POPs in high latitudes and secondary sources, such as re-volatilization from soil surfaces are playing a minor role.

## 4 Conclusions

The present study shows that in the current balance of primary and secondary sources of OCPs in European temperate regions, soils with low SOM and relatively warm climate are a seasonal secondary source of HCH and DDT to the atmosphere, while in colder climates and regions with high SOM will act as a net sink, because soil

## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



concentrations are still under-saturated with respect the atmospheric concentrations and diffuse primary sources may still be playing an important role. Whereas biodegradation completely removes OCPs from the environment, volatilization from secondary sources leads to the re-cycling of OCPs between environmental compartments mediated by atmospheric re-mobilization. This volatilization from low latitudes may have an important effect on levels and inputs in other regions such as the high latitude compartments.

**Supplementary material related to this article is available online at:**  
**[http://www.atmos-chem-phys-discuss.net/11/25937/2011/](http://www.atmos-chem-phys-discuss.net/11/25937/2011/acpd-11-25937-2011-supplement.pdf)**  
**[acpd-11-25937-2011-supplement.pdf](http://www.atmos-chem-phys-discuss.net/11/25937/2011/acpd-11-25937-2011-supplement.pdf).**

*Acknowledgements.* This research project was funded by the European Commission under the Global Change and Ecosystems (FP6) Water Cycle and Soil Related Aspects (AQUATERRA, Project number 505428 GOCE and from the Spanish Ministry of Science and Innovation through the SCARCE Consolider project. A. C. acknowledges a Ph.D. pre-doctoral fellowship from the Spanish Ministry of Education and Science.

## References

- Aulagnier, F. and Poissant, L.: Some Pesticides Occurrence in Air and Precipitation in Québec, Canada, *Environ. Sci. Technol.*, 39, 2960–2967, 2005.
- Barber, J. L., Sweetman, A. J., van Wijk, D., and Jones, K. C.: Hexachlorobenzene in the global environment: Emissions, levels, distribution, trends and processes, *Sci. Total Environ.*, 349, 1–44, 2005.
- Blais, J. M., Schindler, D. W., Muir, D. C. G., Kimpe, L. E., Donald, D. B, and Rosenberg, B.: Accumulation of persistent organochlorine compounds in mountains of western Canada, *Nature.*, 395, 585–588, 1998.
- Bidleman T. F.: Atmospheric transport and air-surface exchange of pesticides, *Water Air Soil Pollut.*, 115, 115–166, 1999.

ACPD

11, 25937–25965, 2011

## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Bidleman, T. F. and Leone, A. D.: Soil-air exchange of organochlorine pesticides in the Southern United States, *Environ Pollut.*, 128, 49–57, 2004.
- Bidleman, T. F., Jantunen, L. M., Harner, T., Wiberg, K., Wideman, J. L., Brice, K., Su, K., Falconer, R. L., Aigner, E. J., Leone, A. D., Ridal, J. J., Kerman, B., Finizio, A., Alegria, H., Parkhurst, W. J., and Szeto, S. Y.: Chiral pesticides as tracers of air-surface exchange, *Environ Pollut.*, 102, 43–49, 1998.
- Borghini, F., Grimalt, J. O., Sanchez-Hernandez, J. C., Barra, R., García, C. J. T., and Forcardi, S.: Organochlorine compounds in soils and sediments of the mountain Andean Lakes, *Environ Pollut.*, 136, 253–266, 2005.
- Buehler, S. S., Basu, I., and Hites, R. A.: Causes of Variability in Pesticide and PCB Concentrations in Air near the Great Lakes, *Environ. Sci. Technol.*, 38, 414–422, 2003.
- Cabrerizo, A., Dachs, J., and Barcelo, D.: Development of a Soil Fugacity Sampler for Determination of Air-Soil Partitioning of Persistent Organic Pollutants under Field Controlled Conditions, *Environ. Sci. Technol.*, 43, 8257–8263, 2009.
- Cabrerizo, A., Dachs, J., Moeckel, C., Ojeda, M.-J., Caballero, G., Barcelo, D., and Jones, K. C.: Ubiquitous Net Volatilization of Polycyclic Aromatic Hydrocarbons from Soils and Parameters Influencing Their Soil-Air Partitioning, *Environ. Sci. Technol.*, 45, 4740–4747, 2011a.
- Cabrerizo, A., Dachs, J., Moeckel, C., Ojeda, M.-J., Caballero, G., Barcelo, D., and Jones, K. C.: Factors Influencing the Soil-Air Partitioning and the Strength of Soils as a Secondary Source of Polychlorinated Biphenyls to the Atmosphere, *Environ. Sci. Technol.*, 45, 4785–4792, 2011b.
- Cortes, D. R., Basu, I., Sweet, C. W., Brice, K. A., Hoff, R. M., and Hites, R. A.: Temporal Trends in Gas-Phase Concentrations of Chlorinated Pesticides Measured at the Shores of the Great Lakes, *Environ. Sci. Technol.*, 32, 1920–1927, 1998.
- Dalla Valle, M., Jurado, E., Dachs, J., Sweetman, A. J., and Jones, K. C.: The maximum reservoir capacity of soils for persistent organic pollutants: implications for global cycling, *Environ Pollut.*, 134, 153–164, 2005.
- Finizio, A., Bidleman, T. F., and Szeto, S. Y.: Emission of chiral pesticides from an agricultural soil in the fraser Valley, british columbia, *Chemosphere.*, 36, 345–355, 1998.
- Gioia, R., Sweetman, A. J., and Jones, K. C.: Coupling Passive Air Sampling with Emission Estimates and Chemical Fate Modeling for Persistent Organic Pollutants (POPs):? A Feasibility Study for Northern Europe, *Environ. Sci. Technol.*, 41, 2165–2171, 2007.
- Gouin, T., Thomas, G. O., Cousins, I., Barber, J., Mackay, D., and Jones, K. C.: Air-surface

exchange of polybrominated diphenyl ethers and polychlorinated biphenyls, Environ. Sci. Technol., 38, 1426–1434, 2002.

Gregor, D. J. and Gummer, W. D.: Evidence of atmospheric transport and deposition of organochlorine pesticides and polychlorinated biphenyls in Arctic snow, Environ. Sci. Technol., 23, 1528–1531, 1989.

Grimalt, J. O., Fernandez, P., Berdie, L., Vilanova, R. M., Catalan, J., Psenner, R., Hofer, R., Appleby, P. G., Rosseland, B. O., Lien, L., Massabuau, J. C., and Battarbee, R. W.: Selective Trapping of Organochlorine Compounds in Mountain Lakes of Temperate Areas, Environ. Sci. Technol., 35, 2690–2697, 2001.

Grimalt, J. O., van Drooge, B. L., Ribes, A., Vilanova, R. M., Fernandez, P., and Appleby, P.: Persistent organochlorine compounds in soils and sediments of European high altitude mountain lakes, Chemosphere, 54, 1549–1561, 2004.

Halse, A. K., Schlabach, M., Eckhardt, S., Sweetman, A., Jones, K. C., and Breivik, K.: Spatial variability of POPs in European background air, Atmos. Chem. Phys., 11, 1549–1564, doi:10.5194/acp-11-1549-2011, 2011.

Harner, T., Shoeib, M., Diamond, M., Stern, G., and Rosenberg, B.: Using Passive Air Samplers To Assess Urban-Rural Trends for Persistent Organic Pollutants. 1. Polychlorinated Biphenyls and Organochlorine Pesticides, Environ. Sci. Technol., 38, 4474–4483, 2004.

Haugen, J.-E., Wania, F., Ritter, N., and Schlabach, M.: Hexachlorocyclohexanes in Air in Southern Norway. Temporal Variation, Source Allocation, and Temperature Dependence, Environ. Sci. Technol., 32, 217–224, 1998.

Hayward, S. J., Gouin, T., and Wania, F.: Levels and Seasonal Variability of Pesticides in the Rural Atmosphere of Southern Ontario, J. Agric. Food Chem., 58, 1077–1084, 2003.

Hinckley, D. A., Bidleman, T. F., Foreman, W. T., and Tuschall, J. R.: Determination of vapor pressures for nonpolar and semipolar organic compounds from gas chromatographic retention data, J. Chem. Eng. Data., 35, 232–237, 1999.

Kaupp, H., Dörr, G., Hippelein, M., McLachlan, M. S., and Hutzinger, O.: Baseline contamination assessment for a new resource recovery facility in Germany Part IV: Atmospheric concentrations of polychlorinated biphenyls and hexachlorobenzene, Chemosphere, 32, 2029–2042, 1996.

Kurt-Karakus, P. B., Bidleman, T. F., Staebler, R. M., and Jones, K. C.: Measurement of DDT Fluxes from a Historically Treated Agricultural Soil in Canada, Environ. Sci. Technol., 40, 4578–4585, 2006.

ACPD

11, 25937–25965, 2011

## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Lee, R. G. M., Burnett, V., Harner, T., and Jones, K. C.: Short-Term Temperature-Dependent Air-Surface Exchange and Atmospheric Concentrations of Polychlorinated Naphthalenes and Organochlorine Pesticides, *Environ. Sci. Technol.*, 34, 393–398, 1999.
- Liu, X., Zhang, G., Li, J., Yu, L.-L., Xu, Y., Li, X.-D., Kobara, Y., and Jones, K. C.: Seasonal Patterns and Current Sources of DDTs, Chlordanes, Hexachlorobenzene, and Endosulfan in the Atmosphere of 37 Chinese Cities, *Environ. Sci. Technol.*, 43, 1316–1321, 2009.
- Mackay, D., Paterson, S., and Schroeder, W. H.: Model describing the rates of transfer processes of organic chemicals between atmosphere and water, *Environ. Sci. Technol.*, 20, 810–813, 1986.
- Meijer, S. N., Shoeib, M., Jones, K. C., and Harner, T.: Air-Soil Exchange of Organochlorine Pesticides in Agricultural Soils. 2. Laboratory Measurements of the Soil-Air Partition Coefficient, *Environ. Sci. Technol.*, 37, 1300–1305, 2003a.
- Meijer, S. N., Ockenden, W. A., Sweetman, A., Breivik, K., Grimalt, J. O., and Jones, K. C.: Global Distribution and Budget of PCBs and HCB in Background Surface Soils: Implications for Sources and Environmental Processes, *Environ. Sci. Technol.*, 37, 667–672, 2003b.
- Nam, J. J., Gustafsson, O., Kurt-Karakus, P., Breivik, K., Steinnes, E., and Jones, K. C.: Relationships between organic matter, black carbon and persistent organic pollutants in European background soils: Implications for sources and environmental fate, *Environ. Pollut.*, 156, 809–817, 2008.
- NOAA ARL HYSPLIT Model, available at: <http://ready.arl.noaa.gov/HYSPLIT.php>, 2011.
- Ockenden, A. W., Prest, H. F., Thomas, G. O., Sweetman, A. J., and Jones, K. C.: Toward an understanding of a global distribution of persistent organic Pollutants: The use of semipermeable membrane devices as time-integrated passive samplers, *Environ. Sci. Technol.*, 32, 1538–1543, 1998.
- Ockenden, W. A., Breivik, K., Meijer, S. N., Steinnes, E., Sweetman, A. J., and Jones, K. C.: The global re-cycling of persistent organic pollutants is strongly retarded by soils, *Environ. Pollut.*, 121, 75–80, 2003.
- Poissant, L. and Koprivnjak, J.-F.: Fate and Atmospheric Concentrations of  $\alpha$ - and  $\gamma$ -Hexachlorocyclohexane in Québec, Canada, *Environ. Sci. Technol.*, 30, 845–851, 1996.
- Ribes, A., Grimalt, J. O., Torres García, C. J., and Cuevas, E.: Temperature and Organic Matter Dependence of the Distribution of Organochlorine Compounds in Mountain Soils from the Subtropical Atlantic (Teide, Tenerife Island), *Environ. Sci. Technol.*, 36, 1879–1885, 2002.
- Růžicková, P., Klánová, J., Čupr, P., Lammel, G., and Holoubek, I.: An Assessment of Air-Soil

## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Exchange of Polychlorinated Biphenyls and Organochlorine Pesticides Across Central and Southern Europe, Environ. Sci. Technol., 42, 179–185, 2007.

Shoeib, M. and Harner, T.: Using measured octanol-air partition coefficients to explain environmental partitioning of organochlorine pesticides, Environ. Toxicol. Chem., 21, 984–990, 2002.

Schuster, J. K., Gioia, R., Sweetman, A. J., and Jones, K. C.: Temporal Trends and Controlling Factors for Polychlorinated Biphenyls in the UK Atmosphere (1991–2008), Environ. Sci. Technol., 44, 8068–8074, 2010.

Schuster, J. K., Gioia, R., Moeckel, C., Agarwal, T., Bucheli, T. D., Breivik, K., Steinnes, E., and Jones, K. C.: Has the Burden and Distribution of PCBs and PBDEs Changed in European Background Soils between 1998 and 2008? Implications for Sources and Processes, Environ. Sci. Technol., 45, 7291–7297, 2011.

Su, Y., Hung, H., Blanchard, P., Patton, G. W., Kallenborn, R., Konoplev, A., Fellin, P., Li, H., Geen, C., Stern, G., Rosenberg, B., and Barrie, L. A.: Spatial and Seasonal Variations of Hexachlorocyclohexanes (HCHs) and Hexachlorobenzene (HCB) in the Arctic Atmosphere, Environ. Sci. Technol., 40, 6601–6607, 2006.

Thomas, G. O., Sweetman, A. J., Parker, C. A., Kreibich, H., and Jones, K. C.: Development and validation of methods for the trace determination of PCBs in biological matrices, Chemosphere, 36, 2447–2459, 1998.

Wang, X., Ren, N., Qi, H., Ma, W., and Li, Y.: Levels, distributions, and source identification of organochlorine pesticides in the topsoils in Northeastern China, J. Environ. Sci., 21, 1386–1392, 2009.

Yao, Y., Tuduri, L., Harner, T., Blanchard, P., Waite, D., Poissant, L., Murphy, C., Belzer, W., Aulagnier, F., Li, Y.-F., and Sverko, E.: Spatial and temporal distribution of pesticide air concentrations in Canadian agricultural regions, Atmos. Environ., 40, 4339–4351, 2006.

Zhang, J.-J., Wen, B., Shan, X.-Q., Zhang, S., and Khan, S. U.: Temporal change in the distribution patterns of hexachlorobenzene and dichlorodiphenyltrichloroethane among various soil organic matter fractions, Environ. Pollut., 150, 234–242, 2007.

Zhang, N., Yang, Y., Tao, S., Liu, Y., and Shi, K.-L.: Sequestration of organochlorine pesticides in soils of distinct organic carbon content, Environ. Pollut., 159, 700–705, 2011.

# Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures




Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Fig. 1.** Ambient air concentration ( $\text{Log Ca}$  in  $\text{ng m}^{-3}$ ) regressed against  $1/T$  for selected HCH isomers, DDT metabolites and HCB.



**Fig. 2.** Concentrations in soil (Log Cs in  $\text{ng gdw}^{-1}$ ) versus soil organic carbon (Log  $f_{\text{OC}}$ ).

## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion







**Fig. 3.** Regression of soil-air partition coefficient ( $\text{Log } K_{SA}$ ) versus temperature ( $1/T$ ) for selected HCH isomers and DDT metabolites.

## Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



# Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.



**Fig. 4.** Soil to air fugacity ratio ( $f_s/f_a$ ) in logarithmic scale for the different sampling campaigns in Ebro (Spain) and UK. Dash lines indicates equilibrium (range +0.23, -0.52).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



# Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.



**Fig. 5.** Examples of soil to air fugacity ratio ( $\text{Log } f_s/f_a$ ) regressed against  $1/T$ .

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Soil-air exchange of  
organochlorine  
pesticides

A. Cabrerizo et al.



**Fig. 6.** Fugacity in soil ( $f_s$ ) versus fugacity in air ( $f_a$ ) for all HCH ( $\alpha$ -HCH,  $\gamma$ -HCH,  $\beta$ -HCH,  $\delta$ -HCH) isomers and all DDT metabolites (p,p'-DDE, p,p'-DDD, o,p'-DDT and p,p'-DDT) for sampling sites in Spain (figures on the left) and sampling sites in UK (figures on the right).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



# Soil-air exchange of organochlorine pesticides

A. Cabrerizo et al.



**Fig. 7.** Soil-air partition coefficient ( $\text{Log } K_{SA}$ ) regressed versus the soil-air fugacity ratio ( $\text{Log } f_s/f_a$ ) for selected OCPs.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

