

**Satellite-derived  
Arctic tropospheric  
BrO columns**

S. Choi et al.

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

# Analysis of satellite-derived Arctic tropospheric BrO columns in conjunction with aircraft measurements during ARCTAS and ARCPAC

S. Choi<sup>1</sup>, Y. Wang<sup>1</sup>, R. J. Salawitch<sup>2</sup>, T. Canty<sup>2</sup>, J. Joiner<sup>3</sup>, T. Zeng<sup>1</sup>,  
T. P. Kurosu<sup>4,\*</sup>, K. Chance<sup>4</sup>, A. Richter<sup>5</sup>, L. G. Huey<sup>1</sup>, J. Liao<sup>1</sup>, J. A. Neuman<sup>6,7</sup>,  
J. B. Nowak<sup>6,7</sup>, J. E. Dibb<sup>8</sup>, A. J. Weinheimer<sup>9</sup>, G. Diskin<sup>10</sup>, T. B. Ryerson<sup>7</sup>,  
A. da Silva<sup>3</sup>, J. Curry<sup>1</sup>, D. Kinnison<sup>9</sup>, S. Tilmes<sup>9</sup>, and P. F. Levelt<sup>11,12</sup>

<sup>1</sup>Georgia Institute of Technology, Atlanta, GA, USA

<sup>2</sup>University of Maryland College Park, College Park, MD, USA

<sup>3</sup>NASA Goddard Space Flight Center, Greenbelt, MD, USA

<sup>4</sup>Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, USA

<sup>5</sup>Institute of Environmental Physics, University of Bremen, Bremen, Germany

<sup>6</sup>Cooperative Institute for Research in Environmental Sciences (CIRES), University of Colorado Boulder, Boulder, CO, USA

<sup>7</sup>NOAA Earth System Research Laboratory, Boulder, CO, USA

<sup>8</sup>University of New Hampshire, Durham, NH, USA

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Satellite-derived  
Arctic tropospheric  
BrO columns**

S. Choi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

<sup>9</sup> National Center for Atmospheric Research, Boulder, CO, USA

<sup>10</sup> NASA Langley Research Center, Hampton, VA, USA

<sup>11</sup> Royal Netherlands Meteorological Institute (KNMI), De Bilt, The Netherlands

<sup>12</sup> University of Technology Eindhoven, Eindhoven, The Netherlands

\* now at: NASA Jet Propulsion Laboratory, Pasadena, CA, USA

Received: 17 August 2011 – Accepted: 12 September 2011 – Published: 21 September 2011

Correspondence to: S. Choi (sungyeon.choi@eas.gatech.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

## Abstract

We derive estimates of tropospheric BrO column amounts during two Arctic field campaigns in 2008 using information from the satellite UV nadir sensors Ozone Monitoring Instrument (OMI) and the second Global Ozone Monitoring Experiment (GOME-2) as well as estimates of stratospheric BrO columns from a model simulation. The sensitivity of the satellite-derived tropospheric BrO columns to various parameters is investigated using a radiative transfer model. We conduct a comprehensive analysis of satellite-derived tropospheric BrO columns including a detailed comparison with aircraft in-situ observations of BrO and related species obtained during the field campaigns. In contrast to prior expectation, tropospheric BrO, when present, existed over a broad range of altitudes. Our results show reasonable agreement between tropospheric BrO columns derived from the satellite observations and columns found using aircraft in-situ BrO. After accounting for the stratospheric contribution to total BrO column, several events of rapid BrO activation due to surface processes in the Arctic are apparent in both the OMI and GOME-2 based tropospheric columns. The wide orbital swath of OMI allows examination of the evolution of tropospheric BrO on about hourly time intervals near the pole. Low pressure systems, strong surface winds, and high planetary boundary layer heights are associated with the observed tropospheric BrO activation events.

## 1 Introduction

Bromine plays an important role in tropospheric ozone chemistry and the resulting oxidation capacity of the polar boundary layer. Bromine radicals catalytically destroy ozone, leading to nearly complete removal near the surface that is termed an ozone ( $O_3$ ) depletion event (ODE). Once  $O_3$  is depleted, high levels of reactive halogen species including atomic bromine (e.g. Br) become the primary oxidants for many species, including methane ( $CH_4$ ) and mercury (Hg) (e.g. Simpson et al., 2007b;

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion




Schroeder et al., 1998). During mercury depletion events (MDEs), Br and BrO are thought to oxidize elemental mercury to more reactive gaseous mercury that deposit to the polar ecosystem (Schroeder et al., 1998; Lu et al., 2001; Ariya et al., 2004; Douglas et al., 2005; Holmes et al., 2010).

5 A primary source of active bromine in the polar boundary layer is thought to be bromide ( $\text{Br}^-$ ) in the condensed phase, which is transformed to gaseous molecular bromine ( $\text{Br}_2$ ) by heterogeneous chemistry. Potential sources of bromide are sea salt aerosols (Fan and Jacob, 1992), surfaces of first year sea ice (Simpson et al., 2007a; Wagner et al., 2007) or newly formed sea ice (Jones et al., 2006), frost flowers  
10 (Kaleschke et al., 2004), or blowing snow triggered by strong winds (Yang et al., 2008, 2010; Jones et al., 2009; A. E. Jones et al., 2010; Begoin et al., 2010; Theys et al., 2011; Toyota et al., 2011).  $\text{Br}_2$  released to the atmosphere is rapidly photolyzed to yield bromine radicals (Br), i.e.



The bromine radicals then lead to catalytic ozone loss, i.e.



20 Barrie et al. (1988) found that filterable bromine is strongly anti-correlated with the abundance of  $\text{O}_3$  in the polar boundary layer. Following this work, many additional measurements of bromine species along with  $\text{O}_3$  have been made in polar environments (e.g. Bottenheim et al., 1990; Barrie et al., 1994; Hausmann and Platt, 1994; Tuckermann et al., 1997; Martinez et al., 1999; Ridley et al., 2003). The association

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





of elevated BrO and depleted surface O<sub>3</sub> during Arctic spring is well established (e.g. Platt and Hönninger, 2003, and references therein).

Bromine monoxide (BrO), an intermediate in the catalytic loss of ozone, is the most commonly observed active bromine species. BrO absorbs ultraviolet (UV) radiation, which enables measurement using remote sensing techniques. Reported in-situ and remotely-sensed ground-, balloon-, and aircraft-based BrO measurements include: (1) boundary layer mixing ratio with Long Path-Differential Optical Absorption Spectroscopy (LP-DOAS) (Hausmann and Platt, 1994; Tuckermann et al., 1997; Martinez et al., 1999); (2) mixing ratio by aircraft in-situ Chemical-Ionization Mass Spectrometry (CIMS) (Neuman et al., 2010; Liao et al., 2011a,b); (3) boundary layer and free tropospheric BrO column using DOAS on aircraft (McElroy et al., 1999); (4) total, stratospheric, and tropospheric columns using ground-based zenith sky and direct sun DOAS (Frieß et al., 1999; Schofield et al., 2004, 2006; Hendrick et al., 2007, 2008; Theys et al., 2007); (5) boundary-layer column with ground-based MAX-DOAS (Hönninger, 2004; Simpson et al., 2007a; Donohoue et al., 2010); (6) balloon profiles using a variety of spectroscopic methods (Fitzenberger et al., 2000; Pfeilsticker et al., 2000; Pundt et al., 2002; Dorf et al., 2008). Despite the many measurements of BrO and related species obtained by various techniques, significant uncertainties remain regarding the importance of very short lived source compounds on the stratospheric bromine budget (e.g. Sect. 1.3.3.3 of WMO, 2011) as well as the magnitude of the global, ubiquitous, background level of tropospheric BrO (e.g. Sect. 5.3 of Theys et al., 2011).

Space-based observation of BrO offers an excellent tool for studying Arctic polar bromine chemistry. Satellite observations provide global coverage, far superior to the spatial coverage available from ground- and aircraft-based measurements. Limb-sounding observations from the Microwave Limb Sounder (MLS) (Livesey et al., 2006; Kovalenko et al., 2007) provide quantification of BrO profiles in the upper and middle stratosphere. The MLS observations of BrO imply a significant contribution to stratospheric bromine from sources other than long-lived CH<sub>3</sub>Br and halons. Limb

**Satellite-derived  
Arctic tropospheric  
BrO columns**

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 observations from the SCanning Image Absorption SpectroMeter for Atmospheric CHartographY (SCIAMACHY) (Sinnhuber et al., 2005; Sioris et al., 2006; Rozanov et al., 2011) extend into the lowermost stratosphere (LMS) and nadir measurements from SCIAMACHY constrain total column BrO, including contributions from the strato-  
10 sphere and troposphere. The retrievals of SCIAMACHY BrO by Rozanov et al. (2011) and Sinnhuber et al. (2005) imply a limited role for supply of bromine by very short lived (VSL) compounds to the LMS and a considerable burden of global, ubiquitous, background BrO whereas the retrieval of SCIAMACHY BrO described by Sioris et al. (2006) implies a larger role for supply of stratospheric Br<sub>y</sub> by VSL compounds and a  
15 much smaller level for background tropospheric BrO (e.g. Sect. 2.5.2.1 of WMO, 2007).

Nadir-viewing instruments on polar-orbiting satellites provide multiple daily observations of total column BrO at high latitude. The high surface albedo of polar regions provides good sensitivity to tropospheric BrO, including that near the surface (Wagner and Platt, 1998; Theys et al., 2011). Chance (1998) and Hegels et al. (1998) retrieved BrO  
20 total vertical column densities (VCD) from nadir radiances in the UV obtained by Global Ozone Monitoring Experiment (GOME) (GOME Users Manual, 1995) on the European Space Agency (ESA) European Remote Sensing 2 (ERS-2) satellite. The early GOME observations showed large enhancements over Hudson Bay during spring 1997, which was attributed to bromine release from the surface (Chance, 1998). Estimates of BrO  
25 total column amount have been subsequently derived from other nadir-viewing satellite sensors including the Ozone Monitoring Instrument (OMI) (Levelt et al., 2006) on the US National Aeronautics and Space Administration (NASA) Aura satellite, SCIAMACHY (Bovensmann et al., 1999) on the ESA Environmental Satellite (EnviSat), and the second Global Ozone Monitoring Experiment (GOME-2) instruments (Munro et al., 2006) that are flying on the series of European Meteorological Operational Satellites (EuMetSat MetOp). Retrievals of VCD BrO from different satellite instruments and by various groups tend to be in fairly close agreement. All satellite instruments report levels considerably larger than predicted by models that account only for stratospheric supply from the decomposition of methyl bromide and halons. The focus of the

**Satellite-derived  
Arctic tropospheric  
BrO columns**

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



scientific community since publication of early GOME observations has been quantifying the degree to which this “excess BrO” resides in the troposphere, the stratosphere, or perhaps both regions of the atmosphere.

Tropospheric BrO columns can be derived from satellite observations using the residual method, in which an estimate of the stratospheric BrO column is subtracted from the satellite-derived VCD of BrO. Wagner and Platt (1998) and Richter et al. (1998) estimated tropospheric VCD of BrO using this approach from GOME observations, based on simple assumptions for stratospheric BrO. These studies and many others assumed that stratospheric BrO is zonally symmetric and that bromine was supplied to the stratosphere solely by the decomposition of long-lived organic compounds. Theys et al. (2009) and Salawitch et al. (2010) showed that the stratospheric VCD of BrO can exhibit strong gradients with respect to longitude at high latitude during spring and Salawitch et al. (2010) questioned prior estimates of residual tropospheric BrO found assuming that the stratospheric burden was zonally symmetric. Theys et al. (2011) derived tropospheric BrO columns from GOME-2 spectra with a model-based stratospheric BrO climatology (Theys et al., 2009) similar to that used here and described below.

Many studies related to bromine and ozone chemistry have been conducted using satellite-derived tropospheric BrO columns. Wagner and Platt (1998) reported elevated regions of BrO vertical column density in the Arctic and Antarctic regions. They noted these enhancements were likely due to increased abundance of tropospheric BrO, rather than a stratospheric disturbance, based on a variety of factors including the correlation between enhanced columns of BrO and the O<sub>2</sub>-O<sub>2</sub> collision complex. Wagner et al. (2001) showed that elevated BrO column amounts observed by GOME were correlated with low ozone in the boundary layer observed in-situ at Ny-Ålesund (Spitsbergen), Norway. In other studies, spatial and temporal features of ODEs have been simulated using 3-dimensional regional chemical transport models and GOME-derived tropospheric BrO columns (Zeng et al., 2003, 2006). Connections between BrO-rich air masses and first-year sea ice have been indicated with back-trajectory analyses

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

using SCIAMACHY data (Wagner et al., 2007). Transport of a large BrO plume near the North Pole is also reported by Begoin et al. (2010). A back trajectory study using satellite-derived tropospheric BrO columns indicated that ODEs can be differentiated into locally activated and transport driven events (Koo et al., 2011).

5 Despite the numerous studies of tropospheric polar bromine chemistry using satellite BrO observations, estimation and interpretation of tropospheric BrO information from space presents ongoing challenges. To properly estimate tropospheric BrO column amounts, the stratospheric contribution to the satellite-derived total column must be accurately represented (e.g. Theys et al., 2009; Salawitch et al., 2010). The global, ubiquitous background tropospheric level of BrO inferred from the satellite record is sensitive to the amount of inorganic bromine ( $\text{Br}_y$ ) delivered to the stratosphere by VSL bromocarbons (Salawitch et al., 2005). Furthermore, low solar elevation angles in the early polar spring lead to large uncertainties in satellite total BrO column retrievals (see below). The presence of clouds further complicates the retrieval of tropospheric BrO from satellite observations. Theys et al. (2011) have recently addressed many of these issues. They showed maps of tropospheric BrO columns derived using a method similar to that described below, and evaluated these columns using ground-based measurements of the tropospheric and stratospheric contributions to the total column. Our study builds upon the work of Theys et al. (2011) by further exploring the issues that affect quantification of tropospheric BrO columns and by using aircraft measurements of BrO to evaluate the satellite-derived tropospheric columns.

15 Salawitch et al. (2010) provided an initial analysis of BrO observations obtained by instruments aboard the NASA and NOAA aircraft (Neuman et al., 2010; Liao et al., 2011b) and the OMI satellite instrument (Kurosu and Chance, 2011) during spring 2008. They simulated the stratospheric BrO column assuming VSL bromocarbons supply between 5 and 10 ppt of  $\text{Br}_y$  to the stratosphere in the form of product gas injection (PGI), in addition to the  $\sim 2$  ppt of  $\text{Br}_y$  supplied by source gas injection (SGI) of the VSL species  $\text{CH}_2\text{Br}_2$ . On the other hand, Theys et al. (2011) simulated the stratospheric burden of BrO assuming 1 ppt of PGI and 5 ppt of SGI (all represented as

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

CH<sub>2</sub>Br<sub>2</sub>). The stratospheric BrO burden found by Salawitch et al. (2010) for their “best case” simulation of 7 ppt from PGI and 2 ppt from SGI is considerably larger (~27% overall difference, with quite a bit of geographic variability) than the stratospheric BrO burden used by Theys et al. (2011). Salawitch et al. (2010) concluded that high column amounts of BrO derived from satellite observations could, in some cases, be attributed to compression of stratospheric air to low altitudes (high pressures) and that this condition, not previously recognized in the analysis of the satellite BrO record, should be considered to properly relate the satellite record to surface events. The magnitude of the stratospheric enhancement is sensitive to the supply of stratospheric Br<sub>y</sub> from VSL bromocarbons as well as the pathways of this transport (i.e. PGI versus SGI; see Chapter 2 of WMO (2003) for a detailed discussion of these pathways). Salawitch et al. (2010) showed agreement, to within uncertainties, between total column BrO measured by OMI and the sum of modeled stratospheric and aircraft-measured tropospheric BrO partial columns. However, their treatment of the tropospheric column did not explicitly account for tropospheric air mass factors (AMFs). Our study builds on this prior work by using a radiative transfer model to calculate tropospheric AMFs and also by considering the effects of clouds, surface reflectivity, and viewing geometry on the evaluation of the bromine budget for Arctic spring 2008.

Validation of satellite-derived BrO columns and their relationship to ODEs was a goal of two Arctic field campaigns conducted in 2008 under the auspices of the International Polar Year (IPY). The Arctic Research of the Composition of the Troposphere from Aircraft and Satellite (ARCTAS) mission was conducted in April and June–July 2008 by NASA (D. J. Jacob et al., 2010) and the US National Oceanic and Atmospheric Administration (NOAA) Aerosol, Radiation, and Cloud Processes affecting Arctic Climate (ARCPAC) mission took place in April 2008 (Brock et al., 2011). Various in-situ aircraft measurements of trace chemicals, including O<sub>3</sub> and reactive bromine species (BrO, Br<sub>2</sub>, and soluble bromide), were made during these campaigns. Neuman et al. (2010) and Liao et al. (2011a) provide an overview of the in-situ measurements of BrO and related species obtained during ARCTAS and ARCPAC.

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Satellite-derived  
Arctic tropospheric  
BrO columns**

S. Choi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

In this paper, we use a detailed approach to derive tropospheric BrO column densities from space-based observations. The satellite and other data sets used here are described in Sect. 2. Calculation of residual tropospheric columns using total columns retrieved from OMI and GOME-2 and the stratospheric column of Salawitch et al. (2010) and accounting appropriately for instrument sensitivity is presented in Sect. 3. Quantitative evaluation of our satellite-derived tropospheric BrO vertical columns using the aircraft in-situ observations of active bromine species collected during ARCTAS and ARCPAC is described in Sect. 4. We examine the results in the context of the meteorological conditions. Finally, we discuss the characteristics of rapid BrO enhancement events (“BrO explosions”) inferred from OMI and GOME-2, including potential relationships with meteorological parameters. Conclusions are given in Sect. 5.

## 2 Data description

We use a variety of aircraft and satellite measurements and model outputs to calculate and evaluate satellite-derived tropospheric BrO columns. Tropospheric BrO columns are inferred from aircraft measurements of in-situ BrO (henceforth referred to as “in-situ columns”, as described in Sect. 2.1). We derive tropospheric vertical column densities (VCDs) using BrO slant column densities (SCD) retrieved from OMI and GOME-2 (see Sect. 2.2) and a model simulation of the stratospheric BrO column as detailed in Sect. 2.3. OMI rotational Raman scene pressures and MODIS cloud products (see Sect. 2.4) are used to assess cloud effects on the derived tropospheric BrO columns. Data from a global reanalysis, described in Sect. 2.5, are used to examine relationships between observed BrO enhancement events and meteorological conditions.

### 2.1 Aircraft in-situ measurements

Aircraft in-situ measurements of BrO and various other trace gas concentrations were made from the NASA DC-8 aircraft during ARCTAS and the NOAA WP-3D aircraft



during ARCPAC. An overview of the instruments, flights, and mission goals and accomplishments are provided in the ARCTAS (D. J. Jacob et al., 2010) and ARCPAC (Brock et al., 2011) overview papers. BrO was measured using Chemical Ionization Mass Spectrometer (CIMS) instruments during both experiments (Neuman et al., 2010; Liao et al., 2011b).

DC-8 CIMS measurements of BrO were made once every 30 s, and WP-3D CIMS measurements were acquired once every 2 s. Measurement uncertainties for BrO are  $\pm 40\%$  with a detection limit of 3 pptv for WP-3D data and  $\pm 40\%$  and a detection limit of 2–5 pptv for DC-8 data (Neuman et al., 2010; Liao et al., 2011b). In this study, we use 1 min averaged data for both DC-8 and WP-3D measurements. The spatial resolution of the 1 min averaged aircraft data is approximately 10 km.

We estimate tropospheric BrO columns from the aircraft BrO mixing ratio measurements for each ascent and descent. Figure 1 shows the flight tracks and locations of the 16 (29) profiles made from the DC-8 (WP-3D). BrO profiles are available for 16 and 17 April 2008 from the DC-8 and 12, 15, 18, 19 and 21 April 2008 from the WP-3D. To calculate tropospheric columns from the aircraft data, the profiles were binned on an altitude grid with 500 m resolution from the surface to 7.5 km (BrO measurements are available only up to 7.5 km altitude). For each bin, the median value was selected as the BrO mixing ratio at that altitude. If the median value was less than 0, then we assumed the mixing ratio was 0.

Figure 2a shows the entire collection of DC-8 BrO measurements and the median DC-8 (henceforth referred to as the composite) profile. Figure 2b shows samples of different types of BrO profiles collected during April 2008. Among the 16 DC-8 BrO profiles, 8 have shapes similar to the DC-8 composite profile, while the others show elevated BrO near the surface and/or in layers at altitudes from 2 to 4 km. WP-3D measurements of BrO exhibit larger variability than those from the DC-8 (Fig. 2c, d). The DC-8 and WP-3D instruments reported much lower mixing ratios of BrO near the surface than have been reported in the past by DOAS instruments in the springtime Arctic boundary layer (Hausmann and Platt, 1994; Tuckermann et al., 1997; Martinez

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

et al., 1999). Surprisingly, the highest mixing ratios of BrO tended to be observed above the boundary layer Neuman et al. (2010); Salawitch et al. (2010).

To estimate the tropospheric BrO column from aircraft measurements, we must first make assumptions about mixing ratios between the surface and the lowest altitude sampled by the aircraft. We only take profiles with aircraft minimum altitudes less than 500 m. Then, we assume that the BrO mixing ratio in the lowest bin (surface to 500 m) is the median BrO mixing ratio between the lowest aircraft altitude and 500 m. When the aircraft did not sample up to the tropopause, we made assumptions about mixing ratios between the highest aircraft altitude and the tropopause. Here, we use the upper part of the DC-8 composite profile to fill empty upper bins. However, the DC-8 composite profile only goes up to 7.5 km, the highest altitude where BrO is sampled by the aircraft. We assume that the BrO mixing ratio between 7.5 km and the tropopause is zero. Aircraft data suggest that BrO mixing ratios at these altitudes are very small for the sampled air masses (see Fig. 2). It is possible that stratospheric to tropospheric transport of air could supply BrO to the upper troposphere, particularly along the western flank of Arctic low pressure systems (i.e. after low altitude tropopause systems pass over a region) (Salawitch et al., 2010). This idea is speculative and is not considered below. Finally, layer column BrO amounts (BrO column amounts for vertical bins) are estimated using pressures and temperatures from aircraft measurements and integrated from the surface to the tropopause, to provide a tropospheric column BrO abundance. We use the tropopause height from the Modern Era Retrospective-Analysis for Research and Application (MERRA) data set (see Sect. 2.5).

Tables 1, 2, and 2 provide information related to the aircraft BrO profiles including the bottom and top altitudes of the aircraft in-situ profiles ( $Alt_B$  and  $Alt_T$ , respectively), the MERRA tropopause height ( $H_{TRP}$ ), and the in-situ tropospheric BrO column ( $Col_{IN-SITU}$ ) obtained by integrating the aircraft profiles. Tables 3, 4, and 4 provide solar zenith angle (SZA), OMI reflectivity at 331 nm ( $R_{331\text{nm}}$ ), the difference between terrain and OMI cloud pressure ( $\Delta P_C$ , explained in Sect. 2.4), the satellite-derived tropospheric columns from OMI ( $Col_{OMI}$ ) and GOME-2 ( $Col_{GOME-2}$ ) corresponding to the

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



in-situ aircraft BrO profiles, and the ratios of satellite-derived tropospheric BrO columns to the in-situ BrO columns ( $\text{Ratio}_{\text{OMI}}$  and  $\text{Ratio}_{\text{GOME-2}}$ ).

In addition to BrO, other aircraft trace gas measurements are useful for inferring halogen chemistry and air mass characteristics. For example,  $\text{Br}_2$  was measured by CIMS instruments from both aircraft. Laboratory studies revealed that the  $\text{Br}_2$  signals include contributions from HOBr that reacted on inlet surfaces. Therefore,  $\text{Br}_2$  represents the lower limit of HOBr +  $\text{Br}_2$  (Neuman et al., 2010). Henceforth, we refer to this measurement as “active bromine”. Soluble bromide was also measured from the DC-8 using mist chamber/ion chromatography. Soluble bromide may include inorganic bromine compounds such as HBr, HOBr, particulate bromine, and possibly BrO and  $\text{Br}_2$  (Ridley et al., 2003; Dibb et al., 2010; Neuman et al., 2010; Liao et al., 2011b). Measurements of  $\text{O}_3$  mixing ratios by chemiluminescence were made from both aircraft. CO was measured using tunable diode laser absorption spectroscopy on the DC-8 and UV fluorescence on the WP-3D (D. J. Jacob et al., 2010; Brock et al., 2011).

## 2.2 Total BrO slant column density from OMI and GOME-2

OMI is a nadir-viewing ultraviolet and visible (UV/Vis) sensor (Levelt et al., 2006) aboard the NASA Aura satellite that is in a sun-synchronous orbit with an overpass of 01:38 p.m. local time. The OMI spectral resolution in the UV-2 channel used to retrieve BrO columns is approximately 0.5 nm. The OMI swath width is about 2600 km. The UV-2 pixel size is approximately  $13 \times 24 \text{ km}^2$  at the swath center and significantly larger at the swath edges. With its wide swath, OMI provides multiple daily observations at high latitudes in spring and daily global coverage at low and middle latitudes. An obstruction outside the instrument that produces radiance errors (known as the “row anomaly”) reduced the swath coverage mainly after May 2008; it does not significantly affect the observations shown here (Claas et al., 2010).

OMI BrO SCDs are retrieved by directly fitting backscattered UV radiances to absorption cross-sections of BrO (the target gas),  $\text{NO}_2$ , HCHO, and  $\text{SO}_2$  as well as inelastic rotational-Raman scattering (also known as the Ring effect) using a non-linear least-squares approach (Chance, 1998). The spectral fitting window for the OMI algorithm

### Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



is 319 to 347.5 nm. BrO cross sections from Wilmouth et al. (1999) are used. The total fitting uncertainty of OMI BrO total column typically ranges from 15 to 51 % (Salawitch et al., 2010; Kurosu and Chance, 2011).

GOME-2 is a 4 channel UV/Vis nadir viewing instrument operating on Metop-A since January 2007 (Callies et al., 2000). It has a local equator crossing time of 9:30 am in the descending node, a swath width of 1920 km, and a spatial resolution of  $40 \times 80 \text{ km}^2$ . For the BrO SCD retrieval used here (Begoin et al., 2010), measurements in the window 336 to 347 nm are used, where GOME-2 has a spectral resolution of about 0.3 nm. SCDs of BrO are retrieved using the standard DOAS approach. Absorption due to BrO (Wahner et al., 1988),  $\text{O}_3$ ,  $\text{NO}_2$ , and the effects of rotational-Raman scattering are included. The uncertainty of the GOME-2 BrO total SCDs is 10 to 30 % depending on solar zenith angle (SZA) and surface albedo and the uncertainty has both random and systematic contributions from spectral interferences and the cross-sections. As a result of throughput loss of the UV channels, random errors have increased since launch. However, this effect is not significant during the time period studied here (Dikty et al., 2011).

All of the analyses conducted in this study are based on the assumption that the magnitude of satellite-derived total column is correct. A detailed analysis of the errors involved in deriving satellite-based tropospheric columns of BrO is beyond the scope of this paper. There are potentially large uncertainties associated with the derivation of both slant and vertical absolute column BrO related to the choice of spectral fitting windows and polynomial parameters used in retrievals as well as assumed BrO vertical profiles (G. Mount, private communication, 2011; DLR, 2009).

For all analyses conducted in this study, we use level 2 (time-ordered) satellite data. Level 2 data provide the best temporal match between satellite observations and aircraft in-situ and ground-based observations.

### 2.3 Stratospheric BrO column

A model simulation of stratospheric bromine species is used to estimate the spatial structure of the stratospheric BrO column. Our approach is similar to that described

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



by Salawitch et al. (2010), but differs in that they showed results for a fixed local solar time of 01:30 p.m. whereas here we calculate BrO along each OMI or GOME-2 orbit, which spans a range of local solar times near the pole. A photochemical steady state (PSS) model is constrained to profiles of temperature, O<sub>3</sub>, NO<sub>y</sub>, Cl<sub>y</sub>, H<sub>2</sub>O, CH<sub>4</sub>, sulfate aerosol surface area, etc. output from a run of Whole Atmosphere Community Climate Model (WACCM) (Garcia et al., 2007) conducted using meteorological fields for spring 2008. Salawitch et al. (2010) provide a description of this WACCM run, which was conducted to support the Stratosphere-Troposphere Analyses of Regional Transport 2008 campaign.

We specify the vertical distribution of Br<sub>y</sub> (the total bromine content of all inorganic bromine species) input to the PSS model, based on a relation between Br<sub>y</sub> and CFC-12. Profiles of CFC-12 are from the NASA Global Modeling and Assimilation Office (GMAO) Goddard Earth Observing System Data Assimilation System Version 5 (GEOS-5) (Rienecker et al., 2007) assimilation for spring 2008. Salawitch et al. (2010) evaluated the sensitivity of stratospheric partial column BrO to the amount of stratospheric bromine supplied by VSL bromocarbons, termed VSL Br<sub>y</sub>. The value of VSL Br<sub>y</sub> is quite uncertain: Tables 1–9 of (WMO, 2011) gives a range of 1 to 8 ppt for this quantity. We use VSL Br<sub>y</sub> = 7 ppt because this value gives best overall agreement between OMI total column BrO and the sum of tropospheric partial column BrO inferred from ARCTAS and ARCPAC aircraft profiles and the simulated stratospheric BrO burden (see Auxiliary Material of Salawitch et al. (2010) for a discussion of the plausibility of VSL Br<sub>y</sub> = 7 ppt).

The PSS model is used to find BrO, at the local solar time of each OMI or GOME-2 orbit, assuming production and loss of all species are in balance over a 24 h period of time for a stationary air parcel. Then, BrO abundance is integrated from the pressure of the tropopause (WMO definition of the thermal tropopause, based on GEOS-5 temperature) to 0.01 hPa, yielding stratospheric column BrO. The diurnal variation of the stratospheric column BrO must be considered in estimating tropospheric BrO column, especially for the high SZAs of the level 2 data examined below. Further details of

**Satellite-derived  
Arctic tropospheric  
BrO columns**

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the PSS model are provided in Sect. 6.3.2 of SPARC CCMVal (2010), and references therein.

As noted above, the  $Br_y$  versus CFC-12 relation used here consists of a baseline value of  $Br_y$  plus 7 ppt (representing stratospheric injection of bromine in the form of product gases of VSL bromocarbons). Baseline  $Br_y$  is set to zero at the tropopause and represents, above the tropopause, supply of bromine from the decomposition of methyl bromide ( $CH_3Br$ ), halons, as well as dibromethane ( $CH_2Br_2$ ). Other groups place  $CH_2Br_2$  into the definition of VSL  $Br_y$ . We consider  $CH_2Br_2$  to be part of baseline  $Br_y$ , however, because this compound is observed above the tropical tropopause (e.g. Wamsley et al., 1998). The total stratospheric  $Br_y$  burden of 26 ppt used here is at the upper end of the present range of uncertainty (e.g. WMO, 2011). This formulation (7 ppt for VSL  $Br_y$  plus baseline that includes  $CH_2Br_2$ ) was chosen because it results in best agreement between OMI total column BrO and the sum of stratospheric (modeled) and tropospheric (measured) partial column BrO. A wide range of values for VSL  $Br_y$  yields “reasonable agreement” between these two terms when all uncertainties are considered (Fig. 6, Auxiliary Material, Salawitch et al., 2010). Despite the considerable uncertainty in stratospheric  $Br_y$  and the impact of this parameter on tropospheric BrO column inferred from satellite observations of total column BrO, our approach of using a value for stratospheric  $Br_y$  near the upper end is supported by the generally close quantitative agreement between inferences of tropospheric BrO column from the satellites and the in-situ data, described below. As noted previously, the fields of stratospheric BrO used here are approximately 27 % larger than those reported by Theys et al. (2011) (with percentage differences that vary considerably with respect to location).

## 2.4 Cloud parameters

Optically thick clouds shield the underlying atmosphere from satellite sensors. The OMI rotational Raman (RR) cloud product (Vasilkov et al., 2008) is used to infer information about the shielding effects of clouds over snow and ice (Vasilkov et al., 2010). This product provides an estimate of the scene (combined cloud and surface) pressure

### Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



over snow and ice surfaces. The Near-real-time SSM/I EASE-grid daily global Ice and snow concentration and Snow Extent (NISE) data set (Nolin et al., 1998) is used to identify snow and ice-covered pixels. When the difference between scene and terrain pressure ( $\Delta P_c$ ) exceeds 250 hPa we infer that clouds have led to significant shielding of tropospheric BrO from the satellite sensor (Vasilkov et al., 2010), as discussed below.

We use level 3 (gridded) cloud optical thickness and cloud-top pressure retrievals from the MODerate-resolution Imaging Spectroradiometer (MODIS) (Platnick et al., 2003) on the NASA Aqua satellite as a secondary check on our detection of shielding clouds. Analysis of cloud effects on the satellite-derived tropospheric BrO columns is presented in Sect. 3.2.

## 2.5 Meteorological data sets

Tropopause heights are inferred along the flight tracks using tropopause pressure and geopotential height profiles from MERRA, a reanalysis based on GEOS-5 system (Rienecker et al., 2007) provided by NASA Global Modeling and Assimilation Office (GMAO). These tropopause heights are used to integrate the aircraft BrO profiles and are given in Tables 1, 2 and 2. MERRA also provides meteorological parameters including sea level pressure, wind speed, and planetary boundary layer height. These meteorological data are used to infer relationships between BrO enhancements and the meteorological conditions. MERRA parameters are provided at  $0.5^\circ$  latitude  $\times$   $0.667^\circ$  longitude resolution. All parameters except geopotential height profiles are provided hourly; geopotential height is given every 6 h.

## 3 Derivation of tropospheric BrO vertical column density

### 3.1 Residual method to obtain tropospheric BrO vertical column density

Tropospheric column BrO can be obtained from satellite total column BrO retrievals using the residual method (e.g. Theys et al., 2011, and references therein). Here, as in

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Theys et al. (2011), we calculate tropospheric BrO vertical column densities ( $VCD_{Trop}$ ) for each OMI or GOME-2 pixel using the derived BrO total slant column ( $SCD_{Total}$ ), an estimate of stratospheric BrO vertical column ( $VCD_{Strat}$ ), stratospheric air mass factors ( $AMF_{Strat}$ ), and an estimate of the tropospheric air mass factor ( $AMF_{Trop}$ ) as follows:

$$VCD_{Trop} = \frac{SCD_{Trop}}{AMF_{Trop}} = \frac{(SCD_{Total} - SCD_{Strat})}{AMF_{Trop}} = \frac{(SCD_{Total} - VCD_{Strat} \cdot AMF_{Strat})}{AMF_{Trop}}. \quad (6)$$

The SCD of a given absorber seen by a satellite sensor is defined as the amount of the absorber along an average light path taken by photons as they travel from the sun, through the atmosphere, and back to the sensor. The SCD is affected by scattering and absorption within the atmosphere as well as reflection off the surface and clouds.

For a given altitude range (denoted by a subscript  $z$ ), The air mass factor ( $AMF_z$ ) is used to convert  $SCD_z$  to  $VCD_z$  as follows:

$$AMF_z = SCD_z / VCD_z. \quad (7)$$

The sensitivity of UV radiance measurements to the BrO layer amounts varies with altitude. This variation depends on viewing geometry (mainly solar zenith angle, SZA, and view zenith angle, VZA), surface albedo, cloud effects, and the vertical BrO profile. We must account for this varying sensitivity in the AMF. Using the optically thin absorber assumption, the  $AMF_z$  can be formulated as

$$AMF_z = SCD_z / VCD_z = \frac{\int W(z)N(z)dz}{\int N(z)dz}, \quad (8)$$

(Palmer et al., 2001; Theys et al., 2011), where  $z$  is altitude,  $N(z)$  is the number density profile of the absorber, and  $W(z)$  is the weighting function profile that represents all the parameters influencing the AMF except the vertical profile of the absorber.

Here, we use estimates of  $AMF_{Strat}$  provided in the OMI and GOME-2 total BrO column products.  $AMF_{Strat}$  for GOME-2 is computed using the SCIATRAN radiative transfer model (Rozanov et al., 2005). The OMI algorithm uses a different approach

**Satellite-derived Arctic tropospheric BrO columns**

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to calculate  $AMF_{Strat}$ ; BrO VCDs are obtained by applying wavelength- and albedo-dependent AMFs to the BrO absorption cross sections prior to fitting. It is assumed that all BrO resides in the stratosphere in this procedure. An effective AMF is then defined as the ratio of the SCD (derived independently as described in Sect. 2.2) to this VCD. We assume that  $AMF_{Strat}$  is represented by this effective air mass factor.

We compute the tropospheric  $AMF_{Trop}$  at 344.6 nm using the Linearized Discrete Ordinate Radiative Transfer (LIDORT) model (Spurr et al., 2001). We use a clear scene assumption in the AMF calculation. The DC-8 composite profile shown in Fig. 2a is taken as the default tropospheric BrO profile in the  $AMF_{Trop}$  calculation. However, we use individual aircraft profiles of BrO, from both the DC-8 and WP-3D, to calculate  $AMF_{Trop}$  for point-to-point comparisons between aircraft in-situ and satellite-derived tropospheric columns.

We have generated a look-up table of  $AMF_{Trop}$  for various representative surface albedos and viewing geometries. A  $5^\circ$  interval for SZA and a  $2.5^\circ$  interval for VZA are used. Because the weighting functions are highly dependent on surface albedo, we use step-widths of 0.1 in the range [0,1] for surface albedo.

The calculation of  $AMF_{Trop}$  uses the derived OMI reflectivity at 331 nm for each pixel, from the OMI Total Ozone Mapping Spectrometer (TOMS) total  $O_3$  product, as a proxy for surface albedo. We thus avoid using a surface albedo climatology, which can lead to AMF errors when the climatology differs from the actual surface albedo. Such errors can occur when the snow or sea ice distribution differs from the climatological mean, which is of particular concern at high latitudes during spring.

### 3.2 Sensitivity of the derived tropospheric column to SZA, surface albedo, and clouds

Here, we investigate the sensitivity of the satellite-derived tropospheric BrO retrieval to SZA, surface albedo, and clouds. They et al. (2011) analyzed the dependence of the weighting function on surface albedo and clouds for  $SZA = 45^\circ$ . Here, we extend this analysis to a wider SZA range ( $40^\circ$ – $90^\circ$ ) relevant to polar observing conditions;



at high latitudes ( $>60^\circ\text{N}$ ) during the day in the early spring SZA is  $>50^\circ$ . Figure 3a shows that when the sun is relatively high in the sky ( $\text{SZA} \leq 60^\circ$ ), OMI and GOME-2 should have good sensitivity to tropospheric BrO for surface albedos  $>0.5$ . Sensitivity is significantly reduced for darker surfaces (e.g. albedo = 0.1). Sensitivity to tropospheric BrO decreases with increasing SZA; there is significantly lower sensitivity at  $80^\circ$  even for a surface albedo of 0.5. Figure 3b shows that at high surface albedo (0.9), there is increased sensitivity to BrO near the surface in addition to good overall tropospheric sensitivity for SZA up to  $80^\circ$ .

To evaluate the effect of optically thick clouds on the sensitivity to tropospheric BrO, we show a weighting function profile for a cloud with an extinction coefficient of 300 at 2.5 km in Fig. 3b. Similar to the results of Theys et al. (2011), this weighting function is the same as the clear-sky case above the cloud, and drops to zero below the cloud. Hence, OMI and GOME-2 are not sensitive to BrO below optically thick clouds.

The effect of optically thin clouds and aerosols on the sensitivity of OMI and GOME-2 to tropospheric BrO is dependent on surface reflectivity and viewing geometry. According to Vasilkov et al. (2010), the UV cloud shielding effect is generally reduced over bright surfaces as compared to dark surfaces for nadir-viewing satellite sensors. For example, there is good sensitivity to tropospheric absorbers below thin clouds over bright surfaces near the nadir swath positions for SZAs  $\leq 67^\circ$ . On the other hand, their simulation shows that there can be UV cloud shielding at the swath edge at high solar zenith angles, even for a thin cloud ( $\tau \sim 0.5$ ). The clear scene assumption in the AMF calculation is appropriate over bright surfaces (i.e. snow/ice) when the derived scene pressure is close to the terrain pressure, which indicates that any particles along the line of sight must either be a thin, non-shielding cloud or an aerosol layer (the so-called Arctic haze) common in polar regions. The optical thickness of Arctic haze during ARCTAS was reported to be up to  $\sim 0.2$  at 354 nm (Shinozuka et al., 2011). Based on the calculations of Vasilkov et al. (2010), OMI and GOME-2 should have good sensitivity to surface BrO in Arctic haze conditions at the near-nadir swath positions, but may have reduced sensitivity at the swath edge.

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



Figure 4a shows computed tropospheric air mass factors. As shown for the weighting functions in Fig. 3, tropospheric air mass factors are higher over brighter surfaces owing to increased near-surface sensitivity. However, tropospheric sensitivity begins to be lost as SZA increases; tropospheric sensitivity almost disappears when  $SZA > 85^\circ$  even for relatively bright surfaces. In this study, we present satellite-derived BrO tropospheric columns only for  $SZA < 85^\circ$ .

The sensitivity of  $AMF_{Trop}$  to the tropospheric BrO profile is assessed in Fig. 4b. For a bright surface (albedo = 0.9), no significant difference exists between tropospheric AMFs computed using the DC-8 composite profile and a profile where all tropospheric BrO is contained below 500 m for  $SZA \leq \sim 60^\circ$ . This is a consequence of good sensitivity at all tropospheric altitudes for bright surfaces and low SZA as shown in Fig. 3. For  $60^\circ < SZA < 75^\circ$ , the sensitivity to middle- and upper-tropospheric BrO is maintained while the near-surface sensitivity drops; this leads to a small sensitivity to the profile shape under these conditions. Profile sensitivity decreases for higher SZAs as the sensitivity to the entire troposphere drops. For a darker surface (albedo = 0.4), we have lower sensitivity to BrO near the surface even when the sun is relatively high ( $SZA \sim 40^\circ$ ), leading to greater profile sensitivity at all SZAs.

We next examine tropospheric column BrO estimated from OMI in the context of the different sensitivities discussed above. Figure 5 shows OMI total, the model stratospheric, and the derived tropospheric BrO columns for OMI orbit 20050 on 22 April 2008 along with several parameters that impact the sensitivity of the radiance observation to the tropospheric BrO column. These parameters include 331 nm reflectivity (a proxy for surface albedo),  $\Delta P_c$ , and SZA.

Nearly zero tropospheric BrO column amounts are obtained over low surface albedo areas (OMI reflectivity < 0.5) such as the Barents Sea (marked with black squares in Fig. 5). Here the retrieved BrO total columns are generally less than the modeled stratospheric column, while the adjacent areas with higher reflectivity have higher total column BrO. The low tropospheric AMFs for these conditions amplify the slant columns in the calculation of the tropospheric column (Eq. 6). Even so, the computed

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tropospheric vertical columns are near zero. This indicates that satellite-derived tropospheric BrO may not be reliable when the surface albedo is low.

In Fig. 5, the cloud shielding effect is shown over Siberia (marked with orange rectangles), where the OMI rotational Raman cloud product reports  $\Delta P_c > 250$  hPa. Aqua MODIS also indicates cloud cover, with cloud top pressures between 450 and 500 hPa (not shown). Our derived tropospheric columns over this region are significantly less than those from adjacent areas without cloud cover. This supports the presence of tropospheric BrO mainly below 450–500 hPa and the ability of OMI and other nadir viewing sensors to capture information about tropospheric BrO over bright polar regions, with SZA up to  $\sim 85^\circ$  when the sky is clear. Theys et al. (2011) also used the cloud shielding effect to infer an amount of BrO in the troposphere, but applied this approach only in the tropics.

For completeness, we include all data in our analysis of aircraft measurements. However, we note aircraft soundings for which observing conditions likely produce large errors in the satellite-derived tropospheric BrO columns. In mapping surface BrO enhancement events with satellite retrievals, we use only data that has good sensitivity to the troposphere:  $SZA < 85^\circ$ ,  $\Delta P_c < 250$  hPa, and surface albedo  $> 0.7$ .

### 3.3 Uncertainty analysis

For a single pixel, we express the uncertainty of the tropospheric VCD,  $\sigma_{VCD_{Trop}}$ , in a simplified form:

$$\sigma_{VCD_{Trop}}^2 = \left( \frac{\sigma_{SCD_{Total}}}{AMF_{Trop}} \right)^2 + \left( \frac{\sigma_{VCD_{Strat}} AMF_{Strat}}{AMF_{Trop}} \right)^2 \quad (9)$$

(Boersma et al., 2004; De Smedt et al., 2008; Theys et al., 2011), where  $\sigma_{SCD_{Total}}$  is the uncertainty of total slant column,  $\sigma_{SCD_{Strat}}$  is the uncertainty of stratospheric slant column, and  $\sigma_{AMF_{Trop}}$  is the uncertainty of tropospheric air mass factor. This representation assumes that the different types of uncertainties are uncorrelated with each other.

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



We take the fitting uncertainty derived from observed minus fitted radiances for each pixel as  $\sigma_{\text{SCD}_{\text{Total}}}$ . This assumes that the  $\text{SCD}_{\text{Total}}$  error has a zero mean. The average fitting uncertainty at latitudes greater than  $60^\circ \text{N}$  is about 18 %.

We estimate the uncertainty of the stratospheric slant column ( $\sigma_{\text{SCD}_{\text{Strat}}}$ ) by multiplying the uncertainty of the stratospheric column BrO by the stratospheric AMF. The uncertainty of the stratospheric column BrO results from a root-sum-squares combination of three terms: the uncertainty in chemical kinetics that govern the BrO to Br<sub>y</sub> ratio, the uncertainty in the dynamics that govern CFC-12 (and hence Br<sub>y</sub> due to Source Gas Injection), and the uncertainty in VSL Br<sub>y</sub>. The uncertainty in chemical kinetics is evaluated by varying the rate constant of individual chemical reactions (including *J* values) by the 1-sigma estimate of uncertainty given by Sander et al. (2006). The most important chemical term is BrO + NO<sub>2</sub> forming BrNO<sub>3</sub>; this rate constant is uncertain by about a factor of 2 at 220 K. The uncertainty due to dynamics is found by repeating calculations for plus and minus 4 % variations in the abundance of CFC-12, because comparison to aircraft observations showed CFC-12 from GEOS-5 was accurate to within  $\pm 4\%$  in the lower stratosphere (Salawitch et al., 2010). Finally, the uncertainty due to VSL Br<sub>y</sub> is set at  $\pm 27\%$ , which represents the mean difference in stratospheric column BrO resulting from our approach to handling this term compared to the approach of Theys et al. (2011).

Here, we neglect the uncertainty in the tropospheric air mass factor as it is relatively small compared with the other error sources. For example, Theys et al. (2011) describe sensitivity tests that show use of a single wavelength for the weighting function profiles leads to an error of less than 5 %. We find that the profile dependence produces error between 7 and 13 % for bright surfaces.

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 4 Results and discussions

### 4.1 Comparisons of aircraft in-situ measurements with satellite retrievals

In this section, comparisons between aircraft and satellite inferences of tropospheric column BrO are shown for cases where agreement is good, agreement is poor but the cause for disagreement is understood, and agreement is poor and not well understood. These comparisons are performed to evaluate the veracity of the satellite-derived tropospheric column BrO product. We present a series of figures for various ARCTAS and ARCPAC flights. For each, we show maps of the satellite-derived total and tropospheric columns and time series of various aircraft measurements, collocated satellite-derived BrO data, and estimated stratospheric BrO amounts with vertical error bars derived using the formulation given in Sect. 3.3. Satellite-derived data are shown for all conditions (e.g. even in the presence of thick clouds that likely shield the surface). We also show plots of various parameters that affect the sensitivity of the satellite measurements to tropospheric BrO and note times and places where satellite observing conditions are not ideal for inferring reliable tropospheric BrO amounts.

#### 4.1.1 Cases of good agreement between satellite and in-situ data

Figure 6a shows maps of the OMI and GOME-2 measurements of total column BrO as well as maps of tropospheric column BrO inferred from each satellite sensor on 17 April 2008. The track of the DC-8 and locations of BrO profiles, for the ARCTAS flight on this date, are marked. The bottom panel shows geographic regions of interest. OMI and GOME-2 orbits closest in time to the ARCTAS flight are used.

The tropospheric column BrO maps have different spatial structure than maps of total column BrO. For example, total column BrO is highest over the north coast of Canada. These local maxima are significantly reduced in the tropospheric columns. A long tail of enhanced values near the North Pole is pronounced in the maps of tropospheric column BrO and is not as prominent in total column BrO. As discussed by Salawitch et

### Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



al. (2010) and Theys et al. (2011), inferring tropospheric BrO from the total BrO column requires an accurate estimate of the stratospheric contribution to the total column, because variability induced by stratospheric dynamics is comparable in magnitude to variability induced by the surface release of bromine. Figure 6a is a perfect illustration of this point, which as noted above has been overlooked in many prior studies.

Figure 6b shows time series plots of DC-8 flight data and collocated satellite measurements on 17 April 2008. DC-8 data include aircraft altitude, in-situ O<sub>3</sub>, CO, BrO, active bromine and soluble bromide. Satellite data include OMI-derived BrO columns (total and tropospheric),  $\Delta P_c$ , OMI 331 nm reflectivity, and Aqua MODIS cloud optical thickness. Estimated errors of satellite-derived BrO columns are presented as vertical error bars. The tropopause height, as given in the MERRA data set, is at approximately ~7 km during the flight (not shown). CO and O<sub>3</sub> data suggest that there were no stratospheric intrusions for any of the collected tropospheric BrO profiles. Obvious signatures of stratospheric air (low CO and high O<sub>3</sub>) were occasionally seen, but only when the aircraft was flying near or above the tropopause.

Satellite-derived BrO tropospheric columns (orange line) are ~zero near Fairbanks (see top panel of Fig. 6b). As discussed in Sects. 3.1 and 3.2, tropospheric BrO information is not reliably retrieved when surface reflectivities are less than ~0.4, which is the case near Fairbanks (see the fourth panel of Fig. 6b).

Both total and tropospheric column BrO show a slight dip near the pole, where the OMI and MODIS cloud products indicate shielding clouds (middle of Fig. 6b). Here, the low values of derived tropospheric BrO over optically thick clouds likely result from the cloud shielding effect (Theys et al., 2011). Such clouds may also reduce tropospheric BrO columns by slowing photochemical production of active bromine.

In-situ tropospheric column BrO, marked as black dots in the time series plot (the first panel of Fig. 6b), are obtained by integrating in-situ BrO mixing ratio profiles as described in Sect. 2.1. Uncertainties in the in-situ columns are shown as black vertical error bars. The horizontal bars indicate the distance covered by corresponding flight ascents/descents. OMI tropospheric BrO columns for point-to-point comparisons

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



with aircraft data, shown as red squares, are obtained by averaging pixels over the corresponding flight segment.

The OMI tropospheric BrO columns exhibit a magnitude and variability similar to that of the DC-8 in-situ columns. For example, profile #12 reports high BrO mixing ratios from the surface to ~4 km; the OMI tropospheric BrO column (as well as GOME-2, shown in Fig. 6a but not in Fig. 6b) is also relatively high at the same location.

However, some discrepancies appear, perhaps because the aircraft captures small scale features that the satellite observations cannot resolve. For example, in-situ profile #13 shows a relatively large value for troposphere column BrO, caused by enhanced BrO mixing ratios in a layer near 2 km. This could be a local event as the nearby aircraft column #14 does not show such an enhancement. The satellite retrieval agrees well with profile #14 but underestimates the abundance of BrO measured in profile #13. Overall, OMI tropospheric column BrO quantitatively agrees with the DC-8 in-situ column BrO to within the respective estimated uncertainties. The comparison between satellite and in-situ tropospheric column BrO for this flight is further quantified by Liao et al. (2011b). These comparisons demonstrate the ability of OMI and GOME-2 to capture the magnitude and spatial distribution of tropospheric column BrO over bright surfaces for clear conditions.

#### 4.1.2 Examination of differences between satellite-derived and in-situ tropospheric columns of BrO

We next describe two cases of apparent inconsistency between satellite and in-situ observations. Figures 7a and 7b show data corresponding to the DC-8 flight on 5 April 2008. In-situ measurements did not detect ozone depletion or high bromine over Hudson Bay, whereas OMI and GOME-2 both report high total column BrO (Salawitch et al., 2010). BrO measurements were not available to produce profiles for this flight, but active bromine and soluble bromide were reported.

Figure 7b shows that the derived tropospheric BrO column from OMI is low over Hudson Bay, where the total column is high. Similar results are found for GOME-2 (Fig. 7a). The OMI measurement of total column BrO shows very similar magnitude

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and structure as the calculated stratospheric column BrO. This suggests that the elevated total column BrO over Hudson Bay could be a consequence of high stratospheric columns. Furthermore, the in-situ CO and O<sub>3</sub> measurements indicate no evidence of stratospheric air below 6 km altitude in ascending or descending flight tracks near Hudson Bay.

Figures 8a and 8b show similar maps and time series, respectively, for the DC-8 flight on 8 April 2008. GOME-2 observations during this flight are not available. Aircraft in-situ BrO measurements are also not available for this flight. This makes it difficult to evaluate the validity of the OMI-derived BrO tropospheric columns. The aircraft observed a severe ozone depletion event ( $[O_3] < 1$  ppbv) and high levels of active bromine and soluble bromide near Alert (marked as black square), whereas the OMI total column BrO is relatively low and the inferred OMI tropospheric column BrO is at or near background levels. The ODE observed by the DC-8 on 8 April 2008 is thus not apparent in the satellite measurement of BrO on this date.

As described in Eqs. (3), (4), and (5), BrO catalyzes the loss of ozone and, at the same time, BrO is produced by reaction of Br with O<sub>3</sub>. This suggests that high BrO concentrations cannot be maintained if O<sub>3</sub> is completely depleted, even though plenty of active bromine may be available. This explanation is consistent with other observations (e.g. Hausmann and Platt, 1994; Tuckermann et al., 1997; Neuman et al., 2010) that show high BrO concentrations only when ozone is partially depleted. The presence of a severe ozone depletion event is therefore indicative of an expectation of little or no enhancement of tropospheric BrO, despite the presence of large amounts of active bromine and soluble bromide. Liao et al. (2011a) also discuss the relationship between O<sub>3</sub> and BrO; little BrO exists when O<sub>3</sub> is low due to the partitioning between Br and BrO.

OMI tropospheric column BrO shows enhancements in the vicinity of the 8 April 2008 ozone depletion event, during the 36 h time period prior to this flight (not shown). This enhancement is likely connected with the nearly complete removal of ozone observed by the DC-8 aircraft.

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



### 4.1.3 Cases of disagreement between satellite and in-situ data

Figure 9a shows maps of satellite-derived BrO columns and the flight track of the NOAA ARCPAC WP-3D on 19 April 2008. The OMI (shown) and GOME-2 (not shown) data both indicate enhanced total and tropospheric columns of BrO over the Chukchi Sea.

5 The aircraft flew into the area of elevated satellite-derived tropospheric column BrO at the northernmost extent of the track.

Figure 9b shows time series plots for the flight of 19 April 2008. Two types of disagreement between in-situ and satellite tropospheric column BrO are observed. First, satellite-derived tropospheric column BrO is considerably less than the in-situ column  
10 over surfaces with moderate reflectivity in the presence of clouds. For example, the OMI estimate is less than the WP-3D columns (#19 and #23) near Fairbanks. These observations were obtained in a region where the surface albedo is between 0.4 and 0.6, in the presence of low level clouds. This type of disagreement may be expected from the sensitivity studies described in Sect. 3.2. It is a challenge for satellite sensors to accurately retrieve tropospheric BrO over a moderately reflective surface in the  
15 presence of clouds.

The second type of disagreement during this flight is illustrated by profile #21, where both OMI and GOME-2 tropospheric column BrO are significantly enhanced (only OMI is shown in the first panel of Fig. 9b), whereas the in-situ BrO column does not show  
20 evidence of enhancement. Artifacts from clouds or surface albedo are not the likely explanation for this disagreement, as the surface was bright and no clouds are evident in  $\Delta P_c$  (fourth panel, Fig. 9b). The highest level of active bromine is observed from the aircraft sensor at the same location where tropospheric column BrO from both satellite sensors show enhancements.  $O_3$  mixing ratios of  $\sim 10$  ppbv suggest the aircraft sampled a partial ozone depletion event (e.g. Ridley et al., 2003). Prior DOAS observations in the Arctic suggest that BrO should still be present for  $O_3$  levels of  $\sim 10$  ppbv (Hausmann and Platt, 1994; Tuckermann et al., 1997). The disagreement between satellite  
25 and aircraft BrO could be related to the different spatial scales, and/or vertical coverage

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





spanned by the respective instruments: the aircraft does not observe the complete profile and the field of view encompassed by the satellite covers a larger area than that sampled by the airplane. Timing may also be a possible explanation for the disagreement, as the aircraft flew near the location of the enhancement about five hours after the OMI overpass and about three hours after that of GOME-2.

Our analysis indicates that the large enhancement in total column BrO seen by OMI and GOME-2 over the Chukchi Sea on 19 April 2008 was due to a tropospheric perturbation. The WP-3D aircraft recorded highly elevated active bromine and partially depleted  $O_3$  near the surface at this precise location, indicating recent association with elevated BrO. It is possible the satellite perturbation was caused by the presence of BrO at higher altitudes than those sampled by the aircraft at this location. By the time the aircraft reached 3 km altitude, it had left the region of highly elevated satellite BrO (terminal point of profile #23). Salawitch et al. (2010) presented an analysis of ground-based Max DOAS observations of BrO for April 2008 that, together with ARCTAS and ARCPAC BrO profiles, showed important contributions to column BrO often originate within the troposphere from altitudes above the top of the planetary boundary layer.

Figure 10a and 10b shows maps of the BrO columns and time series plots for the NOAA WP-3D flight of 21 April 2008, respectively. The maps show enhanced OMI total column BrO over the north shore of Canada and Canadian Archipelago. The stratospheric model shows only a slight enhancement for these regions; the subtraction of the stratospheric model from the OMI column shows significant levels of tropospheric column BrO for much of the flight segment. Time series plots indicate that WP-3D in-situ columns #25, #26, and #27 are lower than the OMI tropospheric column, but the differences are not significant given the uncertainties in OMI and in-situ tropospheric columns. There is, however, good agreement between in-situ column #28 and the satellite-based estimate.

WP-3D profiles #26, #27, and #28 were collected in close proximity (see Fig. 1), but show a large variation in the column amounts. As noted above, the profile #28 column agrees well with the satellite-based column whereas the other two in-situ based BrO

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



columns are lower than the satellite-based estimates. As discussed above for other flights, this result may be explained by the fact that the aircraft captures small-scale spatial features while the satellite observes larger scales.

The overestimation of the tropospheric column BrO by OMI, compared to in-situ columns for profile #25, #26, and #27, could potentially be explained by a stratospheric column that is not fully removed from the total column. One factor not considered in our analysis is irreversible, cross-tropopause exchange of BrO from the stratosphere to the troposphere (STE). Such transport events occur on the western flank of Arctic low pressure systems (Salawitch et al., 2010, and references therein). If the satellite signal were to originate from STE of BrO, the associated BrO may not have been sampled by the WP-3D, since the maximum altitudes of the WP-3D profiles #25, #26 and #27 (6 km) is well below the height of the tropopause (10.5 km). Future measurement of profiles of BrO from near the surface to the lowermost stratosphere, in the footprint of a satellite sensor after passage of an airmass with a low altitude tropopause, are needed to assess the importance of STE of bromine on the interpretation of the satellite record.

## 4.2 BrO explosions observed with satellites

Events of rapid enhancement of tropospheric column BrO are apparent from OMI and GOME-2 observations after adjustment for the stratospheric burden of BrO, for the time period mid-March to late April 2008. The NASA DC-8 aircraft flew into an area of enhanced tropospheric column BrO near the North Pole on 17 April 2008 (Fig. 6a) and the NOAA WP-3D aircraft flew near another tropospheric BrO enhancement on 19 April 2008 (Fig. 9a). Here we examine the event near the North Pole on 17 April 2008 and similar events in more detail using only OMI retrievals. The wide orbital swath and high spatial resolution of OMI, in addition to its frequent observations at high latitudes, provide a unique view of the temporal evolution of these events.

Figure 11 shows the evolution of a tropospheric BrO enhancement event (“BrO explosion”) observed from 16 to 18 April 2008. Here, we only show observations when the following conditions are met, to provide reliable tropospheric BrO information as

### Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



discussed in Sect. 3.2:  $SZA < 85^\circ$ , reflectivity  $> 0.7$ , and  $\Delta P_c < 250$  hPa. The stratospheric column has been removed, as discussed in Sect. 3.1, using photochemical model output for the VSL  $Br_y = 7$  ppt simulation, for the local solar time of each OMI pixel. The major activation of BrO starts at  $\sim 22:00$  UTC on 16 April and lasts for  $\sim 30$  h.

5 We see activations near the North Pole and Canadian Archipelago. The spatial features of the elevated tropospheric column BrO change rapidly, with significant variations over the course of a day.

We have ruled out a stratospheric origin for the enhanced total column BrO observations near the North Pole and Canadian Archipelago on these days. Since stratospheric BrO and  $O_3$  columns exhibit a significant correlation (Theys et al., 2009; Salawitch et al., 2010), the OMI measurement of total  $O_3$  column can be used as a proxy for the spatial pattern of stratospheric column BrO. The comparison of the OMI tropospheric BrO column with the OMI total  $O_3$  column shown in Fig. 12 indicates lack of correlation. Thus, the elevated region of total column BrO is not of stratospheric origin.

15 Figure 12 also shows few shielding clouds near the areas with high BrO columns.

Figure 13 shows three tropospheric BrO enhancement events during April 2008 along with maps of sea level pressure, wind speed at 2 m altitude, and planetary boundary layer height from MERRA for the closest synoptic hour. For these events, the locations of enhanced BrO columns are coincident with high near-surface wind speeds (3rd column) that are geostrophically consistent with localized low pressure systems. Neuman et al. (2010) also note high wind speed along the flight paths in conjunction with high concentrations of active bromine. Our analysis is consistent with the suggestion that strong surface winds associated with low pressure systems can trigger bromine activation via blowing snow (Yang et al., 2008, 2010; Jones et al., 2009; A. E. Jones et al., 2010).

25 Figure 13 also shows that the spatial structure of high tropospheric column BrO is similar to that of the planetary boundary layer (PBL) height, although there is not always a precise alignment of these features. At high latitudes where the meteorological analysis is driven primarily by satellite data, the MERRA fields may contain displacement or

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



other errors, particularly in near-surface fields. The following discussion should therefore be considered somewhat speculative in light of these uncertainties. We provide an animated visualization as a Supplement to depict evolution of enhanced tropospheric column BrO, sea level pressure, wind speed at 2 m and planetary boundary layer height from 16 to 18 April 2008.

Neuman et al. (2010) reported a temperature inversion at 500 m during the BrO activation event of 19 April 2008, more or less consistent with the MERRA PBL height. Bromine chemistry may be related to PBL height as follows: BrO concentrations are controlled in large part by  $O_3$ ; a sufficient amount of  $O_3$  is necessary to maintain high BrO concentrations as explained in Sec. 4.1.2. Ozone is quickly consumed in a shallow boundary layer during bromine activation (Lehrer et al., 2004; Anderson and Neff, 2008). When  $O_3$  is substantially depleted, production of BrO will cease and BrO will be destroyed by various reactions, including those in Eq. (4). On the other hand,  $O_3$  may not become completely depleted in a deeper boundary layer, allowing concentrations of BrO to remain high.

The second row of Fig. 13 illustrates an event at 00:30 UTC on 18 April 2008 and the third row is for an event at 22:40 UTC on 19 April 2008. The two OMI orbits presented are closest in time to airplane flights into the elevated BrO and active bromine layers (profile #12 from DC-8 and #21 from WP-3D, respectively). The locations of the profiles are shown as black diamonds on the maps of tropospheric BrO. The aircraft profiles of temperature, BrO mixing ratio, and active bromine mixing ratio are presented in the last column. Aircraft measurements show enhanced active bromine in the near-surface layer for both flights and enhanced amounts of BrO for the 18 April flight. Satellite-derived tropospheric BrO enhancements are closely related to near-surface parameters including sea level pressure, wind speed at 2 m, and PBL height. This observation suggests that the BrO activations originate at the surface, which of course is consistent with prior expectation. For the event on 18 April, a tail of enhanced BrO column exists parallel to the wind direction inferred from sea level pressure. Bergoin et al. (2010) suggested long range transport of tropospheric BrO for plumes in

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

which recycling of bromine from condensed to gas phase sustains elevated BrO. The profile for BrO measured on 18 April by the DC-8 instrument (Fig. 13) also shows enhancements in the free troposphere, above the top of the PBL height (~500 m). Simultaneous profiles of CO and O<sub>3</sub> (not shown here, see Fig. 6b) imply no stratospheric influence. The existence of elevated BrO above the top of the PBL could be due to vigorous convection over ice leads driven by warm exposed water, with BrO then dispersed horizontally by prevailing winds (Simpson et al., 2007b; Salawitch et al., 2010). Our results are consistent with a surface origin of elevated BrO as well as transport of BrO enriched air parcels away from the source of origin. Indeed, an animation of the panels in Fig. 11 (see Supplement) provides compelling evidence for this behavior.

## 5 Conclusions

We have estimated tropospheric column BrO for April 2008 from OMI and GOME-2 retrieved total column BrO retrievals and a model simulation of the stratospheric column BrO. The sensitivity of satellite radiances to tropospheric BrO depends on various parameters, such as surface albedo, solar zenith angle (SZA), viewing zenith angle (VZA), the shape of the BrO profile, and the presence of clouds. We have quantified these dependences with a radiative transfer model. A detailed comparison is provided between tropospheric column BrO found from OMI and GOME-2 measurements of total column BrO and tropospheric column BrO computed from in-situ observations of BrO and related species acquired by instruments aboard the DC-8 and WP-3D aircraft during the ARCTAS and ARCPAC field campaigns.

Our analysis shows that retrievals of total column BrO from OMI and GOME-2, combined with a model estimate of stratospheric column BrO, can be used to retrieve realistic estimates of the magnitude and spatial variations in tropospheric column BrO provided that proper observing conditions are met. These conditions include bright surfaces (albedos  $\geq \sim 0.8$ ), low SZAs ( $\leq \sim 80^\circ$ ), and the absence of optically thick clouds. For the flight of 17 April 2008, satellite-derived tropospheric column BrO shows

### Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



enhancements where the DC-8 reports intense bromine activation and moderate ozone depletion. The satellite and aircraft based measurements of tropospheric column BrO are in close quantitative agreement on 17 April 2008, particularly for DC-8 flight segments over bright surfaces in clear conditions. In contrast to prior expectation based on DOAS measurements, the tropospheric burden of BrO based on the aircraft profiles was distributed over a broad range of altitudes and was not restricted to the planetary boundary layer. For the DC-8 flight of 5 April 2008, regions of enhanced total column BrO observed by OMI and GOME-2 near Hudson Bay are attributed to compression of stratospheric air associated with a low altitude tropopause, as neither the aircraft nor the satellite based measurements of tropospheric column BrO show enhancements. The regions of enhanced tropospheric column BrO inferred from OMI and GOME-2 on 5 and 17 April 2008 are not readily apparent in maps of total column BrO because stratospheric variability imparts variations in column BrO comparable in magnitude to the tropospheric signal. Use of a zonally symmetric representation of stratospheric column BrO, common in many prior studies, prevents the proper identification of the tropospheric BrO burden (e.g. Theys et al., 2009; Salawitch et al., 2010; Theys et al., 2011).

Satellite and aircraft measurements of tropospheric column BrO do not always exhibit good agreement, at times for reasons that seem well understood and at other times for reasons that are unclear but may be related to differences in the timing or spatial coverage of the respective observations. For the severe ozone depletion event observed by the DC-8 near Alert on 8 April 2008, neither OMI total column BrO column nor OMI tropospheric column BrO were elevated. Atmospheric conditions were favorable for the remote sensing from space of tropospheric BrO near Alert on this date (e.g. clear skies, high surface reflectivity). We believe the lack of a signal for OMI tropospheric column BrO near Alert on 8 April is consistent with our understanding of bromine chemistry: the production of BrO diminishes when ozone is severely depleted (e.g. Hausmann and Platt, 1994; Tuckermann et al., 1997). Hence, the association of elevated tropospheric column BrO and depleted O<sub>3</sub> is expected to be much

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

stronger for a partial ozone depletion event (ODE) than a major ODE. Tropospheric column BrO from OMI does show an enhancement, near Alert, 36 h prior to the major ODE. The aircraft may have been capturing very small scale variability (several nearby profiles showed large differences) compared to space-based observations. Finally, aircraft observations generally did not sample the altitude region where stratosphere to troposphere transport of BrO could potentially be affecting the satellite measurement of column BrO.

We examined several events of rapid enhancement of tropospheric column BrO observed by OMI. Observations at high latitudes from this polar-orbiting, sun-synchronous satellite provide a unique illustration of the rapid temporal evolution of BrO activation because orbital periods are about 100 min apart. Our analysis indicates that BrO activation events occurring on 16, 17, and 18 April 2008 are related to near-surface parameters such as low pressure systems, strong surface winds, and/or high planetary boundary layer. The satellite measurements reveal horizontal transport of activated BrO away from the source of origin, and the aircraft measurements show disbursement of BrO within the free troposphere. The strong quantitative agreement between OMI and aircraft tropospheric column BrO on 17 April 2008 supports the validity of the rapid time evolution, on synoptic scales, revealed by the OMI tropospheric BrO product. The events on 17, 18, and 19 April suggest bromine activation via high winds over snow (Yang et al., 2008; Jones et al., 2009; Yang et al., 2010; A. E. Jones et al., 2010) as well as long range transport of BrO by surface winds (Begoin et al., 2010).

**Supplementary material related to this article is available online at:**  
**[http://www.atmos-chem-phys-discuss.net/11/26173/2011/  
acpd-11-26173-2011-supplement.zip](http://www.atmos-chem-phys-discuss.net/11/26173/2011/acpd-11-26173-2011-supplement.zip)**

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





*Acknowledgements.* This work was supported by the NASA IPY, ARCTAS, and Aura science team programs, NOAA ARCPAC program, GEST GSSP program, and Smithsonian Institution. The authors are grateful to the NOAA NCEP and NASA GMAO reanalysis processing teams as well as the GOME-2, OMI, and MODIS data processing teams for providing data sets. The authors also thank Nicolas Theys for sharing his calculations of stratospheric column BrO and for many helpful email exchanges.

## References

- Anderson, P. S. and Neff, W. D.: Boundary layer physics over snow and ice, *Atmos. Chem. Phys.*, 8, 3563–3582, doi:10.5194/acp-8-3563-2008, 2008. 26204
- Ariya, P., Dastoor, A. P., Amyot, M., Schroeder, W. H., Barrie, L., Anlauf, K., Raofie, F., Ryzhkov, A., Davignon, D., Lalonde, J., and Steffen, A.: The Arctic: a sink for mercury, *Tellus*, 56B, 397–403, 2004. 26176
- Barrie, L. A., Bottenheim, J. W., Schnell, R. C., Crutzen, P. J., and Rasmussen, R. A.: Ozone destruction and photochemical reactions at polar sunrise in the lower Arctic atmosphere, *Nature*, 334, 138–141, 1988. 26176
- Barrie, L. A., Bottenheim, J. W., and Hart, W. R.: Polar Sunrise Experiment 1992 (PSE 1992): Preface, *J. Geophys. Res.*, 99, 25313–25314, 1994. 26176
- Begoin, M., Richter, A., Weber, M., Kaleschke, L., Tian-Kunze, X., Stohl, A., Theys, N., and Burrows, J. P.: Satellite observations of long range transport of a large BrO plume in the Arctic, *Atmos. Chem. Phys.*, 10, 6515–6526, doi:10.5194/acp-10-6515-2010, 2010. 26176, 26180, 26186, 26204, 26207
- Boersma, K. F., Eskes, H. J., and Brinksma, E. J.: Error analysis for tropospheric NO<sub>2</sub> retrieval from space, *J. Geophys. Res.*, 109, D04311, doi:10.1029/2003JD003962, 2004. 26194
- Bottenheim, W., Barrie, L. A., Atlas, E., Heidt, L. E., Niki, H., Rasmussen, R. A., and Shepson, P. B.: Depletion of lower tropospheric ozone during Arctic spring: The polar sunrise experiment 1988, *J. Geophys. Res.*, 95, 18555–18568, 1990. 26176
- Bovensmann, H., Burrows, J. P., Buchwitz, M., Frerick, J., Noël, S., Rozanov, V. V., Chance, K. V., and Goede, A. P. H.: SCIAMACHY: Mission Objectives and Measurement Modes, *J. Atmos. Sci.*, 56, 127–150, 1999. 26178
- Brock, C. A., Cozic, J., Bahreini, R., Froyd, K. D., Middlebrook, A. M., McComiskey, A., Brioude,

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



J., Cooper, O. R., Stohl, A., Aikin, K. C., de Gouw, J. A., Fahey, D. W., Ferrare, R. A., Gao, R.-S., Gore, W., Holloway, J. S., Hbler, G., Jefferson, A., Lack, D. A., Lance, S., Moore, R. H., Murphy, D. M., Nenes, A., Novelli, P. C., Nowak, J. B., Ogren, J. A., Peischl, J., Pierce, R. B., Pilewskie, P., Quinn, P. K., Ryerson, T. B., Schmidt, K. S., Schwarz, J. P., Sodemann, H., Spackman, J. R., Stark, H., Thomson, D. S., Thornberry, T., Veres, P., Watts, L. A., Warneke, C., and Wollny, A. G.: Characteristics, sources, and transport of aerosols measured in spring 2008 during the aerosol, radiation, and cloud processes affecting Arctic Climate (ARCPAC) Project, *Atmos. Chem. Phys.*, 11, 2423–2453, doi:10.5194/acp-11-2423-2011, 2011. 26181, 26183, 26185

Callies, J., Corpaccioli, E., Hahne, A., and Lefebvre, A.: GOME-2 – Metop's second- generation sensor for operational ozone monitoring, *ESA Bulletin-European Space Agency*, 102, 28–36, 2000. 26186

Chance, K.: Analysis of BrO Measurements from the Global Ozone Monitoring Experiment, *Geophys. Res. Lett.*, 25, 3335–3338, 1998. 26178, 26185

Claas, J., Braak R., and Kroon, M.: Row anomaly: introduction and flagging, OMI Science Team Meeting, De Bilt, The Netherlands, June 2010. 26185

De Smedt, I., Müller, J.-F., Stavrou, T., van der A, R., Eskes, H., and Van Roozendael, M.: Twelve years of global observations of formaldehyde in the troposphere using GOME and SCIAMACHY sensors, *Atmos. Chem. Phys.*, 8, 4947–4963, doi:10.5194/acp-8-4947-2008, 2008. 26194

Dibb, J. E., Ziemba, L. D., Luxford, J., and Beckman, P.: Bromide and other ions in the snow, firn air, and atmospheric boundary layer at Summit during GSHOX, *Atmos. Chem. Phys.*, 10, 9931–9942, doi:10.5194/acp-10-9931-2010, 2010. 26185

Dikty, S., Richter, A., Weber, M., Noel, S., Wittrock, F. Bovensmann, H., Munro, R., Lang, R., and Burrows, J. P.: GOME-2 optical degradation as seen in level 2 data time series (2007–2010; BrO, NO<sub>2</sub>, HCHO, H<sub>2</sub>O, and O<sub>3</sub>), EGU General Assembly, Vienna, Austria, available at: [http://www.doas-bremen.de/posters/egu\\_2011\\_dikty.pdf](http://www.doas-bremen.de/posters/egu_2011_dikty.pdf), April 2011. 26186

Deutsches Zentrum für Luft und Raumfahrt e.V.: Algorithm Theoretical Basis Document for GOME-2 Total Column Products of Ozone, Minor Trace Gases, and Cloud Properties, Oberpfaffenhofen, Germany, 2009. 26186

Donohoue, D., Carlson, D., and Simpson, W. R.: Rapid methods for inversion of MAX-DOAS elevation profiles to surface-associated box concentrations, visibility, and heights: application to analysis of Arctic BrO events, *Atmos. Meas. Tech. Discuss.*, 3, 4645–4674,

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


doi:10.5194/amtd-3-4645-2010, 2010. 26177

Dorf, M., Butz, A., Camy-Peyret, C., Chipperfield, M. P., Kritten, L., and Pfeilsticker, K.: Bromine in the tropical troposphere and stratosphere as derived from balloon-borne BrO observations, *Atmos. Chem. Phys.*, 8, 7265–7271, doi:10.5194/acp-8-7265-2008, 2008. 26177

5 Douglas, T. A., Sturm, M., Simpson, W. R., Brooks, S., Lindberg, S., and Perovich, D.: Elevated mercury measured in snow and frost flowers near Arctic sea ice leads, *Geophys. Res. Lett.*, 32, L04502, doi:10.1029/2004GL022132, 2005. 26176

Fan, S.-M. and Jacob, D. J.: Surface ozone depletion in Arctic spring sustained by bromine reactions on aerosols, *Nature*, 359, 522–524, 1992.

10 Fitzenberger, R., Bösch, H., Camy-Peyret, C., Chipperfield, M. P., Harder, H., Platt, U., Sinnhuber, B.-M., Wagner, T., and Pfeilsticker, K.: First profile measurements of tropospheric BrO, *Geophys. Res. Lett.*, 27, 2921–2924, 2000. 26177

Frieß, U., Chipperfield, M. P., Harder, H., Otten, C., Platt, U., Pyle, J., Wagner, T., and Pfeilsticker, K.: Intercomparison of measured and modelled BrO slant column amounts for the Arctic winter and spring 1994/95, *Geophys. Res. Lett.*, 26, 1861–1864, 1999. 26177

15 Garcia, R. R., Marsh, D. R., Kinnison, D. E., Boville, B. A., and Sassi, F.: Simulations of secular trends in the middle atmosphere, 1950–2003, *J. Geophys. Res.*, 112, D09301, doi:10.1029/2006JD007485, 2007. 26187

GOME Users Manual, ESA Special Publication SP-1182, ESTEC, Noordwijk, The Netherlands, 1995. 26178

20 Hausmann, M. and Platt, U.: Spectroscopic measurement of bromine oxide and ozone in the high Arctic during Polar Sunrise Experiment 1992, *J. Geophys. Res.*, 99, 25399–25413, 1994. 26176, 26177, 26183, 26199, 26200, 26206

Hegels, E., Crutzen, P. J., Klupfel, T., Perner, D., and Burrow, J. P.: Global distribution of atmospheric bromine-monoxide from GOME on earth observing satellite ERS-2, *Geophys. Res. Lett.*, 25, 3127–3130, 1998. 26178

25 Hendrick, F., Van Roozendaal, M., Chipperfield, M. P., Dorf, M., Goutail, F., Yang, X., Fayt, C., Hermans, C., Pfeilsticker, K., Pommereau, J.-P., Pyle, J. A., Theys, N., and De Mazière, M.: Retrieval of stratospheric and tropospheric BrO profiles and columns using ground-based zenith-sky DOAS observations at Harestua, 60° N, *Atmos. Chem. Phys.*, 7, 4869–4885, doi:10.5194/acp-7-4869-2007, 2007. 26177

30 Hendrick, F., Johnston, P. V., De Mazière, Fayt, C., Hermans, C., Kreher, K., Theys, N., Thomas, A., and Van Roozendaal, M.: One-decade trend analysis of stratospheric BrO

**Satellite-derived  
Arctic tropospheric  
BrO columns**

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



over Harestua (60°N) and Lauder (45°S) reveals a decline, *Geophys. Res. Lett.*, 35, L14801, doi:10.1029/2008GL034154, 2008. 26177

Holmes, C. D., Jacob, D. J., Corbitt, E. S., Mao, J., Yang, X., Talbot, R., and Slemr, F.: Global atmospheric model for mercury including oxidation by bromine atoms, *Atmos. Chem. Phys.*, 10, 12037–12057, doi:10.5194/acp-10-12037-2010, 2010. 26176

Hönninger and Platt, U.: Observations of BrO and its vertical distribution during surface ozone depletion at Alert, *Atmos. Environ.*, 36, 2481–2489, 2002.

Hönninger, G., Leser, H., Sebastián, O., and Platt, U.: Ground-based measurements of halogen oxides at the Hudson Bay by longpath DOAS and passive MAX-DOAS, *Geophys. Res. Lett.*, 31, L04111, doi:10.1029/2003GL018982, 2004b. 26177

Jones, A. E., Anderson, P. S., Wolff, E. W., Turner, J., Rankin, A. M., and Colwell, S. R.: A role for newly forming sea ice in springtime polar tropospheric ozone loss? Observational evidence from Halley station, Antarctica, *J. Geophys. Res.*, 111, D08306, doi:10.1029/2005JD006566, 2006. 26176

Jones, A. E., Anderson, P. S., Begoin, M., Brough, N., Hutterli, M. A., Marshall, G. J., Richter, A., Roscoe, H. K., and Wolff, E. W.: BrO, blizzards, and drivers of polar tropospheric ozone depletion events, *Atmos. Chem. Phys.*, 9, 4639–4652, doi:10.5194/acp-9-4639-2009, 2009. 26176, 26203, 26207

Jones, A. E., Anderson, P. S., Wolff, E. W., Roscoe, H. K., Marshall, G. J., Richter, A., Brough, N., and Colwell, S. R.: Vertical structure of Antarctic tropospheric ozone depletion events: characteristics and broader implications, *Atmos. Chem. Phys.*, 10, 7775–7794, doi:10.5194/acp-10-7775-2010, 2010. 26176, 26203, 26207

Jacob, D. J., Crawford, J. H., Maring, H., Clarke, A. D., Dibb, J. E., Emmons, L. K., Ferrare, R. A., Hostetler, C. A., Russell, P. B., Singh, H. B., Thompson, A. M., Shaw, G. E., McCauley, E., Pederson, J. R., and Fisher, J. A.: The Arctic Research of the Composition of the Troposphere from Aircraft and Satellites (ARCTAS) mission: design, execution, and first results, *Atmos. Chem. Phys.*, 10, 5191–5212, doi:10.5194/acp-10-5191-2010, 2010. 26181, 26183, 26185

Kaleschke, L., Richter, A., Burrows, J., Afe, O., Heygster, G., Notholt, J., Rankin, A. M., Roscoe, H. K., Hollwedel, J., Wagner, T., and Jacobi, H.-W.: Frost flowers on sea ice as a source of sea salt and their influence on tropospheric halogen chemistry, *Geophys. Res. Lett.*, 31, L16114, doi:10.1029/2004GL020655, 2004. 26176

Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha,

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)




[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


- S., White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *B. Am. Meteorol. Soc.*, 77, 437–470, 1996.
- 5 Koo, J.-H., Wang, Y., Choi, S., Chance, K., Kurosu, T. P., Oltmans, S. J., Weinheimer, A. J., Ryerson, T. B., Hair, J. W., Fenn, M. A., and Shaw, M.: Spatial heterogeneity of ozone depletion in the Arctic troposphere, in preparation, 2011. 26180
- Kovalenko, L. J., Livesey, N. J., Salawitch, R. J., Camy-Peyret, C., Chipperfield, M. P., Cofield, R. E., Dorf, M., Drouin, B. J., Froidevaux, L., Fuller, R., A., Goutail, F., Jarnot, R. F., Jucks, K., Knosp, B. W., Lambert, A., MacKenzie, I. A., Pfeilsticker, K., Pommereau, J.-P., Read, 10 W. G., Santee, M. L., Schwartz, M. J., Snyder, W. V., Stachnik, R., Stek, P. C., Wagner, P. A., and Waters, J. W.: Validation of Aura Microwave Limb Sounder BrO observations in the stratosphere, *J. Geophys. Res.*, 112, D24S41, doi:10.1029/2007JD008817, 2007. 26177
- Kreher, K., Johnston, P. V., Wood, S. W., Nardi, B., and Platt, U.: Ground-based measurements of tropospheric and stratospheric BrO at Arrival Heights, Antarctica, *Geophys. Res. Lett.*, 24, 3021–3024, 1997.
- 15 Kurosu, T. and Chance, K.: OMBRO Readme file, [https://www.cfa.harvard.edu/~tkurosu/SatelliteInstruments/OMI/PGEReleases/READMEs/OMBRO\\_v300\\_README.pdf](https://www.cfa.harvard.edu/~tkurosu/SatelliteInstruments/OMI/PGEReleases/READMEs/OMBRO_v300_README.pdf), 2011. 26180, 26186
- Lehrer, E., Hönninger, G., and Platt, U.: A one dimensional model study of the mechanism of halogen liberation and vertical transport in the polar troposphere, *Atmos. Chem. Phys.*, 4, 2427–2440, doi:10.5194/acp-4-2427-2004, 2004. 26204
- 20 Levelt, P. F., van den Oord, G. H. J., Dobber, M. R., Mälkki, A., Visser, H., de Vries, J., Stammes, P., Lundell, J., and Saari, H.: The Ozone Monitoring Instrument, *IEEE T. Geosci. Remote Sens.*, 44, 1093–1101, doi:10.1109/TGRS.2006.872333, 2006. 26178, 26185
- 25 Liao, J., Sihler, H., Huey, L. G., Neuman, J. A., Tanner, D. J., Friess, U., Platt, U., Flocke, F. M., Orlando, J. J., Shepson, P. B., Beine, H. J., Weinheimer, A. J., Sjostedt, S. J., Nowak, J. B., Knapp, D. J., Staebler, R. M., Zheng, W., Sander, R., Hall, S. R., and Ullmann, K.: A comparison of Arctic BrO measurements by chemical ionization mass spectrometry and long path-differential optical absorption spectroscopy, *J. Geophys. Res.-Atmos.*, 116, D00r02, doi:10.1029/2010jd014788, 2011. 26177, 26181, 26199
- 30 Liao, J., Huey, L. G., Scheuer, E., Dibb, J. E., Stickel, R. E., Tanner, D. J., Neuman, J. A., Nowak, J. B., Choi, S., Wang, Y., Joiner, J., Salawitch, R. J., Canty, T., Weinheimer, A. J., Shetter, R. E., Fried, A., Brune, W., Anderson, B., Zhang, W., Chen, G., Crawford, J., and Hecobian, A.:

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Characterization of soluble bromide measurements and a case study of BrO observations during ARCTAS, *Atmos. Chem. Phys. Discuss.*, submitted, 2011. 26177, 26180, 26183, 26185, 26198

Livesey, N. J., Kovalenko L. J., Salawitch, R. J., MacKenzie, I. A., Chipperfield, M. P., Read, W. G., Jarnot, R. F., and Waters, J. W.: EOS Microwave Limb Sounder observations of upper stratospheric BrO: Implications for total bromine, *Geophys. Res. Lett.*, 33, L20817, doi:10.1029/2006GL026930, 2006. 26177

Lu, J. Y., Schroeder, W. H., Barrie, L. A., Steffen, A., Welch, H. E., Martin, K., Lockhart, L., Hunt, R. V., Boila, G., and Richter, A.: Magnification of atmospheric mercury deposition to polar regions in springtime: the link to tropospheric ozone depletion chemistry, *Geophys. Res. Lett.*, 28, 3219–3222, 2001. 26176

Martinez, M., Arnold, T., and Perner, D.: The role of bromine and chlorine chemistry for arctic ozone depletion events in Ny-Ålesund and comparison with model calculations, *Ann. Geophys.*, 17, 941–956, doi:10.1007/s00585-999-0941-4, 1999. 26176, 26177, 26183

McElroy, C. T., McLinden, C. A., and McConnell, J. C.: Evidence for bromine monoxide in the free troposphere during the Arctic polar sunrise, *Nature*, 397, 338–341, 1999. 26177

Munro, R., Eisinger, M., Anderson, C., Callies, J., Corpaccioli, E., Lang, R., Lefebvre, A., Livschitz, Y., and Albinana, A. P.: GOME-2 on MetOp, in: *Proc. of The 2006 EUMETSAT Meteorological Satellite Conference*, Helsinki, Finland, 12–16 June 2006, EUMETSAT, p. 48, 2006. 26178

Neuman, J. A., Nowak, J. B., Huey, L. G., Burkholder, J. B., Dibb, J. E., Holloway, J. S., Liao, J., Peischl, J., Roberts, J. M., Ryerson, T. B., Scheuer, E., Stark, H., Stickel, R. E., Tanner, D. J., and Weinheimer, A.: Bromine measurements in ozone depleted air over the Arctic Ocean, *Atmos. Chem. Phys.*, 10, 6503–6514, doi:10.5194/acp-10-6503-2010, 2010. 26177, 26180, 26181, 26183, 26184, 26185, 26199, 26203, 26204

Nolin, A., Armstrong, R. L., and Maslanik, J.: Near Real-Time SSM/I EASE-Grid Daily Global Ice Concentration and Snow Extent, Jan to Mar 2004 (updated daily), Boulder, CO, USA: National Snow and Ice Data Center, Digital media, 1998. 26189

Palmer, P. I., Jacob, D. J., Chance, K., Martin, R. V., Spurr, R. J. D., Kurosu, T. P., Bey, I., Yantosca, R., Fiore, A., and Li, Q.: Air-mass factor formulation for spectroscopic measurements from satellites: application to formaldehyde retrievals from GOME, *J. Geophys. Res.*, 106(D13), 14539–14550, doi:10.1029/2000JD900772, 2001. 26190

Pfeilsticker, K., Sturges, W. T., Bosch, H., Camy-Peyret, C., Chipperfield, M. P., Engel, A.,

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)




[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Fitzenberger, R., Muller, M., Payan, S., and Sinnhuber, B. M.: Lower stratospheric organic and inorganic bromine budget for the Arctic winter 1998/99, *Geophys. Res. Lett.*, 27, 3305–3308, 2000 26177

Platnick, S., King, M. D., Ackerman, S. A., Menzel, W. P., Baum, B. A., Ri/edi, J. C., and Frey, R. A.: The MODIS cloud products: algorithms and examples from Terra, *IEEE T. Geosci. Remote*, 41, 459–473, 2003. 26189

Platt, U. and Hönninger, G.: The role of halogen species in the troposphere, *Chemosphere*, 52, 325–338, 2003 26177

Pundt, I., Pommereau, J. P., Chipperfield, M. P., Van Roozendaal, M., and Goutail, F.: Climatology of the stratospheric BrO vertical distribution by balloon-borne UV-visible spectrometry, *J. Geophys. Res.*, 107, 4806, doi:10.1029/2002JD002230, 2002. 26177

Richter, A., Wittrock, F., Eisinger, M., and Burrows, J. P.: GOME Observations of Tropospheric BrO in Northern Hemispheric Spring and Summer 1997, *Geophys. Res. Lett.*, 25, 2683–2686, 1998. 26179

Ridley, B. A., Atlas, E. L., Montzka, D. D., Browell, E. V., Cantrell, C. A., Blake, D. R., Blake, N. J., Cinquini, L., Coffey, M. T., Emmons, L. K., Cohen, R. C., DeYoung, R. J., Dibb, J. E., Eisele, F. L., Flocke, F. M., Fried, A., Grahek, F. E., Grant, W. B., Hair, J. W., Hannigan, J., Heikes, B. J., Lefer, B. L., Mauldin, R. L., Moody, J. L., Shetter, R. E., Snow, J. A., Talbot, R. W., Thornton, J. A., Walega, J. G., Weinheimer, A. J., Wert, B. P., and Wimmers, A. J.: Ozone Depletion Events Observed in the High Latitude Surface Layer During the TOPSE Aircraft Program, *J. Geophys. Res.*, 108, 8356, doi:10.1029/2001JD001507, 2003. 26176, 26185, 26200

Rienecker, M. M., Suarez, M. J., and Todling, R.: The GEOS-5 data assimilation system Documentation of versions 5.0.1, 5.1.0, and 5.2.0. NASA Tech. Memo. 2007-104606, vol. 27, edited by: Suarez, M. J., 2007. 26187, 26189

Rozanov, A., Rozanov, V., Buchwitz, M., Kokhanovsky, A., and Burrows, J. P.: SCIATRAN 2.0 a new radiative transfer model for geophysical applications, *Adv. Space Res.*, 36, 1015–1019, 2005. 26190

Rozanov, A., Köhl, S., Doicu, A., McLinden, C., Pukite, J., Bovensmann, H., Burrows, J. P., Deutschmann, T., Dorf, M., Goutail, F., Grunow, K., Hendrick, F., von Hobe, M., Hrechanyy, S., Lichtenberg, G., Pfeilsticker, K., Pommereau, J. P., Van Roozendaal, M., Stroh, F., and Wagner, T.: BrO vertical distributions from SCIAMACHY limb measurements: comparison of algorithms and retrieval results, *Atmos. Meas. Tech.*, 4, 1319–1359, doi:10.5194/amt-4-



1319-2011, 2011. 26178

5 Saiz-Lopez, A., Mahajan, A. S., Salmon, R. A., Bauguitte, S. J.-B., Jones, A. E., Roscoe, H. K., and Plane, J. M. C.: Boundary layer halogens in coastal Antarctica, *Science*, 317, 348–351, doi:10.1126/science.1141408, 2007.

10 Salawitch, R. J., Weisenstein, D. K., Kovalenko, L. J., Sioris, C. E., Wennberg, P. O., Chance, K., Ko, M. K. W., and McLinden, C. A.: Sensitivity of ozone to bromine in the lower stratosphere, *Geophys. Res. Lett.*, 32, L05811, doi:10.1029/2004GL021504, 2005. 26180

15 Salawitch, R., Canty, T., Kurosu, T., Chance, K., Liang, Q., da Silva, A., Pawson, S., Nielsen, J. E., Rodriguez, J. M., Bhartia, P. K., Liu, X., Huey, L. G., Liao, J., Stickel, R. E., Tanner, D. J., Dibb, J. E., Simpson, W. R., Donohoue, D., Weinheimer, A., Flocke, F., Knapp, D., Montzka, D., Neuman, J. A., Nowak, J. B., Ryerson, T. B., Oltmans, S., Blake, D. R., Atlas, E. L., Kinnison, D. E., Tilmes, S., Pan, L. L., Hendrick, F., Van Roozendael, M., Kreher, K., Johnston, P. V., Gao, R. S., Johnson, B., Bui, T. P., Chen, G., Pierce, R. B., Crawford, J. H., and Jacob, D. J.: A new interpretation of total column BrO during Arctic spring, *Geophys. Res. Lett.*, 37, L21805, doi:10.1029/2010GL043798, 2010. 26179, 26180, 26181, 26182, 26184, 26186, 26187, 26188, 26195, 26196, 26198, 26201, 26202, 26203, 26205, 26206

20 Sander, S. P., Friedl, R. R., Golden, D. M., Kurylo, M. J., Moortgat, G. K., Keller-Rudek, H., Wine, P. H., Ravishankara, A. R., Kolb, C. E., Molina, M. J., Finlayson-Pitts, B. J., Huie, R. E., and Orkin, V. L.: Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies – Evaluation Number 15, NASA JPL Publication 06-2, Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, 2006. 26195

Schroeder, W. H., Anlauf, K. G., Barrie, L. A., Lu, J. Y., Steffen, A., Schneeberger, D. R., and Berg, T.: Arctic springtime depletion of mercury, *Nature*, 394, 331–332, 1998. 26176

25 Schofield, R., Kreher, K., Connor, B. J., Johnston, P. V., Thomas, A., Shooter, D., Chipperfield, M. P., Rodgers, C. D., and Mount, G. H.: Retrieved tropospheric and stratospheric BrO columns over Lauder, New Zealand, *J. Geophys. Res.*, 109, D14304, doi:10.1029/2003JD004463, 2004. 26177

30 Schofield, R., Johnston, P. V., Thomas, A., Kreher, K., Connor, B. J., Wood, S., Shooter, D., Chipperfield, M. P., Richter, A., von Glasow, R., and Rodgers, C. D.: Tropospheric and stratospheric BrO columns over Arrival Heights, Antarctica, 2002, *J. Geophys. Res.*, 111, D22310, doi:10.1029/2005JD007022, 2006. 26177

Shinozuka, Y., Redemann, J., Livingston, J. M., Russell, P. B., Clarke, A. D., Howell, S. G., Freitag, S., O'Neill, N. T., Reid, E. A., Johnson, R., Ramachandran, S., McNaughton, C. S.,

ACPD

11, 26173–26243, 2011

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Satellite-derived  
Arctic tropospheric  
BrO columns**

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Kapustin, V. N., Brekhovskikh, V., Holben, B. N., and McArthur, L. J. B.: Airborne observation of aerosol optical depth during ARCTAS: vertical profiles, inter-comparison and fine-mode fraction, *Atmos. Chem. Phys.*, 11, 3673–3688, doi:10.5194/acp-11-3673-2011, 2011. 26192
- 5 Simpson, W. R., Carlson, D., Hönninger, G., Douglas, T. A., Sturm, M., Perovich, D., and Platt, U.: First-year sea-ice contact predicts bromine monoxide (BrO) levels at Barrow, Alaska better than potential frost flower contact, *Atmos. Chem. Phys.*, 7, 621–627, doi:10.5194/acp-7-621-2007, 2007a. 26176, 26177
- Simpson, W. R., von Glasow, R., Riedel, K., Anderson, P., Ariya, P., Bottenheim, J., Burrows, J., Carpenter, L. J., Frieß, U., Goodsite, M. E., Heard, D., Hutterli, M., Jacobi, H.-W., Kaleschke, L., Neff, B., Plane, J., Platt, U., Richter, A., Roscoe, H., Sander, R., Shepson, P., Sodeau, J., Steffen, A., Wagner, T., and Wolff, E.: Halogens and their role in polar boundary-layer ozone depletion, *Atmos. Chem. Phys.*, 7, 4375–4418, doi:10.5194/acp-7-4375-2007, 2007b. 26175, 26205
- 10 Sinnhuber, B.-M., Rozanov, A., Sheode, N., Afe, O. T., Richter, A., Sinnhuber, M., Wittrock, F., Stiller, G. P., von Clarmann, T., Linden, A., and Burrows, J. P.: Global observations of stratospheric bromine monoxide from SCIAMACHY, *Geophys. Res. Lett.*, 32, L20810, doi:10.1029/2005GL023839, 2005. 26178
- Sioris, C. E., Kovalenko, L. J., McLinden, C. A., Salawitch, R. J., Van Roozendael, M., Goutail, F., Dorf, M., Pfeilsticker, K., Chance, K., von Savigny, C., Liu, X., Kurosu, T. P., Pommereau, J.-P., Boesch, H., and Frerick, J.: Latitudinal and vertical distribution of bromine monoxide in the lower stratosphere from Scanning Imaging Absorption Spectrometer for Atmospheric Chartography limb scattering measurements, *J. Geophys. Res.*, 111, D14301, doi:10.1029/2005JD006479, 2006. 26178
- 20 SPARC Report on the Evaluation of Chemistry-Climate Models, edited by: Eyring, V., Shepherd, T. G., and Waugh, D. W., SPARC Report No. 5, WCRP-132, WMO/TD-No. 1526, <http://www.atmos.physics.utoronto.ca/SPARC>, 2010. 26188
- Spurr, R. J. D., Kurosu, T. P., and Chance, K.: A linearized discrete ordinate radiative transfer model for atmospheric remote sensing retrieval, *J. Quant. Spectrosc. Ra.*, 68, 689–735, 2001. 26191
- 30 Theys, N., Van Roozendael, M., Hendrick, F., Fayt, C., Hermans, C., Baray, J.-L., Goutail, F., Pommereau, J.-P., and De Mazière, M.: Retrieval of stratospheric and tropospheric BrO columns from multi-axis DOAS measurements at Reunion Island (21° S, 56° E), *Atmos. Chem. Phys.*, 7, 4733–4749, doi:10.5194/acp-7-4733-2007, 2007. 26177

Theys, N., Van Roozendael, M., Errera, Q., Hendrick, F., Daerden, F., Chabrilat, S., Dorf, M., Pfeilsticker, K., Rozanov, A., Lotz, W., Burrows, J. P., Lambert, J.-C., Goutail, F., Roscoe, H. K., and De Mazière, M.: A global stratospheric bromine monoxide climatology based on the BASCOE chemical transport model, *Atmos. Chem. Phys.*, 9, 831–848, doi:10.5194/acp-9-831-2009, 2009. 26179, 26180, 26203, 26206

Theys, N., Van Roozendael, M., Hendrick, F., Yang, X., De Smedt, I., Richter, A., Begoin, M., Errera, Q., Johnston, P. V., Kreher, K., and De Mazière, M.: Global observations of tropospheric BrO columns using GOME-2 satellite data, *Atmos. Chem. Phys.*, 11, 1791–1811, doi:10.5194/acp-11-1791-2011, 2011. 26176, 26177, 26178, 26179, 26180, 26181, 26188, 26189, 26190, 26191, 26192, 26194, 26195, 26197, 26206

Toyota, K., McConnell, J. C., Lupu, A., Neary, L., McLinden, C. A., Richter, A., Kwok, R., Semeniuk, K., Kaminski, J. W., Gong, S.-L., Jarosz, J., Chipperfield, M. P., and Sioris, C. E.: Analysis of reactive bromine production and ozone depletion in the Arctic boundary layer using 3-D simulations with GEM-AQ: inference from synoptic-scale patterns, *Atmos. Chem. Phys.*, 11, 3949–3979, doi:10.5194/acp-11-3949-2011, 2011. 26176

Tuckermann, M., Ackermann, R., Golz, C., Lorenzen-Schmidt, H., Senne, T., Stutz, J., Trost, B., Unold, W., and Platt, U.: DOAS-observation of halogen radical-catalysed Arctic boundary layer ozone destruction during the ARCTOC-campaigns 1995 and 1996 in Ny-Ålesund, Spitsbergen, *Tellus*, 49B, 533–555, 1997. 26176, 26177, 26183, 26199, 26200, 26206

Van Roozendael, M., Wagner, T., Richter, A., Pundt, I., Arlander, D. W., Burrows, J. P., Chipperfield, M., Fayt, C., Johnston, P. V., Lambert, J. C., Kreher, K., Pfeilsticker, K., Platt, U., Pommereau, J. P., Sinnhuber, B. M., Tornkvist, K. K., and Wittrock, F.: Intercomparison of BrO measurements from ERS-2 GOME, ground-based and balloon platforms, in: *Remote Sensing Of Trace Constituents In The Lower Stratosphere, Troposphere And The Earth's Surface: Global Observations, Air Pollution And The Atmospheric Correction*, vol. 29, *Adv. Space Res.*, 11, 1661–1666, 2002.

Vasilkov, A. P., Joiner, J., Spurr, R. J. D., Bhartia, P. K., Levelt, P., and Stephens, G.: Evaluation of the OMI cloud pressures derived from rotational Raman scattering by comparisons with other satellite data and radiative transfer simulations, *J. Geophys. Res.*, 113, D15S19, doi:10.1029/2007JD008689, 2008. 26188

Vasilkov, A. P., Joiner, J., Haffner, D., Bhartia, P. K., and Spurr, R. J. D.: What do satellite backscatter ultraviolet and visible spectrometers see over snow and ice? A study of clouds and ozone using the A-train, *Atmos. Meas. Tech.*, 3, 619–629, doi:10.5194/amt-3-619-2010,

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


2010. 26188, 26189, 26192

Wagner, T. and Platt, U.: Satellite mapping of enhanced BrO concentrations in the troposphere, *Nature*, 395, 486–490, 1998. 26178, 26179

Wagner, T., Leue, C., Wenig, M., Pfeilsticker, K., and Platt, U.: Spatial and temporal distribution of enhanced boundary layer BrO concentrations measured by the GOME instrument aboard ERS-2, *J. Geophys. Res.*, 106, 24225–24235, 2001. 26179

Wagner, T., Ibrahim, O., Sinreich, R., Frieß, U., von Glasow, R., and Platt, U.: Enhanced tropospheric BrO over Antarctic sea ice in mid winter observed by MAX-DOAS on board the research vessel Polarstern, *Atmos. Chem. Phys.*, 7, 3129–3142, doi:10.5194/acp-7-3129-2007, 2007. 26176, 26180

Wahner A., Ravishankara, A. R., Sander S. P., Friedl R. R.: Absorption cross-section of BrO between 312 and 385 nm at 298 and 223-K, *Chem. Phys. Lett.*, 152, 507–512, 1988. 26186

Wamsley, P. R., Elkins, J. W., Fahey, D. W., Dutton, G. S., Volk, C. M., Myers, R. C., Montzka, S. A., Butler, J. H., Clarke, A. D., Fraser, P. J., Steele, L. P., Lucarelli, M. P., Atlas, E. L., Schauffler, S. M., Blake, D. R., Rowland, F. S., Sturges, W. T., Lee, J. M., Penkett, S. A., Engel, A., Stimpfle, R. M., Chan, K. R., Weisenstein, D. K., Ko, M. K. W., and Salawitch, R. J.: Distribution of halon-1211 in the upper troposphere and lower stratosphere and the 1994 bromine budget, *J. Geophys. Res.*, 103, 1513–1526, doi:10.1029/97JD02466, 1998. 26188

Wilmouth, D. M., Hanisco, T. F., Donahue, N. M., and Anderson, J. G.: Fourier transform ultraviolet spectroscopy of the A (2)Pi(3/2) ← X (II3/2)-I-2 transition of BrO, *J. Phys. Chem. A*, 103, 8935–8945, 1999. 26186

WMO (World Meteorological Organization), Scientific Assessment of Ozone Depletion: 2002, Global Ozone Research and Monitoring Project – Report No. 47, Geneva, Switzerland, 2003. 26181

WMO (World Meteorological Organization), Scientific Assessment of Ozone Depletion: 2006, Global Ozone Research and Monitoring Project – Report No. 50, Geneva, Switzerland, 2007. 26178

WMO (World Meteorological Organization), Scientific Assessment of Ozone Depletion: 2010, Global Ozone Research and Monitoring Project – Report No. 52, 516 pp., Geneva, Switzerland, 2011. 26177, 26187, 26188

Yang, X., Pyle, J. A., and Cox, R. A.: Sea salt aerosol production and bromine release: Role of snow on sea ice, *Geophys. Res. Lett.*, 35, L16815, doi:10.1029/2008GL034536, 2008. 26176, 26203, 26207

Yang, X., Pyle, J. A., and Cox, R. A.: Sea salt aerosol production and bromine release: Role of snow on sea ice, *Geophys. Res. Lett.*, 35, L16815, doi:10.1029/2008GL034536, 2008. 26176, 26203, 26207

5 Zeng, T., Wang, Y., Chance, K., Browell, E. V., Ridley, B. A., and Atlas, E. L.: Widespread persistent near-surface ozone depletion at northern high latitudes in spring, *Geophys. Res. Lett.*, 30, 2298, doi:10.1029/2003GL018587, 2003. 26179

Zeng, T., Wang, Y., Chance, K., Blake, N., Blake, D., and Ridley, B.: Halogen-driven low-altitude O<sub>3</sub> and hydrocarbon losses in spring at northern high latitudes, *J. Geophys. Res.*, 111, D17313, doi:10.1029/2005JD006706, 2006. 26179

ACPD

11, 26173–26243, 2011

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 1.** Information for NASA DC-8 BrO profiles shown in Fig. 1. “Day” means day of April 2008. Alt<sub>B</sub> (Alt<sub>T</sub>) means the bottom (top) altitude of the profile. “Col.” stands for BrO vertical column density (VCD). Units of BrO VCD are 10<sup>13</sup> molecules cm<sup>-2</sup>. “Col.<sub>ADDED</sub>” is the estimated amount of BrO that was added to the measured in situ column to provide an estimate of the complete tropospheric column.

No.	Day	UTC	Lat. [°]	Lon. [°]	Alt <sub>B</sub> [m]	Alt <sub>T</sub> [km]	H <sub>TRP</sub> [km]	Col. IN-SITU	Col. ADDED
1	16	20:52	69.7	-145.3	95	4.2	7.6	4.36	5.3 %
2	16	21:17	71.1	-147.4	82	4.2	7.0	2.85	8.1 %
3	16	22:17	73.2	-159.0	89	4.0	6.9	3.97	5.8 %
4	16	22:34	72.1	-161.9	105	3.6	7.3	2.14	27.4 %
5	16	23:44	66.3	-165.5	206	4.1	7.0	3.04	7.6 %
6	16	24:06	65.2	-165.2	387	4.2	9.3	2.87	8.0 %
7	17	21:34	70.3	-148.7	75	5.2	8.0	2.68	3.9 %
8	17	21:53	71.0	-148.2	83	6.3	8.1	2.18	22.4 %
9	17	22:51	77.1	-148.4	99	6.3	7.4	3.86	0.3 %
10	17	23:29	80.3	-148.4	90	5.5	8.2	3.44	1.1 %
11	17	24:13	84.9	-148.5	112	5.4	7.3	3.04	3.4 %
12	17	24:40	87.2	-148.4	112	5.9	6.3	4.35	0.8 %
13	17	26:58	78.3	-156.5	89	7.0	7.7	5.20	3.7 %
14	17	27:15	77.1	-156.6	89	4.5	7.5	2.57	9.0 %
15	17	28:15	71.8	-157.5	90	4.1	6.6	2.11	23.2 %
16	17	28:35	71.3	-156.8	75	5.8	8.2	1.88	10.6 %

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

**Table 2a.** Similar to Table 1 but for NOAA WP-3D profiles of BrO, measured on 12 to 15 April 2008.

No.	Day	UTC	Lat. [°]	Lon. [°]	Alt <sub>B</sub> [m]	Alt <sub>T</sub> [km]	H <sub>TRP</sub> [km]	Col. IN-SITU	Col. ADDED
1	12	21:14	65.8	-148.1	289	4.6	7.4	2.55	6.5 %
2	12	23:06	70.6	-152.7	261	5.2	7.3	1.31	8.0 %
3	12	23:39	72.6	-153.3	107	4.1	7.2	1.96	37.7 %
4	12	24:00	74.0	-153.3	77	3.3	7.0	1.27	43.8 %
5	12	24:27	75.1	-151.2	80	5.0	7.2	3.34	3.1 %
6	12	24:54	75.1	-144.5	104	5.0	6.7	0.15	67.7 %
7	12	26:09	74.0	-139.7	99	3.4	6.5	1.41	39.4 %
8	15	19:17	64.8	-149.5	166	3.5	8.2	0.62	62.9 %
9	15	22:24	66.4	-165.0	135	4.6	5.9	0.73	22.6 %
10	15	22:48	67.8	-165.0	135	4.3	5.9	0.81	28.5 %
11	15	23:12	69.1	-165.1	77	3.2	6.1	1.71	32.5 %
12	15	24:30	71.9	-161.7	93	6.6	6.7	0.10	2.9 %
13	15	25:00	71.7	-161.8	100	6.5	6.6	0.07	17.9 %
14	15	25:45	69.6	-157.2	207	4.9	6.3	1.60	15.2 %
15	15	26:50	65.4	-148.6	203	5.3	7.3	1.05	10.0 %

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

**Table 2b.** Similar to Table 1 but for NOAA WP-3D profiles of BrO, measured on 18 to 21 April 2008.

No.	Day	UTC	Lat. [°]	Lon. [°]	Alt <sub>B</sub> [m]	Alt <sub>T</sub> [km]	H <sub>TRP</sub> [km]	Col. IN-SITU	Col. ADDED
16	18	22:39	72.5	-141.9	92	7.1	8.5	4.36	-0.0%
17	18	23:24	73.0	-137.7	107	6.8	8.4	0.85	0.4%
18	18	24:51	70.9	-144.2	146	5.2	9.6	1.92	5.5%
19	19	22:18	65.5	-148.5	219	4.8	11.7	4.49	13.8%
20	19	24:06	71.5	-156.5	188	4.7	8.7	2.44	6.8%
21	19	26:51	74.0	-160.0	151	3.8	6.6	1.79	22.0%
22	19	27:39	72.6	-153.8	151	7.4	8.1	2.12	0.0%
23	19	29:04	66.7	-149.1	304	5.4	11.3	2.29	4.6%
24	21	18:31	66.0	-145.6	287	6.1	11.0	4.38	0.3%
25	21	21:21	72.7	-127.3	316	6.3	8.4	1.43	6.6%
26	21	23:27	69.6	-136.7	159	4.6	10.6	0.96	17.3%
27	21	24:10	69.5	-136.7	164	4.7	10.5	1.31	12.7%
28	21	25:21	69.3	-137.6	162	5.9	10.3	2.97	17.3%
29	21	26:24	66.1	-146.0	222	5.4	11.0	1.58	6.6%

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

**Table 3.** Satellite-derived information corresponding to DC-8 BrO profile locations shown in Fig. 1. Units and abbreviations are as used in Tables 1, 2 and 2. “Ratio<sub>OMI</sub>” means ratio of the OMI-derived tropospheric column to the in-situ column. “Ratio<sub>GOME-2</sub>” is the same but for GOME-2.

No.	Day	SZA [°]	$R_{331\text{nm}}$	$\Delta P_c$ [hPa]	Col. OMI	Col. GOME-2	Ratio OMI	Ratio GOME-2
1	16	60	0.80	19.9	0.98	1.08	0.23	0.25
2	16	61	0.88	46.7	4.12	3.91	1.52	1.44
3	16	63	0.91	41.2	2.99	3.67	0.78	0.96
4	16	61	0.89	44.2	2.35	2.78	1.17	1.39
5	16	56	0.85	23.4	2.15	2.33	0.74	0.80
6	16	55	0.83	49.3	1.97	2.26	0.72	0.82
7	17	59	0.85	15.8	2.70	1.80	1.03	0.79
8	17	60	0.86	21.3	2.47	2.65	1.17	1.26
9	17	67	0.88	31.1	3.10	2.31	0.81	0.60
10	17	70	0.90	46.3	2.21	1.54	0.65	0.45
11	17	74	0.87	82.7	3.11	2.81	1.05	0.95
12	17	77	0.87	146.9	4.69	5.08	1.08	1.17
13	17	74	0.90	39.6	2.85	2.16	0.55	0.42
14	17	75	0.90	32.9	2.81	2.05	1.15	0.84
15	17	78	0.86	33.5	2.48	2.82	1.26	1.43
16	17	80	0.85	38.6	2.32	2.17	1.25	1.16

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

**Table 4a.** Similar to Table 3 but for NOAA WP-3D profiles of BrO, measured on 12 to 15 April 2008.

No.	Day	SZA [°]	$R_{331\text{ nm}}$	$\Delta P_c$ [hPa]	Col. OMI	Col. GOME-2	Ratio OMI	Ratio GOME-2
1	12	57	0.44	11.9	-0.46	–	-0.18	–
2	12	62	0.89	85.9	1.59	–	1.22	–
3	12	65	0.89	80.4	1.24	1.43	0.63	0.73
4	12	67	0.90	89.5	2.00	2.31	1.58	1.82
5	12	69	0.90	80.7	1.58	1.95	0.47	0.58
6	12	72	0.89	70.7	1.04	2.00	6.70	12.94
7	12	77	0.90	76.5	1.23	1.21	0.87	0.86
8	15	62	0.45	155.9	-1.39	-0.42	-2.23	-0.67
9	15	57	0.85	79.7	2.09	2.88	2.85	3.93
10	15	58	0.86	92.9	1.63	2.20	2.01	2.71
11	15	59	0.86	92.7	2.19	2.45	1.28	1.44
12	15	64	0.90	104.3	2.07	3.56	19.74	34.07
13	15	65	0.91	106.1	2.33	3.21	32.63	44.95
14	15	67	0.94	106.7	1.97	2.22	1.24	1.39
15	15	74	0.46	77.8	0.28	-0.58	0.27	-0.55

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

**Table 4b.** Similar to Table 3 but for NOAA WP-3D profiles of BrO, measured on 18 to 21 April 2008.

No.	Day	SZA [°]	$R_{331\text{nm}}$	$\Delta P_c$ [hPa]	Col. OMI	Col. GOME-2	Ratio OMI	Ratio GOME-2
16	18	62	–	–	4.08	2.71	0.94	0.62
17	18	65	–	–	4.20	2.81	4.97	3.32
18	18	67	–	–	3.76	2.14	1.96	1.12
19	19	54	0.61	279.7	0.91	0.33	0.20	0.07
20	19	62	0.78	176.6	1.97	2.55	0.81	1.04
21	19	72	0.83	142.0	4.23	6.00	2.37	3.36
22	19	77	0.79	179.7	1.22	1.80	0.58	0.85
23	19	86	0.72	263.6	0.78	0.95	0.34	0.42
24	21	63	0.47	266.0	0.12	0.25	0.03	0.06
25	21	61	0.83	110.5	3.03	3.54	2.13	2.48
26	21	62	0.80	198.1	3.14	3.29	3.27	3.43
27	21	64	0.80	205.6	3.02	3.45	2.32	2.64
28	21	69	0.79	193.5	2.40	2.87	0.81	0.97
29	21	71	0.47	260.5	0.12	0.29	0.07	0.18

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Satellite-derived  
Arctic tropospheric  
BrO columns

S. Choi et al.



**Fig. 1.** Flight tracks and the locations of BrO profiles measured by instruments on board the (a) NASA DC-8 aircraft during ARCTAS and (b) NOAA WP-3D aircraft during ARCPAC.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.




**Fig. 2.** (a) Collection of DC-8 measurements of BrO (blue dots) and DC-8 composite BrO profile (black line) and (b) DC-8 BrO profiles #7, #10, #11 and #12, (c) similar to (a) but for WP-3D measurements of BrO, (d) WP-3D BrO profiles #5, #22, and #23.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.



**Fig. 3.** BrO weighting functions at 344.6 nm for various solar zenith angles (SZA) and surface albedos for **(a)** clear sky conditions and **(b)** clear and cloudy sky conditions, where the cloudy case is for an optically thick cloud (cloud extinction coefficient = 300).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.




**Fig. 4.** (a) Tropospheric air mass factors (AMFs) at 344.6 nm computed using the DC-8 composite BrO profile for various surface albedos as a function of SZA; (b) similar to (a) but also showing AMFs found assuming all of the BrO is below 500 m (dotted lines).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)




**Satellite-derived Arctic tropospheric BrO columns**

S. Choi et al.




**Fig. 5.** Various parameters and estimated quantities for OMI orbit 20050 at 03:23 UTC on 22 April 2008 highlighting reduced tropospheric BrO over low albedo surfaces (black box) and reduced BrO in the presence of shielding clouds (orange box). See text for more details and discussion.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



**Satellite-derived Arctic tropospheric BrO columns**

S. Choi et al.



**Fig. 6a.** Maps on 17 April 2008 of (top left) OMI total vertical column density (VCD) of BrO, (top right) OMI tropospheric VCD of BrO, (middle left) GOME-2 total VCD of BrO, and (middle right) GOME-2 tropospheric VCD of BrO column. Negative values are included as dark blue. The flight track of the DC-8 on 17 April 2008 is indicated by black lines; aircraft profile locations where BrO was measured are marked with black diamonds. (Bottom) Geographic regions of interest described in the paper.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.



**Fig. 6b.** Time series plots for NASA ARCTAS DC-8 flight of 17 April 2008 with line colors corresponding to the same color y axis. The top panel shows various estimates of VCD of BrO along the DC-8 flight track, including OMI total column, the modeled stratospheric column, and tropospheric columns based on aircraft in-situ measurements (black dots) as well as our analysis of OMI retrievals (red squares). The error bars of the tropospheric columns are described in the text. The two middle panels show various aircraft in-situ measurements along with the flight altitude. The bottom panel shows OMI reflectivity at 331 nm and cloud retrieved parameters from OMI and MODIS, as described in the text. The measurement accuracy of DC-8 BrO measurements is  $\pm 40\%$ . Soluble bromide is not plotted when the mixing ratio is below the detection limit (1–6 pptv).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Satellite-derived  
Arctic tropospheric  
BrO columns**

S. Choi et al.



**Fig. 7a.** Similar to Fig. 6a but for the DC-8 flight on 5 April 2008.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.




**Fig. 7b.** Similar to Fig. 6b but for the DC-8 flight on 5 April 2008. Aircraft in situ BrO measurements are not available for this flight.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.



**Fig. 8a.** Similar to Fig. 6a but for the DC-8 flight on 8 April 2008. GOME-2 data are not available for this flight. The location of severe ozone depletion and high active bromine is marked with a black square.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Satellite-derived  
Arctic tropospheric  
BrO columns

S. Choi et al.




**Fig. 8b.** Similar to Fig. 7b but for the DC-8 flight on 8 April 2008.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



**Satellite-derived  
Arctic tropospheric  
BrO columns**

S. Choi et al.



**Fig. 9a.** Similar to Fig. 6a but for the WP-3D flight of 19 April 2008.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion





**Satellite-derived  
Arctic tropospheric  
BrO columns**

S. Choi et al.



**Fig. 10a.** Similar to Fig. 6a but for the WP-3D flight of 21 April 2008.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 10b.** Similar to Fig. 9b but for the WP-3D flight on 21 April 2008, with tropopause pressure added.

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Satellite-derived  
Arctic tropospheric  
BrO columns

S. Choi et al.



**Fig. 11.** OMI tropospheric BrO VCD for different orbits (UTC time as indicated) from 16 to 18 April 2008.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



Satellite-derived  
Arctic tropospheric  
BrO columns

S. Choi et al.



**Fig. 12.** (a) OMI tropospheric BrO VCD; (b) OMI total ozone column; (c) terrain minus cloud pressure for OMI orbit 19976 on 17 April 2008.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version

Interactive Discussion







**Fig. 13.** First column: OMI tropospheric BrO VCD; Second column: MERRA sea level pressure; third column: MERRA wind speed at 2 m altitude; fourth column: MERRA planetary boundary layer height; fifth column: aircraft in-situ BrO (red) and active bromine mixing ratio profiles (blue) and temperature profile (purple) corresponding to OMI orbit 19979 on 17 April 2008 (top row), orbit 19990 on 18 April 2008 (middle row), and orbit 20018 on 19 April 2008 (bottom row).

## Satellite-derived Arctic tropospheric BrO columns

S. Choi et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

