

Abstract

Detailed investigations of the chemical and microphysical properties of atmospheric aerosol particles were performed at the puy-de-Dôme (pdD) research station (1465 m) in autumn (September and October 2008), winter (February and March 2009), and summer (June 2010) using a Time-of-Flight Aerosol Mass Spectrometer. Over the three campaigns, the average mass concentrations of the non-refractory submicron particles ranged from $10 \mu\text{g m}^{-3}$ up to $27 \mu\text{g m}^{-3}$. Highest nitrate and ammonium mass concentrations were measured during the winter and during periods when marine modified airmasses were arriving at the site, whereas highest concentrations of organic particles were measured during the summer and during periods when continental air-masses arrived at the site. The measurements reported in this paper show that atmospheric particle composition is strongly influenced by both the season and the origin of the airmass. The total organic mass spectra were analysed using positive matrix factorisation to separate individual organic components contributing to the overall organic particle mass concentrations. These organic components include a low volatility oxygenated organic aerosol particle (LV-OOA) and a semi-volatile organic aerosol particle (SV-OOA). Correlations of the LV-OOA components with fragments of m/z 60 and m/z 73 (mass spectral markers of wood burning) during the winter campaign suggest that wintertime LV-OOA are related to aged biomass burning emissions, whereas organic aerosol particles measured during the summer are likely linked to biogenic sources. Equivalent potential temperature calculations, gas-phase, and LIDAR measurements define whether the research site is in the planetary boundary layer (PBL) or in the free troposphere (FT)/residual layer (RL). We observe that SV-OOA and nitrate particles are associated with air masses arriving from the PBL where as particle composition measured from RL/FT airmasses contain high mass fractions of sulphate and LV-OOA. This study provides unique insights into the effects of season and airmass variability on regional aerosol particles measured at an elevated site.

Seasonal characterisation of aerosol particles at the puy de Dôme site

E. J. Freney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Aerosol particles play important roles in affecting the radiative balance of the earth and air quality. For this reason, detailed and long-term analysis of the physicochemical properties of atmospheric aerosol particles is crucial to understand their processes and lifetime in the atmosphere. A network of both high (>2000 m) and low (<1000 m) elevation measurement sites have provided the opportunity to study the long-term variability of aerosol particle properties in Europe. These urban and rural research sites are most commonly equipped with size distribution measurements and filters for offline particle composition measurements (Nyeki et al., 1998; Weingartner et al., 1999; Gelencser et al., 2007; Venzac et al., 2009).

Results arising from observations made at these research sites offer information on the seasonal and chemical variability of aerosol particles measured in different atmospheric environments. However, the coarse time and size resolution of these measurements limits our knowledge on the physical and chemical properties of aerosol particles. In the last decade, online aerosol mass spectrometers have been used to provide highly time resolved quantitative data on submicron inorganic and organic aerosol species. There have recently been several reports summarizing aerosol mass spectrometry measurements made in the Northern Hemisphere (Zhang et al., 2005; Jimenez et al., 2009; Ng et al., 2010), showing that aged organic compounds are the largest contributor to the submicron particle mass. In Europe, Lanz et al. (2010) overviewed the results of several campaigns carried out in the alpine region over a number of different seasons, providing detailed information on the aerosol particle chemical and physical properties at both high- and low-altitude sites and during several different seasons of the year. Similar to previous observations they showed that organic compounds were the most abundant species in the submicron particle fraction, with biomass burning organic aerosol particles being important during the winter and low-volatility organic aerosol (LV-OOA) dominating during the spring and summer. Although these studies provide information on the background composition during dif-

Seasonal characterisation of aerosol particles at the pyu de Dôme site

E. J. Freney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ferent seasons they are unable to provide information on the seasonal variability at the same site.

The puy-de-Dôme (pdD) research site is classified as a mid-altitude site (1465 m) where at night the research site is commonly in the residual layer (RL) or in the free troposphere (FT), but during the day it is exposed to airmasses arriving from the planetary boundary layer (PBL) (Venzac et al., 2009). Therefore, at the pdD site, measured particle composition during the day can be representative of boundary layer air masses, whereas at night particle composition is representative of regional air masses. These features make the pdD research site an ideal place to characterize both regional and local boundary layer pollution conditions through examining the diurnal variability of aerosol particle properties. A Time-of-Flight Aerosol Mass Spectrometer (ToF-AMS) was deployed at the pdD site during three different seasons of the year as part of the European monitoring and evaluation program (EMEP). The aim of EMEP is to characterize the physical and chemical properties of the aerosol population in Europe through carrying out simultaneous aerosol composition measurements at several different research sites around Europe. In this paper, we report the chemical composition, concentrations, and temporal variations of the non-refractory particle components at the pdD site. The focus of this study was to characterize the seasonal variability in the chemistry of both long-range transported and less aged PBL airmasses with high-time resolution, and thereby offering insights into the sources and properties of aerosol particles arriving at an elevated background site.

2 Experimental

2.1 Sampling site

The pdD research station is located in central France (45°46 N 2°57 E). The station is surrounded by agricultural and forest areas. The city of Clermont-Ferrand (150 000 inhabitants) is located 16 km east of the station at 396 m. A detailed description of

Seasonal characterisation of aerosol particles at the puy de Dôme site

E. J. Freney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Seasonal characterisation of aerosol particles at the py de Dôme site

E. J. Freney et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the site is provided by Venzac et al. (2009). The pdD research site contains a number of different instruments capable of resolving the physical, chemical, optical, and hygroscopic properties of aerosol particles. The site is in cloud approximately 50 % of the time between November and March. For this reason the site is equipped with two different types of stainless steel inlets; a whole air inlet capable of sampling cloud droplets together with interstitial (non-activated) particles (whole air inlet (WAI)), and a second inlet capable of sampling only interstitial particles. The WAI inlet has a 50 % cut-off at 30 μm and the interstitial inlet has a 50 % cut-off at 1.6 μm . These inlets do not allow for isokinetic sampling. Meteorological parameters (including wind direction, wind speed, relative humidity (RH), pressure, and temperature), aerosol physical and chemical properties (particle size, black carbon mass), and gas-phase species (SO_2 , CO , CO_2 , O_3 , NO , and NO_2) are measured continuously throughout the year. Particle size distributions between 50 and 300 nm were measured using a scanning mobility particle sizer (SMPS), and black carbon (BC) mass concentration was measured using a Multi Angle Absorption Photometer (MAAP 5012). The SMPS and MAAP were sampling behind the WAI, whereas the AMS was situated behind the interstitial inlet.

2.2 Backward trajectories

In order to determine the transport pathways of the aerosol particles prior to arriving at the pdD site (1465 m), the Hybrid Single Particle Lagrangian Trajectory (HYSPPLIT) model was used. Seventy-two hour backward trajectories were computed for every hour for each day of the campaign. Trajectories were classified according to their predominant transport direction prior to measurement as either continental (C), marine (M), marine modified (Mod), and Mediterranean (Med) (Fig. S1). If air mass trajectories show a combination of transport direction during the 72-h period the air mass is classified as mixed.

2.3 LIDAR and temperature profile measurements

Using temperature, pressure, and relative humidity data from measurements made simultaneously at the Observatoire de Physique du Globe de Clermont-Ferrand (OPGC) ground based research station (425 m a.s.l., 11 km east from the pdD Research Station) and at the pdD site, the differential equivalent potential temperature (θ_e) was calculated as a function of elevation using the Boltons approximation (Bolton, 1980). The calculated (θ_e) is used to identify if the measurements at the pdD site were performed in the PBL, FT, or RL. Fig. 1, shows the correlation of θ_e at the OPGC with those at the pdD site coloured by hour of day for the three different campaigns. The largest difference in θ_e between the two sites occurs during the evening hours. This difference is most pronounced during the winter campaign when the pdD is most often in the free troposphere. During the summer campaign, LIDAR measurements were also available (Fig. 1d). The spatial raw resolution of the detected signals is 15 m. The LIDAR instrument is positioned on the roof of the OPGC. There was no cloud cover over the OPGC or pdD during the LIDAR measurements. The boundary layer height was determined by a wavelength covariance technique (Brooks, 2003), and exhibit a weak diurnal variation in boundary layer height. However, these measurements show that during the summer, the pdD site was always sampling in the PBL or RL.

2.3.1 Time-of-flight aerosol mass spectrometer

The chemical composition and mass concentration of the non-refractory submicron particulate matter (NR-PM₁) was measured with an Aerodyne ToF-AMS (Drewnick et al., 2005; Canagaratna et al., 2007). In order to extract chemically resolved mass concentrations of individual aerosol species, the AMS raw data are evaluated with standard assumptions as described by Allan et al. (2004). Filter measurements, in which the AMS sampled filtered air, were made approximately every 48 hrs. The resolved mass concentrations include ammonium, sulphate, nitrate, organic, and chloride species. For the organic mass spectra additional analysis was carried out using posi-

Seasonal characterisation of aerosol particles at the puy de Dôme site

E. J. Freney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tive matrix factorisation (PMF). The ToF-AMS was sampling behind an interstitial inlet. For this reason, when the site was advected by clouds, the ToF-AMS sampled only those particles not activated into clouds. For the purpose of this study only periods when the site was not in cloud are taken into account.

5 The calculation of the mass concentrations from the ToF-AMS mass spectra requires that a collection efficiency (CE) be applied to the data. The CE is defined as the fraction of the particle mass that is measured by the ToF-AMS, relative to what would have been measured if all particles were spherical and particle bounce was negligible. Recently, it has been demonstrated that particles containing high ammonium nitrate concentra-
10 tions are more efficiently sampled by the AMS than other inorganic and organic species (Middlebrook et al., 2011). Ammonium nitrate particles tend to be spherical and retain water even at very low RH values and are therefore efficiently focused by the particle beam and are not subjected to particle losses from particle bounce (Middlebrook et al., 2011). Middlebrook et al. (2011) illustrated, using a series of equations, how
15 a composition dependent CE can be applied to particle mass concentrations. In this study, a similar composition dependent CE is applied to the data. When ammonium nitrate mass concentrations contribute less than 0.25 to the total aerosol particle mass a CE of 0.5 is used. As the fraction of ammonium nitrate increases to a maximum of 0.78 (Pure NH_4NO_3) the applied CE approaches 1. Since the majority of aerosol
20 particles arriving at the pdD site are thought to be internally mixed, this composition dependent CE was applied to all species sampled by the AMS. In order to assess how accurate our chosen CE was at quantifying the mass of aerosol particles in the atmosphere, we compared the total volume of aerosol particles sampled by the AMS and BC with that sampled by a SMPS. The AMS volume was calculated by dividing
25 the mass concentrations of organic and inorganic species by their respective densities ($1.72 \mu\text{g cm}^{-3}$ for SO_4 , NH_4 , NO_3 , and $1.2 \mu\text{g cm}^{-3}$ for organics). During this study, the SMPS was sampling behind a WAI inlet, where as the AMS was sampling behind the interstitial inlet and hence the AMS did not sample particles with the same efficiency as the SMPS. In addition the SMPS was not sampling in the same building at the time of

Seasonal characterisation of aerosol particles at the py de Dôme site

E. J. Freney et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the respective campaigns. The resulting slopes and r^2 for each comparison were 0.76 and 44 % (SMPS:AMS = 0.95), 0.51 and 71 % (SMPS:AMS = 1.61), and 0.68 and 44 % (SMPS:AMS = 0.85) for the autumn, winter, and summer, respectively (Fig. S2). Periods with largest discrepancies were observed during episodes when increased particle numbers were sampled by the SMPS. The size distributions of all particles during all seasons arriving at pdD are characteristic of aged particles showing a major accumulation mode peaking at an aerodynamic diameter of 600 nm (Supplement Fig. S3). This monomodal peak is expected since the site is not exposed to fresh emissions but rather to particles transported long distances and thus considerably aged.

2.4 Positive matrix factorisation

The unit mass resolution organic mass spectra from the ToF-AMS instrument are analysed using positive matrix factorisation (PMF). PMF is a statistical technique in which the organic mass spectral time series (X) are represented as a linear combination of a set of characteristic factor profiles (F) and their time-dependant intensities (G) (Paatero and Tapper, 1994; Paatero, 1997). PMF is therefore described by the matrix equation $X = GF + E$, where E is the residual matrix, defined as the difference between the data matrix X and the fitted solution GF . The PMF analysis was conducted using PMF2 software package (P. Paatero, University of Helsinki, Finland), together with the PMF analysis and evaluation tool (Ulbrich et al., 2009). The error matrix calculated in the Squirrel software (version 1.51) was modified following the recommendations of Ulbrich et al. (2009) and references therein. A minimum error of 1 ion was applied throughout the X matrix, and the organic peaks calculated as the fraction of the CO_2^+ ion (m/z 44, 18, 17, and 16) were down weighted as described in Ulbrich et al. (2009). Matrix rotations were explored by varying the f peak parameter from -3 to $+3$. The f peak parameter allows the user to explore different PMF solutions that give the same fit to the data (Ulbrich et al., 2009). The number of PMF factors and f peak values were determined by minimising the scaled residuals and factor correlations with external tracers (includ-

Seasonal characterisation of aerosol particles at the pyu de Dôme site

E. J. Freney et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ing CO, O₃, NO_x, SO₂, NO₃⁻, and SO₄²⁻) and comparing the component mass spectra with reference mass spectra available on the AMS mass spectral database (Lanz et al., 2007; Ulbrich et al., 2009). An optimum peak value of 0 was applied for each dataset. Correlations of the resolved mass spectral profiles with reference mass spectra and correlations of the organic time series with the time series of other particulate and gas phase species are supplied in the Supplement (Table S1).

3 Results and discussion

3.1 Characterisation of the submicron particle composition

A time series of sulphate (SO₄), nitrate (NO₃), ammonium (NH₄), organics (Org), chloride (Cl), and black carbon BC-bearing particles is shown in Fig. 2. Meteorological parameters are listed in Table 1. Average values for each aerosol particle species measured by the AMS, during clear sky periods are listed in Table 2. The weather conditions during this study were characterized by high RH (86 ± 12 %) and temperatures ranging from 0.1 °C to 18 °C (average = 9 ± 4 °C), -8 °C to 13 °C (average = 1 ± 4 °C), and 2 °C to 25 °C (average = 14 ± 5 °C), during the autumn, winter, and summer campaigns, respectively. Highest average aerosol mass concentrations were observed during the summer campaign (27 µg m⁻³) and lowest during the winter campaign (10 µg m⁻³), in agreement with seasonal variability observed at the site for particle number concentrations (Venzac et al., 2009) and particle mass concentrations measured at other high-altitude stations in Europe (Kasper and Puxbaum, 1998; Weingartner et al., 1999).

On the basis of campaign averages and in the order of the campaigns (autumn 2008, winter 2009, summer 2010), the organic mass contributed 32 % (2.52 µg m⁻³), 23 % (1.24 µg m⁻³), and 56 % (15.59 µg m⁻³) to the NR-PM₁. Sulphate particles contributed between 30% (2.4 µg m⁻³), 22 % (1.28 µg m⁻³), and 20 % (5.45 µg m⁻³), and nitrate contributed 15% (1.14 µg m⁻³), 31 % (1.71 µg m⁻³), and 9 % (2.33 µg m⁻³). Ammonium contributed 18 % (1.36 µg m⁻³), 20 % (1.08 µg m⁻³), and 13% (3.69 µg m⁻³). Chloride

Seasonal characterisation of aerosol particles at the py de Dôme site

E. J. Freney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



for NO₃ bearing particles because of the high contributions of organic aerosol particles to the submicron mass. This might indicate different origins like biogenic precursors for secondary organic aerosols versus anthropogenic precursors for the inorganic components.

3.2 PMF analysis of organic fraction

In order to fully characterize the organic fraction collected at pdD, PMF analysis was performed separately on each of the datasets, however the resolved PMF factors were similar during all seasons allowing for a direct comparison of the datasets. Two prominent types of organic species were resolved for each of the campaigns. Mass spectra and time series of each of the different components resolved during each campaign are illustrated in Figs. 3 and S4, respectively. A full list of correlations of all PMF components with reference mass spectra and with gas and particle phase species is available in the Supplement (Table S1). The average mass concentrations and fractional contributions of the different organic components are shown in Table 2. Only oxidised organic aerosol (OOA) particles were resolved from ToF-AMS measurements made at the pdD research site. This OOA could be separated into two different types of OOA; OOA1 and OOA2. OOA1 contributed approximately $65 \pm 5\%$ to the total organic aerosol during all campaigns and had a similar mass spectral signature to OOA measured in Europe and in the United States (Ng et al., 2010, 2011) (Fig. 3). The mass spectral properties of this OOA are defined by having a dominant peak at m/z 44 and are generally associated with organic aerosol particles that are aged and have a low volatility. They are therefore termed a low-volatility oxidised organic aerosol particle (LV-OOA).

For the summer campaign the time series of the LV-OOA component showed strongest correlations with that of O₃ ($r^2 = 0.52$), CO ($r^2 = 0.33$), and BC ($r^2 = 0.44$) (Supplement Table S1). Ozone has been shown to be a good marker for the production of secondary organic aerosols in some campaigns (Cabada et al., 2004), but other studies show that this is not always the case (Zhang et al., 2005). In our three campaigns the strongest correlation of LV-OOA with ozone is observed for the summer

Seasonal characterisation of aerosol particles at the py de Dôme site

E. J. Freney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



when high organic mass concentrations were arriving at the pdD site, weaker correlations were observed during autumn and winter ($r^2 < 0.01$ and 0.10 , respectively) possibly suggesting a change in the precursor sources to the formation of LV-OOA depending on the season.

5 Results from the CARBOSOL (study of the present and retrospective state of the organic versus inorganic aerosol over Europe) project (Gelencser et al., 2007), involving organic, levoglucosan, and black carbon measurements from several different European sites show that during the winter months there is a high contribution of biomass burning to secondary organic aerosol particles (Gelencser et al., 2007; Puxbaum et al.,
10 2007; Lanz et al., 2008). We show a high contribution of secondary organic aerosol particles similar to that of fulvic acid during all campaigns, but during the winter campaign the LV-OOA mass spectra also contains marker peaks of wood burning ((Alfarra et al., 2007) (m/z 60 (contributing 0.9% to the total mass spectra) and m/z 73 (contributing 0.7% to the total mass spectra)) and correlates strongly with the time series
15 of m/z 73 and m/z 60. These peak signals suggest that the LV-OOA measured during the winter are possibly related to aged wood burning particles and due to secondary organic aerosols from wood burning emissions which can also contain significant contributions of m/z 60 (Heringa et al., 2011). The LV-OOA component had average mass concentrations of $\sim 0.4 \mu\text{g m}^{-3}$ for most of the winter campaign, but during an isolated
20 4-h period on 26 February, its mass concentrations increased up to $2 \mu\text{g m}^{-3}$. Organic particle mass concentrations measured during this period showed little correlation with gas phase species or other inorganic compounds suggesting that this is an aged aerosol particle type. The residuals measured from the PMF analysis during this period of the winter campaign are higher than during the rest of the campaign.
25 However, a pure primary wood burning factor could not be extracted even when PMF analysis was performed only on this isolated period.

The second type of organic aerosol resolved during each of the campaigns closely resembled a less aged oxidized organic aerosol. The less oxidized organic aerosol particle measured had dominant mass spectral peaks at m/z 43, 44, 55, and 57, and

Seasonal characterisation of aerosol particles at the pyu de Dôme site

E. J. Freney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

fraction of m/z Org43 to the total organics (f43) (Fig. 4). All of our points and PMF components from each campaign cluster within the well-defined triangular regions as outlined by Ng et al. (2010). The calculated f44 and f43 from each of the PMF components for each campaign is also plotted on the graph. During the autumn, winter, and summer campaign the f44 for LV-OOA organics ranges from 0.19 to 0.24 which is similar to values reported by Ng et al. (2010) (0.10 to 0.25). SV-OOA during the autumn, winter, and summer ranged from 0.02 to 0.14.

From Fig. 4 it is apparent that aerosol particles measured during the summer are situated on the right side of the graph when it is likely to have more biogenic activity whereas those measured during the winter, when we expect to have the highest contribution from biomass burning, are located on the left side of the triangle. Organic mass spectra measured during the autumn are in the middle. Laboratory studies on secondary organic aerosol formation from biogenic precursors show the resulting f44, f43 on the right side of the Ng triangle (Jimenez et al., 2009; Ng et al., 2010), whereas those from biomass burning emissions tend to be situated on the left side of the triangle (Heringa et al., 2011). These observations indicate that the formation of organic aerosol particle is strongly influenced by different processes and precursor sources occurring at different times of the year. The seasonal segregation shown in our measurements of the organic mass spectra suggests that the triangle is not only descriptive for the aging of organic aerosol particles but to some extent can also describe the source of the secondary organic aerosol particles.

3.3 Diurnal variations and air mass source

Backward trajectory calculations show that during the autumn and summer campaigns air masses at pdD arrived mainly from the continental sector (39 % Autumn and 80 % Summer), whereas marine airmasses (Marine (29 %)+Marine modified (43 %)) contributed 72 % during the winter. Mediterranean air masses were only measured during the autumn campaign where they contributed 13 % to the total airmasses arriving during the sampling period. This air mass pattern is typical of that observed at the pdD station

Seasonal characterisation of aerosol particles at the py de Dôme site

E. J. Freney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Seasonal
characterisation of
aerosol particles at
the puy de Dôme site**

E. J. Freney et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

(Venzac et al., 2009). The fractional contribution of each species is averaged during continental, marine, and marine modified air masses, for a given season (Fig. 5). High aerosol mass concentrations are observed during continental air masses ($3.3 \mu\text{g m}^{-3}$), while lowest concentrations are measured in marine and marine modified air masses ($1.2 \mu\text{g m}^{-3}$ and $2.0 \mu\text{g m}^{-3}$ respectively), in agreement with Bourcier et al. (2011). Figure 5 illustrates that within a given air mass type the aerosol composition changes significantly with season. Hence, seasonal variability is not solely driven by changes in air mass. Highest fractions of organic and nitrate aerosol particles are observed for continental and marine modified airmasses, organic and sulphate particles are dominant in marine airmasses, particularly during the winter campaign. Despite variability, the organic aerosol particles contribute a large fraction to the overall particle mass whatever the air mass type (20 to 51 %) or season (23 to 50 %).

One of the conclusions of Lanz et al. (2010) was that the time of year and location of the sampling site could largely explain the variability in chemical composition for different campaigns and was found to be a better description of the aerosol composition than the type of site (rural vs urban). Here we show that although the time of year has an important influence on the aerosol particle composition, the source of the airmasses still has significant effects on the submicron particle composition.

In earlier studies, the seasonal variability of aerosol properties were shown to be linked to the frequency at which the pdD site was located in the PBL or FT/RL Venzac et al. (2009). The pdD site is mostly in the PBL during the day and FT/RL during the night time hours, particularly during the winter. Fig. 6 shows the diurnal variability of the mass concentration of the inorganic and organic species during each campaign. Particle number concentration (N_{tot}) measurements made at the site over several years show that in general the typical diurnal variability is characterized by an increase in N_{tot} between 12:00 and 14:00 UTC (Venzac et al., 2009). In this study we observe that the diurnal variability in N_{tot} was strongest during the autumn and winter, and weakest during the summer campaign (Table 1). AMS measurements show strongest diurnal variability in particle mass concentrations during the winter (Fig. 6), when the pdD

site is expected to spend the majority of the time in the free troposphere and only be advected by boundary layer airmasses for several hours during the day.

During autumn and winter we observe increases in the mass concentration of all species during the day and decreases at night. However, the relative contributions of all species do not change. For the summer campaign, mass concentrations of Org and SO_4^{2-} increase while those of NO_3 decrease during the daytime hours (Fig. 6). These changes are presumably a result of photochemical secondary aerosol production of organics and sulphates during the day, rather than changes from BL to FT. Summer night time increases in NO_3 are likely related to the lower mass concentrations of sulphate at night, allowing excess NH_3 to react with nitrate (Seinfeld and Pandis, 1998), this process is also favoured by the lower temperatures and higher RH during the night time. A closer look at the different organic components determined from PMF analysis show that the organic components day time increases are evenly attributed to both LV-OOA and SV-OOA. During the summer campaign the diurnal temperature difference was approximately 5°C , however during autumn and winter it was less than 2°C .

The chemical composition of aerosol particles were also examined relative to the age of the air masses. Since NO_y measurements were not available at the pdD site during these measurements, the ratio of CO/NO_x was used as a tracer for major pollution sources and for photochemical processing. According to Morgan et al. (2010), a value between 10 and 50 for the CO/NO_x ratio, is classified as near-source conditions. During all measurement campaigns the CO/NO_x ratio range from 7 to greater than 400 with an average value of 93 ± 43 , confirming that most of the airmasses arriving at the pdD are representative of aged regional emissions. Plotting the fraction of SO_4 , NO_3 , LV-OOA, and SV-OOA to the total aerosol mass concentration against the CO/NO_x ratio (Fig. S5), shows that LV-OOA is associated with aged air masses while highest nitrate concentrations favour low CO/NO_x ratios arising presumably from boundary layer sources.

The age of the air masses are likely connected to the BL/FT segregation. Hence, on the basis of both thermodynamical and chemical indicators we used a combination

Seasonal characterisation of aerosol particles at the py de Dôme site

E. J. Freney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

of θ_e measurements and gas-phase measurements to characterize the properties of aerosol particles likely associated with BL/FT air masses. Plotting the difference in θ_e ($\Delta\theta_e$) against the CO/NO_x ratio, we observe that there is not a strong linear relationship between θ_e and the CO/NO_x ratio. However, lowest values of θ_e are related to low CO/NO_x ratios (Fig. 7) indicating that, overall, the thermodynamical indicator of BL/FT air masses, θ_e , is coherent with the chemical aging indicator, CO/NO_x. These plots suggest that boundary layer air masses can be characterized by low θ_e and low CO/NO_x ratios, whereas airmasses arriving from the free troposphere or residual layer are characterized by high θ_e and high CO/NO_x ratios (Fig. 7). If these plots are coloured by particle composition, we observe that high concentrations of NO₃ and SV-OOA are most often related to airmasses mostly associated to BL air, whereas highest concentrations of LV-OOA are related to air masses associated to residual layer or free tropospheric air.

4 Conclusions

This is the first time that the ToF-AMS has been deployed during three seasons at an altitude site, providing a unique opportunity to study the seasonal variability of regional airmasses. Through investigating the chemical composition of submicron aerosol composition during autumn 2008, winter 2009, and summer 2010, we show that although the time of year has a strong influence on the chemical properties of the air mass, we also observe that the source (FT/BL) and transport direction of airmasses influence the properties of aerosol particle. Examining particle mass concentrations together with gas-phase measurements allows us to associate nitrate and less aged organics with airmasses arriving from the BL whereas airmasses arriving from the FT/RL are correlated with aged organics and sulphate particles. Detailed analysis of the organic component showed that two different types of organic aerosol particles were measured at the site: a LV-OOA particle and a SV-OOA particle. The lack of a clear sign of primary contributions to organic mass is similar as previously found in Crete (Hildebrandt

Seasonal characterisation of aerosol particles at the puy de Dôme site

E. J. Freney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



et al., 2010). Small changes in mass spectral properties between seasons as well as correlations with different gas and particle phase species suggest that there are different precursor sources contributing to the formation of secondary organic aerosols during different seasons of the year. From the analysis of the resolved organic mass spectra we conclude that biomass burning emissions contribute to the formation of secondary organic aerosols during the winter and biogenic emissions contribute during the summer. Considering seasonal influences on the formation of SOA is important in modelling and quantifying organic mass concentrations in the atmosphere. These measurements show that through a combination of continuous aerosol particle measurements, we can understand the influence of different air mass sources (FT/BL) and transport directions on the aging and formation of the aerosol particle distribution, particularly the organic fraction.

Supplementary material related to this article is available online at:
[http://www.atmos-chem-phys-discuss.net/11/27139/2011/
acpd-11-27139-2011-supplement.pdf](http://www.atmos-chem-phys-discuss.net/11/27139/2011/acpd-11-27139-2011-supplement.pdf).

Acknowledgements. We thank Manjula Canagaratna for helpful discussions. This work was performed in the framework of the Research Infrastructure Action under the FP6 Structuring the European Research Area Programme, EUSAAR Contract No. RIII3-CT-2006-026140. The work has been partly funded by FP6 Integrated project EUCAARI, Contract No. 36833. E. J. Freney is funded by an AXA postdoctoral fellowship (ACI-AMS) and by a Marie-Curie international reintegration grant.



The publication of this article is financed by CNRS-INSU.
27156

ACPD

11, 27139–27170, 2011

Seasonal characterisation of aerosol particles at the py de Dôme site

E. J. Freney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

- Alfarra, M. R., Prevot, A. S. H., Szidat, S., Sandradewi, J., Weimer, S., Lanz, V. A., Schreiber, D., Mohr, M., and Baltensperger, U.: Identification of the mass spectral signature of organic aerosols from wood burning emissions, *Environ. Sci. Technol.*, 41, 5770–5777, doi:10.1021/es062289b, 2007. 27150
- Allan, J. D., Delia, A., Coe, H., Bower, K., Alfarra, M., Jimenez, J., Middlebrook, A., Drewnick, F., Onasch, T., Canagaratna, M., Jayne, J., and Worsnop, D.: A generalised method for the extraction of chemically resolved mass spectra from aerodyne aerosol mass spectrometer data, *J. Aerosol Sci.*, 35, 909–922, doi:10.1016/j.jaerosci.2004.02.007, 2004. 27144
- Bahreini, R., Keywood, M., Ng, N., Varutbangkul, V., Gao, S., Flagan, R., Seinfeld, J., Worsnop, D., and Jimenez, J.: Measurements of secondary organic aerosol from oxidation of cycloalkenes, terpenes, and m-xylene using an Aerodyne aerosol mass spectrometer, *Environ. Sci. Technol.*, 39, 5674–5688, doi:10.1021/es048061a, 2005. 27151
- Bolton, D.: The computation of equivalent potential temperature, *Mon. Weather Rev.*, 108, 1046–1053, doi:10.1175/1520-0493(1980)108<1046:TCOEPT>2.0.CO;2, 1980. 27144
- Bourcier, L., Sellegri, K., Jaffrezo, J.-L., Chausse, P., Pichon, J. M., Aymoz, G., Michaud, V., and Laj, P.: Seasonal variation of aerosol chemistry at the puy de Dome station (1465 m a.s.l), France, in preparation, 2011. 27153
- Brooks, I.: Finding boundary layer top: Application of a wavelet covariance transform to lidar backscatter profiles, *J. Atmos. Ocean. Tech.*, 20, 1092–1105, doi:10.1175/1520-0426(2003)020<1092:FBLTAO>2.0.CO;2, 2003. 27144
- Cabada, J., Pandis, S., Subramanian, R., Robinson, A., Polidori, A., and Turpin, B.: Estimating the secondary organic aerosol contribution to PM_{2.5} using the EC tracer method, *Aerosol Sci. Tech.*, 38, 140–155, doi:10.1080/02786820390229084, 2004. 27149
- Canagaratna, M. R., Jayne, J. T., Jimenez, J. L., Allan, J. D., Alfarra, M. R., Zhang, Q., Onasch, T. B., Drewnick, F., Coe, H., Middlebrook, A., Delia, A., Williams, L. R., Trimborn, A. M., Northway, M. J., DeCarlo, P. F., Kolb, C. E., Davidovits, P., and Worsnop, D. R.: Chemical and microphysical characterization of ambient aerosols with the aerodyne aerosol mass spectrometer, *Mass Spectrom. Rev.*, 26, 185–222, doi:10.1002/mas.20115, 2007. 27144
- Drewnick, F., Hings, S., DeCarlo, P., Jayne, J., Gonin, M., Fuhrer, K., Weimer, S., Jimenez, J., Demerjian, K., Borrmann, S., and Worsnop, D.: A new time-of-flight aerosol mass spectrometer (ToF-AMS) – Instrument description and first field deployment, *Aerosol Sci. Tech.*, 39,

Seasonal characterisation of aerosol particles at the puy de Dôme site

E. J. Freney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



637–658, doi:10.1080/02786820500182040, 2005. 27144

Gelencser, A., May, B., Simpson, D., Sanchez-Ochoa, A., Kasper-Giebl, A., Puxbaum, H., Caseiro, A., Pio, C., and Legrand, M.: Source apportionment of PM_{2.5} organic aerosol over Europe: Primary/secondary, natural/anthropogenic, and fossil/biogenic origin, *J. Geophys. Res.-Atmos.*, 112, D23S04, doi:10.1029/2006JD008094, 2007. 27141, 27150

Heringa, M. F., DeCarlo, P. F., Chirico, R., Tritscher, T., Dommen, J., Weingartner, E., Richter, R., Wehrle, G., Prvt, A. S. H., and Baltensperger, U.: Investigations of primary and secondary particulate matter of different wood combustion appliances with a high-resolution time-of-flight aerosol mass spectrometer, *Atmos. Chem. Phys.*, 11, 5945–5957, doi:10.5194/acp-11-5945-2011, 2011. 27150, 27152

Hildebrandt, L., Engelhart, G. J., Mohr, C., Kostenidou, E., Lanz, V. A., Bougiatioti, A., DeCarlo, P. F., Prevot, A. S. H., Baltensperger, U., Mihalopoulos, N., Donahue, N. M., and Pandis, S. N.: Aged organic aerosol in the Eastern Mediterranean: the Finokalia Aerosol Measurement Experiment 2008, *Atmos. Chem. Phys.*, 10, 4167–4186, doi:10.5194/acp-10-4167-2010, 2010. 27155

Hock, N., Schneider, J., Borrmann, S., Römpp, A., Moortgat, G., Franze, T., Schauer, C., Pöschl, U., Plass-Dülmer, C., and Berresheim, H.: Rural continental aerosol properties and processes observed during the Hohenpeissenberg Aerosol Characterization Experiment (HAZE2002), *Atmos. Chem. Phys.*, 8, 603–623, doi:10.5194/acp-8-603-2008, 2008. 27148

Hueglin, C., Gehrig, R., Baltensperger, U., Gysel, M., Monn, C., and Vonmont, H.: Chemical characterisation of PM_{2.5}, PM₁₀ and coarse particles at urban, near-city and rural sites in Switzerland, *Atmos. Environ.*, 39, 637–651, doi:10.1016/j.atmosenv.2004.10.027, 2009. 27148

Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dunlea, E. J., Huffman, J. A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of Organic

Seasonal characterisation of aerosol particles at the pyu de Dôme site

E. J. Freney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Seasonal
characterisation of
aerosol particles at
the puy de Dôme site**

E. J. Freney et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Real-Time Methods for Estimating Organic Component Mass Concentrations from Aerosol Mass Spectrometer Data, *Environ. Sci. Technol.*, 45, 910–916, doi:10.1021/es102951k, <http://pubs.acs.org/doi/abs/10.1021/es102951k>, 2011. 27149, 27151, 27166

5 Nyeki, S., Li, F., Weingartner, E., Streit, N., Colbeck, I., Gaggeler, H., and Baltensperger, U.: The background aerosol size distribution in the free troposphere: An analysis of the annual cycle at a high-alpine site, *J. Geophys. Res.-Atmos.*, 103, 31749–31761, doi:10.1029/1998JD200029, 1998. 27141

Paatero, P.: Least squares formulation of robust non-negative factor analysis, *Chemometr. Intell. Lab.*, 37, 23–35, doi:10.1016/S0169-7439(96)00044-5, 3rd International Conference on Environmetrics and Chemometrics, LAS VEGAS, NV, 11–13 Sep 1995, 1997. 27146

10 Paatero, P. and Tapper, U.: Positive matrix factorization- A nonnegative factor model with optimal utilization of error-estimates of data values, *Environmetrics*, 5, 111–126, doi:10.1002/env.3170050203, 1994. 27146

Puxbaum, H., Caseiro, A., Sanchez-Ochoa, A., Kasper-Giebl, A., Claeys, M., Gelencser, A., Legrand, M., Preunkert, S., and Pio, C.: Levoglucosan levels at background sites in Europe for assessing the impact of biomass combustion on the European aerosol background, *J. Geophys. Res.-Atmos.*, 112, doi:10.1029/2006JD008114, 2007. 27150

Seinfeld, J. and Pandis, S.: Atmospheric Chemistry and Physics, from Air Pollution to Climate Change, *J. Atmos. Chem.*, 37, 212–214, doi:10.1023/A:1006483708571, 1998. 27154

20 Slowik, J. G., Stroud, C., Bottenheim, J. W., Brickell, P. C., Chang, R. Y.-W., Liggio, J., Makar, P. A., Martin, R. V., Moran, M. D., Shantz, N. C., Sjostedt, S. J., van Donkelaar, A., Vlasenko, A., Wiebe, H. A., Xia, A. G., Zhang, J., Leaitch, W. R., and Abbatt, J. P. D.: Characterization of a large biogenic secondary organic aerosol event from eastern Canadian forests, *Atmos. Chem. Phys.*, 10, 2825–2845, doi:10.5194/acp-10-2825-2010, 2010. 27151

25 Ulbrich, I. M., Canagaratna, M. R., Zhang, Q., Worsnop, D. R., and Jimenez, J. L.: Interpretation of organic components from Positive Matrix Factorization of aerosol mass spectrometric data, *Atmos. Chem. Phys.*, 9, 2891–2918, doi:10.5194/acp-9-2891-2009, 2009. 27146, 27147

Venzac, H., Sellegri, K., Villani, P., Picard, D., and Laj, P.: Seasonal variation of aerosol size distributions in the free troposphere and residual layer at the puy de Dme station, France, *Atmos. Chem. Phys.*, 9, 1465–1478, doi:10.5194/acp-9-1465-2009, 2009. 27141, 27142, 27143, 27147, 27153

Weingartner, E., Nyeki, S., and Baltensperger, U.: Seasonal and diurnal variation of aerosol

size distributions ($10 < D < 750$ nm) at a high-alpine site (Jungfrauoch 3580 m a.s.l.), J. Geophys. Res.-Atmos., 104, 26809–26820, doi:10.1029/1999JD900170, Joint Symposium on Atmospheric Chemistry and Climate Change, Univ. Washington, Seattle, Washington, 19–25 Aug, 1998, 1999. 27141, 27147

- 5 Zhang, Q., Alfarra, M., Worsnop, D., Allan, J., Coe, H., Canagaratna, M., and Jimenez, J.: Deconvolution and quantification of hydrocarbon-like and oxygenated organic aerosols based on aerosol mass spectrometry, Environ. Sci. Technol., 39, 4938–4952, doi:10.1021/es048568l, 2005. 27141, 27149

**Seasonal
characterisation of
aerosol particles at
the puy de Dôme site**

E. J. Freney et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Seasonal characterisation of aerosol particles at the puy de Dôme site

E. J. Freney et al.

Table 1. Average meteorological parameters and particle number concentration measured at the pdD site for each of the campaigns. The errors represent one standard deviation.

Season	RH%	Wind speed (ms ⁻¹)	Temp °C	UV	Particle concentration (SMPS) (cm ⁻³)	
					Day	Night
Autumn	9 ± 4	86 % ± 18	6 ± 4	5 ± 7	3018 ± 1969	2003 ± 1259
Winter	1 ± 4	86 % ± 21	7 ± 4	5 ± 7	1557 ± 1443	1301 ± 1160
Summer	14 ± 5	87 % ± 12	2.2 ± 1.2	9 ± 11	3405 ± 1156	3364 ± 1108

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Seasonal characterisation of aerosol particles at the py de Dôme site

E. J. Freney et al.

Table 2. The fractional contribution of each species to the total submicron particle mass (%) and their corresponding average mass concentrations ($\mu\text{g m}^{-3}$). The value shown for each of the separated organic species (LVOOA and SV-OOA) is the fractional contribution of these species to the total organics.

Season		SO ₄	NO ₃	NH ₄	Chl	Org	BC	LV-OOA	SV-OOA
Autumn	%	30	15	18	0.3	32	5	73	27
	$\mu\text{g m}^{-3}$	2.4	1.14	1.36	0.02	2.52	0.38	1.27	0.48
Winter	%	23	31	20	1	23	2	52	48
	$\mu\text{g m}^{-3}$	1.28	1.71	1.08	0.07	1.24	0.2	0.412	0.418
Summer	%	20	9	13	0.2	56	2	68	32
	$\mu\text{g m}^{-3}$	5.45	2.33	3.69	0.06	15.59	0.474	7.82	3.76

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Seasonal characterisation of aerosol particles at the puy de Dôme site

E. J. Freney et al.




Fig. 1. Equivalent potential temperature (θ_e) determined from meteorological parameters measured at pD and at the OPGC for the **(a)** autumn, **(b)** winter, and **(c)** summer campaign coloured by hour of day (UTC). **(d)** Lidar image, coloured by range-corrected signal at 355 nm (arbitrary units) taken during the summer campaign showing that the pD station was always in the boundary layer (values greater than 3×10^5). The dark dotted line represents to the height of the pD (1465 m a.s.l.).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Seasonal characterisation of aerosol particles at the puy de Dôme site

E. J. Freney et al.




Fig. 2. Mass concentration of organic and inorganic particles measured by the AMS, and BC measured by MAAP during autumn, winter, and summer. Pie charts illustrate the average particle composition measured during each campaign. Coloured bars at the top of each graph indicate the source of the airmass. When the bars are multicoloured it indicates that the air-masses were mixed during this period of sampling.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Seasonal characterisation of aerosol particles at the py de Dôme site

E. J. Freney et al.




Fig. 3. Average mass spectra of organic species extracted from the total organic mass spectra for the (a) Autumn, (b) winter, and (c) Summer campaign. Smaller sticks overlaid on each mass spectral pattern represent those of Ng et al. (2011).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 4. Plot of the fraction of Org44 to total Organics (f_{44}) against the fraction of Org43 to the total organics (f_{43}) for each campaign. Each triangular point represents data averaged over a 60-minute period. The star shaped points show f_{44} and f_{43} of each of the different organic species extracted from the total organic signal using PMF. The black dotted lines show the boundaries set by Ng et al. (2010).

Seasonal characterisation of aerosol particles at the puy de Dôme site

E. J. Freney et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Seasonal characterisation of aerosol particles at the puy de Dôme site

E. J. Freney et al.




Fig. 5. Particle composition averaged over season and airmass type.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 6. Diurnal variation of mass concentration of BC and inorganic and organic particles for autumn, (a) and (b), winter, (c) and (d), and summer, (e) and (f).

**Seasonal
characterisation of
aerosol particles at
the puy de Dôme site**

E. J. Freney et al.




Fig. 7. Difference in thetaE vs. the CO/NO_x ratio during winter 2009. Plots are coloured by **(a)** SO₄, **(b)** NO₃, **(c)** LV-OOA, and **(d)** SV-OOA.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion