

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

# Atmospheric mercury observations from Antarctica: seasonal variation and source and sink region calculations

K. Aspmo Pfaffhuber<sup>1</sup>, T. Berg<sup>1,\*</sup>, D. Hirdman<sup>1,\*\*</sup>, and A. Stohl<sup>1</sup>

<sup>1</sup>Norwegian Institute for Air Research (NILU), Norway

\* now at: Department of Chemistry, Norwegian University of Science and Technology, Norway

\*\* now at: Centre for Environmental and Climate Research (CEC), Lund University, Sweden

Received: 11 October 2011 – Accepted: 13 October 2011 – Published: 28 October 2011

Correspondence to: K. Aspmo Pfaffhuber (kap@nilu.no)

Published by Copernicus Publications on behalf of the European Geosciences Union.

## Atmospheric mercury observations from Antarctica

K. Aspmo Pfaffhuber  
et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

## Abstract

Long term atmospheric mercury measurements in the Southern Hemisphere are scarce and in Antarctica completely absent. Recent studies have shown that the Antarctic continent plays an important role in the global mercury cycle. Therefore, long term measurements of gaseous elemental mercury (GEM) were initiated at the Norwegian Antarctic Research Station, Troll (TRS) in order to improve our understanding of atmospheric transport, transformations and removal processes of GEM. GEM measurements started in February 2007 and are still ongoing, and this paper presents results from the first four years. The mean annual GEM concentration was  $0.93 \pm 0.19 \text{ ng m}^{-3}$  and is in good agreement with other recent southern hemispheric measurements. Measurements of GEM were combined with the output of the Lagrangian particle dispersion model FLEXPART, for a statistical analysis of GEM source and sink regions. It was found that the ocean is a source of GEM to TRS year round, especially in summer and fall. None of the Southern Hemisphere continents contribute significantly to the direct transport of GEM to TRS, but they are important for determining the overall GEM load in the Southern Hemisphere and for the mean GEM concentration at TRS. Further, the sea ice and marginal ice zones are GEM sinks in spring as also seen in the Arctic, but the Antarctic oceanic sink seems weaker. Contrary to the Arctic, a strong summer time GEM sink was found, when air originates from the Antarctic Plateau, which shows that the summertime removal mechanism of GEM is completely different and is caused by other chemical processes than the springtime atmospheric mercury depletion events. The results were corroborated by an analysis of ozone source and sink regions.

## 1 Introduction

Antarctica is the most remote of all continents and is usually perceived as an isolated and hostile place and a symbol of the last great wilderness untouched by human

ACPD

11, 29117–29139, 2011

## Atmospheric mercury observations from Antarctica

K. Aspö Pfaßhuber  
et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



disturbance. However, like other remote regions on Earth, it is not escaping the impact of local pollutant emissions due to increasing human presence and pollution imported from other continents (Priddle, 2002; Bargagli, 2005; Bergstrom et al., 2006; Stohl and Sodemann, 2010). Mercury behaves exceptional for a metal in the environment; it has a very complex biogeochemical cycle and exists in a variety of forms in the atmosphere, such as gaseous elemental mercury (GEM), gaseous oxidized mercury (GOM) and particulate bound mercury (PBM). The discovery of a unique scavenging process called atmospheric mercury depletion events (AMDEs), where the concentration of atmospheric mercury drops precipitously within hours (Schroeder et al., 1998), resulted in intensified research on atmospheric mercury. Mercury is now being monitored at many sites, although mainly in the Northern Hemisphere. Measurements in the Southern Hemisphere are generally scarce and particularly the Antarctic regions have not been extensively monitored and mainly sporadic measurements have been made. Efforts have been initiated to study AMDEs at coastal sites (Ebinghaus et al., 2002; Temme et al., 2003a; Sprovieri et al., 2002; Brooks et al., 2008a) and more recently the Antarctic Plateau has become a new focus of attention (Brooks et al., 2008b). These efforts show us that we currently underestimate the role of this continent in the global mercury cycle (Dommergue et al., 2010). Intensive measurement campaigns with extensive measurement programs provide large amounts of information that apply to a short time period, however, care should be taken when extrapolating to annual budgets and large geographical areas. Long term monitoring can provide valuable information on seasonal and annual variation as well as long term concentration trends, and hence the global mercury cycle. Combining such large data sets with corresponding atmospheric transport model output allows a statistical analysis of the sources and sinks of mercury in large regions around the measurement site (Hirdman et al., 2009) by investigating the origin of high and, respectively, low measured mercury concentrations.

A long-term monitoring program of GEM was initiated in 2007 at the Norwegian Antarctic monitoring station Troll in order to extend the global mercury database and

## Atmospheric mercury observations from Antarctica

K. Aspö Pfaßhuber et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

to improve the current understanding of the atmospheric transport, transformation and removal processes of GEM. The data from Troll is to our knowledge the longest times series of GEM from the Antarctic continent and measurements are still ongoing. In this study we present seasonal and annual GEM concentration variations and present a statistical analysis of the source- and sink regions for GEM observed at Troll.

## 2 Methods

### 2.1 Site description

The Norwegian Antarctic Troll Research Station (TRS) is located in Queen Maud Land at 72°01' S, 2°32' E at an elevation of 1275 m and approximately 220 km from the Antarctic coast. TRS is situated on snow-free bedrock and accessible by air-transport during Antarctic summer, facilitated by a blue-ice airfield on the glacier 7 km north of the main station. After TRS was turned from a summer into an all-year station in 2005, the *Norwegian Institute for Air Research* (NILU) deployed a container-housed atmospheric observatory in February 2007. TRS is one of the few stations located on the slope of the Antarctic ice sheet (and the only one manned year-round). It is exposed to a varying influence from both the Antarctic Plateau and the Southern Ocean, while the environment at other long-term activity stations either is of coastal or high-elevation continental nature. A detailed description of TRS can be found in Hansen et al. (2009) along with meteorological conditions, instrument set-up and some first results.

### 2.2 Experimental

Measurements of GEM were initiated in February 2007. A Tekran gas phase analyser (Model 2537A, Tekran Inc) was installed to collect and determine GEM concentrations in air. The instrument was programmed to sample air at a flow rate of 1.5 l min<sup>-1</sup> with a 5 min time resolution. For data analysis, 1 h averages were used, unless otherwise

## Atmospheric mercury observations from Antarctica

K. Aspö Pfaßhuber et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



stated. Auto calibrations were performed every 25 h using the instrument's internal calibration source. The internal calibration source was checked against manual injections once per year. The quality assurance and control protocol applied on all data left a data coverage of more than 94 % during the entire sampling period. The sample inlet is located approximately 6 m above surface. More details on the instrumental set-up and quality control can be found in Hansen et al. (2009), Berg et al. (2003) and Aspmo et al. (2005). Additionally, atmospheric in-situ concentrations of ozone were recorded by UV absorption spectrometry (API 400). Ozone ( $O_3$ ) concentrations were measured with 1-min time resolution, and 1 h averages were used for data analysis. More details on  $O_3$  measurements can be found in Hansen et al. (2009). For the analysis of source regions, GEM and  $O_3$  data were averaged to 3 h to fit with the time resolution of our transport model output.

### 2.3 Source and sink region analysis

To identify sources and sinks of measured GEM and  $O_3$ , 3-hourly backward simulations from TRS with the Lagrangian particle dispersion model FLEXPART (Stohl et al., 1998, 2005) were used. FLEXPART was driven with 3-hourly operational meteorological data from the European Centre for Medium-Range Weather Forecasts with  $1^\circ \times 1^\circ$  resolution. During every 3-hour interval, 60 000 particles were released at the measurement point and followed backward for 20 days to calculate emission sensitivity ( $S$ ) on a  $1^\circ \times 1^\circ$  grid, under the assumption that removal processes can be neglected.  $S$  (in units of  $s\ m^{-3}$ ) in a particular grid cell is proportional to the particle residence time in that cell and measures the simulated concentration at the receptor that a source of unit strength ( $1\ kg\ s^{-1}$ ) in the cell would produce. The  $S$  distribution in a 100 m layer adjacent to the surface (so-called footprint layer) was used as input to the statistical analysis of surface sources and sinks.

A statistical method described in detail in Hirdman et al. (2009, 2010) was used to identify possible source and sink regions of GEM. The method is similar to older methods based on trajectory calculations (Ashbaugh, 1983; Ashbaugh et al., 1985)

## Atmospheric mercury observations from Antarctica

K. Aspmo Pfaffhuber et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



but takes advantage of the superior quality of Lagrangian particle dispersion model output compared to simple trajectories, which ignore turbulence and convection. Every one of  $M$  measurements were related to a modeled footprint  $S$  field, and the average footprint  $S_T$  was calculated as

$$S_T(i, j) = \frac{1}{M} \sum_{m=1}^M S(i, j, m) \quad (1)$$

where  $i, j$  are grid indices of  $S$ . Then, the subset of the data with the  $L = M/10$  highest 10% (or, respectively, lowest 10%) of measured GEM concentrations was selected to calculate

$$S_P(i, j) = \frac{1}{L} \sum_{l=1}^L S(i, j, l) \quad (2)$$

where the suffix P indicates the percentile (10% or 90%). The ratio

$$R_P(i, j) = \frac{L}{M} \frac{S_P(i, j)}{S_T(i, j)} \quad (3)$$

can then be used for identifying grid cells that are likely sources (or sinks) of GEM. If air mass transport patterns were the same for the data subset and for the full data set, one would expect  $R_P(i, j) = 0.1$  for all  $i, j$ . Information on sources and sinks of GEM are contained in the deviations from this expected value. For the top decile of the data, for instance,  $R_{90}(i, j) > 0.1$  means that high measured GEM concentrations are associated preferentially with high  $S$  values in grid cell  $(i, j)$ , indicating a likely source, whereas  $R_{90}(i, j) < 0.1$  indicates a possible sink or at least the absence of a source. Conversely, when using the lowest decile of the data,  $R_{10}(i, j) > 0.1$  indicates a likely sink in grid cell  $(i, j)$ , and  $R_{10}(i, j) < 0.1$  a source.

Not all features of  $R_P$  are statistically significant. Therefore, calculation of  $R_P$  was limited to grid cells where  $S_T > 5 \times 10^{-9} \text{ s m}^{-3}$  and a bootstrap resampling analysis

## Atmospheric mercury observations from Antarctica

K. Aspö Pfaffhuber et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



was employed to ensure statistical significance. For more details on the method, see Hirdman et al. (2009, 2010).

### 3 Results and discussion

#### 3.1 Mercury concentrations at TRS compared to Zeppelin (Spitsbergen, Arctic) and measurements in the Southern Hemisphere

The arithmetic mean GEM concentration during more than four years of continuous measurements at TRS was  $0.93 \pm 0.19 \text{ ng m}^{-3}$ , whereas the median concentration was  $0.97 \text{ ng m}^{-3}$ . This is, to the best of the authors' knowledge, the longest time series of GEM from the Antarctic continent and the only annual time series covering more than a year. The mean GEM concentration observed at TRS is in good agreement with other recent observations from high latitudes in the Southern Hemisphere (Ebinghaus et al., 2002; Xia et al., 2010; Temme et al., 2003b). Figure 1a shows a box and whisker plot comparing NILUs Antarctic GEM measurements at TRS and Arctic GEM measurements obtained at Zeppelin station, Spitsbergen ( $78^{\circ}54' \text{ N}$ ,  $11^{\circ}52' \text{ E}$ ). As shown in Fig. 1a the annual concentration variability is rather similar at the two locations, whereas the distribution is shifted towards lower concentrations as TRS. The GEM concentrations observed at TRS are only about two-thirds of the GEM concentrations at Zeppelin ( $1.57 \text{ ng m}^{-3}$ , annual average 2007–2010). The mean concentration at Zeppelin falls within the range of what is being considered the northern hemispheric background concentration,  $1.5\text{--}1.7 \text{ ng m}^{-3}$  (Slemr et al., 2003). GEM concentrations observed in the Southern Hemisphere are lower because most of the mercury emission sources are located on the Northern Hemisphere (Pacyna et al., 2006) and in the atmosphere the lifetime of GEM (Schroeder and Munthe, 1995) is not long enough for homogeneous mixing on a global scale. At Cape Point, South Africa, Slemr et al. (2008) observed a small significant decrease in atmospheric mercury concentrations from  $1.29 \text{ ng m}^{-3}$  in 1996 to  $1.19 \text{ ng m}^{-3}$  in 2004. This decline in concentration

## Atmospheric mercury observations from Antarctica

K. Aspö Pfaßhuber  
et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Atmospheric mercury observations from Antarctica

K. Aspö Pfaßhuber  
et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



has continued and Brunke et al. (2010) reported GEM concentrations from Cape Point from 2007 and 2008 to be  $0.94 \pm 0.16 \text{ ng m}^{-3}$ . Slemr et al. (2003) suggested a small negative concentration gradient of GEM towards high southerly latitudes, as the median GEM concentration observed at the German Antarctic research station Neumayer in 2000 and 2001 was  $1.10 \text{ ng m}^{-3}$  (Ebinghaus, 2002). However, this southward decreasing concentration gradient is not apparent in the data presented here as the concentrations at Cape Point and TRS are practically equal. Our time series is not long enough for trend analysis, but the data presented here does not show any change in annual mean or median GEM concentration from February 2007 to June 2011. As argued by Cole et al. (2011) the Polar Regions are different, compared to temperate latitudes, with respect to how they respond to mercury emissions reductions.

### 3.2 Seasonal variation in mercury concentrations at TRS

Figure 1b shows, through monthly box plots of all data collected, the seasonal variation of measured GEM concentrations. The median GEM concentration shows a maximum in winter and a minimum in summer, contrary to observations from Cape Point for which Slemr et al. (2008) and Brunke et al. (2010) hypothesized that the seasonality predominantly was driven by mercury emissions. At TRS the winter maximum is most likely caused by the lack of photochemical oxidation processes during the polar night (from May to July).

From late fall through winter (April–July), GEM concentrations remains at a virtually constant level of  $1.00 \pm 0.07 \text{ ng m}^{-3}$ , and the median concentration equals the mean. In spring and summer (August–February) GEM concentrations are highly variable ranging from 0.02 to  $6.04 \text{ ng m}^{-3}$  and with mean concentration of  $0.86 \pm 0.24 \text{ ng m}^{-3}$ . The high variability suggests that exchange processes at the surface (source and/or sinks) must be more active at high southern latitudes in spring and summer than during winter. In spring (August, September and October), the median concentration is higher than the mean concentration, indicating that the springtime mean is influenced by episodic low concentrations caused by AMDEs as also observed by Ebinghaus et al. (2002).



In summer (November, December and January), the median concentration is lower compared to the mean indicating that the mean is influenced by episodic high concentrations. These observations are in good agreement with measurements performed at the German Antarctic research station Neumayer in 2000 and 2001 (Ebinghaus et al., 2002).

Diurnal variation in the GEM concentrations was not observed at any time of the year at TRS. The reason for this can be that there are possibly no sources or sinks for GEM with a diurnal cycle in the vicinity of this site. The ground surrounding TRS is mainly snow free bedrock year round, a type of surface that does not lead to a great deal of deposition followed by solar radiation induced re-emission. Additionally, the sample inlet is located 6 m above ground, a height that perhaps attenuates minor diurnal variation that might be observed closer to the surface.

### 3.3 AMDEs at TRS

From the end of August until the end of October, AMDEs were observed at TRS in strong positive correlation with  $O_3$  (correlation coefficients  $r$  up to 0.79). AMDEs are often operationally defined as GEM concentrations below  $1.0 \text{ ng m}^{-3}$  for AMDEs observed in the Arctic. Using the same relative decrease in GEM concentrations for the Antarctic, AMDEs at TRS would occur at GEM concentrations below  $0.6 \text{ ng m}^{-3}$ . This occurs in 5% of the springtime observations as in contrast to 21% of the observations at Zeppelin in the Arctic (Berg et al., 2011). As seen in the Fig. 2, periodic GEM concentrations below  $0.6 \text{ ng m}^{-3}$  also occur after the end of October, however these GEM depletions are not caused by AMDE chemistry as GEM is anti-correlated with  $O_3$  in these cases, as also discussed in Temme et al. (2003a) (see also following discussion).

## Atmospheric mercury observations from Antarctica

K. Aspömp Pfaffhuber et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

### 3.4 Statistical analysis of source and sink regions

Figure 3 shows  $R_p$  fields for both the highest and lowest 10 % of all GEM data divided into the different seasons; winter, spring, summer and fall. Winter is represented as the months May–July, spring is represented by September–October, summer is represented by November–January and fall is represented by February. March, April and August data are not included in the analysis, because these months turned out to be transition periods not falling clearly into one of the above defined seasons. The  $R_{90}$  plots, representing the highest 10 % of all GEM measurements show that the open ocean is to a varying degree a source region of GEM to TRS all year round (Fig. 3 a–d). Especially in summer and fall the open ocean is a strong source of GEM (Fig. 3c–d). The highest  $R_{90}$  values are found over the high-latitude seas surrounding Antarctica. This is similar to the Arctic situation, where Hirdman et al. (2009) also found the Arctic Ocean to be a source of GEM in summer, after being a strong sink in spring. They suggested that some of the mercury lost in spring from the Arctic atmosphere during AMDEs could be re-emitted in summer. In the Antarctic case, the spring-time oceanic sink appears weaker (Fig. 3f). This may in part be caused by the location of TRS not being exposed directly to depletion events but rather to transport of mercury-depleted air, as can also be seen by the lower frequency of AMDEs at TRS compared to Zepelin (see above). However, the strong oceanic emissions in summer (Fig. 3c) may be a result of spring-time deposited mercury or evasion resulting from a sea ice free ocean.

The figure also shows that emissions of mercury from the Southern Hemisphere continents do not contribute significantly to direct transport of GEM to TRS. South Africa, one of the largest emitters of mercury globally (Pacyna et al., 2010), does not appear as an identifiable source region for GEM at TRS. Meridional transport from South Africa to Antarctica typically takes more than 20–30 days, especially in summer (Stohl and Sodemann, 2010) and does not occur frequently enough on the 20 day timescale of the FLEXPART calculations to be represented in the statistics. Even when extending the

## Atmospheric mercury observations from Antarctica

K. Aspö Pfaßhuber et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



calculations to 30 days we did not register a clear signal because on this time scale the sampled air masses are a mixture of air originating from many different regions at the same time and do not cause clear concentration variability at the measurement station. In addition, on these timescales modeled transport also becomes very uncertain. Correspondingly, the other continents do not appear directly as a source regions for GEM at TRS in our analysis. However, these emissions are certainly important for determining the overall GEM loading in the Southern Hemisphere atmosphere and, thus, also for the mean concentrations measured at TRS. Indeed, especially for spring (Fig. 3b) the analysis provides a hint that transport from lower-latitude regions (including the tip of South America) is frequently associated with the highest GEM concentrations, suggesting mid-latitude sources of the observed high GEM concentrations, although the particular source regions could not be resolved.

Considering the  $R_{10}$  plots, representing the lowest 10% of all GEM measurements, for winter (Fig. 3e), the  $R_{10}$  values are almost everywhere below 0.15 and mostly below 0.1. This indicates that air masses associated with low GEM concentrations avoid surface contact and therefore often descend from above the boundary layer. In spring (Fig. 3f), however, the highest  $R_{10}$  values are found in conjunction with sea ice and marginal ice zones, indicating removal at the surface. This is the area where AMDEs are known to occur (Kaleschke et al., 2005; Simpson et al., 2007; Steffen et al., 2008). Additionally, high  $R_{10}$  values are found over the Antarctic Plateau where there are no emission sources. In summer (Fig. 3g) on the other hand, when the sea ice extent is at an absolute minimum, no surface removal is observed in this area, and the highest  $R_{10}$  values are found exclusively over the Antarctic Plateau. In fall (Fig. 3h), the area with high  $R_{10}$  values over the plateau is small, but the Ross Sea appears as an additional sink.

In spring,  $O_3$  and GEM chemistry is closely tied together both in the Arctic and Antarctic atmosphere (Schroeder et al., 1998; Ebinghaus et al., 2002). To support the validity of the interpretation of the GEM plots, the statistical analysis was repeated for  $O_3$ . The  $R_{10}$  plots for  $O_3$  (Fig. 4 e–h) shows that air masses over the Antarctic continent

## Atmospheric mercury observations from Antarctica

K. Aspö Pffaffhuber  
et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

and the seas surrounding it containing low concentrations of  $O_3$  have above average surface contact ( $R_{10} > 0.1$ ), thus  $O_3$  is destroyed at the surface. In spring (Fig. 4f) low  $O_3$  concentrations mainly originate from sea ice dense areas surrounding Queen Maud Land as also seen for GEM (Fig. 3f), though the extent of low GEM values is smaller than for  $O_3$ . The  $R_{90}$  plots for  $O_3$  (Fig. 4a–d) show that air masses containing high concentrations of  $O_3$  have in general little surface contact. This probably indicates descent from the stratosphere. Only air masses originating from low-latitudes (north of  $30^\circ$  S) are associated with high  $O_3$  concentrations, probably indicating transport of photochemically formed  $O_3$  from lower latitudes even though the actual continental source regions cannot be resolved. The  $R_{90}$  summer and fall plots (Fig. 4c–d) shows  $O_3$  rich air originating from the Antarctic Plateau. This is exactly the same area from where air low in GEM comes from.

In contrast to the Arctic where the periodic low concentration GEM episodes end with the onset of snowmelt (Lindberg et al., 2002), episodic low GEM concentrations are observed at TRS throughout the summer (see Fig. 2). Sprovieri et al. (2003) and Brooks et al. (2008a) argue that halogen chemistry causes the summertime depletion of GEM such as during the springtime AMDEs. Contrary to that, Sprovieri et al. (2003) who carried out Hg species measurements at the Italian Antarctic research base, Terra Nova Bay, close to the Ross Sea, put forward a hypothesis that the summer time low level GEM and the coinciding high level GOM could be due to pure meteorological factors rather than atmospheric chemistry processes, such as stratification of the planetary boundary layer. The height of the atmospheric mixed layer changes over time, which also leads to changes of its chemical composition. This is particularly valid for species emitted from the ground or sea surface, the dilution of which depends on the mixed layer height. However, our plots and data show that the summertime low GEM concentrations are not caused by either of the two explanations; it must be a different mechanism than AMDEs because transport from remaining sea ice regions (e.g., in the Weddell Sea) is not associated with low GEM concentrations at TRS in summer and fall (Fig. 3g–h) and it cannot be stratification of the planetary boundary layer because

**Atmospheric mercury observations from Antarctica**

K. Aspömf Pfaffhuber et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



GEM and O<sub>3</sub> anti-correlates. As proposed by Temme et al. (2003a), AMDEs in Antarctica end abruptly in the beginning of November and another oxidation mechanism takes over. They suggested that GEM oxidation had already occurred before air parcels were advected to their measurement location. The Antarctic Plateau mixed-layer represents a highly oxidizing environment during summer resulting from low temperatures typically not exceeding -25 °C leading to frequent cases of strong near surface temperature inversions in combination with continuous sunlight giving rise to non-stop photochemical reactions both within the snowpack and the atmospheric boundary layer (Crawford et al., 2001; Davis et al., 2008). These conditions result in an efficient release of NO<sub>x</sub> from the snowpack to the atmosphere, leading to extremely high NO mixing ratios. NO rapidly converts abundant HO<sub>2</sub> into OH, enhancing the OH concentration and O<sub>3</sub> production. Summertime OH concentrations over the Antarctic Plateau (~ 10<sup>6</sup> cm<sup>-3</sup>) are considerably higher than coastal OH concentrations (~ 10<sup>5</sup> cm<sup>-3</sup>) (Bloss et al., 2007) in part due to the low mixing layer at the South Pole but also the effect of snowpack emissions of NO<sub>x</sub> (Davis et al., 2004). Temme et al. (2003a) proposed OH, HO<sub>2</sub>, O(1D) as possible GEM oxidants. The high concentrations of GOM observed at the coastal sites by Temme et al. (2003a) and Sprovieri et al. (2003) must have been transported from the Antarctic Plateau to the measurement location. This would be consistent with our finding that in summer the lowest GEM concentrations are associated with transport from the Plateau (Fig. 3g), suggesting an oxidative sink there.

Figure 4c–d shows that the same transport pathway from the Antarctic Plateau, which causes low GEM concentrations at TRS, is associated with the highest O<sub>3</sub> concentrations at TRS in summer and fall. This shows clearly that the mechanism removing the GEM must be different from the AMDE chemistry.

## 4 Conclusions

More than four years of atmospheric mercury measurements from the Antarctic Research Station Troll are presented in this paper. The mean concentration of GEM was

29129

## Atmospheric mercury observations from Antarctica

K. Aspömp Pfaffhuber et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



0.93 ± 0.19 ng m<sup>-3</sup>, which is in good agreement with recent GEM measurements at other high latitude sites in the Southern Hemisphere. Significant long-term decreases in GEM concentrations are observed at many monitoring sites both in the Northern Hemisphere and at Cape Point, South Africa as a response to decreasing mercury emissions. However, no such change in the annual mean or median concentration could be detected in the rather short time series from TRS.

A seasonal concentration variation was observed at TRS, with a maximum in winter and minimum in summer. The wintertime maximum is caused by absent photochemical removal processes which allows for a buildup of atmospheric mercury. Spring and summer shows highly variable GEM concentrations indicating extremely active surface exchange processes at this time of the year. No diurnal variation was observed throughout the year, which is likely caused by the nature of the surrounding surfaces being not ideal for deposition followed by radiation induced re-emission. AMDEs were observed every spring in strong correlation with O<sub>3</sub>; however the depletions seem weaker when compared to the Arctic as seen in both the oceanic springtime sink and the AMDE occurrence frequency.

Statistical analysis of source and sink regions show that the ocean is a source of mercury to TRS all year round, and especially in summer and fall. This is likely caused by oceanic emission of springtime deposited GEM or evasion resulting from a sea ice free ocean. None of the southern hemispheric continents are a direct source of mercury to TRS, but they do contribute significantly to the overall mercury loading in the Southern Hemisphere atmosphere. Sinks for GEM in winter are lacking indicating that air masses containing low GEM concentrations avoid surface contact and often descend from above the boundary layer. In spring, removal of GEM is observed at the surface in conjunction with sea ice and marginal ice zones, caused by AMDEs. In summer, no oceanic sink is observed, however the interior of the Antarctic Plateau appears as a strong sink, indicating that the frequent low GEM concentration episodes observed during summer are caused by a different mechanism than AMDEs.

## Atmospheric mercury observations from Antarctica

K. Aspö Pfaﬀhuber  
et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Atmospheric mercury observations from Antarctica

K. Aspö Pffaffhuber  
et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A statistical analysis of  $O_3$  showed that the Antarctic continent and the surrounding seas are a sink, thus  $O_3$  is destroyed at the surface. In spring this sink is mainly located within sea ice dense areas surrounding Queen Maud Land, the same area as where the springtime GEM sink is found. In summer on the other hand the situation is different; the oceanic  $O_3$  sink is maintained, but the GEM sink region is found over the Antarctic Plateau where air rich in  $O_3$  is coming from. The Antarctic Plateau is a highly oxidizing environment in summer; resulting in efficient release of  $NO_x$  from the snowpack leading to high  $NO/NO_2$  cycling rapidly enhances radical concentrations, such as OH, which further leads to increased  $O_3$  production. It is suggested that the increase in radical concentration due to snowpack emissions of  $NO_x$  possibly also leads to strong GEM oxidation. The results show that the same transport pathway from the Antarctic Plateau causing low GEM concentrations at TRS is associated with the highest  $O_3$  concentrations at TRS in summer and fall. This shows that the elevated summertime concentration of oxidized mercury species observed by others at coastal locations results from different chemical processes than those causing AMDEs.

*Acknowledgements.* Financial support to sustain measurements is given through the Norwegian Antarctic Research Expeditions (NARE) program administered by the Norwegian Polar Institute (NPI). We are very grateful for the technical support offered by the overwintering NPI staff at TRS, and we also want to thank our excellent technician and instrument care taker Jan H. Wasseng. Funding by the Norwegian Research Council in the framework of CLIMSLIP supported the model analyses.

## References

- Ashbaugh, L. L.: A statistical trajectory technique for determining air pollution source regions, J. Air Poll. Contr. Ass., 33, 1096–1098, 1983.
- Ashbaugh, L. L., Malm, W. C., and Sadeh, W. Z.: A residence time probability analysis of sulfur concentrations at Grand Canyon National Park, Atmos. Environ., 19, 1263–1270, 1985.

**Atmospheric mercury observations from Antarctica**K. Aspmo Pfaffhuber  
et al.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Aspmo, K., Gauchard, P. A., Steffen, A., Temme, C., Berg, T., Bahlmann, E., Banic, C., Domergue, A., Ebinghaus, R., Ferrari, C., Pirrone, N., Sprovieri, F., and Wibetoe, G.: Measurements of atmospheric mercury species during an international study of mercury depletion events at Ny-Alesund, Svalbard, spring 2003. How reproducible are our present methods?, *Atmos. Environ.*, 39, 7607–7619, 2005.
- Bargagli, R., Agnorelli, C., Borghini F., and F. Monaci.: Enhanced deposition and bioaccumulation of mercury in Antarctic terrestrial ecosystems facing a coastal polynya, *Environ. Sci. Technol.*, 39, 8150–8155, 2005.
- Berg, T., Sekkesaeter, S., Steinnes, E., Valdal, A. K., and Wibetoe, G.: Springtime depletion of mercury in the European Arctic as observed at Svalbard, *Sci. Total Environ.*, 304, 43–51, 2003.
- Berg, T., Pfaffhuber, K. A., Cole, A., and Steffen, A.: Long-term atmospheric mercury measurements in the Norwegian Arctic: trend analysis, in preparation, 2011.
- Bergstrom, D. M., Convey P., and Huiskes, A. H. L.: *Trends in Antarctic Terrestrial and Limnetic Ecosystems*, Springer, Dordrecht, 369 pp., 2006.
- Bloss, W. J., Lee, J. D., Heard, D. E., Salmon, R. A., Bauguitte, S. J.-B., Roscoe, H. K., and Jones, A. E.: Observations of OH and HO<sub>2</sub> radicals in coastal Antarctica, *Atmos. Chem. Phys.*, 7, 4171–4185, doi:10.5194/acp-7-4171-2007, 2007.
- Brooks, S., Lindberg, S., Southworth, G., and Arimoto, R.: Springtime atmospheric mercury speciation in the McMurdo, Antarctica coastal region, *Atmos. Environ.*, 42, 2885–2893, 2008a.
- Brooks, S., Arimoto, R., Lindberg, S., and Southworth, G.: Antarctic polar plateau snow surface conversion of deposited oxidized mercury to gaseous elemental mercury with fractional long-term burial, *Atmos. Environ.*, 42, 2877–2884, 2008b.
- Brunke, E.-G., Labuschagne, C., Ebinghaus, R., Kock, H. H., and Slemr, F.: Gaseous elemental mercury depletion events observed at Cape Point during 2007–2008, *Atmos. Chem. Phys.*, 10, 1121–1131, doi:10.5194/acp-10-1121-2010, 2010.
- Cole, A., Steffen, A., Poissant, L., Pilote, M., Pfaffhuber, K. A., and Berg, T.: Trends in atmospheric mercury concentrations in the northern hemisphere: Why is the arctic different? Abstract submitted to the 10th International Conference on Mercury as a Global Pollutant, 7 Halifax, Canada, 24–29 July 2011, RS4-P5, 2011.
- Crawford, J. H., Davis, D. D., Chen, G., Buhr, M., Oltmans, S., Weller, R., Mauldin, L., Eisele, F., Shetter, R., Lefer, B., Arimoto, R., and Hogan A.: Evidence for photochemi-



## Atmospheric mercury observations from Antarctica

K. Aspmo Pfaffhuber  
et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

cal production of ozone at the South Pole surface, *Geophys. Res. Lett.*, 28, 3641–3644, doi:10.1029/2001GL013055, 2001.

Davis, D., Chen, G., Buhr, M., Crawford, J., Lenschow, D., Lefer, B., Shetter, R., Eisele, F., Mauldin, L., and Hogan, A.: South Pole  $\text{NO}_x$ , chemistry: an assessment of factors controlling variability and absolute levels, *Atmos. Environ.*, 38, 5375–5388, 2004.

Davis, D. D., Seelig, J., Huey, G., Crawford, J., Chen, G., Wang, Y., Buhr, M., Helmig, D., Neff, W., Blake, D., Arimoto, R., and Eisele, F.: Areassessment of Antarctic Plateau reactive nitrogen based on ANTCI 2003 airborne ground based measurements, *Atmos. Environ.*, 42, 2831–2848, 2008.

Dommergue, A., Sprovieri, F., Pirrone, N., Ebinghaus, R., Brooks, S., Courteaud, J., and Ferrari, C. P.: Overview of mercury measurements in the Antarctic troposphere, *Atmos. Chem. Phys.*, 10, 3309–3319, doi:10.5194/acp-10-3309-2010, 2010.

Ebinghaus, R., Kock, H. H., Temme, C., Einax, J. W., Lowe, A. G., Richter, A., Burrows, J. P., and Schroeder, W. H.: Antarctic springtime depletion of atmospheric mercury, *Environ. Sci. Technol.*, 36, 1238–1244, 2002.

Hansen, G., Aspmo, K., Berg, T., Edvardsen, K., Fiebig, M., Kallenborn, R., Krognes, T., Lunder, C., Stebel, K., Schmidbauer, N., Solberg, S., and Yttri, K. E.: Atmospheric monitoring at the Norwegian Antarctic Station Troll: measurement programme and first results, *Polar Res.*, 28, 353–363, 2009.

Hirdman, D., Aspmo, K., Burkhart, J. F., Eckhardt, S., Sodemann, H., and Stohl, A.: Transport of mercury in the Arctic atmosphere: Evidence of a spring-time sink and summer-time source, *Geophys. Res. Lett.*, 36, L12814, doi:10.1029/2009GL038345, 2009.

Hirdman, D., Sodemann, H., Eckhardt, S., Burkhart, J. F., Jefferson, A., Mefford, T., Quinn, P. K., Sharma, S., Ström, J., and Stohl, A.: Source identification of short-lived air pollutants in the Arctic using statistical analysis of measurement data and particle dispersion model output, *Atmos. Chem. Phys.*, 10, 669–693, doi:10.5194/acp-10-669-2010, 2010.

Kaleschke, L., Richter, A., Burrows, J., Afe, O., Heygster, G., Notholt, J., Rankin, A. M., Roscoe, H. K., Hollwedel, J., Wagner, T., and Jacobi, H.-W.: Frost flowers on sea ice as a source of sea salt and their influence on tropospheric halogen chemistry, *Geophys. Res. Lett.*, 31, L16114, doi:10.1029/2004GL020655, 2004.

Lindberg, S. E., Brooks, S., Lin, C. J., Scott, K. J., Landis, M. S., Stevens, R. K., Goodsite, M., and Richter, A.: Dynamic oxidation of gaseous mercury in the Arctic troposphere at polar sunrise, *Environ. Sci. Technol.*, 39, 1245–1256, 2002.

**Atmospheric mercury observations from Antarctica**K. Aspmo Pfaffhuber  
et al.[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Pacyna, E. G., Pacyna, J. M., Steenhuisen, F., and Wilson, S.: Global anthropogenic mercury emission inventory for 2000, *Atmos. Environ.*, 40, 4048–4063, 2006.
- Pacyna, E. G., Pacyna, J. M., Sundseth, K., Munthe, J., Kindbom, K., Wilson, S., Steenhuisen, F., and Maxon, P.: Global emission of mercury to the atmosphere from anthropogenic sources and projection to 2020, *Atmos. Environ.*, 44, 2487–2499, 2010.
- Schroeder, W. H. and Munthe, J.: Atmospheric mercury – an overview, *Atmos. Environ.*, 32, 809–822, 1995.
- Schroeder, W. H., Anlauf, K. G., Barrie, L. A., Lu, J. Y., Steffen, A., Schneeberger, D. R., and T. Berg: Arctic springtime depletion of mercury, *Nature*, 394, 331–332, 1998.
- Simpson, W. R., Carlson, D., Hönninger, G., Douglas, T. A., Sturm, M., Perovich, D., and Platt, U.: First-year sea-ice contact predicts bromine monoxide (BrO) levels at Barrow, Alaska better than potential frost flower contact, *Atmos. Chem. Phys.*, 7, 621–627, doi:10.5194/acp-7-621-2007, 2007.
- Slemr, F., Brunke, E.-G., Ebinghaus, R., Temme, C., Munthe, J., Wangberg, I., Schroeder, W., Steffen, A., and Berg, T.: Worldwide trend of atmospheric mercury since 1977, *Geophys. Res. Lett.*, 30, 1516, doi:10.1029/2003GL016954, 2003.
- Slemr, F., Brunke, E.-G., Labuschagne, C., and Ebinghaus, R.: Total gaseous mercury concentration at the Cape Point GAW stations and their seasonality, *Geophys. Res. Lett.*, 35, L11807, doi:10.1029/2008GL033741, 2008.
- Sprovieri, F., Pirrone, N., Hedgecock, I. M., Landis, M. S., and Stevens, R. K.: Intensive atmospheric mercury measurements at Terra Nova Bay in Antarctica during November and December 2000, *J. Geophys. Res.*, 107, 4722, doi:10.1029/2002JD002057, 2002.
- Steffen, A., Douglas, T., Amyot, M., Ariya, P., Aspmo, K., Berg, T., Bottenheim, J., Brooks, S., Cobbett, F., Dastoor, A., Dommergue, A., Ebinghaus, R., Ferrari, C., Gardfeldt, K., Goodsite, M. E., Lean, D., Poulain, A. J., Scherz, C., Skov, H., Sommar, J., and Temme, C.: A synthesis of atmospheric mercury depletion event chemistry in the atmosphere and snow, *Atmos. Chem. Phys.*, 8, 1445–1482, doi:10.5194/acp-8-1445-2008, 2008.
- Stohl, A. and Sodemann, H.: Characteristics of atmospheric transport into the Antarctic troposphere, *J. Geophys. Res.*, 112, D02305, doi:10.1029/2009JD012536, 2010.
- Stohl, A., Hittenberger, M., and Wotawa, G.: Validation of the Lagrangian particle dispersion model FLEXPART against large scale tracer experiment data, *Atmos. Environ.*, 32, 4245–4264, 1998.
- Stohl, A., Forster, C., Frank, A., Seibert, P., and Wotawa, G.: Technical note: The Lagrangian

particle dispersion model FLEXPART version 6.2, Atmos. Chem. Phys., 5, 2461–2474, doi:10.5194/acp-5-2461-2005, 2005.

5 Temme, C., Einax, J. W., Ebinghaus, R., and Schroeder, W. H.: Measurements of atmospheric mercury species at a coastal site in the Antarctic and over the South Atlantic Ocean during polar summer, Environ. Sci. Technol., 37, 22–31, 2003a.

Temme, C., Slemr, F., Ebinghaus, R., and Einax, J. W.: Distribution of mercury over the Atlantic Ocean in 1996 and 1999–2001, Atmos. Environ., 37, 1889–1897, 2003b.

10 Priddle, J. (Ed.): Regionally based assessment of persistent toxic substances - Antarctica regional report, United Nations Environmental Programme – Chemicals, Geneva, Switzerland, 2002

Xia, C., Xie, Z., and Sun, L: Atmospheric mercury in the marine boundary layer along a cruise path from Shanghai, China to Prydz Bay, Antarctica, Atmos. Environ., 44, 1815–1821, 2010.

---

**Atmospheric mercury observations from Antarctica**

K. Aspö Pfaßhuber et al.

---

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Atmospheric mercury observations from Antarctica**

K. Aspö Pffaffhuber et al.




**Fig. 1.** (a) Box and whisker plot presenting the concentration distribution of GEM measurements at TRS (Antarctic) and Zeppelin (Arctic). (b) Box and whisker plot presenting the monthly GEM concentration distribution from all the data collected at TRS. The middle line in the box shows the median concentration, the box indicates the range between the 25th and the 75th percentile. Whiskers above and below the box indicate the 90th and 10th percentile, respectively. The filled square shows the arithmetic mean.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Atmospheric mercury observations from Antarctica**

K. Aspö Pfaßhuber et al.




**Fig. 2.** Comparison of the time series of GEM concentrations from TRS in the Antarctic and Zeppelin in the Arctic. The time series from Zeppelin is shifted 182 days as compared to the TRS time series such that the seasons coincide.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)


## Atmospheric mercury observations from Antarctica

K. Aspö Pfaffhuber et al.



**Fig. 3.** Fields of  $R_{90}$  (a–d) and  $R_{10}$  (e–h) for GEM measurements at TRS from 2007 through June 2010. The location of TRS is marked with a white asterisk. Areas where  $S_7$  is below the threshold are plotted white.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

**Atmospheric mercury  
observations from  
Antarctica**K. Aspö Pfaffhuber  
et al.

**Fig. 4.** Fields of  $R_{90}$  (a–d) and  $R_{10}$  (e–h) for ozone measurements at TRS from 2007 through June 2010. The location of TRS is marked with a white asterisk. Areas where  $S_7$  is below the threshold are plotted white.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)