Atmos. Chem. Phys. Discuss., 11, 5867–5933, 2011 www.atmos-chem-phys-discuss.net/11/5867/2011/ doi:10.5194/acpd-11-5867-2011 © Author(s) 2011. CC Attribution 3.0 License.



This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

# The Pasadena Aerosol Characterization Observatory (PACO): chemical and physical analysis of the Western Los Angeles Basin aerosol

S. P. Hersey<sup>1</sup>, J. S. Craven<sup>2</sup>, K. A. Schilling<sup>2</sup>, A. R. Metcalf<sup>1</sup>, A. Sorooshian<sup>3</sup>, M. N. Chan<sup>1</sup>, R. C. Flagan<sup>1,2</sup>, and J. H. Seinfeld<sup>1,2</sup>

 <sup>1</sup>Division of Engineering and Applied Science, California Institute of Technology, Pasadena, CA, USA
 <sup>2</sup>Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena, CA, USA
 <sup>3</sup>Department of Chemical and Environmental Engineering, University of Arizona, Tucson, AZ, USA

Received: 3 February 2011 - Accepted: 4 February 2011 - Published: 18 February 2011

Correspondence to: J. H. Seinfeld (seinfeld@caltech.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Discussion Pa	<b>ACPD</b> 11, 5867–5933, 2011							
per   Di	Pasadena, CA aerosol characterization							
iscussion	S. P. Hersey et al.							
Pane	Title Page							
	Abstract	Introduction						
7	Conclusions	References						
iscuss	Tables	Figures						
ion P	I	►I.						
aner	•	•						
_	Back	Close						
Disci	Full Scre	en / Esc						
Ission	Printer-frier	dly Version						
Рар	Interactive	Discussion						
Ð		$\bigcirc$						

# Abstract

5

The Pasadena Aerosol Characterization Observatory (PACO) represents the first major aerosol characterization experiment centered in the Western/Central Los Angeles Basin. The sampling site, located on the campus of the California Institute of Technology in Pasadena, was positioned to sample a continuous afternoon influx of transported urban aerosol with a photochemical age of 1–2 h and generally free from major local contributions. Sampling spanned 5 months during the summer of 2009, which were broken into 3 regimes on the basis of distinct meteorological conditions. Regime I was characterized by a series of low pressure systems, resulting in high humidity and rainy periods with clean conditions. Regime II typified early summer meteorology,

and rainy periods with clean conditions. Regime II typified early summer meteorology, with significant morning marine layers and warm, sunny afternoons. Regime III was characterized by hot, dry conditions with little marine layer influence.

Organic aerosol (OA) is the most significant constituent of Los Angeles aerosol (42, 43, and 55% of total submicron mass in regimes I, II, and III, respectively), and that

- the overall oxidation state remains relatively constant on timescales of days to weeks  $(O:C = 0.44 \pm 0.08, 0.55 \pm 0.05, and 0.48 \pm 0.08$  during regimes I, II, and III, respectively), with no difference in O:C between morning and afternoon periods. Periods characterized by significant morning marine layer influence followed by photochemically favorable afternoons displayed significantly higher aerosol mass and O:C ratio,
- <sup>20</sup> suggesting that aqueous processes may be important in the generation of secondary aerosol and oxidized organic aerosol (OOA) in Los Angeles.

Water soluble organic mass (WSOM) reaches maxima near 14:00–15:00 local time (LT), but the percentage of AMS organic mass contributed by WSOM remains relatively constant throughout the day. Sulfate and nitrate reside predominantly in accumulation

<sup>25</sup> mode aerosol, while afternoon SOA production coincides with the appearance of a distinct fine mode dominated by organics. Particulate  $NH_4NO_3$  and  $(NH_4)_2SO_4$  appear to be  $NH_3$ -limited in regimes I and II, but a significant excess of particulate  $NH_4^+$  in the hot, dry regime III suggests less marine  $SO_4^{2-}$  and the presence of organic amines.



Positive Matrix Factorization (PMF) analysis of C-ToF-AMS data resolved three factors, corresponding to a hydrocarbon-like OA (HOA), semivolatile OOA (SV-OOA), and low-volatility OOA (LV-OOA). HOA appears to be a periodic plume source, while SV-OOA exhibits a strong diurnal pattern correlating with ozone. Peaks in SV-OOA con-

centration correspond to peaks in DMA number concentration and the appearance of a fine organic mode. LV-OOA appears to be an aged accumulation mode constituent that may be associated with aqueous-phase processing, correlating strongly with sulfate and representing the dominant background organic component.

Filter analysis revealed a complex mixture of species during periods dominated by SV-OOA and LV-OOA, with LV-OOA periods characterized by shorter-chain dicarboxylic acids (higher O:C ratio), as well as appreciable amounts of nitrate- and sulfatesubstituted organics. Phthalic acid was ubiguitous in filter samples, suggesting that

PAH photochemistry may be an important SOA pathway in Los Angeles. Water uptake characteristics indicate that hygroscopicity is largely controlled by or-

<sup>15</sup> ganic mass fraction (OMF). The hygroscopicity parameter  $\kappa$  averaged 0.31±0.08, approaching 0.5 at low OMF and 0.1 at high OMF, with increasing OMF suppressing hygroscopic growth and increasing critical dry diameter for CCN activation ( $D_d$ ).

Finally, PACO will provide context for results forthcoming from the CalNex field campaign, which involved ground sampling in Pasadena during the spring and summer of 2010.

# 1 Introduction

20

25

Over half the world's population lives in urban areas, and that fraction is expected to increase in coming decades (Bremner et al., 2009). There have been a number of recent coordinated studies focused on characterizing particulate air quality in major urban areas, such as Pittsburgh, PA (Pittsburgh Air Quality Study, PAQS; e.g., Pekney et al., 2006; Bein et al., 2006; Wittig et al., 2004; Cabada et al., 2004; Modey et al., 2004), Los Angeles, CA (Southern California Air Quality Study, SCAQS; e.g., Elder-



ing et al., 1994; Watson et al., 1994; Chow et al., 1994; Turpin and Huntzicker, 1991; Southern California Ozone Study 1997, SCOS97-NARSTO; e.g., Croes and Fujita, 2003; Liu et al., 2000; Pastor et al., 2003; Hughes et al., 2002; Liu et al., 2000; Secondary Organic Aerosol in Riverside, SOAR; e.g., Docherty et al., 2008; Eatough et al.,

- <sup>5</sup> 2008; Denkenberger et al., 2007), Mexico City, Mexico (Megacity Initiative: Local and Global Research Observations, MILAGRO; e.g., DeCarlo et al., 2008; Stone et al., 2008; Aiken et al., 2009), and Beijing, China (Campaign of Air Quality Research in Beijing, CAREBEIJING; e.g., Matsui et al., 2009; Yue et al., 2009; van Pinxteren et al., 2009).
- <sup>10</sup> The Los Angeles metropolitan area, with a population of over 17 million people, has long been considered a laboratory for air quality. Once the quintissential example of classic photochemical smog, the Los Angeles area has, as a result of decades of aggressive emission controls, reached a point where 8-h US National Ambient Air Quality Standard ozone exceedances have decreased from over 200 days/year in the
- <sup>15</sup> 1980s to 120 in 2008 (AQMD, 2010). Nonetheless, this area remains one in which particulate matter levels continue to regularly exceed air quality standards, with annual PM<sub>2.5</sub> averages exceeding the national standard every year from 2000–2008 in Los Angeles County (CARB, 2010).

The Los Angeles Basin is generally characterized as source-rich in its western area,
 with prevailing west and southwest winds transporting emissions downwind toward the east and northeast (Lu and Turco, 1995). Three major coordinated studies have been undertaken to characterize Los Angeles air quality, in addition to ongoing work to describe facets of Los Angeles aerosol (Table 1). The majority of previous studies in the Los Angeles Basin have been centered in the downwind, eastern part of the basin, roughly 60 km east of downtown Los Angeles. The aerosol in this region is predomi-

nantly aged, with local contributions as well.

A major source of gaseous and particulate pollution in the Los Angeles area is mobile source emissions. The Los Angeles aerosol undergoes dramatic evolution from nighttime/morning periods to the photochemically active afternoon, characterized by



changes in both its microphysical properties and composition. Overall, the submicron aerosol is dominated by organic aerosol (OA), which is predominantly secondary in origin (SOA), and nitrate (Docherty et al., 2008), with smaller amounts of sulfate and a minor chloride contribution. Because of strict  $SO_2$  controls in the basin, sulfate is primarily the result of marine sources (Watson et al., 1994).

5

10

The Pasadena Aerosol Characterization Observatory experiment (PACO) was conducted from May–August 2009 in Pasadena, CA, in the north-central/north-western portion of the Los Angeles Basin, 16 km NE of downtown Los Angeles. Aerosol physics, composition, and hygroscopicity were measured during 54 days that spanned three distinct regimes of the annual meteorological cycle. A major forest fire occurred at the end of the sampling period, starting on 26 August and consuming over 160 000

acres of the Angeles National Forest before containment on 16 October. Named the "Station Fire", its southernmost extent was roughly 8 km from the sampling site, and emissions from the fire impacted the sampling site with periodic heavy smoke, soot, and ash. Results from PACO sampling during the Station Fire will be presented in subsequent work.

PACO represents the first prolonged experiment aimed at characterizing the aerosol near the source-rich western/central part of the Los Angeles basin; comparisons with previous studies in downwind areas will help determine the rate of aerosol generation

from Los Angeles sources, as well as the nature of aerosol transformations with downwind transport. Further, the PACO study was well-timed to precede the 2010 CalNex field campaign (http://www.esrl.noaa.gov/csd/calnex/).

This paper presents an overview of the PACO experiment. Aerosol number and volume distributions measured by a differential mobility analyzer (DMA) exhibit dis-

tinct diurnal patterns, and composition data from an Aerodyne Compact-Time-of-Flight Aerosol Mass Spectrometer (C-ToF AMS) indicate that these patterns are associated largely with the production of semivolatile oxidized organic species during photochemically active daytime periods. AMS-based Positive Matrix Factorization (PMF) results indicate that the character of OA in Los Angeles is overwhelmingly oxidized and



secondary, and the high time resolution data allow for investigation of the daily evolution of OA components. Ultra Performance Liquid Chromatography/Electrospray Ionization Time-of-Flight High-Resolution Mass Spectrometry (UPLC/ESI-TOFMS) analysis of high-volume filter samples provides further evidence for diurnal changes in the charac-

- ter of OA, while thermal-optical analysis of low-volume, denuded filter samples reveals the magnitude of organic and elemental carbon in morning versus afternoon sampling periods. A particle-into-liquid sampler with offline ion chromatography analysis (PILS-IC) provides quantitative water-soluble inorganic ion composition during PACO, while a PILS coupled with online total organic carbon analysis (PILS-TOC) measured total
- <sup>10</sup> water soluble organic carbon measurements during the third regime. Finally, trends in aerosol hygroscopicity indicate the degree to which changes in the magnitude and character of OA affect particle water uptake. PACO results are compared with previous aerosol characterization experiments in the Los Angeles Basin, with major differences attributed to regional variations in both source profiles and degree of aging.

#### 15 2 Methods

Sampling was conducted on the Caltech campus in Pasadena, at 34.138° N, 118.124° W. Caltech is located in the western/central portion of the Los Angeles Basin, 8 km south of the San Gabriel Mountains and 16 km NE of downtown Los Angeles in the San Gabriel Valley. The 210 freeway is located 1.5 km north of the site, with
heavy traffic between the hours of 08:00 and 09:00 LT, and then again between 15:00 and 19:00 LT. Prevailing daytime winds are from the W or SW, minimizing the impact of primary emissions from the 210 freeway at the sampling site, but overnight NW winds occasionally bring freeway emissions to Caltech. Other than this periodic impact from local primary emissions, the Pasadena aerosol is representative of transported urban particulate pollution from near the source-rich downtown area of the Los Angeles

<sup>25</sup> ban particulate pollution from near the source-rich downtown area of the Los Angeles Basin.



PACO sampling took place between May and August of 2009, spanning the transition from spring to summer meteorological regimes. PACO was separated into three distinct analysis periods on the basis of meteorology, as summarized in Table 2.

With the exception of a high-volume filter sampler located outdoors on the roof, instruments sampled from inlets connected to a main sampling line drawing in ambient air at a 16 liters per minute (L min<sup>-1</sup>) laminar flow rate. Sample flow to each instrument was supplemented with pumps to be isokinetic with flow in the main sampling line, and sample lines were designed to draw from the center of flow through the main line. Ambient air was dried by nafion driers upstream of each instrument.

# 10 2.1 Meteorology

Hourly meteorological data were downloaded from the Remote Automatic Weather Stations (RAWS) archive from the Western Regional Climate Center at the Desert Research Institute (DRI, Reno, Nevada). Data are from the Santa Fe Dam station, approximately 16 km east of the sampling site in the San Gabriel Valley, and are available online: http://www.raws.dri.edu/cgi-bin/rawMAIN.pl?caCSFD.

# 2.2 Gas-phase data

25

Hourly  $O_3$ ,  $NO_x$ , and CO data were obtained from the California Air Resources Board's Pasadena (South Wilson) monitoring site located on the Caltech campus. Data are available online at http://www.arb.ca.gov/adam/hourly/hourly1.php.

# 20 2.3 Differential Mobility Analyzer

Particle size distribution measurements were performed with a cylindrical scanning differential mobility analyzer (TSI Model 3081) upstream from a condensation particle counter (TSI Model 3760). A logarithmic scan from a mobility diameter of 15 nm to 815 nm was completed every 237 s. The sheath and excess flows of  $2.5 L \text{ min}^{-1}$  were used, with a 10:1 flow rate ratio of sheath-to-aerosol.



# 2.4 Aerosol Mass Spectrometer

An Aerodyne compact time-of-flight aerosol mass spectrometer (C-ToF-AMS; Drewnick et al., 2005; Canagaratna et al., 2007) measured aerosol composition and size information for 54 days during PACO. The AMS collects sub-micron particles via an aero-

- <sup>5</sup> dynamic lens into a particle sizing chamber, after which particles are vaporized and ionized by a heater and filament assembly. Aerosol fragments are then orthogonally extracted into an ion time-of-flight chamber where they are detected and interpreted as mass spectra. Adjustments to the AMS fragmentation table were made regularly, based on 21 filter sample periods over the course of PACO (Allan et al., 2004b). Ad-
- ditionally, thirteen ionization efficiency calibrations were made using ammonium nitrate solutions. The AMS sulfate measurement was compared to sulfate concentrations from the PILS-IC, resulting in an AMS collection efficiency (CE) of 0.5. This CE is in agreement with previous ambient studies (Huffman et al., 2005; Drewnick et al., 2003; Allan et al., 2004a). AMS data were processed with the ToF-AMS Analysis Toolkit in Igor Pro 6. (Wayamatrias, Lake Oswage, OR)
- <sup>15</sup> 6 (Wavemetrics, Lake Oswego, OR).

20

#### 2.5 Positive Matrix Factorization analysis

The dominant fraction of aerosol is organic in most urban areas (Zhang et al., 2007). OA comprises a multitude of compounds, such that characterization by identification of each component is not currently possible (Schauer et al., 1996). Positive Matrix Factorization (PMF) has been used to deconvolve AMS mass spectra in order to identify classes of organic compounds that constitute OA (Paatero and Tapper, 1994; Paatero, 1997; Ulbrich et al., 2009). PMF solutions represent linear, positive combinations of mass spectra that describe variability in data, and AMS-based PMF results from most sites resolve factors whose mass spectra have characteristics of hydrocarbon-like OA

(HOA) and oxygenated OA (OOA), with other factors such as biomass burning OA (BBOA) occurring in some areas (Zhang et al., 2007). OOA can often be further deconvolved into low-volatility OOA (LV-OOA) and semivolatile OOA (SV-OOA), broadly



representing aged, transported, and highly oxidized OA and fresher, local, and less oxidized OA, respectively (Ng et al., 2010). AMS measurements, in conjunction with PMF analysis, allow for investigation of the diurnal evolution of OA and rapid changes in OA occurring during photochemically intense periods. Further, the combination of PMF solutions with analysis of the relative abundance of mass-to-charge (m/z) ratios

<sup>5</sup> PMF solutions with analysis of the relative abundance of mass-to-charge (m/z) ratios 43 and 44 can be used as an aerosol aging diagnostic that allows comparison between locations (Ng et al., 2010).

For the PACO dataset, bulk AMS organic aerosol was prepared and then processed using the PMF2.exe algorithm (Paatero and Tapper, 1994). Solutions from the PMF algorithm were compared to meteorological variables and chemical tracers, as well as previously published mass spectra, using the PMF Evaluation Tool (PET Ulbrich et al., 2009). The strategic procedure for probing the PMF solution space is provided in Appendix A.

# 2.6 PILS-IC

10

- A particle-into-liquid sampler coupled with off-line ion chromatography (PILS-IC, Sorooshian et al., 2006) sampled during 4-h periods (07:00–11:00 and 15:00–19:00 LT), providing quantitative measurements of inorganic and organic ions. The PILS samples ambient air through a 1 μm cut size impactor and three denuders (URG and Sunset Laboratory) designed to remove inorganic and organic gases that might bias aerosol measurements. Sampled air is then exposed to high water supersaturation in a steam chamber, where particles grow sufficiently large to be collected by inertial impaction before being delivered to vials held on a rotating carousel. Samples were collected every 30 min, and were stored at 2 °C until analysis by a dual IC system (Dionex ICS-2000). PILS-IC data from PACO are used primarily to determine collection officiency corrections for the AMS and are not presented in detail here.
- <sup>25</sup> efficiency corrections for the AMS and are not presented in detail here.



# 2.7 PILS-TOC

Water-soluble organic carbon (WSOC) was quantified during 12-h periods (07:00–19:00 LT) using a PILS (Brechtel Manufacturing Inc.) coupled to a Sievers Model 800 Total Organic Carbon (TOC) Analyzer (PILS-TOC). Details of this technique are described in detail by Sullivan et al. (2004, 2006). Briefly, particles smaller than 2.5 µm in diameter are sampled by the PILS and passed immediately through an organic carbon denuder (Sunset Laboratory Inc.) to remove organic vapors. Particles are grown into droplets, collected by inertial impaction, and delivered to a TOC analyzer for quantification of WSOC, with data reported every 6 min. To account for dilution of the PILS liquid
wash flow to the TOC analyzer owing to collected drops and condensation on the PILS droplet impactor, a constant dilution factor of 1.15 is assumed. This factor is based on measurements with an independent PILS system coupled to ion chromatography, a technique that can more precisely quantify the dilution factor. The reported WSOC levels are the difference between the measured and background concentrations. Con-

tamination in the MQ water was the main component of the background levels, which were assumed to be constant between periodic background measurements. The overall measurement uncertainty is estimated to be approximately 10%. The PILS-TOC was employed during the final, photochemically active regime of PACO, from 10 July to 4 August.

#### 20 2.8 Filter sampling

25

Both high- and low-volume filter samples were collected during 4-h sampling periods (07:00–11:00 and 15:00–19:00 LT). High-volume ( $1250 L min^{-1}$ ) samples were collected with a Tisch TE-6001 sampler fitted with a PM<sub>2.5</sub> fractionater, using Whatman Quartz Microfibre Filters (cat. #1851-865). Low volume ( $15.9 L min^{-1}$ ) samples were collected on 47 mm PALL Life Sciences Pallflex membrane filters, downstream from a Sunset Laboratory parallel-plate carbon filter denuder. Sampling was carried out according to the IMPROVE\_A protocol, except that oven limitations forced filters to be



pre-baked at 300 °C before sampling. Despite the lower pre-baking temperature, subsequent analysis of pre-baked filters revealed  $\leq 0.01 \,\mu g$  of organic carbon (OC) and elemental carbon (EC). After sampling, filters were wrapped in two layers of aluminum foil and stored at -20 °C until analysis.

- <sup>5</sup> Low- and high- volume samples were analyzed for OC and EC via thermal-optical analysis (TOA) with a Sunset OC-EC TOA analyzer, following the IMPROVE\_A temperature and calibration protocols. High volume quartz filters were also analyzed by Ultra Performance Liquid Chromatography/Electrospray Ionization Time-of-Flight High-Resolution Mass Spectrometry (UPLC/ESI-TOFMS). One-eighth of each filter was ex-
- tracted with high-purity methanol under ultrasonication for 45 min. Filter extracts were filtered through a PALL Life Sciences Acrodisc CR 25-mm syringe filter (PTFE membrane, 0.2-mm pore size) into a scintillation vial. The filtered extracts were blown dry under a gentle nitrogen stream at ambient temperature. The residue was reconstituted with 50:50 v/v methanol with 0.1% acetic acid and water. Blank filters were extracted
- and treated in the same manner as the field samples. Extracts were analyzed by a Waters ACQUITY ultra performance liquid chromatography (UPLC) system, coupled with a Waters LCT Premier TOF mass spectrometer equipped with an electrospray ionization (ESI) source, allowing for accurate mass measurements (i.e., determination of molecular formulas) to be obtained for each observed ion. Extracts were analyzed by
- <sup>20</sup> UPLC/ESI TOFMS operated in both negative and positive ion modes. Details of operation protocols, including column information and chromatographic method for the UPLC/ESI-TOFMS are given by Surratt et al. (2008).

# 2.9 Hygroscopicity

Hygroscopicity measurements were carried out with the Differential Aerosol Sizing and
 Hygroscopicity Spectrometer Probe (DASH-SP, Brechtel Mfg), which is described by
 Sorooshian et al. (2008). Briefly, ambient particles pass through a nafion dryer before
 size-selection by a cylindrical, single-classification differential mobility analyzer (DMA).
 The resulting monodisperse aerosol is split into five separate flows – one providing



a redundant measurement of total particle concentration at the DMA-selected size with a water condensation particle counter (TSI Model 3831), and the remaining four channels consisting of parallel nation humidification chambers (Perma Pure, LLP, Model MD-070-24FS-4), followed by correspondingly humidified custom optical particle coun-

- <sup>5</sup> ters (OPCs). In the OPC sample volume, particles pass through a focused laser beam ( $\lambda = 532$  nm, World Star Technologies, Model TECGL-30) and scatter light in proportion to size ( $D_p$ ) and refractive index (RI). Forward-scattered light is collected and focused on a photomultiplier tube, and the resulting electrical pulse is recorded by a high-speed data acquisition computer. An iterative data processing algorithm, based on laboratory
- <sup>10</sup> calibrations with salts of known refractive indices, is used to determine the best fit on a solution surface relating electrical pulse height, size, and refractive index. The hygroscopic growth factor (GF =  $D_{p,wet}/D_{p,dry}$ ) is corrected for the RI change caused by particulate liquid water at elevated RH.

In the current study, the DASH-SP sampled for 4-h periods (07:00–11:00 and 15:00– 19:00 LT). Hygroscopicity was measured at dry sizes corresponding to *D*<sub>p,em</sub> of 150, 175, 200, and 225 nm. Multiple RH sensors in the nafion tubes and OPCs controlled RHs to dry (≤ 8%), 74%, 85%, and 92%, with RH uncertainty of 1.5%. Because calculated RI varied little during sampling periods, GF calculations were made with a single RI for each sampling period, representing the average RI calculated for the entire period. The error introduced by this assumption is small, as light scattering in the DASH

is far more sensitive to size than RI. Overall uncertainty in GF calculations is calculated to be 4.5%.

#### 3 Results and discussion

# 3.1 Atmospheric conditions

<sup>25</sup> Figure 1a-c is a compilation of meteorological data, as well as gas- and aerosolphase composition for regimes I, II, and III, respectively. Wind speed and direction



display distinct diurnal patterns, with stagnant or very light winds from the N/NE in the overnight to early morning hours (20:00–06:00 LT) changing direction to come from the W/SW shortly after sunrise (06:00–07:00 LT). Between 06:00 and 16:00 LT, wind speed increases from 3–4 km h<sup>-1</sup> to 13–15 km h<sup>-1</sup> out of the W/SW before decreasing back to
very light or stagnant out of the N/NE by 20:00 LT. One effect of these wind patterns is to allow local emissions to build up in the Los Angeles Basin during the nighttime and early morning hours, leading to significant carryover of aged aerosol in the background Los Angeles air, as proposed by Blumenthal et al. (1978). When sea breezes develop, fresh emissions in source-rich areas are gradually transported from the Western Los
Angeles Basin toward downwind areas in the E/NE. Situated 16 km from downtown

- Los Angeles, Pasadena can be considered as a receptor site that first receives transported pollution between 10:00 and 11:00 LT, after roughly 4–5 h of transport. During the afternoon hours (13:00–18:00 LT), the sampling site receives a steady influx of air from the W/SW with photochemical age of 1–2 h. There are no significant freeways or
- point sources of particulate pollution in close W/SW proximity to the sampling site, and so it is expected that air sampled during PACO is representative of transported urban air in Los Angeles. A major freeway with significant diesel and gasoline mobile sources is located 1.5 km to the north of the sampling site, and so overnight N/NE winds may have periodically impacted the site with primary particulate emissions. Daytime varia-
- tion in gas- and aerosol-phase composition is expected to result from three sources:
   (1) temporal changes in emissions from source-rich areas, (2) diurnal variations in the intensity of photochemistry, and (3) the age of transported pollution.

A clear diurnal trend is observed in RH and solar radiation, with RH increasing to near 100% on most nights and remaining at or near saturation until sunrise. The incidence

of nighttime and morning marine cloudcover was common during regimes I and II, while regime III was typically drier, with clear nighttime skies more common. Periods of more significant marine layer influence are indicated by longer periods of saturated RH and more gradual decrease in RH (and slower increase in solar radiation) after sunrise. Regime I has been classified as "springtime meteorology" due to the impact



of several low pressure systems in Southern California, bringing periodic drizzle and leading to several days of cloudcover and significantly diminished solar radiation (i.e. 6 to 12 June). Daytime RH rarely fell below 50% during regime I. Regime II exhibited the strongest summertime marine layer influence, with heavy cloudcover persisting
until late morning (10:00 to 12:00 LT) nearly every day before making way for sunny, hazy afternoons with daytime RH between 25 and 40%. Regime III observed the least marine layer influence, with many mornings of clear skies and daytime RH typically between 10 and 30%. Despite differences in meteorology, the diurnal wind patterns described above were robust, with stagnant morning winds giving way to afternoon
sea breezes and pollutant transport from the W/SW.

Gas-phase data give a sense of the atmospheric chemical environment during sampling. O<sub>3</sub> exhibits a strong diurnal pattern correlated with solar radiation and serves as a tracer for photochemical activity. During periods of relatively dry atmospheric conditions, the O<sub>3</sub> concentration peak increases in magnitude over subsequent days (see 17 to 19 July and 27 July to 2 August in regime III), suggesting that either pollutants accumulate or photochemical activity increases during periods of consistent, stable atmospheric conditions, resulting in photochemical episodes. CO and NO<sub>x</sub> display more random behavior, exhibiting periodic peaks in concentration indicative of plumes of fresher emissions.

#### 20 3.2 Aerosol composition

OC and EC mass concentrations, in addition to OC:EC ratios from representative days in each regime, are presented in Table 3. OC is higher in afternoon periods when compared with mornings in each regime (27, 18, and 4% higher in regimes I, II, and III, respectively), while EC is lower during afternoons (7, 35, and 22% lower in regimes I, 1, and III, respectively). Further, OC:EC ratios are significantly higher during afternoon periods. These diurnal trends in OC are expected, given the importance of photochemical SOA production, and the percentage increase in OC mass between morning and afternoon periods is similar to that observed by the AMS (15, 17, and 10% higher in



regime I, II, and III, respectively). Discrepancies may arise due to the different size ranges measured by the AMS and filter sampling (upper limits of measurement are 1.0 and 2.5  $\mu$ m for the AMS and filters, respectively), in addition to potential volitalization of semivolatile organics relating to the thermo-optical OC/EC analysis technique.

<sup>5</sup> The modest increase in OC between morning and afternoon periods during the photochemically intense regime III may indicate that significant OC existed in the background aerosol during this period. The decrease in EC observed between morning and afternoon periods suggests that primary organics are a more significant portion of the Los Angeles aerosol in the mornings, and the relatively low concentrations of EC suggest that the dominant source of organic aerosol in Los Angeles is secondary.

DMA data are presented in panel 8 of Fig. 1a–c. Diurnal patterns are evident in both DMA volume and number in all regimes. DMA volume tends to reach a maximum in the mid-to-late morning (10:00, 11:00, 08:00 LT in regime I, II, and III, respectively), when RH is still high and particles have presumably undergone cloud processing. The

- early maxima in regime III underscore the absence of significant marine layer influence persisting into the late morning. On average, DMA number concentration reaches a maximum between 14:00 and 15:00 LT during each regime, roughly 3 h after transported pollution is expected to first arrive at the sampling site (Fig. 2). This indicates that increases in fine particle mass at the PACO sampling site are not the result of
- transport alone, but also affected by photochemical production in transported, polluted air from source-rich areas. There was no significant difference in wind patterns between regimes, so the slightly earlier peak in DMA number in regime II suggests that marine moisture and aqueous-phase processes (most prevalent in regime II) may hasten secondary aerosol production. Figure 2 indicates that appreciable concentrations
- of fine particles (< 100 nm) are present in late nights and early mornings (particularly in regimes II and III), possibly comprised of fresh emissions, nighttime-generated NO<sub>3</sub><sup>-</sup>, or fine particle carryover residing in a shallow mixed layer.

The third and fourth panels of Fig. 1a–c show time traces of bulk AMS aerosol mass concentration and mass fraction, respectively, while the sixth panel shows a time trace



of approximate O:C ratio, based on Aiken et al. (2008), as well as WSOC concentration from the PILS-TOC (regime III). Organics dominated the aerosol sampled during PACO, and except for short episodes in regime I, the approximate O:C ratio remained relatively constant during each regime ( $0.44 \pm 0.08$ ,  $0.55 \pm 0.05$ , and  $0.48 \pm 0.08$  during  $^{5}$  regimes I, II, and III, respectively). Organic aerosol in regime II appears to be more oxidized than I and III, suggesting that the marine moisture and presumed aqueous processing may enhance the rate of organic oxidation in Los Angeles. From bulk AMS time traces it is evident that organics increase in both magnitude and mass fraction each afternoon, particularly during the more photochemically active regimes II and III. It is noteworthy that there are no distinct trends in approximate O:C between mornings and afternoons (morning approximate O:C =  $0.44 \pm 0.08$ ,  $0.54 \pm 0.08$ , and  $0.49 \pm 0.07$ 

for regimes I, II, and III, respectively, while afternoon approximate  $O:C = 0.44 \pm 0.08$ ,  $0.53 \pm 0.06$ ,  $0.50 \pm 0.08$  for regimes I, II, and III, respectively). This suggests that there is little change in overall oxidation state of Los Angeles aerosol, even on timescales of <sup>15</sup> weeks. It is unlikely that all organics are uniformly oxidized at all times in Los Angeles,

but rather that there exists a continuum of OA oxidation, with more reduced and more oxidized ends of that continuum maintaining a relatively constant ratio for extended periods of time.

Trends in aerosol mass fraction are presented in Fig. 4. Organics dominate the Los Angeles submicron aerosol, contributing 42, 43, and 55% of aerosol mass in regimes I, II, and III, respectively. Diurnal trends are evident in Fig. 4, with organic mass fraction increasing significantly from morning (07:00–11:00 LT) to midday (11:00–15:00 LT) to afternoon (15:00–19:00 LT) in each regime. Sulfate is a significant component in marine/moisture-influenced regimes I and II, and is a more minor component in regime

III, suggesting that cloud processing and marine influence may be an important source of sulfate aerosol in Los Angeles. Nitrate contributes equal mass fractions in each regime, with a minimum in the afternoon and a maximum in the morning. Ammonium contributes a remarkably constant mass fraction to the aerosol, despite significantly less inorganic sulfate in regime III, suggesting an ammonium surplus for aerosol





sampled in regime III. The significant peak in AMS masses during the early morning of 5 July (regime II) corresponds to a total submicron mass of over  $200 \,\mu g \,m^{-3}$ , and resulted from a myriad of fireworks displays in the area surrounding the sampling site. This plume has been removed from further analysis.

- Diurnal trends in bulk composition are highlighted in the top panel of Fig. 3. It is 5 noteworthy that during the significantly marine-layer-impacted regime II, bulk organic, sulfate, nitrate, and ammonium masses are all enhanced. This is further evidence that aqueous processing associated with marine layer influence may be an important factor in generating aerosol mass in Los Angeles. Typically, organics increase in magnitude
- to a maximum near 14:00–15:00 LT in each regime, corresponding to a maximum in 10 ozone, solar radiation, and DMA number – evidence of photochemical SOA production. Sulfate exhibits slight enhancements during periods of peak photochemical activity in all three regimes, but as suggested by Kleeman et al. (1999), it appears to be predominantly a background marine species. Nitrate mass is highest in the nighttime
- and morning periods, likely resulting from ozone-NO<sub>v</sub> production of HNO<sub>3</sub> at night and 15 HO<sub>x</sub>-NO<sub>x</sub> production of HNO<sub>3</sub> after rush hour in the mornings. As mentioned in Sect. 3.1, there were periods of high  $O_3$  observed during PACO, and during several of these periods there also appeared to be significantly elevated levels of aerosol nitrate during nighttime sampling (e.g. 30 July to 3 August). Further, morning peaks in aerosol nitrate varied significantly in magnitude, suggesting that there may be significant variation in 20
- the amount of gas-phase HNO<sub>3</sub> production from day to day.

Ammonium mass generally follows the combined trends of  $SO_4^{2-}$  and  $NO_3^{-}$ , but appears to be in excess in regime III. Defining an ammonium ratio as:

(moles of  $NH_4^+$  measured by AMS)

(moles of  $NH_4^+$  required to neutralize  $SO_4^{2-}$  and  $NO_3^-$ )

aerosol sampled in regimes I and II was found to be generally neutralized (average 25 ammonium ratio =  $0.96 \pm 0.21$  and  $0.94 \pm 0.17$  for regimes I and II, respectively), while aerosol from regime III was characterized by significant ammonium excess (ammonium 5883



ratio = 1.28  $\pm$  0.30). These results suggest that aerosol phase NH<sub>4</sub>NO<sub>3</sub> was ammonialimited during regimes I and II, but ammonia may have been in excess during regime III. Investigating molar ratios of inorganic species, it is apparent that the role of  $SO_{A}^{2-}$ is significantly diminished in regime III, corresponding to the absence of marine layer <sup>5</sup> influence. The ratio of  $NO_3^-$  to  $NH_4^+$  remains relatively constant across regimes (0.32, 0.29, and 0.34 in regimes I, II, and III, respectively), while the ratio of  $SO_4^{2-}$  to  $NH_4^+$ exhibits a marked decrease in regime III (0.42, 0.41, and 0.24 in regimes I, II, and III, respectively). The AMS has difficulty distinguishing whether NH,NH<sub>2</sub>, and NH<sub>2</sub><sup>+</sup> (m/z 15, 16, and 17, respectively) are contributed by inorganics or organic amines. It is hypothesized that ammonium is chiefly inorganic during the moister, marine-layer-10 influenced regimes I and II, serving primarily to neutralize  $SO_4^{2-}$  and  $NO_3^{-}$ , while there may be significant amounts of organic amines in the absence of marine layer influence during regime III. It is also possible that these organic amines are present in all regimes, which would imply that inorganic  $SO_4^{2-}$  and  $NO_3^{-}$  are not fully neutralized by  $NH_4^+$  during

regimes I and II. 15

> The bottom panel of Fig. 3 shows size-resolved PToF-AMS composition for a representative morning and afternoon period in each regime, with 9 June, 19 June, and 19 July representing regimes I, II, and III, respectively. Size-resolved data indicate that aerosol in regime I is typically internally-mixed, with the mode centered at  $\sim 400-$

- 500 nm. Afternoon PToF data from regime I indicate that photochemical organic pro-20 duction typically results in organics condensing on existing accumulation mode aerosol. Regime II is characterized by significantly higher aerosol mass and more complex size distributions. While there is a significant internally-mixed accumulation mode present and centered at ~ 600-700 nm, there is also significant aerosol mass at sizes smaller
- than 300 nm in both morning and afternoon sampling periods, though the mass at 25 smaller sizes does not comprise a distinct mode. Afternoon PToF data from regime Il indicate that a fine mode appears coincident with photochemistry, suggesting that fine mode aerosol is predominately comprised of secondary species. Nitrate, sulfate, and ammonium are generally confined to accumulation mode aerosol in regime II.



Regime III PToF data reveal an aerosol that is dominated by organics at all sizes and is distinctly bimodal. In addition to an internally mixed accumulation mode centered at ~ 500–600 nm, distributions from regime III also reveal a significant fine mode centered at ~ 100–200 nm. The accumulation mode tends to be comprised of sulfate, nitrate, and ammonium (in addition to organics) in the morning periods, with less nitrate in afternoons. Afternoon PToF data from regime III, like those from regime II, reveal

- fine-mode aerosol growth resulting from production of secondary organic species. Numerous studies have shown that the critical species for new particle formation in the atmosphere is  $H_2SO_4$ . The extent to which the observed fine mode aerosol might have
- originated from the nucleation of gas-phase species during PACO is unclear. However, given the relatively large amount of pre-existing aerosol surface area, as well as the relatively low concentration of SO<sub>2</sub>, the precursor to H<sub>2</sub>SO<sub>4</sub>, it is unlikely that new particle formation by nucleation is an important process in Los Angeles. Instead, the growth of fine-mode organics is proposed to occur by condensation of SOA on existing ultrafine
- primary aerosol. This tentative conclusion is contrasted with results from Pittsburgh, in which Stanier et al. (2004) found new particle formation to be an important process, strongly correlated with SO<sub>2</sub> concentrations. Whether condensing on nucleated sulfate or existing ultrafine primary aerosol, growth of fine mode particles in Los Angeles is predominantly due to SOA.

Figure 5 shows the hourly average for WSOC mass (top panel, representing WSOC organic mass) and % of AMS organic accounted for by WSOC (bottom panel) for regime III. A factor of 1.8 was used to convert water soluble carbon (WSOC) mass to water soluble organic mass (WSOM), within the range of factors presented by Turpin and Lim (2001), and matching the factor used by Docherty et al. (2008) for aerosol in the Los Angeles Basin. Averaged hourly data reveal a trend in WSOM, increasing from a minimum in the late evening and early morning to a maximum at 15:00 LT, cor-

responding to maxima in  $O_3$  concentration, solar radiation, AMS organic, and DMA number concentration. This suggests that photochemical production of SOA in air transported from source-rich areas is a significant source of WSOM in the Los Angeles



Basin. While there is a diurnal trend in WSOM, the fraction of AMS organic accounted for by WSOM is relatively constant at  $0.44 \pm 0.16$  during regime III, with a slight enhancement between 15:00 and 19:00 LT. This suggests that while significant WSOM is photochemically generated during the course of a day, water soluble organics comprise

a relatively constant percentage of organic aerosol in Los Angeles. WSOC has been found to correlate well with oxidation state of organic aerosol (e.g., Kondo et al., 2007). Given the relatively constant approximate O:C ratio observed in regime III and steady afternoon influx of polluted air on the order of 1–2 h old, the constant WSOM:AMS organic ratio is not unexpected.

#### 10 3.3 PMF results

15

AMS data from PACO were combined into a single continuous dataset for analysis by the PMF method. PMF produced several non-unique solutions that were compared with known mass spectra and external time traces in order to choose final solutions. Figure 6 presents a summary of PMF factors, while panel 6 of Fig. 1a–c represents time traces for PMF factors for regimes I, II, and III, respectively. Details of the PMF method and solution selection are presented in Appendix A.

Three factors were resolved from the PMF method, and comparisons with previously published spectra indicate that they correspond to a hydrocarbon-like organic aerosol (HOA) and semivolatile- and low-volatility-oxygenated organic aerosol (SV-OOA and

- <sup>20</sup> LV-OOA, respectively). The HOA factor is closely correlated with previously published spectra from periods dominated by reduced, primary organics (Zhang et al., 2005; Lanz et al., 2007; Ulbrich et al., 2009), and has significant signal at *m/z* 41 and 57. The LV-OOA factor is closely correlated with previously-published OOA-1 factors and OOA-like factors observed in highly oxidizing environments (Zhang et al., 2005; Lanz
- et al., 2007; Ulbrich et al., 2009; Alfarra et al., 2004). The SV-OOA factor is also correlated with previously-published OOA factors, but has a higher degree of correlation with previously-published factors expected to represent less-aged, less-oxidized, semivolatile oxygenated organic aerosol, or SV-OOA (Zhang et al., 2005; Lanz et al.,



2007; Ulbrich et al., 2009; Bahreini et al., 2005; Alfarra et al., 2004). The LV-OOA factor has a larger ratio of *m/z* 44 to *m/z* 43, an indication of a more aged, oxidized organic component (Ng et al., 2010). The SV-OOA factor, on the other hand, has a smaller *m/z* 44 to *m/z* 43 ratio. PACO PMF factors correlated closely with "standard" HOA,
<sup>5</sup> SV-OOA, and LV-OOA factors reported by Ng et al. (2011), based on an average of results from 15 sites.

As seen in Fig. 1a–c, the HOA factor has the character of a periodic plume source, with low background concentrations punctuated by sudden, significant increases in concentration that remain high for several hours at a time. These increases in HOA concentration occurred primarily during regime I and are typically correlated with or just following peaks in CO concentration (panel 3 in Fig. 1a–c), though the low timeand concentration-resolution of CO measurements made statistical analysis difficult. HOA is typically associated with primary organic aerosol (POA) (e.g., Ulbrich et al., 2009; Ng et al., 2010), and the only significant local source of POA is expected to be

- <sup>15</sup> the 210 freeway, 1.5 km north of the sampling site. There appears to be a trend of HOA plumes coinciding with or immediately following periods of light winds out of the N, which suggests that the HOA-like factor may represent primary freeway emissions. It is unclear why this HOA factor does not appear during every period of light winds out of the N. Low correlation between the HOA factor and ozone (Pr = 0.39) suggests
- that HOA production is not a photochemical process, and low correlations with inorganic AMS traces (Pr = 0.18, 0.15, 0.21, and 0.02 for sulfate, nitrate, ammonium, and chloride, respectively) suggest that HOA production is independent of formation of inorganic aerosol, and that HOA may comprise an externally mixed mode, separate from the accumulation mode aerosol that is typically comprised of inorganic constituents.

<sup>25</sup> The more dominant LV-OOA factor is most significantly correlated with inorganic AMS traces (Pr = 0.75, 0.46, 0.65, and 0.46 for sulfate, nitrate, ammonium, and chloride, respectively). As discussed in Sect. 3.2, inorganics are most prevalent in accumulation mode aerosol (rather than fine mode), and their loadings appear to be associated with significant marine layer influence in the Los Angeles Basin (e.g., regimes I and II).



The high degree of correlation between LV-OOA and inorganic constituents suggests that the LV-OOA factor represents an organic component that typically resides in accumulation mode aerosol and may be associated with marine layer/cloud processing influence. The correlation between LV-OOA and ozone is relatively low (Pr = 0.39), indicative of an aged background organic species, as opposed to a rapidly-produced

- Indicative of an aged background organic species, as opposed to a rapidly-produced photochemical product that exhibits a distinct diurnal trend. Examining time traces for LV-OOA in Fig. 1a–c reveals that the LV-OOA factor exhibits relatively high background levels in the mornings and at night, while periodically showing moderate increases in magnitude during midday periods of photochemistry. These periodic photochemical
- trends in LV-OOA correspond to periods of intense photochemistry (e.g., 18–19 July), characterized by midday ozone concentrations near 80–100 ppb. This suggests that while LV-OOA appears to be the major constituent of the aged background aerosol discussed in Sect. 3.2, it can be photochemically produced on relatively short timescales during periods of intense photochemical activity. Its strong correlation with SO<sub>4</sub><sup>2-</sup>, con-
- sistent presence during humid overnight and morning hours, coincidence with accumulation mode aerosol, and greater prevalence in moisture-impacted regimes I and II suggest that there may be significant aqueous processing sources for LV-OOA in Los Angeles.

Figure 1a–c show a regular diurnal trend in SV-OOA, with overnight minima increasing to maximum concentrations daily at approximately 14:00-15:00 LT, correlating with maxima in solar radiation, AMS organic, and DMA number concentration. The SV-OOA factor exhibits a reasonable degree of correlation with ozone (Pr = 0.53), suggesting that semivolatile organics are products of short-timescale photochemical oxidation of VOCs. Correlations with inorganic traces are lower than those for LV-OOA (Pr = 0.42,

<sup>25</sup> 0.31, 0.45, and 0.19 for sulfate, nitrate, ammonium, and chloride, respectively). As discussed in Sect. 3.2, aerosol size distributions exhibit significant enhancements in fine mode aerosol in afternoon periods, with the fine mode comprised almost entirely of organics (particularly in regimes II and III). Further, SV-OOA correlates strongly with DMA number concentration (Pr = 0.62, vs. 0.00 and 0.33 for HOA and LV-OOA, respectively).



The regular, significant enhancements in SV-OOA during peak photochemical periods, high correlation with DMA number concentration combined with lower correlations with inorganics (which typically comprise accumulation mode aerosol), and the regular appearance of fine organic modes during afternoons suggests that the SV-OOA factor is comprised of early-generation photochemical products that are chiefly responsible for the growth of fine-mode aerosol in the afternoons. Based on wind patterns discussed in Sect. 3.1, SV-OOA is proposed to consist of secondary organic products on the order of 1-2h old.

Ng et al. (2011) applied a simple method for making real-time estimates of OA components, based on an empirical scaling factor that relates HOA and OOA (further seq-10 regated as SV-OOA and LV-OOA) to signal at fragments m/z 57 and m/z 44. HOA is estimated by  $b \times (C_{57} - 0.1 \times C_{44})$ , where b for PACO was found to be 14.0, compared with 13.4 (range = 8.5 - 19.2) for Ng et al. (2011). PACO OOA (combined SV-OOA and LV-OOA) is estimated by  $c \times C_{44}$ , where c was found to be 8.1 for PACO, compared with 6.6 (range = 2.9 - 9.0) by Ng et al. (2011). The PACO SV-OOA and LV-OOA factors are 15

estimated by  $3.3 \times C_{44}$  and  $4.8 \times C_{44}$ , respectively.

Figure 7 shows the mass fraction of organic accounted for by each organic component during each regime. OA in Los Angeles is overwhelmingly oxidized in nature, with LV- and SV-OOA combined contributing 77, 92, and 86% of OA during regimes

- I, II, and III, respectively. HOA contributed the largest fraction to OA during regime I, 20 in large part due to episodically high HOA concentrations during May and early June. HOA is a minor factor during regimes II and III. SV-OOA is a major constituent of OA during periods of photochemistry, with the hot, dry, photochemically intense regime III seeing the most significant contribution of SV-OOA to OA. LV-OOA dominates in
- regimes I and II, which were both periods of significant marine/moisture influence and 25 cloud processing.

# 3.4 Filter analysis

PMF factors are mathematical solutions that return constant mass spectra that are combined linearly to describe variability in AMS data, and it is important to compare



PMF results to physical, independent characterizations of OA. For PACO, UPLC-ESI-(+ and -)-MS analysis of high-volume filter samples was performed on 9 selected filters representing periods of high HOA, SV-OOA, and LV-OOA concentrations, in order to distinguish chemical profiles associated with the PMF factors. Periods distinguished

5 as "high SV- and LV-OOA" were chosen such that those species were the single, dominant component of OA. PACO filter sampling did not include any periods in which HOA was the isolated, dominant species, as periods distinguished as "high HOA" were also impacted by high background levels of LV-OOA.

Functional groups of interest in UPLC-ESI-(+)-TOF-MS analysis were basic nitrogen groups (i.e. amines, imines) as well as oxygen-containing moieties (i.e. carbonyls, 10 alcohols, ethers). The complexity of the samples made complete speciation by this technique difficult, but it was possible to identify the majority of the base peaks. Nonoxygenated nitrogenous compounds likely to be amines were detected in significant concentrations and with some variety of carbon chain-length (C8-C16) in the high HOA + LV-OOA periods, as well as during morning periods in which background LV-15

OOA dominated the organic fraction. This suggests that amines may be an important class of organic compounds in aged, background Los Angeles aerosol. The most prominent class of compounds in all filters was oxygenated organic species, with increased variety of oxygenated species during periods of significant photochemical organic production.

20

UPLC-ESI-(-)-TOF-MS offered better mass spectral resolution and more complete identification of base peaks than (+)-mode, allowing for more complete chemical profiles. Phthalic acid is ubiquitous in appreciable guantities in the extractable fraction of Los Angeles OA. Phthalic acid has been identified as a significant product of polyaromatic hydrocarbon (PAH) photochemistry (Kautzman et al., 2010), so this result 25 suggests that PAHs may be a significant class of precursor VOCs to OA in Los Angeles. The high HOA + LV-OOA samples were characterized by appreciable quantities of nitrogen- and sulfur-containing organics (> 10% of the extractable, identifiable fraction), which is unique to periods in which both HOA and LV-OOA were high in



magnitude. Carboxylic acids were significant during both SV- and LV-OOA periods (> 15% of extractable, identifiable fraction), though the carboxylic acids identified during SV-OOA periods tended to be longer-chain (C7-C10, as opposed to C4-C9 for LV-OOA periods). Additionally, aerosol in LV-OOA periods tended to have a more complex

- <sup>5</sup> chemical profile, with no single class of compounds dominating the extractable fraction identified by UPLC. This suggests that while the SV- and LV-OOA periods contain many of the same chemical compounds, LV-OOA tends to be characterized by smaller organic acids (higher O:C ratio), as well as a more complex profile of organonitrates, organosulfates, and oxidized acids.
- As explained in Ulbrich et al. (2009), aerosol partitioning with a bilinear PMF model is inherently limited in fitting a dynamic organic component. With a dataset as large as that represented by PACO, it is not unexpected that the organic component undergoes numerous and significant changes over the course of a single regime. Thus, two unique, constant mass spectra used to describe variability in a bilinear model may be
- <sup>15</sup> more accurately viewed as two points on a continuum of volatility, rather than two distinct components of differing volatility. This hypothesis is supported by the commonality of compound classes identified by UPLC analysis for SV- and LV-OOA periods. The continuum of oxygenation is further illustrated in Fig. 8, a plot of m/z 44 (CO<sub>2</sub><sup>+</sup>) vs. m/z43 (mostly C<sub>2</sub>H<sub>3</sub>O<sup>+</sup>) proposed by Ng et al. (2010) as a mass spectral diagnostic for
- <sup>20</sup> the age of organic aerosol. It is hypothesized that LV-OOA components tend to have a higher ratio of m/z 44 to m/z 43 and represent a lower-volatility, more-aged organic component with a higher degree of oxidation. Results from PACO indicate that the organic component of Los Angeles aerosol produces a continuum of  $f_{44}$  values during each regime and that the most aged organic fraction may actually be characterized
- by a predominance of longer-chain, oxidized monoacids, complex amines, and nitrateand sulfate- containing organics. It is concluded that for the Los Angeles aerosol, a bilinear PMF model, with mass spectra assumed to be constant over an entire regime, is helpful in qualitatively assessing the degree of aging of the organic component, but represents a broad simplification of a dynamic organic fraction of aerosol.



# 3.5 Hygroscopicity

The intensity of scattered light is a strong function of particle diameter ( $D_p$ ), and since aerosol water uptake largely determines particle size, it is a major variable determining the interaction of particles with radiation. Urban aerosol tends to be dominated by <sup>5</sup> organic material, which is often hydrophobic, leading to overall growth factors (GFs,  $D_{p,wet}/D_{p,dry}$ ) that are smaller than those for pure inorganic aerosol. At RH above the deliquescence RH of common inorganic aerosol constituents, the comparatively low hygroscopicity of organics tends to result in an inverse correlation between organic mass fraction and overall aerosol hygroscopicity (Shinozuka et al., 2009; Quinn et al., 2005; Hersey et al., 2009).

Because urban aerosol is, in part, an external mixture of organic and inorganic components, hygroscopic behavior is often complex, with several GF modes observed (e.g., Cocker et al., 2001; Massling et al., 2005, 2009; Swietlicki et al., 2008; Meier et al., 2009; Tiitta et al., 2010; Rose et al., 2010). The optical sizing method utilized by

- the DASH-SP provides rapid measurements on timescales suitable for aircraft studies, but does not allow resolution of more than two modes (a nonhygroscopic and hygroscopic mode). During PACO, two modes were typically distinguishable: a nonhygroscopic mode (1 < GF < 1.15 at 92% RH) identified during morning sampling (≤ 30% of aerosol), decreasing to very minor significance in the afternoon, in addition to a domi-</p>
- nant hygroscopic mode that will be the focus of the rest of this section. This suggests that aerosol sampled in the morning included an appreciable fraction of nonhygroscopic particles, with afternoon photochemistry and SOA production resulting in an aerosol of more uniform, unimodal hygroscopicity. Since it was not possible to resolve multiple growth modes, GF data represent the overall subsaturated water uptake behavior for hygroscopic particles.

The DASH-SP measured GF at dry particle sizes of 150, 175, 200, and 225 nm. There was no statistically significant difference between GF at these dry sizes, and so data were averaged to give GFs representative for particles between 150 and 225



nm dry diameter. Particles sampled during morning periods (07:00–11:00 LT) were significantly more hygroscopic at 74 and 92% than those sampled during afternoon periods (15:00–19:00 LT); morning GF averaged 1.14  $\pm$  0.13, 1.37  $\pm$  0.10, and 1.97  $\pm$  0.45 at 74, 85, and 92% RH, respectively, while afternoon GF averaged 1.06 $\pm$  0.04,

<sup>5</sup> 1.37±0.07, and 1.74±0.20 at 74, 85, and 92% RH, respectively. Suppressed GF at 74 and 92% during afternoon sampling but constant GF at 85% suggests that the hygroscopic growth curve (GF vs. RH) for morning-sampled aerosol is one with a higher GF at lower RHs, combined with a steep ascending section at RH > 85%, suggestive of a predominantly inorganic aerosol with a small hygroscopic organic component and little nonhygroscopic aerosol.

These GF values are significantly higher than those measured during the 1987 SCAQS study in the eastern Los Angeles Basin  $(1.23 \pm 0.08 \text{ for } 200 \text{ nm particles at } 90 \pm 3\% \text{ RH}$ ; Zhang and McMurray, 1993), and are more consistent with the most hygroscopic mode measured in Pasadena in 1999 (1.6 at 89% RH; Cocker et al., 2001).

Figure 9 shows GF at 74, 85, and 92% plotted against organic mass fraction (OMF), with markers colored by sampling date and time. PACO data show a clear trend of suppressed GF with increasing organic mass fraction (Pr = -0.56, -0.78, -0.71 for 74, 85, and 92%, respectively). The lowest GFs and highest values of organic mass fraction occur late in PACO, during the dry and photochemically intense regime III characterized by significant photochemical production of SV-OOA.

An increasingly popular representation of aerosol hygroscopicity is the  $\kappa$  parameter, developed by Petters and Kreidenweis (2007). As outlined in Shinozuka et al. (2009), one can use size-resolved GF data at sub-saturated RH to solve the equation:

$$S(D) = \frac{D^3 - D_d^3}{D^3 - D_d^3(1 - \kappa)} \exp\left[\frac{4\sigma M_w}{RT\rho_w D}\right]$$

for  $\kappa$ , where *S* is the saturation ratio over an aqueous droplet (0.74, 0.85, or 0.92), *D* and  $D_d$  are the humidified and dry diameters, respectively,  $\sigma$  is the surface tension at the interface of air and pure water (0.072 J m<sup>-2</sup>),  $M_w$  is the molecular weight of water



(2)

(18 g mol<sup>-1</sup>), *R* is the gas constant (8.3 J mol<sup>-1</sup>K<sup>-1</sup>), *T* is the temperature (298 K), and  $\rho_w$  is the density of water (1 g cm<sup>-3</sup>). Because the effect of organics on surface tension is not explicitly accounted for in this model, the  $\kappa$  here may be regarded an "effective hygroscopicity parameter", which includes both solute-induced water activity changes and surface tension effects (Petters and Kreidenweis, 2007; Rose et al., 2010).

Using the calculated  $\kappa$  values, one may estimate critical dry diameter ( $D_d$ ), or the diameter above which aerosol particles activate as cloud consenation nuclei (CCN) at a given supersaturation ( $S_c$ ), using the following equations:

$$\kappa = \frac{4A^3}{27D_d^3 \ln^2 S_c}$$

10 where:

$$A = \frac{4\sigma M_{\rm w}}{RT\rho_{\rm w}}$$

Following Shinozuka et al. (2009) and assuming  $S_c$  of 0.2%, the relationship between  $\kappa$  and  $D_d$  becomes:

 $D_{\rm d} = \kappa^{(-1/3)} \cdot 70 \,\rm nm$ 

<sup>15</sup> Values of  $\kappa$  and critical dry diameter at 0.2% supersaturation were calculated based on GFs at 85% RH and dry particle diameter of 200 nm. Uncertainties are estimated to be 25%, based on sizing uncertainties in the DASH-SP and RH variation in the instrument (Sorooshian et al., 2008). Figure 10 shows  $\kappa$  (left axis) and critical dry diameter (right axis;  $D_d$ ) plotted against organic mass fraction (OMF), colored by time and date. Kappa values range from 0.15 to 0.51, with an average of 0.31 ± 0.08, while critical diameters ranged from 87 to 131 nm, with an average of 105 ± 10 nm. As with GF data, there is a clear trend of decreasing  $\kappa$  and increasing critical dry diameter with increasing organic mass fraction, with the lowest hygroscopicity and CCN activity



(3)

(4)

(5)

and higher RH exhibited more hygroscopic behavior. At low values of organic mass fraction,  $\kappa$  approaches that of nitrate and sulfate salts (0.53–0.67), while approaching 0.1 at high organic fraction. Cubison et al. (2008) found that CCN concentrations in eastern Los Angeles were best modeled by treating organics (including the oxidized

- <sup>5</sup> fraction) as nonhygroscopic. GF- and CCN- derived  $\kappa$  values for hygroscopic organics range from 0.01 to 0.5 (Petters and Kreidenweis, 2007), with only the most hygroscopic organics (i.e. malonic acid) approaching the high end of that range. Results from PACO suggest that organics are moderately hygroscopic, with  $\kappa$  values on the lower end of the range cited by Petters and Kreidenweis (2007).
- <sup>10</sup> Shinozuka et al. (2009) presented paramaterizations for calculating as a function of organic mass fraction for 100 nm  $D_p$ , dry aerosol in Central Mexico and the North American West Coast ( $\kappa = 0.34-0.20 \cdot \text{OMF}$  and  $\kappa = 0.47-0.43 \cdot \text{OMF}$ , respectively), and these parameterizations are shown in Fig. 10. Overall, aerosol sampled during PACO are more hygroscopic at a given OMF than those presented in Shinozuka et al.
- <sup>15</sup> (2009), and are parameterized as  $\kappa = 0.50 0.29 \cdot \text{OMF}$ . This suggests that the organic fraction measured by Shinozuka et al. (2009) inhibits hygroscopicity significantly more than in Los Angeles. It is possible that the organics measured in Central Mexico and off the N. American West Coast exhibit more surface-active behavior, inhibiting hygroscopic activity, or that those organics are predominantly nonhygroscopic.

#### 20 4 Conclusions

25

A major ambient sampling experiment was conducted at a site in Pasadena, CA positioned to sample a constant afternoon influx of transported Los Angeles pollution with a photochemical age of roughly 1–2 h and generally free from the influence of local point sources. Sampling spanned 4 months during the summer of 2009, and was characterized by three distinct meteorological regimes: a "springtime" period characterized by clean conditions coincident with low pressure systems (regime I), a typical early-



photochemically active afternoons (regime II), and a late-summer period characterized by little marine layer influence, hot and dry conditions, and elevated photochemical activity (regime III).

Chemical analysis of aerosol revealed that OC dominates particulate carbon in Los
 Angeles (OC:EC average of 8.04), and is enhanced relative to EC by an average of 47% in the afternoon, suggesting significant SOA production. High RH during marine-layer-influenced morning sampling periods resulted in aqueous processing of aerosol particles and high aerosol volume, while drier afternoon periods saw increased aerosol number and volume due to SOA production. Maxima in aerosol volume occured in the mid-late morning, reflecting a balance between RH-induced hygroscopic growth and SOA production. Maxima in aerosol number 14:00–15:00 LT,

indicating that daytime photochemistry tended to generate a fine aerosol mode. Organics are the most significant constituent of transported Los Angeles aerosol, contributing an average of 42, 43, and 55% of total submicron mass in regimes I, II,

- and III, respectively. Interestingly, there was no significant difference between morning and afternoon O:C ratios, suggesting that the constant influx of 1–2 h old aerosol results in an overall organic oxidation state at the sampling site varying little despite significant SOA production. Regime II, characterized by significant morning marine layer influence followed by photochemically active afternoons, displayed significantly higher
- aerosol mass and higher average O:C ratio, suggesting that aqueous processes may be important in the generation of secondary aerosol and OOA in Los Angeles. Afternoons during regime II and III are characterized by the appearance of a fine organic mode, suggesting that SOA may be responsible for growth of fine particles. It is unclear whether these fine particles are the result of sulfate nucleation and subsequent SOA
   growth or SOA condensation on existing primary aerosol.

Water soluble organic mass (WSOM) reaches maxima near 14:00–15:00 LT, coincident with markers for photochemistry and suggesting significant SOA contribution to WSOM. Sulfate is predominantly an accumulation mode constituent, and its magnitude appears to depend on the degree of marine influence. Nitrate is similarly



a predominantly accumulation mode species, reaching maxima after morning rush hour periods. Nighttime nitrate production is observed during periods in which maximum ozone levels indicated significant photochemical events. An ammonium ratio, defined as the moles of ammonium measured divided by the moles of ammonium

required to neutralize sulfate and nitrate as (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> and NH<sub>4</sub>NO<sub>3</sub>, respectively, revealed that ammonia limits particulate (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> and NH<sub>4</sub>NO<sub>3</sub> in regimes I and II. An ammonium ratio of 1.28±0.30 in regime III suggests that ammonium was in excess and that organic amines may have been an important constituent of submicron aerosol during this hot, dry, photochemically intense sampling period with reduced marine sulfate
 influence.

PMF analysis of C-ToF-AMS data resolved three factors, corresponding to HOA, SV-OOA, and LV-OOA, and correlating well with the "standard" factors presented in Ng et al. (2011). The HOA factor appears to be a periodic plume source, generally appearing during or shortly after peaks in CO concentration. SV-OOA displays a strong diurnal pattern, correlating strongly with ozone and the appearance of a fine mode

- diurnal pattern, correlating strongly with ozone and the appearance of a fine mode (< 100 nm) in the aerosol size distribution, suggesting that these SOA products are generated on timescales of 1–2 h and tend to comprise a fine aerosol mode. LV-OOA is correlated with inorganic nitrate and sulfate and appears to be an aged carryover organic component that resides in accumulation mode aerosol, but may be generated
- on short timescales (1–2 h) under extremely oxidizing conditions. Its prevalence during humid overnight and morning hours and enhanced importance in moisture-influenced regimes I and II suggest that there may be significant aqueous-phase sources for LV-OOA. Overall the organic fraction of Los Angeles aerosol is overwhelmingly oxidized, with LV- and SV-OOA contributing 77, 92, and 86% of OA in regimes I, II, and III, respectively.

Filter analysis by both (+) and (-) mode UPLC-ESI-TOF-MS reveals a complex mixture of organic compounds dominated by oxygenated species. Long (C8-C16), non-oxygenated nitrogenous compounds likely to be amines are detected during periods impacted by high concentrations of background LV-OOA, suggesting that amines



may be an important constituent of aged organic aerosol in Los Angeles. Phthalic acid represents a ubiquitous, appreciable fraction of extractable OA, suggesting that PAH photochemistry may be an important pathway for SOA production in Los Angeles. Organo-nitro-sulfate species are observed during periods when aerosol appears to be

a mixture of aged background species and freshly emitted particles. Carboxylic acids appear to be major contributers to both SV- and LV-OOA, with LV-OOA tending to be associated with shorter (C4-C9) carboxylic acids and thereby higher O:C ratio.

The organic fraction of transported Los Angeles aerosol is dynamic and undergoes numerous and significant changes on timescales of days to weeks, despite exhibiting

- <sup>10</sup> consistency in the overall degree of oxidation. Many compounds are common to periods identified as distinct by PMF analysis, and PACO data suggest that Los Angeles OA is best described as a complex, dynamic mixture of compounds that represent a continuum of oxidation. Thus, a bilinear PMF model that assumes constant mass spectra representative of classes of OA is a useful tool in qualitatively assessing the degree of OA aging and oxidation, but is a dramatic simplification of a complex organic
- fraction.

Finally, aerosol water uptake characteristics measured by the DASH-SP indicate that a bimodal distribution becomes more unimodal during periods of photochemical SOA production. The degree of hygroscopicity of the main hygroscopic mode is highly anticorrelated with organic mass fraction, as indicated by decreasing GF and  $\kappa$ , as well as increasing critical diameter with increased organic fraction. Values for  $\kappa$  averaged

0.31±0.08, approaching 0.5 at low OMF and 0.1 at high OMF. It appears that organics in Los Angeles inhibit hygroscopicity less than those measured in Central Mexico and off the N. American West Coast, potentially owing to less surface-active behavior or an
 organic fraction comprised of more water soluble species.

20

PACO represents the first major aerosol characterization in the Western Los Angeles Basin, and comparison with SOAR will help identify spatial variation in Los Angeles aerosol and determine the nature of aerosol evolution as it is transported from sourcerich (western) to downwind (eastern) parts of Los Angeles. Further, its timing will serve



to contextualize and compare forthcoming results from the CalNex field campaign, part of which involved ground sampling at the PACO site.

# **Appendix A**

## **5 PMF analysis**

PMF was initiated using 50 seed values, and then again using a range of fpeak values from -1 to 1 varying by 0.1. Figures A1 and A2 show the  $Q/Q_{expected}$  values for the different solutions. The PMF results were sorted by regime, number of factors, and seed (only where it caused the solution to vary) and compared to previously published mass spectra in the AMS online Spectral Database explained in Ulbrich et al. (2009) (Ulbrich, I. M., Lechner, M., and Jimenez, J. L.: AMS Spectral Database, available at: http://cires.colorado.edu/jimenez-group/AMSsd/). Table A1 shows Pearson's *r* values comparing the PMF mass spectral profile solutions to existing mass spectra. Table A2 shows Pearson's *r* values comparing the PMF time series solutions to gas phase trac-15 ers, AMS inorganic tracers, PILS tracers, and WSOC. Figure A3 shows total residuals

ers, AMS inorganic tracers, PILS tracers, and WSOC. Figure A3 shows total residuals of all masses over time for solutions with different factor numbers. Figure A4 shows the scaled residuals at each mass for the presented 3 factor solution. The 2 and 4 factor solution profiles and time series contributions are shown in Fig. A5a, b.

The difference in the total residuals was noticeable between the 1 and 2 factor solution and the 2 and 3 factor solution, but negligible between the 3 and 4 factor solution. In the 2 factor solution there is evidence of an HOA-like factor and OOA-like factor from the mass spectral comparison, but the when compared to the 3 factor solution, the time series correlations with AMS inorganic and certain gas phase tracers is improved. In the 3 factor case, 3 solutions were explored, and were representative of the 50 possible

solutions from the PMF results. Regarding the mass spectral profile, the seed = 6 solution was chosen since it has very strong correlations with the real mass spectra, and each of the profiles are least similar to one another (see the right hand side of Table A2)



when compared to the two other 3 factor solutions. When considering the time series comparison, the seed = 6 LV-OOA like factor (factor 1) has the highest correlation with the AMS inorganic species. The p-ToF data suggests the organic profile of LV-OOA has similar particle size distributions to the AMS inorganic species (this was investigated

- in areas of known LV-OOA like factor and low SV-OOA like factor, since PMF was not performed on pToF data). When the afternoon sun reaches a maximum, there is another, smaller mode that is believed to be SV-OOA, without the strong inorganic signal. Also, the seed = 6 SV-OOA factor correlates best with ozone when compared to the two other solutions. The tracers for HOA were limited in this study since the resolution
- for the CO and NO<sub>x</sub> were poor and the *r* values are very low. However, when comparing the seed = 6 and seed = 1 solutions, the correlations of the HOA-like factor to CO and NO<sub>x</sub> are slightly larger. Additionally, the HOA-like factor was compared to m/z57, which is higher for the seed = 6 solutions. The HOA correlations for the seed = 4 solutions are slightly higher than in the seed = 1 and seed = 6 solutions, but this solu-
- tion is not as strong mass spectrally, and when comparing the tracers for the two other factors. The four factor situation provided two different solutions, representative of the 50 seeds. In addition to the residuals not changing dramatically, in the mass spectral comparison, for both solutions, two of the factors have very similar *r* values (factor 1 and 4 for seed 1 and factor 2 and 4 for seed 4). Also, the time series correlations are
   worse in the four factor solutions, when compared the the presented 3 factor solution.

Acknowledgements. Funding was provided by the Electric Power Research Institute. The authors would like to thank Jason Surratt for his helpful comments and direction. We would also like to thank Eric Edgerton and the SEARCH network for guidance on high-volume filter sampling and for the use of a sampler during PACO.



#### References

- Aiken, A. C., DeCarlo, P. F., Kroll, J. H., Worsnop, D. R., Huffman, J. A., Docherty, K. S., Ulbrich, I. M., Mohr, C., Kimmel, J. R., Sueper, D., Sun, Y., Zhang, Q., Trimborn, A., Northway, M., Ziemann, P. J., Canagaratna, M. R., Onsach, T. B., Alfarra, M. R., Prevot, A. S. H., Dommen, J., Duplissy, J., Metzger, A., Baltensperger, U., and Jimenez, J. L.: O/C and OM/OC ratios of primary, secondary, and ambient organic aerosols with high-resolution time-of-flight aerosol mass spectrometry, Environ. Sci. Technol., 42, 4478–4485, 2008. 5882
  Aiken, A. C., Salcedo, D., Cubison, M. J., Huffman, J. A., DeCarlo, P. F., Ulbrich, I. M., Docherty, K. S., Sueper, D., Kimmel, J. R., Worsnop, D. R., Trimborn, A., Northway, M., Stone, E. A., Schauer, J. J., Volkamer, R. M., Fortner, E., de Foy, B., Wang, J., Laskin, A., Shutthanandan, V., Zheng, J., Zhang, R., Gaffney, J., Marley, N. A., Paredes-Miranda, G.,
- Arnott, W. P., Molina, L. T., Sosa, G., and Jimenez, J. L.: Mexico City aerosol analysis during MILAGRO using high resolution aerosol mass spectrometry at the urban supersite (T0) Part 1: Fine particle composition and organic source apportionment, Atmos. Chem. Phys., 9, 6633–6653, doi:10.5194/acp-9-6633-2009, 2009, 5870
- Alfarra, M., Coe, H., Allan, J., Bower, K., Boudries, H., Canagaratna, M., Jimenez, J., Jayne, J., Garforth, A., Li, S., and Worsnop, D. R.: Characterization of urban and rural organic particulate in the Lower Fraser Valley using two aerodyne aerosol mass spectrometers, Atmos. Environ., 38, 5745–5758, 2004. 5886, 5887, 5921
- Allan, J., Bower, K., Coe, H., Boudries, H., Jayne, J., Canagaratna, M., Millet, D., Goldstein, A., Quinn, P., Weber, R., and Worsnop, D.: Submicron aerosol composition at Trinidad Head, California, during ITCT 2K2: its relationship with gas phase volatile organic carbon and assessment of instrument performance, J. Geophys. Res. Atmos., 109, D23S24, doi:10.1029/2003JD004208, 2004a. 5874
- Allan, J., Delia, A., Coe, H., Bower, K., Alfarra, M., Jimenez, J., Middlebrook, A., Drewnick, F., Onasch, T., Canagaratna, M., Jayne, J., and Worsnop, D.: A generalised method for the extraction of chemically resolved mass spectra from aerodyne aerosol mass spectrometer data, J. Aerosol Sci., 35, 909–922, doi:10.1016/j.jaerosci.2004.02.007, 2004b. 5874
   AQMD: Historic ozone air quality trends, available at: http://www.aqmd.gov/smog/o3trend.html
   (last access: August 2010), 2010. 5870
  - Bahreini, R., Keywood, M., Ng, N., Varutbangkul, V., Gao, S., Flagan, R., Seinfeld, J., Worsnop, D., and Jimenez, J.: Measurements of secondary organic aerosol from oxidation





5902

of cycloalkenes, terpenes, and m-xylene using an Aerodyne aerosol mass spectrometer, Environ. Sci. Technol., 39, 5674–5688, 2005. 5887, 5921

- Bein, K. J., Zhao, Y. J., Pekney, N. J., Davidson, C. I., Johnston, M. V., and Wexler, A. S.: Identification of sources of atmospheric PM at the Pittsburgh Supersite – Part II: Quantitative
- comparisons of single particle, particle number, and particle mass measurements, Atmos. Environ., 40, S424–S444, 2006. 5869

Blumenthal, D., White, W., and Smith, T.: Anatomy of a Los Angeles smog episode: pollutant transport in the daytime sea breeze regime, Atmos. Environ., 12, 893–907, 1978. 5879

Bremner, J., Haub, C., Lee, M., Mather, M., and Zuehlke, E.: World Population Highlights:

- <sup>10</sup> Key Findings From PRB's 2009 World Population Data Sheet, Population Reference Bureau, available at: www.prb.org, 2009. 5869
  - Cabada, J. C., Rees, S., Takahama, S., Khlystov, A., Pandis, S. N., Davidson, C. I., and Robinson, A. L.: Mass size distributions and size resolved chemical composition of fine particulate matter at the Pittsburgh supersite, Atmos. Environ., 38, 3127–3141, 2004. 5869
- <sup>15</sup> Canagaratna, M., Jayne, J., Jimenez, J., Allan, J., Alfarra, M., Zhang, Q., Onasch, T., Drewnick, F., Coe, H., Middlebrook, A., Worsnop, D. R., and Jimenez, J. L.: Chemical and microphysical characterization of ambient aerosols with the aerodyne aerosol mass spectrometer, Mass Spectrom. Rev., 26, 185–222, 2007. 5874

CARB: iADAM Air Quality Data Statistics, California Air Resources Board, available at: http://www.arb.ca.gov/adam/, 2010. 5870

//www.arb.ca.gov/adam/, 2010. 5870
 Chow, J. C., Fujita, E. M., Watson, J. G., Lu, Z. Q., Lawson, D. R., and Asbaugh, L. L.: Evaluation of filter-based aerosol measurements during the 1987 Southern California -Quality

Study, Environ. Monit. Assess., 30, 49–80, 1994. 5870
Cocker, D. R., Whitlock, N. E., Flagan, R. C., and Seinfeld, J. H.: Hygroscopic properties of Pasadena, California aerosol, Aerosol Sci. Technol., 35, 637–647, 2001. 5892, 5893, 5909

- Pasadena, California aerosol, Aerosol Sci. Technol., 35, 637–647, 2001. 5892, 5893, 5909 Croes, B. E. and Fujita, E. M.: Overview of the 1997 Southern California Ozone Study (SCOS97-NARSTO), Atmos. Environ., 37, S3–S26, 2003. 5870
  - Cubison, M. J., Ervens, B., Feingold, G., Docherty, K. S., Ulbrich, I. M., Shields, L., Prather, K., Hering, S., and Jimenez, J. L.: The influence of chemical composition and mixing state of
- Los Angeles urban aerosol on CCN number and cloud properties, Atmos. Chem. Phys., 8, 5649–5667, doi:10.5194/acp-8-5649-2008, 2008. 5895
  - DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T., Docherty, K. S., Worsnop, D. R., and Jimenez, J. L.:



Field-deployable, high-resolution, time-of-flight aerosol mass spectrometer, Anal. Chem., 78, 8281–8289, 2006. 5909

- DeCarlo, P. F., Dunlea, E. J., Kimmel, J. R., Aiken, A. C., Sueper, D., Crounse, J., Wennberg, P. O., Emmons, L., Shinozuka, Y., Clarke, A., Zhou, J., Tomlinson, J.,
- <sup>5</sup> Collins, D. R., Knapp, D., Weinheimer, A. J., Montzka, D. D., Campos, T., and Jimenez, J. L.: Fast airborne aerosol size and chemistry measurements above Mexico City and Central Mexico during the MILAGRO campaign, Atmos. Chem. Phys., 8, 4027–4048, doi:10.5194/acp-8-4027-2008, 2008. 5870

Denkenberger, K. A., Moffet, R. C., Holecek, J. C., Rebotier, T. P., and Prather, K. A.: Real-time,

- single-particle measurements of oligomers in aged ambient aerosol particles, Environ. Sci. Technol., 41, 5439–5446, 2007. 5870, 5909
  - Docherty, K. S., Stone, E. A., Ulbrich, I. M., DeCarlo, P. F., Snyder, D. C., Schauer, J. J., Peltier, R. E., Weber, R. J., Murphy, S. M., Seinfeld, J. H., Grover, B. D., Eatough, D. J., and Jimenez, J. L.: Apportionment of primary and secondary organic aerosols in Southern
- <sup>15</sup> California during the 2005 Study of Organic Aerosols in Riverside (SOAR-1), Environ. Sci. Technol., 42, 7655–7662, 2008. 5870, 5871, 5885, 5909
  - Drewnick, F., Schwab, J., Hogrefe, O., Peters, S., Husain, L., Diamond, D., Weber, R., and Demerjian, K.: Intercomparison and evaluation of four semi-continuous PM<sub>2.5</sub> sulfate instruments, Atmos. Environ., 37, 3335–3350, doi:10.1016/S1352-2310(03)00351-0, 2003. 5874
- <sup>20</sup> Drewnick, F., Hings, S., DeCarlo, P., Jayne, J., Gonin, M., Fuhrer, K., Weimer, S., Jimenez, J., Demerjian, K., Borrmann, S., and Worsnop, D.: A new time-of-flight aerosol mass spectrometer (TOF-AMS) – instrument description and first field deployment, Aerosol Sci. Tech., 39, 637–658, doi:10.1080/02786820500182040, 2005. 5874

Eatough, D. J., Grover, B. D., Woolwine, W. R., Eatough, N. L., Long, R., and Farber, R.: Source apportionment of 1 h semi-continuous data during the 2005 Study of Organic Aerosols in

 apportionment of 1 h semi-continuous data during the 2005 Study of Organic Aerosols in Riverside (SOAR) using positive matrix factorization, Atmos. Environ., 42, 2706–2719, 2008. 5870

- Eldering, A., Cass, G. R., and Moon, K. C.: An Air monitoring network using continuous particlesize distribution monitors – connecting pollutant properties to visibility via mie scattering calculations, Atmos. Environ., 28, 2733–2749, 1994. 5869
- Hersey, S. P., Sorooshian, A., Murphy, S. M., Flagan, R. C., and Seinfeld, J. H.: Aerosol hygroscopicity in the marine atmosphere: a closure study using high-time-resolution, multiple-RH DASH-SP and size-resolved C-ToF-AMS data, Atmos. Chem. Phys., 9, 2543–2554,

30



doi:10.5194/acp-9-2543-2009, 2009. 5892

- Huffman, J., Jayne, J., Drewnick, F., Aiken, A., Onasch, T., Worsnop, D., and Jimenez, J.: Design, modeling, optimization, and experimental tests of a particle beam width probe for the aerodyne aerosol mass spectrometer, Aerosol Sci. Tech., 39, 1143–1163, doi:10.1080/02786820500423782, 2005. 5874
- Hughes, L. S., Allen, J. O., Salmon, L. G., Mayo, P. R., Johnson, R. J., and Cass, G. R.: Evolution of nitrogen species air pollutants along trajectories crossing the Los Angeles area, Environ. Sci. Technol., 36, 3928–3935, 2002. 5870, 5909

Kautzman, K. E., Surratt, J. D., Chan, M. N., Chan, A. W. H., Hersey, S. P., Chhabra, P. S.,

- Dalleska, N. F., Wennberg, P. O., Flagan, R. C., and Seinfeld, J. H.: Chemical composition of gas- and aerosol-phase products from the photooxidation of naphthalene, J. Phys. Chem. A, 114, 913–934, 2010. 5890
  - Kleeman, M., Hughes, L., Allen, J., and Cass, G.: Source contributions to the size and composition distribution of atmospheric particles: Southern California in September 1996, Environ.

<sup>15</sup> Sci. Technol., 33, 4331–4341, 1999. 5883

5

Kondo, Y., Miyazaki, Y., Takegawa, N., Miyakawa, T., Weber, R. J., Jimenez, J. L., Zhang, Q., and Worsnop, D. R.: Oxygenated and water-soluble organic aerosols in Tokyo, J. Geophys. Res., 112, D01203, doi:10.1029/2006JD007056, 2007. 5886

Lanz, V. A., Alfarra, M. R., Baltensperger, U., Buchmann, B., Hueglin, C., and Prévôt, A. S. H.:

- Source apportionment of submicron organic aerosols at an urban site by factor analytical modelling of aerosol mass spectra, Atmos. Chem. Phys., 7, 1503–1522, doi:10.5194/acp-7-1503-2007, 2007. 5886, 5921
  - Liu, D. Y., Prather, K. A., and Hering, S. V.: Variations in the size and chemical composition of nitrate-containing particles in Riverside, CA, Aerosol Sci. Technol., 33, 71–86, 2000. 5870
- Lu, R. and Turco, R.: Air pollutant transport in a coastal environment II. Three-dimensional simulations over Los Angeles Basin, Atmos. Environ., 29, 1499–1518, 1995. 5870
  - Massling, A., Stock, M., and Wiedensohler, A.: Diurnal, weekly, and seasonal variation of hygroscopic properties of submicrometer urban aerosol particles, Atmos. Environ., 39, 3911–3922, 2005. 5892
- Massling, A., Stock, M., Wehner, B., Wu, Z. J., Hu, M., Bruggemann, E., Gnauk, T., Herrmann, H., and Wiedensohler, A.: Size segregated water uptake of the urban submicrometer aerosol in Beijing, Atmos. Environ., 43, 1578–1589, 2009. 5892

Matsui, H., Koike, M., Kondo, Y., Takegawa, N., Kita, K., Miyazaki, Y., Hu, M., Chang, S. Y.,



Blake, D. R., Fast, J. D., Zaveri, R. A., Streets, D. G., Zhang, Q., and Zhu, T.: Spatial and temporal variations of aerosols around Beijing in summer 2006: model evaluation and source apportionment, J. Geophys. Res. Atmos., 114, D00G13, 2009. 5870

Meier, J., Wehner, B., Massling, A., Birmili, W., Nowak, A., Gnauk, T., Brüggemann, E., Her-

rmann, H., Min, H., and Wiedensohler, A.: Hygroscopic growth of urban aerosol particles 5 in Beijing (China) during wintertime: a comparison of three experimental methods, Atmos. Chem. Phys., 9, 6865–6880, doi:10.5194/acp-9-6865-2009, 2009. 5892

Modey, W. K., Eatough, D. J., Anderson, R. R., Martello, D. V., Takahama, S., Lucas, L. J., and Davidson, C. I.: Ambient fine particulate concentrations and chemical composition at two

- sampling sites in metropolitan Pittsburgh: a 2001 intensive summer study, Atmos. Environ., 10 38, 3165-3178, 2004. 5869
  - Na, K., Sawant, A. A., Song, C., and Cocker, D. R.: Primary and secondary carbonaceous species in the atmosphere of Western Riverside County, California Atmos, Environ., 38. 1345-1355, 2004, 5909
- Ng, N. L., Canagaratna, M. R., Zhang, Q., Jimenez, J. L., Tian, J., Ulbrich, I. M., Kroll, J. H., 15 Docherty, K. S., Chhabra, P. S., Bahreini, R., Murphy, S. M., Seinfeld, J. H., Hildebrandt, L., Donahue, N. M., DeCarlo, P. F., Lanz, V. A., Prévôt, A. S. H., Dinar, E., Rudich, Y., and Worsnop, D. R.: Organic aerosol components observed in Northern Hemispheric datasets from Aerosol Mass Spectrometry, Atmos. Chem. Phys., 10, 4625-4641, doi:10.5194/acp-10-4625-2010, 2010. 5875, 5887, 5891
- 20
- Ng, N. L., Canagaratna, M. R., Jimenez, J. L., Zhang, Q., Ulbrich, I. M., and Worsnop, D. R.: Real-time methods for estimating organic component mass concentrations from aerosol mass spectrometer data, Environ. Sci. Technol., 45, 910-916, doi:10.1021/es102951k, 2011. 5887, 5889, 5897, 5921
- Paatero, P.: A weighted non-negative least squares algorithm for three-way "PARAFAC" factor analysis, Chemom. Intell. Lab. Syst., 38, 223-242, 1997. 5874
  - Paatero, P. and Tapper, U.: Positive matrix factorization a nonnegative factor model with optimal utilization of error-estimates of data values, Environmetrics, 5, 111-126, 1994. 5874. 5875
- Pastor, S. H., Allen, J. O., Hughes, L. S., Bhave, P., Cass, G. R., and Prather, K. A.: Ambient single particle analysis in Riverside, California by aerosol time-of-flight mass spectrometry during the SCOS97-NARSTO, Atmos. Environ., 37, S239–S258, 2003. 5870, 5909 Pekney, N. J., Davidson, C. I., Bein, K. J., Wexler, A. S., and Johnston, M. V.: Identification of



sources of atmospheric PM at the Pittsburgh Supersite, Part I: single particle analysis and filter-based positive matrix factorization, Atmos. Environ., 40, S411–S423, 2006. 5869

- Petters, M. D. and Kreidenweis, S. M.: A single parameter representation of hygroscopic growth and cloud condensation nucleus activity, Atmos. Chem. Phys., 7, 1961–1971, doi:10.5194/acp-7-1961-2007, 2007. 5893, 5894, 5895
- van Pinxteren, D., Bruggemann, E., Gnauk, T., Iinuma, Y., Muller, K., Nowak, A., Achtert, P., Wiedensohler, A., and Herrmann, H.: Size- and time-resolved chemical particle characterization during CAREBeijing-2006: different pollution regimes and diurnal profiles, J. Geophys. Res. Atmos., 114, D00G09, doi:10.1029/2008JD010890, 2009. 5870

5

Quinn, P., Bates, T., Baynard, T., Clarke, A., Onasch, T., Wang, W., Rood, M., Andrews, E., Allan, J., Carrico, C., Coffman, D., and Worsnop, D. R.: Impact of particulate organic matter on the relative humidity dependence of light scattering: a simplified parameterization, Geophys. Res. Lett., 32, 22, 2005. 5892

Rose, D., Nowak, A., Achtert, P., Wiedensohler, A., Hu, M., Shao, M., Zhang, Y., Andreae, M. O.,

and Pöschl, U.: Cloud condensation nuclei in polluted air and biomass burning smoke near the mega-city Guangzhou, China – Part 1: Size-resolved measurements and implications for the modeling of aerosol particle hygroscopicity and CCN activity, Atmos. Chem. Phys., 10, 3365–3383, doi:10.5194/acp-10-3365-2010, 2010. 5892, 5894

Sawant, A. A., Na, K., Zhu, X., Cocker, K., Butt, S., Song, C., and Cocker III, D. R.: Character-

- <sup>20</sup> ization of PM<sub>2.5</sub> and selected gas-phase compounds at multiple indoor and outdoor sites in Mira Loma, California, Atmos. Environ., 38, 6269–6278, 2004. 5909
  - Schauer, J. J., Rogge, W. F., Hildemann, L. M., Mazurek, M. A., and Cass, G. R.: Source apportionment of airborne particulate matter using organic compounds as tracers, Atmos. Environ., 30, 3837–3855, 1996. 5874
- Shinozuka, Y., Clarke, A. D., DeCarlo, P. F., Jimenez, J. L., Dunlea, E. J., Roberts, G. C., Tomlinson, J. M., Collins, D. R., Howell, S. G., Kapustin, V. N., McNaughton, C. S., and Zhou, J.: Aerosol optical properties relevant to regional remote sensing of CCN activity and links to their organic mass fraction: airborne observations over Central Mexico and the US West Coast during MILAGRO/INTEX-B, Atmos. Chem. Phys., 9, 6727–6742, doi:10.5194/acp-9-6727-2009, 2009, 5892, 5893, 5894, 5895
- Sorooshian, A., Brechtel, F., Weber, R., Corless, A., Flagan, R., and Seinfeld, J.: Modeling and characterization of a particle-into-liquid sampler (PILS), Aerosol Sci. Technol., 40, 396–409, 2006. 5875



- Sorooshian, A., Hersey, S., Brechtel, F., Corless, A., Flagan, R., and Seinfeld, J.: Rapid, sizeresolved aerosol hygroscopic growth measurements: differential aerosol sizing and hygroscopicity spectrometer probe (DASH-SP), Aerosol Sci. Technol., 42, 445–464, 2008. 5877, 5894
- Stanier, C., Khlystov, A., and Pandis, S.: Nucleation events during the Pittsburgh Air Quality Study: description and relation to key meteorological, gas phase, and aerosol parameters special issue of aerosol science and technology on findings from the fine particulate matter supersites program, Aerosol Sci. Technol., 38, 253–264, 2004. 5885
- Stone, E. A., Snyder, D. C., Sheesley, R. J., Sullivan, A. P., Weber, R. J., and Schauer, J. J.:
   Source apportionment of fine organic aerosol in Mexico City during the MILAGRO experiment 2006, Atmos. Chem. Phys., 8, 1249–1259, doi:10.5194/acp-8-1249-2008, 2008. 5870
   Sullivan, A., Weber, R., Clements, A., Turner, J., Bae, M., and Schauer, J.: A method for on-line measurement of water-soluble organic carbon in ambient aerosol particles: results from an
  - urban site, Geophys. Res. Lett., 31, L13105, doi:10.1029/2004GL019681, 2004. 5876
- <sup>15</sup> Sullivan, A., Peltier, R., Brock, C., De Gouw, J., Holloway, J., Warneke, C., Wollny, A., and Weber, R.: Airborne measurements of carbonaceous aerosol soluble in water over Northeastern United States: method development and an investigation into water-soluble organic carbon sources, J. Geophys. Res., 111, 1–14, 2006. 5876

Surratt, J., Gomez-Gonzalez, Y., Chan, A., Vermeylen, R., Shahgholi, M., Kleindienst, T., Ed-

- ney, E., Offenberg, J., Lewandowski, M., Jaoui, M., Maenhaut, W., Claeys, M., Flagan, R. C., and Seinfeld, J. H.: Organosulfate formation in biogenic secondary organic aerosol, J. Phys. Chem. A, 112, 8345–8378, 2008. 5877
  - Swietlicki, E., Hansson, H., Hameri, K., Svenningsson, B., Massling, A., McFiggans, G., Mc-Murry, P., Petaja, T., Tunved, P., Gysel, M., Topping, D., Weingartner, E., Baltensperger, U.,
- Rissler, J., Wiedensohler, A., and Kulmala, M.: Hygroscopic properties of submicrometer atmospheric aerosol particles measured with H-TDMA instruments in various environments a review, Tellus B, 60, 432–469, 2008. 5892
  - Tiitta, P., Miettinen, P., Vaattovaara, P., Joutsensaari, J., Pet, T., Virtanen, A., Raatikainen, T., Aalto, P., Portin, H., Romakkaniemi, S., Kokkolab, H., Lehtinena, E. J., Markku, K., and
- <sup>30</sup> Laaksonena, A.: Roadside aerosol study using hygroscopic, organic and volatility TDMAs: characterization and mixing state, Atmos. Environ., 44, 976–986, 2010. 5892
  - Turpin, B. J. and Lim, H.-J.: Species contributions to PM<sub>2.5</sub> mass concentrations: revisiting common assumptions for estimating organic mass, Aerosol Sci. Technol., 35, 602–610, 2001.





5885

- Turpin, B. J. and Huntzicker, J. J.: Secondary formation of organic aerosol in the Los-Angeles Basin – a descriptive analysis of organic and elemental carbon concentrations, Atmos. Environ. A, 25, 207–215, 1991. 5870, 5909
- <sup>5</sup> Ulbrich, I. M., Canagaratna, M. R., Zhang, Q., Worsnop, D. R., and Jimenez, J. L.: Interpretation of organic components from Positive Matrix Factorization of aerosol mass spectrometric data, Atmos. Chem. Phys., 9, 2891–2918, doi:10.5194/acp-9-2891-2009, 2009. 5874, 5875, 5886, 5887, 5891, 5921

Watson, J. G., Chow, J. C., Lu, Z. Q., Fujita, E. M., Lowenthal, D. H., Lawson, D. R., and Ash-

- baugh, L. L.: Chemical mass-balance source apportionment of PM<sub>(10)</sub> during the Southern California Air-Quality Study, Aerosol Sci. Technol., 21, 1–36, 1994. 5870, 5871, 5909
  - Wittig, A. E., Anderson, N., Khlystov, A. Y., Pandis, S. N., Davidson, C., and Robinson, A. L.: Pittsburgh air quality study overview, Atmos. Environ., 38, 3107–3125, 2004. 5869
  - Yue, D. L., Hu, M., Wu, Z. J., Wang, Z. B., Guo, S., Wehner, B., Nowak, A., Achtert, P., Wiedensohler, A., Jung, J., Kim, Y. J., and Liu, S.: Characteristics of aerosol size distributions and
- sohler, A., Jung, J., Kim, Y. J., and Liu, S.: Characteristics of aerosol size distributions and new particle formation in the summer in Beijing, J. Geophys. Res. Atmos., 114, D00G12, doi:10.1029/2008JD010894, 2009. 5870
  - Zhang, Q., Alfarra, M., Worsnop, D., Allan, J., Coe, H., Canagaratna, M., and Jimenez, J.: Deconvolution and quantification of hydrocarbon-like and oxygenated organic aerosols based
- on aerosol mass spectrometry, Environ. Sci. Technol., 39, 4938–4952, 2005. 5886, 5921 Zhang, Q., Jimenez, J. L., Canagaratna, M. R., Allan, J. D., Coe, H., Ulbrich, I., Alfarra, M. R., Takami, A., Middlebrook, A. M., Sun, Y. L., Dzepina, K., Dunlea, E., Docherty, K., De-Carlo, P. F., Salcedo, D., Onasch, T., Jayne, J. T., Miyoshi, T., Shimono, A., Hatakeyama, S., Takegawa, N., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demer-
- jian, K., Williams, P., Bower, K., Bahreini, R., Cottrell, L., Griffin, R. J., Rautiainen, J., Sun, J. Y., Zhang, Y. M., and Worsnop, D. R.: Ubiquity and dominance of oxygenated species in organic aerosols in anthropogenically-influenced Northern Hemisphere midlatitudes, Geophys. Res. Lett., 34, L13801, doi:10.1029/2007GL029979, 2007. 5874
- Zhang, X. and McMurray, S.: Mixing characteristics and water content of submicron aerosols measured in Los Angeles and at the Grand Canyon, Atmos. Environ. A, 27, 1593–1607, 1993. 5893



 Table 1. Previous Los Angeles air quality studies and major findings relative to particulate matter.

Study	Date	Major aerosol findings
Southern California Air Quality Study (SCAQS)	Summer/ fall 1987	<ul> <li>SOA/OA ~ 40% in summer afternoon (Turpin and Huntzicker, 1991)</li> <li>Primary vehicle exhaust is the major aerosol component near PACO sampling site (Watson et al., 1994)</li> <li>20–32% of PM<sub>10</sub> is SO<sub>4</sub><sup>2-</sup> and NO<sub>3</sub><sup>-</sup> (Watson et al., 1994)</li> </ul>
Southern California Ozone Study (SCOS97-NARSTO)	Summer 1997	<ul> <li>Submicron organic mode evolves during periods of intense photochemistry, coincident with maximum O<sub>3</sub> (Pastor et al., 2003)</li> <li>NH<sub>4</sub>NO<sub>3</sub> and organics added to submicron aerosol with photochemical age (Pastor et al., 2003)</li> <li>NH<sub>3</sub> sources in eastern basin enhance particulate NH<sub>4</sub>NO<sub>3</sub> (Hughes et al., 2002)</li> </ul>
Secondary Organic Aerosol in River- side (SOAR-1, 2)	Aug, Nov 2005	<ul> <li>Significant diurnal changes in organic aerosol, with species more reduced in the morning and more oxidized in the afternoon (DeCarlo et al., 2006)</li> <li>Oligomers detected in aged, acidic particles (Denkenberger et al., 2007)</li> <li>SOA/OA = 70–90% at midday and ~ 45% during morning commute; SOAR average = 74% (Docherty et al., 2008)</li> </ul>
Individual studies	1999, 2001–2002	<ul> <li>Complex, multi-modal GF data in Pasadena (Cocker et al., 2001)</li> <li>SOA &gt; 50% of total PM<sub>2.5</sub> in eastern basin (Sawant et al., 2004; Na et al., 2004)</li> </ul>



#### Table 2. PACO sampling regimes (2009).

Regime	Dates	Characteristics
Springtime meteorology	22 May to 12 Jun	<ul> <li>Basin impacted by a series of low pressure systems</li> <li>Cutoff low brought rare springtime precipitation and cleaner atmosphere</li> <li>Persistent cloudcover, unstable atmosphere</li> <li>Low daytime temperatures, high RH, and limited photochemistry</li> <li>High visibility with low O<sub>3</sub></li> </ul>
Early summer/ marine influence	18 Jun to 7 Jul	<ul> <li>Limited impact from low pressure systems</li> <li>Morning marine layer eroded several hours after sunrise</li> <li>Higher daytime temperatures, slightly lower RH, and photochemically active afternoons</li> <li>Hazy afternoons with increased O<sub>3</sub></li> </ul>
Photo- chemically active	10 Jul to 4 Aug	<ul> <li>Semi-permanent high pressure over Southern California</li> <li>Little marine moisture and infrequent, quickly-eroded marine layer</li> <li>High daytime temp. (often exceeding 35 °C), very low daytime RH (≤ 15%), photochemically active days</li> <li>Decreased visibility, high afternoon O<sub>3</sub></li> </ul>



	OC		EC		OC:EC	
Regime	AM	PM	AM	PM	AM	PM
Ι	$3.88 \pm 0.32$	$4.92 \pm 1.18$	$0.43 \pm 0.02$	$0.40 \pm 0.27$	9.1	12.4
II	$4.24 \pm 0.46$	$5.01 \pm 1.62$	$0.80 \pm 0.31$	$0.52 \pm 0.40$	5.3	9.6

7.4

9.9

 $6.16 \pm 1.40$   $6.41 \pm 1.44$   $0.83 \pm 0.47$   $0.65 \pm 0.22$ 

**Table 3.** OC/EC concentrations ( $\mu$ g m<sup>-3</sup>) for representative days from each regime.

Ш

iscussion	<b>AC</b> 11, 5867-	ACPD 11, 5867–5933, 2011 Pasadena, CA aerosol characterization S. P. Hersey et al.							
Paper   Discuss	Pasad aei charact S. P. He								
sion Paper	Title	Page							
Discuss	Conclusions Tables	References Figures							
ion Paper	14	►I ►							
Discussi	Back Full Sc Printer-frie	reen / Esc endly Version							
on Paper		Printer-friendly Version							

Table A1. Mass spectral comparison of several PMF solutions.

2. Lanz, V. A., et al., Atmos. Chem. Phys. 2007, 7, 1503-1522. 3. Ulbrich, I.M., et al., Atmos. Chem. and Phys. 2009, 9, 2891-2918.

4. Bahreini, R. et al. ES&T 2005, 39, pgs.5674-5688 5. Alfarra, R., et al. Atmos. Env. 2004, 38, 5745-5758.

										Mass	Spectra	a Comp	arison								
Factor Number (p)	Seed	Brushfire (4)	Manchester high CO low O3 (5)	Pittsburgh HOA (3)	Pittsburgh HOA(1)	Pittsburgh OOA I (3)	Pittsburgh OOA II (3)	Pittsburgh OOA (1)	Urban Vancouver high O3 (5)	Urban Vancouver low O3 (5)	Zurich Summer HOA (2)	Zurich Summer OOA I (2)	Zurich Summer OOA II (2)	Zurich Summer Wood burning (2)	Zurich Winter HOA (2)	Zurich Winter OOA (2)	Zurich Winter Wood burning (2)	Factor 1	Factor 2	Factor 3	Factor 4
1	1	0.96	0.68	0.68	0.51	0.97	0.85	0.97	0.99	0.77	0.44	0.95	0.58	0.64	0.66	0.97	0.81				
1	1	0.94	0.55	0.54	0.35	0.98	0.80	0.98	0.96	0.65	0.30	0.98	0.47	0.62	0.50	0.98	0.76		0.64		
2	1	0.77	0.94	0.97	0.91	0.67	0.78	0.66	0.78	0.96	0.78	0.57	0.76	0.51	0.90	0.61	0.73	0.64			
1	1	0.92	0.50	0.49	0.29	0.98	0.77	0.98	0.94	0.60	0.25	0.98	0.44	0.61	0.42	0.98	0.74		0.56	0.56	
2	1	0.74	0.91	0.95	0.88	0.63	0.73	0.62	0.74	0.93	0.76	0.53	0.71	0.51	0.88	0.56	0.69	0.56		0.94	
3	1	0.74	0.95	0.97	0.93	0.63	0.79	0.63	0.76	0.97	0.79	0.54	0.75	0.48	0.92	0.58	0.70	0.56	0.94		
1	4	0.93	0.66	0.66	0.49	0.93	0.79	0.93	0.93	0.73	0.42	0.89	0.56	0.64	0.55	0.92	0.81		0.92	0.57	
2	4	0.92	0.51	0.50	0.30	0.98	0.78	0.98	0.95	0.61	0.26	0.99	0.45	0.61	0.44	0.98	0.74	0.92		0.39	
3	4	0.59	0.93	0.97	0.96	0.46	0.67	0.45	0.61	0.93	0.83	0.35	0.72	0.39	0.95	0.38	0.58	0.57	0.39		
1	6	0.93	0.52	0.51	0.32	0.98	0.78	0.98	0.95	0.62	0.27	0.98	0.46	0.62	0.45	0.98	0.75		0.41	0.90	
2	6	0.61	0.92	0.96	0.95	0.47	0.65	0.46	0.60	0.92	0.81	0.35	0.71	0.43	0.93	0.39	0.60	0.41		0.73	
3	6	0.94	0.81	0.82	0.68	0.92	0.89	0.92	0.97	0.89	0.58	0.88	0.67	0.62	0.80	0.91	0.82	0.90	0.73		
1	1	0.94	0.57	0.56	0.37	0.98	0.81	0.98	0.97	0.67	0.32	0.98	0.49	0.63	0.54	0.98	0.76		0.89	0.49	0.93
2	1	0.90	0.60	0.59	0.43	0.89	0.74	0.89	0.87	0.66	0.37	0.84	0.54	0.65	0.43	0.87	0.80	0.89		0.53	0.86
3	1	0.61	0.92	0.96	0.94	0.48	0.67	0.48	0.63	0.93	0.81	0.38	0.68	0.41	0.94	0.41	0.58	0.49	0.53		0.56
4	1	0.93	0.69	0.66	0.51	0.95	0.86	0.94	0.96	0.75	0.46	0.92	0.62	0.58	0.60	0.95	0.82	0.93	0.86	0.56	
1	4	0.90	0.60	0.59	0.43	0.89	0.74	0.89	0.87	0.66	0.37	0.85	0.54	0.65	0.43	0.88	0.80		0.89	0.53	0.87
2	4	0.94	0.57	0.56	0.37	0.98	0.81	0.98	0.97	0.67	0.32	0.98	0.49	0.63	0.54	0.98	0.76	0.89		0.49	0.93
3	4	0.61	0.92	0.96	0.94	0.48	0.67	0.48	0.63	0.93	0.81	0.38	0.68	0.41	0.94	0.41	0.58	0.53	0.49		0.56
4	4	0.93	0.69	0.66	0.51	0.95	0.87	0.94	0.96	0.76	0.46	0.92	0.63	0.58	0.60	0.95	0.82	0.87	0.93	0.56	
Refe	ences	•																•			
1. Zh	ang, Q.	., et al. ES	5&T 2005	, 39, 493	8-4952.																

**Discussion** Paper 11, 5867-5933, 2011 Pasadena, CA aerosol characterization **Discussion** Paper S. P. Hersey et al. **Title Page** Abstract Introduction Conclusions References **Discussion** Paper **Tables Figures** 14 ►I. ◄ ► Close Back **Discussion** Paper Full Screen / Esc **Printer-friendly Version** Interactive Discussion

**ACPD** 

		Gas P	hase Ti	racers	AM	6 Inorga	anic Tra	cers	PILS Tracers								
Factor Number (p)	Seed	Ozone	CO	NOX	504	NO3	NH4	ChI	wsoc	org57	NH4	NO3	S04	Factor 1	Factor 2	Factor 3	Factor 4
1	1	0.51	0.38	0.24	0.66	0.45	0.63	0.35	0.58	0.79	0.70	0.67	0.76				
1	1	0.46	0.33	0.14	0.73	0.45	0.64	0.43	0.53	0.54	0.74	0.69	0.79		0.51		
2	1	0.41	0.34	0.30	0.37	0.31	0.44	0.12	0.30	0.91	0.56	0.56	0.63	0.51			
1	1	0.47	0.33	0.16	0.71	0.44	0.63	0.42	0.56	0.53	0.73	0.68	0.79		-0.02	0.73	
2	1	-0.06	0.15	-0.03	0.23	0.18	0.24	0.06	-0.39	0.61	0.22	0.22	0.20	-0.02		0.00	
3	1	0.51	0.34	0.30	0.43	0.32	0.47	0.18	0.50	0.77	0.60	0.59	0.67	0.73	0.00		
1	4	0.07	0.12	-0.14	0.22	0.29	0.30	0.02	-0.19	0.22	0.10	0.25	0.14		-0.19	-0.02	
2	4	0.47	0.32	0.19	0.66	0.39	0.58	0.40	0.58	0.54	0.71	0.66	0.77	-0.19		0.64	
3	4	0.42	0.35	0.30	0.42	0.31	0.45	0.15	0.36	0.93	0.59	0.58	0.65	-0.02	0.64		
1	6	0.39	0.31	0.09	0.75	0.46	0.65	0.46	0.47	0.49	0.76	0.73	0.81		0.05	0.60	
2	6	0.39	0.17	0.04	0.18	0.15	0.21	0.02	-0.29	0.70	0.16	0.17	0.14	0.05		-0.08	
3	6	0.53	0.31	0.28	0.42	0.31	0.45	0.19	0.56	0.62	0.60	0.58	0.67	0.60	-0.08		
1	1	0.43	0.30	0.15	0.71	0.39	0.58	0.43	0.53	0.52	0.73	0.66	0.79		-0.14	0.26	0.56
2	1	0.01	0.15	-0.07	0.16	0.29	0.26	0.01	-0.22	0.07	0.03	0.20	0.04	-0.14		0.06	-0.18
3	1	0.24	0.29	0.22	0.28	0.19	0.30	0.04	0.16	0.94	0.43	0.43	0.46	0.26	0.06		0.44
4	1	0.51	0.28	0.27	0.31	0.34	0.44	0.13	0.39	0.58	0.57	0.58	0.66	0.56	-0.18	0.44	
1	4	0.01	0.15	-0.07	0.16	0.29	0.26	0.01	-0.23	0.07	0.02	0.20	0.04		-0.15	0.06	-0.18
2	4	0.43	0.30	0.15	0.70	0.39	0.58	0.43	0.54	0.52	0.73	0.66	0.79	-0.15		0.26	0.56
3	4	0.24	0.29	0.22	0.27	0.19	0.30	0.04	0.16	0.94	0.43	0.43	0.46	0.06	0.26		0.43
4	4	0.51	0.28	0.27	0.31	0.34	0.44	0.13	0.39	0.58	0.57	0.58	0.66	-0.18	0.56	0.43	

Table A2. Time series comparison of several PMF solutions.





Fig. 1a. Regime I (wind direction marker size proportional to wind speed; max size = 20 km/h).





Fig. 1b. Regime II (wind direction marker size proportional to wind speed; max size = 20 km/h).





Fig. 1c. Regime III (wind direction marker size proportional to wind speed; max size = 20 km/h).





Fig. 2. Average diurnal DMA number concentrations (cm<sup>-3</sup>) for regimes I, II, and III.











**Fig. 4.** AMS bulk mass fractions for AM (07:00–11:00 LT), midday (11:00–15:00 LT), PM (15:00–19:00 LT), and night (19:00–07:00 LT) for regimes I, II, and III.





Fig. 5. WSOC magnitude (squares) and % of AMS organic (circles). Black markers represent the average over regime III.





**Fig. 6.** Mass spectra for the three factors identified in PMF analysis (HOA, SV-OOA, and LV-OOA; left) and Pearson's R correlations of those solution mass spectra with previously reported mass spectra (1, 2, 3, 4, 5, 6: Zhang et al., 2005; Lanz et al., 2007; Ulbrich et al., 2009; Bahreini et al., 2005; Alfarra et al., 2004; Ng et al., 2011, respectively).







5922

III.



**Fig. 8.**  $f_{44}$  vs.  $f_{43}$  for PACO, with color scale corresponding to date. Square markers represent values for HOA, SV-OOA, and LV-OOA factors identified in PACO.







Fig. 9. GF at 74, 85, and 92% RH plotted against organic mass fraction, with color scale corresponding to date.



**Fig. 10.**  $\kappa$  and critical dry diameter plotted against mass fraction organic, with color scale corresponding to date; lines represent empirical parameterizations from PACO, Central Mexico, and N. American West Coast.





Interactive Discussion



5926



**Fig. A2.**  $Q/Q_{exp}$  for varying fpeak values for p = 3. The circled marker is the presented solution.

Interactive Discussion



Fig. A3. Sum of the residuals for 1, 2, 3, and 4 factor soultions.

Discussio		ACPD							
on Pape		11, 5867–5933, 2011							
	aer charact	aerosol characterization							
iscussion	S. P. He	rsey et al.							
Pape	Title	Page							
Ð	Abstract	Introduction							
	Conclusions	References							
iscussio	Tables	Figures							
on Pa		►I							
aper	•	•							
	Back	Close							
Discu	Full Scr	een / Esc							
ssion	Printer-frie	ndly Version							
Paper	Interactive	Discussion							
		BY							



Fig. A4. Scaled residuals for the 3 factor solution (seed = 6).





Fig. A5a. Mass spectral profiles for the 2 factor solution.





Fig. A5b. Mass spectral profiles for the 4 factor solution.





Fig. A6. Time series contributions for the 2-factor solution.

Discussion Pa	<b>ACPD</b> 11, 5867–5933, 2011								
per   Discussior	Pasadena, CA aerosol characterization S. P. Hersey et al.								
Pap	Title	Page							
er	Abstract	Introduction							
	Conclusions	References							
iscuss	Tables	Figures							
ion P	14	۶I							
aper	•	Þ							
_	Back	Close							
Discussion	Full Screen / Esc Printer-friendly Version								
Paper	Interactive Discussion								



Fig. A7. Time series contributions for the 4-factor solution.

$\Box$								
SCIDS	ACPD							
sion F	11, 5867–5	11, 5867–5933, 2011						
aper	Pasade	Pasadena, CA						
_	aero characte	osol erization						
)iscuss	S. P. Her	S. P. Hersey et al.						
sion F								
ap	Title I	Title Page						
Ð,	Abstract	Introduction						
	Conclusions	References						
SCIDS	Tables	Figures						
ion P	14	۶I						
aper	•	•						
	Back	Close						
Discu	Full Scre	en / Esc						
ssion	Printer-frien	dly Version						
Pap	Interactive	Discussion						
D		•						