

***Interactive comment on* “Daytime HONO Vertical Gradients during SHARP 2009 in Houston, TX” by K. W. Wong et al.**

K. W. Wong et al.

jochen@atmos.ucla.edu

Received and published: 17 November 2011

Response to Comment by C. Fittschen

We would like to thank the Dr. Fittschen for the helpful comments on our manuscript. Below is our response (in italics).

The authors discuss in the introduction the possibility of HONO formation from excited NO_2 with H_2O , as reported by Li et al (2008). Even though the authors exclude themselves this reaction as a major source of the observed HONO, we would point out our recent work showing clearly, that the observed OH radicals in the work of Li et al originate from a process involving at least 2 photons: Direct observation

C12014

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



of OH radicals after 565 nm multi-photon excitation of NO₂ in the presence of H₂O
Chemical Physics Letters, Volume 513, Issues 1-3, 6 September 2011, Pages 12-16
Damien Amedro, Alexander E. Parker, Coralie Schoemaeker, Christa Fittschen.

Therefore this reaction is of no importance for the atmosphere.

The comments are well-taken. We have included this study in our paper. The sentence "Recent laboratory study by Amedro et al. (2011) also found that this reaction is not important in daytime HONO formation." was added to the text.

Amedro, D., Parker, A. E., Schoemaeker, C., and Fittschen, C.: Direct observation of OH radicals after 565 nm multi-photon excitation of NO₂ in the presence of H₂O, Chem. Phys. Lett., 513, 12-16, 2011

Interactive comment on Atmos. Chem. Phys. Discuss., 11, 24365, 2011.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

