

***Interactive comment on “Reactions of  
 $\text{H}^+(\text{pyridine})_m(\text{H}_2\text{O})_n$  and  $\text{H}^+(\text{NH}_3)_1(\text{pyridine})_1(\text{H}_2\text{O})_n$   
with  $\text{NH}_3$ : experiments and kinetic modelling  
under tropospheric conditions” by M. J. Ryding  
et al.***

**M. J. Ryding et al.**

pan@chem.gu.se

Received and published: 8 January 2012

Please find our answers as a supplement.

Please also note the supplement to this comment:

<http://www.atmos-chem-phys-discuss.net/11/C13948/2012/acpd-11-C13948-2012-supplement.pdf>

Interactive comment on Atmos. Chem. Phys. Discuss., 11, 24535, 2011.

C13948

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper






Figure 9. Concentration of most abundant cluster ions as a function of pyridine desorption rate coefficient. The values were calculated using Model B with inclusion of evaporation of pyridine. The initial concentrations are given in Table 2 except the concentration of pyridine that was kept at  $9.83 \times 10^7 \text{ cm}^{-3}$ .

Fig. 1. Figure 9.




Figure 10. Concentration of most abundant cluster ions as a function of aerosol concentration. The values were calculated using Model A. The initial concentrations are given in Table 2 except the concentration of pyridine that was kept at  $9.83 \times 10^7 \text{ cm}^{-3}$  and the aerosol concentration that was varied.

Fig. 2. figure 10.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper