

***Interactive comment on* “Primary and secondary sources of formaldehyde in urban atmospheres: Houston Texas region” by D. D. Parrish et al.**

D. D. Parrish et al.

David.D.Parrish@noaa.gov

Received and published: 14 February 2012

We thank Dr. Olaguer for his comment. We do agree that there are direct emissions of formaldehyde (CH_2O) from flares, although they are quite small in the context of other CH_2O sources in the Houston area. The emission rates quoted by Dr. Olaguer (0.2–2.2 kg CH_2O /hr) are not inconsistent either with our estimate of total primary emissions from Houston area industrial facilities (317 kg CH_2O /hr estimated from the emission inventory of Kim et al, 2011) or the direct measurements from three of the major facilities that we cite (45, 17 and 22 kg CH_2O /hr from Mellqvist et al., 2010b). We cannot comment on whether the individual flares investigated by the SHARP campaign are or are not included in any particular emission inventory. However, it is clear that these individual flare emissions of CH_2O are one to two orders of magnitude smaller than the pri-

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

mary CH₂O emissions from individual facilities, and two to three orders of magnitude smaller than the total primary CH₂O emissions from all Houston industrial facilities. Further, these total primary CH₂O industrial emissions are themselves a small fraction ($4\pm 2\%$ in our analysis) of total CH₂O sources in Houston.

Interactive comment on Atmos. Chem. Phys. Discuss., 11, 32601, 2011.

ACPD

11, C15330–C15331,
2012

Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

C15331

