Atmos. Chem. Phys. Discuss., 11, C2657–C2659, 2011 www.atmos-chem-phys-discuss.net/11/C2657/2011/ © Author(s) 2011. This work is distributed under the Creative Commons Attribute 3.0 License.



## Interactive comment on "Application of

## SCIAMACHY and MOPITT CO total column measurements to evaluate model results over biomass burning regions and Eastern China" by C. Liu et al.

C. Liu et al.

thomas.wagner@mpic.de

Received and published: 3 May 2011

Comment to "Application of SCIAMACHY and MOPITT CO total column measurements : : :." by Liu et al.

In section 2.3 the authors mention the instrumental issues that affect the SCIAMACHY CO measurements, as they have also been reported in literature before. They also mention that sophisticated solutions have been developed but that these corrections

C2657

are not perfect. The authors then tried different methods to correct for these problems following a.o. Gloudemans et al. 2005 and Buchwitz et al. 2007. The results are described in Liu et al. 2010 (PhD-thesis) and not shown in the ACPD manuscript here, but they conclude "that part of the problems of the SCIAMACHY CO data could beimproved, at least for limited periods of time, but systematic biases still remained". Having looked at Liu et al. 2010 and the results they obtain by applying the variable offset correction as introduced by Gloudemans et al., 2005, I am surprised that this correction has little or no effect on the time-series over the Sahara. Effectively the resulting 'corrected' time series is very similar to the uncorrected time-series as shown in Fig. 1 in this ACPD-manuscript. For comparison I include the figure below showing the resulting time-series for our SCIAMACHY CO dataset (IMLM) applying our variable offset correction over the same area and time-period as is used by Liu et al. As you can see there is no obvious drop in CO values correlated to the changing ice-layer. This is why we think the correction is guite adeguate -although not perfect- to deal with the ice-layer on the SCIAMACHY detector. Basically I am surprised about the somewhat disappointing results that the authors obtain - as described in Liu et al. 2010- when applying this correction approach to their IMAP CO retrieval.

Author comment: We were also disappointed about the relatively small improvements by applying the different correction procedures. The reasons for these findings are not totally clear to us, but obviously the effects of the ice layer and bad/dead pixels depend on the actual settings of the spectral analysis (as e.g. pointed out by Buchwitz et al., 2007).

The authors also write in Liu et al., 2010 "Here it is interesting to note that the improvement reported by other studies (e.g. Gloudemans et al., 2005) is better than found in our study. The reason for these differences are not clear, but might be caused by the use of different retrieval settings". I think it is important to make such a statement in the current manuscript to make clear that apparently different results are obtained with different SCIAMACHY retrievals applying a similar correction approach. Author comment: We added this information to the revised version.

Interactive comment on Atmos. Chem. Phys. Discuss., 11, 1265, 2011.

C2659