

Supporting information to “Heterogeneous ice nucleation: Exploring the transition from stochastic to singular freezing behavior “

D. Niedermeier¹, R. A. Shaw², S. Hartmann¹, H. Wex¹, T. Clauss¹, J. Voigtländer¹, and F. Stratmann¹

¹Leibniz Institute for Tropospheric Research, 04318 Leipzig, Germany

²Dept. of Physics, Michigan Technological University, Houghton, MI 49931, USA




Fig. 1 The solid lines in the left figure show three different fit curves to the immersion freezing behavior ($T > -38^\circ\text{C}$ indicated by the dashed line) of supercooled droplets each having a single ATD particles immersed (measured with LACIS, Niedermeier et al., 2010). The different parameter combinations, which lead to different time dependencies (see right figure), feature least square differences between fitted curves and experimental data points which are smaller than 10^{-3} . Black curve: $n_{\text{site}} = 1$; $\mu_\theta = 2.125$ rad; $\sigma_\theta = 0.325$ rad; Red curve: $n_{\text{site}} = 4$; $\mu_\theta = 2.31$ rad; $\sigma_\theta = 0.34$ rad; Green curve: $n_{\text{site}} = 7$; $\mu_\theta = 2.48$ rad; $\sigma_\theta = 0.39$ rad.