Atmos. Chem. Phys. Discuss., 11, C5676–C5677, 2011 www.atmos-chem-phys-discuss.net/11/C5676/2011/ © Author(s) 2011. This work is distributed under the Creative Commons Attribute 3.0 License.



ACPD 11, C5676–C5677, 2011

> Interactive Comment

Interactive comment on "Secondary organic aerosol formation from the photooxidation of isoprene, 1,3-butadiene, and 2,3-dimethyl-1,3-butadiene under high NO_x conditions" by K. Sato et al.

K. Sato et al.

sato.kei15@gmail.com

Received and published: 29 June 2011

Thank you very much for your valuable comments. We reply your comments as follows:

"1. At various places in the text the authors seem to be implying that methacrolein is oxidized in the particles by OH and that this leads to oligomer formation. I think this is inconsistent with the figures and also what has been proposed previously."

What we really indicate in the manuscript is that oligomers are produced by "gas-phase methacrolein oxidation with OH radicals followed by particle-phase oligomerizations of



Discussion Paper



methacrolein oxidation products". However, we express this process as "the heterogeneous reaction of methacrolein" at various places in the text. Probably, these expressions are ambiguous and will lead readers' misunderstanding. We revise these expressions adequately at various places in the text.

Interactive comment on Atmos. Chem. Phys. Discuss., 11, 4313, 2011.

Interactive Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper

