

Interactive
Comment

Interactive comment on “TransCom continuous experiment: comparison of ^{222}Rn transport at hourly time scales at three stations in Germany” by S. Taguchi et al.

S. Taguchi et al.

s.taguchi@aist.go.jp

Received and published: 21 September 2011

General comments

The authors would like to thank the reviewer(#3) for the valuable comments that have improved the manuscript. Please find below a point by point answer to the comments. The answers are displayed in italics.

– Specific minor suggestions for the authors –

Section 2.1, page 19260, line 15: I can't follow the equation for the corrected Freiburg values.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



The description is moved to the Appendix. A regression equation between Heidelberg monitor and Freiburg monitor was incorrect. It was corrected in the Appendix and an evidence is shown in Fig.S1. Now you may follow the equation, namely $FRB_new=(0.8FRB_old-0.23)/0.704$.

Section 2.2, page 19262, line 3-5: I am interested in the BLH comparison between radiosonde and ECMWF. Which method did you use to determine the BLH from the radiosonde observations? Why did you think there is no evidence of inconsistency between observations and ECMWF?

We defined BLH_radiosonde as the altitude where potential temperature matches with that at the surface when the lowest layer is in an unstable condition as demonstrated in Fig.S2. Temporal variations of the estimated BLH are shown in Fig.S3. BLH in the ECMWF regional model data set are sampled when the radiosonde observations estimated BLH with horizontal interpolation from 0.5 resolution data. The altitude of BLH estimated from radio sonde and regional model are compared in Fig.S4. Geographical distributions of the stations are shown in Fig.S5. We modified sentences as below: "To assess the diagnosed BLH in the ECMWF dataset, we also estimated BLH from radiosonde observations available in the neighborhood (Nancy, Hanau, Stuttgart, Sigmaringen), defining the BLH as the height at which potential temperature is the same as the surface when the lowest layer is unstable. We found reasonable correlation between the radiosonde and ECMWF BLHs except at Sigmaringen which had relatively few radiosondes available in 2002-2003."

Section 3, page 19262, line 19-22: The reported BLH from 14 models may be determined from different algorithm. Do you think this will have any impacts on your comparison results?

It is difficult to determine how much differences in the definition of BLH in different models could impact our results. We now note this issue in our discussion of Fig.3. "The method used to define BLH in each model may also contribute to the differences

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

from the ECMWF derived BLH.” In respect of Fig.8, it is possible that since we compare against MAX-MIN BLH, i.e. amplitude, then this may reduce some of the impact of differences in BLH definitions.

4.Section 4.3, page 19266, line 1: Please define NSD.

defined

Figure S1. Results of intercomparison conducted at Freiburg in August of 2002.

Figure S2. BLH was estimated when the lowest layer is in an unstable condition. It was defined as the altitude where potential temperature is the same as that at the surface.

Figure S3. BLH estimated from radiosonde (IGRA) data in 2002 and 2003.

Figure S4. BLH estimated from IGRA data set and ECMWF Forecast.

Figure S5. IGRA stations used in this study.

Interactive comment on Atmos. Chem. Phys. Discuss., 11, 19253, 2011.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper






Fig. 1. Results of intercomparison conducted at Freiburg in August of 2002.


Interactive
Comment

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper




prd79011.a/Users/p1871/aist01/pbl_pbl_igra.sh Wed Sep 7 14-23-36 JST 2011

Fig. 2. BLH was estimated when the lowest layer is in an unstable condition. It was defined as the altitude where potential temperature is the same as that at the surface.

[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Interactive
Comment

PBL TOP for 2002 and 2003. x=TYPE2, DATA=IGRA



<http://www.ncdc.noaa.gov/oa/climate/igra/index.php>

prd79011.a/Users/p1871/aist01/pbl_pbl_igra_2y.sh Wed Sep 7 14:28:36 JST 2011

Fig. 3. BLH estimated from radiosonde (IGRA) data in 2002 and 2003.


Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper



[Interactive
Comment](#)

prd79011.a/Users/p1871/aist01/pbl cmp_2blh.sWed Sep 7 14:39:39 JST 2011

Fig. 4. BLH estimated from IGRA data set and ECMWF Forecast.[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)[Discussion Paper](#)

Interactive
Comment




Fig. 5. GRA stations used in this study.

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Discussion Paper