Atmos. Chem. Phys. Discuss., 11, C9836–C9837, 2011 www.atmos-chem-phys-discuss.net/11/C9836/2011/ © Author(s) 2011. This work is distributed under the Creative Commons Attribute 3.0 License.



Interactive comment on "Black carbon fractal morphology and short-wave radiative impact: a modelling study" by M. Kahnert and A. Devasthale

Anonymous Referee #2

Received and published: 5 October 2011

When I started career as an aerosol researcher, I had no choice but to use Mie theory for calculating optical properties of black carbon (soot) particles. I was able to cite rich 1970s era literature to support the approach. However, from the moment I had seen the first microscopic photograph of soot particles I knew Mie theory could not be a reasonable approximation.

Therefore I am glad I can review one of the papers improving the treatment of black carbon aerosol in radiative transfer computations. I believe this is an important manuscript, presenting novel and useful results. It certainly warrants publishing it in ACP with only some minor corrections.

Being more lazy than Reviewer #1, I have a simpler task as there is no need to repeat the same comments. There are however a few (minor) points I would like to make:

C9836

Wherever the authors use phrases like "resulting in a lower TOA radiative impact of spheres" (p. 23118), they should make it clear they mean "lower positive". It may seem redundant and unnecessary but aerosol researchers do assume that "lower radiative impact/forcing" in the case of aerosol means "lower negative" one.

I would not agree with the statement "Multiple scattering becomes increasingly important as the optical depth is increased. Therefore, the compensating eiňĂect of the asymmetry parameter is strongest under more polluted conditions." (bottom of page 23118 and the top of the next one). The shape of the phase function (and therefore the asymmetry parameter) becomes less and less important with each order of scattering as the light field becomes more diffuse. I would therefore argue that this is exactly the reason why with more polluted air (greater optical depth), the effects of lacy and compact aggregates are most similar.

The first name in Henyey-Greenstein has been misspelled in page 23118 (an extremely unlucky page, it seems). I would also suggest citing the H-G 1941 paper directly (L. C. Henyey and J. L. Greenstein, 1941, "Diffuse radiation in the galaxy," Astrophys. J. 93, 70 - 83)

Finally, I do not like the automatically (?) created references which include the number of page the reference was cited it. I find it highly misleading. I do hope it is not a new obligatory format in ACP.

Interactive comment on Atmos. Chem. Phys. Discuss., 11, 23103, 2011.