

Received: 16 February 2012 – Accepted: 10 April 2012 – Published: 27 April 2012

Correspondence to: V. Gros (valerie.gros@lsce.ipsl.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

ACPD

12, 10937–10994, 2012

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Hydroxyl radicals play a central role in the troposphere as they control the lifetime of many trace gases. Measurement of OH reactivity (OH loss rate) is important to better constrain the OH budget and also to evaluate the completeness of measured VOC budget. Total atmospheric OH reactivity was measured for the first time in an European Megacity: Paris and its surrounding areas with 12 million inhabitants, during the MEGAPOLI winter campaign 2010. The method deployed was the Comparative Reactivity Method (CRM). The measured dataset contains both measured and calculated OH reactivity from CO, NO_x and VOCs measured via PTR-MS, GC-FID and GC-MS instruments. The reactivities observed in Paris covered a range from 10 s⁻¹ to 130 s⁻¹, indicating a large loading of chemical reactants. The present study showed that, when clean marine air masses influenced Paris, the purely local OH reactivity (20 s⁻¹) is well explained by the measured species. Nevertheless, when there is a continental import of air masses, high levels of OH reactivity were obtained (120–130 s⁻¹) and the missing OH reactivity measured in this case jumped to 75%. Using covariations of the missing OH reactivity to secondary inorganic species in fine aerosols, we suggest that the missing OH reactants were most likely highly oxidized compounds issued from photochemically processed air masses of anthropogenic origin.

1 Introduction

The hydroxyl radical (OH) represents the most important oxidant in the troposphere. It governs the atmospheric lifetime of most reactive trace species of anthropogenic and biogenic origin and contributes to the self-cleansing capacity of the atmosphere.

Generally, the production processes for OH are relatively well understood. However, the loss processes or the sink term of the OH radicals is still poorly constrained. In most cases the overall sink term of OH radicals is estimated by the sum of the products of concentration and the rate coefficient of all individually measured species reactive to

ACPD

12, 10937–10994, 2012

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Total OH reactivity
measurements in
Paris**C. Dolgorouky et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the OH radical. This method has proved to be rather limited due to the large number of “OH reactive” volatile organic compounds (VOCs) in urban air (Lewis et al., 2000 and Xu et al., 2003). Recently, Goldstein and Galabally (2007) postulated that 10^5 different VOCs have been detected in the Earth’s atmosphere. Moreover, recent studies

5 showed that there are unknown reactive species that contribute to the total OH reactivity (Di Carlo et al., 2004; Sinha et al., 2010; Kato et al., 2011). The necessity of measuring the total sink of OH radicals, also called the OH reactivity (s^{-1}), has therefore become evident. Presently three different types of instruments exist for the direct measurement of the total OH reactivity. The LIF community has two versions depend-

10 ing on how they produce OH in their reactor. The first method is based on the Laser Induced Fluorescence (LIF) technique which records the decay (loss rate) of the OH radicals. Two versions of instruments using LIF technique exist: one that measures the OH reactivity as the inverse of the atmospheric OH lifetime in a reaction flow tube with a movable OH injector (Kovacs and Brune, 2001; Mao et al., 2010; Ingham et al.,

15 2009) the other one uses a laser pump and probe technique (Sadanaga et al., 2005). The second method is called the Comparative Reactivity Method (CRM) and is based on comparative OH reactivity measurement in a flow reactor against the known OH reactivity of an added reagent (e.g. pyrrole, C_4H_5N). CRM has been developed and described in detail by Sinha et al. (2008).

20 Total OH reactivity has been measured in different environments for the last decade. Lou et al. (2010) reported 18 studies conducted in urban, rural and forested areas. The urban areas investigated included studies over New York City (summer 2001 and winter 2004), Mexico City Metropolitan area (spring 2003), Tokyo (summer 2003, winter 2004), Houston (autumn 2000, autumn 2006), Nashville (summer 1999) and Mainz

25 (summer 2008). The ranges of the OH reactivity registered in these urban areas went from $6 s^{-1}$, for the clean atmosphere of the small city of Mainz, Germany (Sinha et al., 2008) to $200 s^{-1}$, for the extremely polluted air of Mexico City (Shirley et al., 2006). More recently, Sinha et al. (2012) report measurements from El Arenosillo in Spain,

where OH reactivity ranged from below detection limit in marine air masses up to 70 s^{-1} in continental air masses.

Studies from forested areas have identified high percentages of missing OH reactivities, obtained from the comparison between the measured and the calculated reactivity, (70 % in the tropical forest of Surinam, Sinha et al., 2008 and 50 % in boreal forest in Finlande, Sinha et al., 2010), but also in rural areas (50 % in a subtropical rural area in PRD of China, Lou et al., 2010). In contrast, missing OH reactivity studies in megacities reported relatively low levels of missing OH reactivity: maximums of 30 % for New York 2004 and Tokyo 2003. If the studies showed that in the case of the forested areas, the “responsible compounds” of the missing OH reactivity were non-measured biogenic compounds, the question is still open for the origin of the missing OH reactivity in an urban area.

This paper presents the OH reactivity measurements by CRM method conducted in January–February 2010 within the framework of the EU project MEGAPOLI (<http://megapoli.dmi.dk>), in the city of Paris and its surrounding region (called “Ile de France”) with almost 12 million inhabitants, and which constitutes one of the few megacities in Europe.

The employed method (CRM) was previously described in only four other studies: Sinha et al. (2008), Sinha et al. (2010), Kim et al. (2011) (but this is a modified version and not strictly our method) and Sinha et al. (2012). All of these measurements were made in environments poor in NO emissions, as interferences due to OH recycling via $\text{NO} + \text{HO}_2$ reactions at $\text{NO} > 5 \text{ ppbv}$ in the setup were reported. This is for the first time that the CRM method has been used to measure OH reactivity measurements in an urban area characterized by high levels of NO emissions, as Paris was identified as a city mainly impacted by traffic emissions (Vardoulakis et al., 2002; Gros et al., 2007). Therefore, the initial part of the article presents the experimental setup and the quantification of interferences in the system used for the MEGAPOLI winter campaign 2010.

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The geographical location of Paris – a relatively small-area basin surrounded by rural areas – makes it a favourable place to study purely local OH reactivity levels emitted by a megacity when air masses come from the clean marine western sector, and to characterize the impact of the European contribution when air masses come from the eastern sector.

Within the MEGAPOLI framework a large number of compounds has been simultaneously measured during this campaign: NO_x (sum of NO and NO₂), carbon monoxide (CO), ozone (O₃), non-methane hydrocarbons (NMHCs) and oxygenated VOCs (OVOCs). We could therefore compare the observed OH reactivity with the calculated values, which were derived using the observed concentrations of the trace species.

2 Experimental

2.1 Experimental setup and field deployment

2.1.1 Experimental setup

The comparative reactivity method for measuring the total OH reactivity is described in detail by Sinha et al. (2008). A brief overview of the principle of the measurement is nevertheless presented here: firstly, a mixture of pyrrole (C₄H₅N) and synthetic dry air is injected into the reactor. The concentration C₁ of the pyrrole is monitored in the flow exiting the reactor by a Proton Transfer Reaction – Mass Spectrometer (PTR-MS). Secondly, the UV rays emitted by a mercury lamp produce OH radicals ([OH] < [pyrrole]) to react with pyrrole ($k_{\text{Pyr}+\text{OH}} = 1.2 \times 10^{-10} \text{ cm}^3 \text{ molecules}^{-1} \text{ s}^{-1}$). The pyrrole concentration registered now is C₂. Next, the synthetic air is replaced with ambient air. The OH reactive species contained in the ambient air compete with pyrrole for the available OH radicals. This causes an increase in the pyrrole monitored concentration from C₂ to C₃. The three measured concentrations, C₁, C₂ and C₃, allow the calculation of the total OH reactivity via the following equation based on competitive kinetics, which

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



has been derived by Sinha et al. (2008).

$$R_{\text{air}} = \frac{C_3 - C_2}{C_1 - C_3} \cdot K_{\text{py+OH}} \cdot C_1 \quad (1)$$

The schematic diagram of the CRM measurement system deployed for the MEGAPOLI campaign is shown in Fig. 1. The glass reactor used for the MEGAPOLI winter campaign is identical in shape and dimension with the one described in Sinha et al. (2008).

To measure C_1 , C_2 and C_3 , three different instrumental modules were used: the “synthetic air” module, the “nitrogen” module and the “ambient air” module (see Fig. 1). The “nitrogen” module is used to produce the OH radical via the UV photolysis of the water molecules imported by the humidified nitrogen bubbled through a pure water bottle. The same method to produce OH radicals has largely been used in the techniques of OH radical measurements (LIF measurements: Kovacs et al. 2003; Mao et al., 2009).

While switching from C_1 to C_2 , the synthetic air entering the reactor is humidified via the “synthetic air” module. Two mass flow controllers (MFC A and MFC B on the Fig. 1) are used and two flows of a 100% humidified and dry zero air are mixed together. Their mixing proportion determines the humidity of the final zero air flow. The relative humidity of C_2 baseline signal during MEGAPOLI was maintained constant at 50%, for a temperature of 24 °C. For more details concerning the correction applied to the data set because of the difference in the humidity of the synthetic air and ambient air, the reader is referred to Sect. 2.2.2, OH reactivity quantifying tests.

When switching from C_2 to C_3 , the “synthetic air” module is switched OFF and the “atmospheric air” module is then connected, via a 3 way valve. This module consists of a Teflon VOC pump (KNF), a Teflon micro valve and 1/4 and 1/16 inch flow reducing Teflon tubing (see Fig. 1).

The used detector located in the exit of the reactor is a high sensitivity PTR-MS, from Ionicon Analytik, Austria. The PTR-MS technology has been firstly introduced in studies of atmospheric chemistry by Lindinger et al. (1998). For this campaign the main monitored compound is pyrrole which is detected at the mass $m/z = 68$ a.m.u. Other

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



studies showed that this detector is ideal for measuring pyrrole, as the protonated ion does not fragment (Sinha et al., 2008, 2009). Also this mass is known to be free of interferences. The explored calibration range of pyrrole concentrations that are measured is from 0.5 to 250 ppbv, but on field measurements were always higher than 10 ppbv. The instrumental precision is of 6 % for 2 standard deviations (2σ). The humidity inside the reaction cell is important for accurate determination of the OH reactivity. To track the humidity into the reactor, the masses of the first two water clusters are also monitored: m/z 37 (cluster $\text{H}_3\text{O}^+\text{H}_2\text{O}$) and m/z 55 (cluster $\text{H}_3\text{O}^+(\text{H}_2\text{O})_2$), an approach described by Amman et al. (2006) and Sinha et al. (2008).

Besides the water cluster measurements that were used as proxies for the humidity into the reactor, an extra sensor for relative humidity and temperature was also installed in the exhaust line of the reactor (see Fig. 1). In contrast to the reactivity setup of Sinha et al. (2008) for the CRM used during the MEGAPOLI winter campaign, no other extra pump has been used at the exhaust line, the reactor being therefore maintained at ambient pressure with no over pressure inside the reactor.

2.1.2 Field deployment

The CRM system was deployed at the Laboratoire d'Hygiène de la Ville de Paris (LHVP) during the MEGAPOLI winter campaign 2010. The site is located in the heart of Paris (the 13th district) and therefore considered to be representative of the background pollution of central Paris.

The OH reactivity system was installed in a room where the temperature was maintained at 24 °C. The ambient air was sampled through a Teflon line of 1/4 inch of diameter and a length of about 10 m. The inlet was situated on the roof of LHVP, at a height of around 14 m above the ground level. In order to avoid any particulate contamination, a 0.45 μm Teflon filter was used at the entrance of the sampling line. OH reactivity studies showed that the OH reactivity measurements are not influenced by the positioning of a Teflon filter in the entrance of the inlet line (Sadanaga et al., 2006).

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Total OH reactivity
measurements in
Paris**

C. Dolgorouky et al.

Ambient air was then sampled at 150 ml min^{-1} in a total flow of 245 ml min^{-1} (dilution factor of the OH reactivity into the reactor of 1.65). The residence time of the compounds in the inlet lines was kept at $<35 \text{ s}$ by using a Teflon VOC sampling pump which draws a flow of 3 l min^{-1} . A 1/4 inch and 1/16 inch flow-reducing Teflon tubes were used downstream to stabilize the ambient air flow into the reactor at 150 ml min^{-1} (see Fig. 1).

Atmospheric OH reactivity measurements were performed between 23 January and 13 February. The time resolution of the measurements was of 2 min. Gaps in the data set were due to regular check-up of the instrument.

2.2 Reactivity tests

2.2.1 Evaluation of the system

At the beginning of the campaign, a series of tests were conducted from 8 to 23 January to characterize the system and the new reaction cell.

To check the accuracy of the CRM OH reactivity quantification, two standards have been used: a propane gas standard (Westfalen A.G.; 29.8 ppm in N_2 ; stated uncertainty 5%) and a standard which was a mixture of 3 compounds in N_2 : acetone, methanol and isoprene (Messer; 5.01 ppm acetone, stated uncertainty 5%; 6.78 ppm methanol, stated uncertainty 5%; 3.90 ppm isoprene, stated uncertainty 10%). The second standard has been analyzed in the laboratory in December 2009 prior to the MEGAPOLI campaign. Different quantities of standard have been introduced into the reactor. Using the definition of the OH reactivity, the reactivity due to the standard has been calculated via the equation:

$$R_{\text{total}} = \sum k_{\text{compound} + \text{OH}} \cdot [\text{compound}] \quad (2)$$

where the $k_{\text{compound} + \text{OH}}$ represents the constant rate of each compound (propane, acetone, methanol and isoprene) with the OH radical ($k_{\text{OH} + \text{propane}} = 1.1 \times 10^{-12} \text{ cm}^3$

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

molecules⁻¹ s⁻¹, $k_{\text{OH} + \text{acetone}} = 1.8 \times 10^{-13} \text{ cm}^3 \text{ molecules}^{-1} \text{ s}^{-1}$, $k_{\text{OH} + \text{methanol}} = 7.65 \times 10^{-13} \text{ cm}^3 \text{ molecules}^{-1} \text{ s}^{-1}$, $k_{\text{OH} + \text{isoprene}} = 1.0 \times 10^{-10} \text{ cm}^3 \text{ molecules}^{-1} \text{ s}^{-1}$, Atkinson et al., 2006) and the [compound] is the concentration of the compound in molecules cm⁻³. Figure 2 presents the measured versus the calculated reactivity. The two standards allowed us to cover a range of OH reactivity from 3 to 50 s⁻¹ for the propane standard, and a dynamic range comprised from 50 up to 300 s⁻¹ for the acetone, methanol and isoprene standard. Corrections using facsimile for deviations from the first order kinetics as described by Sinha et al. (2008) and for dilution of air in the reactor have already been applied.

The total uncertainty of the calculated reactivity is around 20 % and comprises: uncertainty due to the rate coefficient of the standards with the OH radicals (18 %), flow fluctuations (~10 %) and uncertainty due to the accuracy of the standards (5 %). The measured reactivity is obtained with the CRM (vertical axis). The total uncertainty for the measured reactivity is around 18 % and comprises: uncertainty due to the rate coefficient of the pyrrole with the OH radicals (13 %), uncertainty of the standards (5 %), due to flow fluctuations (10 %) and the instrument precision (2σ , ~6 %).

The calibration tests show a good linearity and accountability of the measured reactivity. The estimated detection limit of the ambient OH reactivity measurements within the setup is around 3 s⁻¹.

2.2.2 OH reactivity quantifying tests

As already identified by Sinha et al. (2008), interferences are potentially present in the setup. The penray mercury lamp is used to photolyse water at 185 nm. A fraction of the pyrrole is also photolyzed at 185 nm. The percentage of pyrrole photolysed in our setup is $19 \pm 2 \%$, but this does not influence our measurement because the Eq. (1) takes into consideration the real pyrrole concentration (after photolysis) available for reaction with the OH radicals (C_1 base line).

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Humidity difference between the zero air and the ambient air

Another potential interference is the humidity difference between the zero air and the ambient air. Different humidity conditions for the C_2 and C_3 signals correspond to different yields of OH radicals in the reactor, and therefore can lead to over or under estimation of the total OH reactivity. To avoid this, the humidity of the zero air must be kept the same as the one of the ambient air. In the experimental setup used during the winter 2010 MEGAPOLI campaign, the zero air was humidified to a constant value of 50 %. As the atmospheric humidity varied over the day, corrections were applied on the C_2 base line signal so that the overall OH field stays constant. To do this, a “humidity test” was run. The zero air flowing into the reactor was humidified to different degrees by mixing varying amounts of wet and dry zero air, with the help of the “synthetic air” module presented in Fig. 1. The decrease in the pyrrole baseline (C_2) is then recorded for different relative humidities. During this test the mercury lamp is maintained ON, as the purpose is the quantification of the real OH radical production into the setup. The humidity is tracked via the sensor installed on the exhaust line of the reactor.

As the campaign took place in winter, the relative humidity of the ambient air rarely went below 50 %, usually the relative humidity being between 50 and 90 %. The results of this test are therefore presented for a relative humidity range between 50 and 100 % (see Fig. 3).

A humidity correction has been applied based on this test on the C_2 baseline signal. Note that while this correction may seem significant (\sim up to 30 %), as the ambient air was more humid than zero air, it would only cause underestimation of the true OH reactivity thus leave conclusions of the paper regarding missing OH reactivity robust and intact.

The test revealed that for a relative humidity of between 60–70 %, the setup is much more sensitive to a change in the pyrrole signal than for humidity comprised between 80 and 90 % (the slope being smaller). Therefore, if there is no possibility of regulating the humidity of the C_2 signal to the one of the C_3 signal, as in the present work, it is

ACPD

12, 10937–10994, 2012

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



advised to keep the total relative humidity of the C₂ base line signal in the ranges of 80 to 90 %.

NO interference into the setup

As has been shown previously (Sinha et al., 2008), for high values of the NO into the setup, secondary production of OH radicals is observed due to the recycling of the HO₂ radicals. This interference is non-negligible for measurements made during the MEGAPOLI winter campaign, as the NO atmospheric levels went up to 100 ppb during the morning rush hour (see top panel from Fig. 6). To overcome this potential interference, firstly, the NO effect on the OH reactivity was quantified. Afterwards, a correction factor was applied to the ambient OH reactivity.

For the quantification of the NO interference, different amounts of NO were injected into the reactor and the decrease of the pyrrole signal due to additional OH radicals, recycled via the HO₂ + NO → OH + NO₂ ($k_{\text{HO}_2+\text{NO}} = 8.8 \times 10^{-12} \text{ cm}^3 \text{ molecules}^{-1} \text{ s}^{-1}$, Atkinson et al., 2004) reaction, was analysed. The result of the test is presented in the Fig. 4, where the absolute change in the measured OH reactivity was plotted against the NO values in the setup.

The test was run for an initial OH reactivity of 50 s⁻¹ (using the propane standard described previously) and showed a change in the slope at NO value of 20 ppbv. A secondary NO test was made 2 weeks later with no propane injected into the reactor (OH reactivity null). This secondary test was not presented on Fig. 4, but again, the change in the slope was noted for NO of 20 ppbv into the setup. Two different slopes were therefore considered for correction of the ambient OH reactivity.

The vertical axis uncertainty was 20 % and corresponded to the total OH reactivity uncertainty as quantified for the calibration tests described in Sect. 2.2.1. The total horizontal uncertainty was 11 %, and took into consideration NO flow fluctuations (10 %) and the uncertainty of the NO standard (5 %).

For the MEGAPOLI winter campaign, the OH reactivity data set was corrected for the NO interference as follows: for NO values into the setup smaller than 20 ppbv

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(equivalent to ambient NO concentration of 33 ppbv, for a dilution factor into the reactor of 1.65), the measured OH reactivity was increased with a value equivalent of the product between the 1st slope determined ($1.13 \text{ s}^{-1} \text{ ppbv}^{-1}$) and the measured NO value into the setup. For NO values higher than 20 ppbv, the measured OH reactivity was increased with the $0.46 \cdot \text{NO}$ concentration + 12.51 s^{-1} . 99 % of the OH reactivity dataset corresponds to the first regime ($\text{NO} < 20 \text{ ppbv}$) and only 1 % to the second regime ($\text{NO} > 20 \text{ ppbv}$).

Frequency of the C₀, C₁ and C₂ signal measurements

No C₁ measurements were completed between 23 and 30 January, being replaced by C₀ measurements. These measurements correspond to measurements made in the same conditions as C₁ with the UV lamp turned OFF. The photolytic loss of pyrrole described by the difference between C₀ and C₁ signals is an indicator of the stability of the emission rays of the lamp, and therefore of the constant OH field production into the reactor. The photolytic loss for different C₀s was quantified at the beginning and during the campaign as follows: from 8 to 23 January and after 31 January. The results indicated a constant photolytic loss of $19 \pm 2 \%$ over the entire period. The missing C₁ levels were therefore confidently quantified based on the correspondent C₀ levels.

The CRM method is based on the comparison between the C₂ and C₃ signals and therefore frequent measurements of the C₂ signal are required. During the MEGAPOLI winter campaign, the C₂ baseline signal was determined 2 to 3 times per day. Although not very frequently measured, the variability of the C₂ showed a daily variation of maximum 15 % over the entire campaign. An additional error of 15 % due to extrapolating successive baselines was therefore considered to the total uncertainty of the measured OH reactivity. A total uncertainty of 24 % was finally considered for the measured OH reactivity values during the MEGAPOLI campaign. This result comprises: uncertainty due to the rate coefficient of the pyrrole with the OH radicals (13 %), uncertainty of the standard (5 %), due to flow fluctuations (6 %), the instrument precision (2σ , $\sim 6 \%$) and the uncertainty due to low frequency C₂ measurements (15 %).

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.3 Ancillary measurements at LHVP site during MEGAPOLI winter campaign

Within the MEGAPOLI framework, additional measurements of CO, NO_x, non-methane hydrocarbons (NMHCs) and oxygenated volatile organic compounds (OVOCs), as well as aerosol chemical composition were also performed.

Ozone was measured with an analyzer based on UV absorption (41M, Environment SA) and nitrogen oxides were measured by chemiluminescence with an AC31M analyzer (Environnement SA). CO was measured by an analyzer based on IR absorption (48i-TL instrument, Thermo Electron Corporation, Waltham, MA, USA). More details for these instruments are given in Gros et al. (2011).

2.3.1 Non-methane hydrocarbon on-line measurements by gas chromatography

Two portable gas chromatographs equipped with a flame ionization detector (GCFID, Chromatotec, Saint Antoine, France) were used to measure non-methane hydrocarbons (NMHC) in ambient air. The instruments were located on the roof of the LHVP (~14 m above ground level). The first analyser, ChromaTrap, allowed the measurement of C₂–C₆ hydrocarbons and the second, AirmoBTX, the measurement of C₆–C₁₀ hydrocarbons. As the same instruments were used for atmospheric measurements during the spring 2007 Paris campaign, their technical information (type of columns, sampling flows, preconcentration, desorption-heating temperatures and times, etc.) was already described in details by Gros et al. (2011).

For both GC instruments, the sampling time was 10min and analysis time was 20 min, and therefore measurements were performed with a time resolution of 30 min.

2.3.2 VOC measurements by PTR-MS

HS-PTRMS (High Sensitivity Proton Transfer Reaction Mass Spectrometer, Ionicon Analytic, Lindinger et al., 1998) was deployed during the intensive field campaign in

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



order to quantify VOCs and oxygenated VOCs. The sampling line was installed on the roof of the LHVP building and consists of 10 m Teflon tubing. A primary flow of 10 l min^{-1} was maintained in the sampling line. During the campaign 39 m/z were monitored in the SIM mode with a temporal resolution of 2.5 min. These ions include $m/z = 25, 32, 37, 39, 55, 80$ used to control the PTRMS performances (fragmentation, ionization) and the 33 remaining m/z correspond to individual VOC's or sum of isomers. Within this study 5 VOC's measured by PTRMS were considered: benzene ($m/z = 79$), toluene ($m/z = 93$), acetonitrile ($m/z = 42$), methanol ($m/z = 33$) and acetaldehyde ($m/z = 45$). Transmission of the PTR-MS quadrupole were adjusted by the mean of 18 individual VOCs with certified concentration overlapping the mass range of the VOCs monitored within the field campaign. The 5 selected VOCs were also individually calibrated in the range 0.2–8 ppb at 50 % HR.

2.3.3 Oxygenated compounds measurements by GC – MS

For OVOC measurements, air sampling was performed at 15 ml min^{-1} flow during 30 min by Air server-Unity I (Markes International, Llantrisant, UK). After sampling, the compounds were then thermally desorbed and injected through a transfer line into a high polar column and analyzed by GC (Agilent, Massy, France) equipped with a FID and a Mass Spectrometer. This device allowed the identification and the quantification of 19 OVOC and 11 NMHC from C_2 to C_9 with a time resolution of 1.5 h. Technical precisions are given by Roukos et al. (2009).

2.3.4 Fine aerosol measurements by AMS

A suite of instrumentation for the characterization of the physico-chemical properties of particles was deployed in the garden of the LHVP building (approx. 14 m below the sampling line of the OH-reactivity system). The instruments included a High Resolution Time-of-Flight Aerosol Mass Spectrometer (HR-ToF-AMS), here simply referred to as AMS, Aerodyne Research Inc., DeCarlo et al. (2006). A collection efficiency of 0.4

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



was calculated for the HR-ToF-AMS based on comparison with concurrent TDMPs and particle-into-liquid sampler (PILS) data (Healy et al., 2011). All instruments were connected to the same sampling system consisting of a PM₁₀ inlet located approximately 6 m above ground level directly followed by an automatic aerosol diffusion dryer system maintaining the relative humidity in the line below 30 % Tuch et al. (2009).

The AMS was operated at 5 min time resolution. Due to the 600 °C surface temperature of the vaporizer, the AMS can only measure the non-refractory (NR) part of the particles (ammonium, nitrate, chloride, sulphate, and organic matter). Therefore, based on the aerodynamic lenses transmission efficiency and the detected compounds, AMS results are commonly considered to correspond to the non-refractory PM₁ aerosol (NR-PM₁) Canagaratna et al. (2007).

2.4 Description of the FLEXPART model

The FLEXPART model represents a quantitative dispersion model which takes into consideration turbulence, convection parameterizations, dry and wet deposition to do backwards calculations from a measurement site (receptor) for the volume mixing ratio of a generic inert tracer with infinite lifetime. The generic inert tracer consists of 40 000 particles released in a small box at the measurement location and during the measurement interval (3 h) and followed backward in time for 20 days. The model output unit is called PES, acronym from “potential emission sensitivity” distribution. The unit of the PES is in s kg⁻¹ and its value in a particular grid (geographical place) is proportional to the particle residence time in that cell. It is a measure for the simulated mixing ratio at the receptor that a source of unit (1 kg s⁻¹) in the respective grid cell would produce. For more details on the way that the model works, the reader is sent to the FLEXPART descriptive paper of Stohl et al. (2005).

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3 Results and discussion

In order to understand the OH reactivity results, a short description of the air masses impacting the site and a brief discussion on the pollution levels registered in Paris during the MEGAPOLI winter campaign are presented as follows.

3.1 Air mass origins during MEGAPOLI

OH reactivity measurements were made from 23 January until 13 February. During this time, a wide variety of air masses passed over Paris and its surroundings. In the supplementary information of the article, all of these air masses are synthesized along with their origins (see Appendix A in Supplement), as represented by the 20 days back trajectories calculated with the FLEXPART model.

We have chosen for discussion in this paper three contrasting periods characterising different regimes in the OH reactivity measurements. The corresponding backward trajectories are given in Fig. 5. The first period (hereafter called “period I”) starts with 26 January, 03:00, and ends on 27 January, 21:00. This period corresponded to air masses originating from the eastern part of Europe, passing firstly over the Benelux area and in the second part of this period over Germany.

The second period (hereafter called “period II”) started on 2 February, 09:00, and ended on 4 February, 03:00. In this period, clean air masses originating from the Atlantic Ocean arrived over Paris.

The third period (hereafter called “period III”) started with 9 January, 18:00, and lasted until the end of the campaign, 13 February, 16:00. The air masses originated again from the northern part of the continent, passing over Germany before joining Paris.

In conclusion, periods I and III were therefore characterized as “continental air masses”, while period II was characterized as “oceanic air masses”.

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.2 Air quality in Paris during MEGAPOLI

3.2.1 Ambient levels of VOCs in Paris – comparison to levels observed in other urban environments

The median values of the main gas tracers recorded in Paris over the entire MEGAPOLI winter campaign were: 300 ± 98 ppbv for CO, 27 ± 11 ppbv for NO₂ and 13 ± 8 ppbv for ozone (O₃). NO presented a high variability during this period: a mean value of 10 ppbv and a median of 5.5 ppbv. Previously a campaign has been conducted in Paris during spring 2007 (Gros et al., 2011) and a brief comparison with the pollutant levels is given on the same site as the MEGAPOLI winter campaign 2010. The median values corresponding to CO, NO, NO₂ and O₃ measured during spring 2007 were 248 ± 80 ppbv for CO, 4.4 ± 1.6 ppbv for NO, 11.2 ± 7.4 ppbv for NO₂ and 26 ± 13 ppbv for O₃. It can be seen that higher levels of CO, NO and NO₂ were registered during MEGAPOLI 2010 than for the 2007 campaign, and this is likely because of the higher anthropogenic emissions expected during the cold season. Because of a lower photochemical activity during winter, the O₃ level measured during MEGAPOLI 2010 campaign was lower than in spring 2007.

A brief comparison between the pollution levels reported from several cities where OH reactivity measurements are available is presented in Table 1. These cities were: Houston, New York, Mexico City, Tokyo and Paris. The background and the maximum registered levels of CO, NO_x, O₃ and the sum of non-methane hydrocarbons (NMHC) are summarized. The periods of the year when the measurements were done and the total population of the Metropolitan Areas were also presented. The most populated city was Tokyo with 36 million inhabitants, followed by New York and Mexico City, each of 22 million inhabitants. Paris and its surrounding area, called Ile de France, counted for 12 million inhabitants, representing one of the few megacities in Europe. Among the studied cases, Tokyo (36° N, 140° E) presented measurements in winter, in the same period when the measurements of Paris were realized. As Tokyo is at about 35° N one may expect the winter effect of more emissions to be less pronounced.

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Total OH reactivity
measurements in
Paris**C. Dolgorouky et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The measured levels of the compounds presented in the Table 1 correspond to median daily values. The background values correspond to midnight median levels, while the maximums represent the maximum registered of the median daily values. For more details on the measured data for Houston, New York and Mexico City, the reader is sent to Shirley et al. (2006) and Mao et al. (2009) and references therein. For Tokyo measurements, see Yoshino et al. (2006).

In terms of carbon monoxide, the background levels measured in Paris are similar to values measured in the other cities, though slightly lower. In winter, Tokyo registered a maximum of CO 3 to 4 times higher than the Paris maximum, but 2 times lower than the maximum observed for Mexico-City. Nevertheless, the maximum level of CO in Paris measured in the cold season remains comparable to maximums observed in New York City and Houston, though smaller and for different periods.

In terms of NO_x, the Paris background is slightly higher than the background levels measured in the other cities. Nevertheless, the maximum value remains comparable to the values measured in New York City and Tokyo (in summer and in autumn), but still almost two times lower than the maximum registered for the Mexico City in spring season and Tokyo in winter season. This is a comparison only on levels and does not consider inter annual variation of this pollutant concentration.

Generally, the Paris background level of NMHC sum was two times lower than in New York and Houston, 2000. The maximum is being 30 % to 50 % lower than those of New York and Houston, 2000 respectively. The high values in the sum of the NMHC measured for Houston in 2006 were mainly due to spikes at night correlated to emission events from plants in the Houston Ship Channel (Day et al., 2010). Greater discrepancies were nevertheless noted with respect to Mexico City when both the background and the maximum levels were 10 times higher than in Paris.

This quick comparative analysis allowed us to conclude that Paris presents intermediate levels of gaseous pollutants (CO, NO_x, NMHC), between medium size cities (Houston) and megacities like Tokyo, New York and Mexico City. This is explained by its smaller population and its less industrialized character.

Examining now the speciated VOCs measured during the MEGAPOLI winter campaign, Table 2 gives their mean and range over the measurement period. Some compounds (ex. Benzene, Toluene) were measured by several instruments and measurements from the instrument with the better time resolution were selected. Their corresponding OH reactivity, calculated as the product between the concentration and the equivalent reaction rate constant (Kovacs and Brune, 2001; Kovacs et al., 2003; Sadanaga et al., 2005; Mao et al., 2009, 2010) is also given and will be discussed in Sect. 3.4.

Alkanes, alkenes-alkynes (and isoprene, though usually under the detection limits, because of the cold season), aromatics and the oxygenated compounds accounted for 26, 19, 7 and 48 % respectively of the total volume mixing ratio of the measured VOCs. The oxygenated VOCs contribute a large part to the total concentration of the compounds measured during the campaign. Among the oxygenated species, ethanol by itself accounted for more than half (56 %) to the total OVOC concentration, followed by the sum of methanol and acetaldehyde 30 %, and acetone with 8 %.

VOC Source apportionment analysis was conducted by Gaimoz et al. (2011) for the spring 2007 campaign, allowing the characterization of the local pollution in Paris. The term “local pollution” means pollution from Paris and “Ile de France”. The main source identified for VOC was traffic emission (fuel exhaust and evaporation) contributing to 50 %. Other contributing sources included natural gas emissions, background input and biogenic emissions.

3.2.2 VOC variability: influence of local sources versus long-range transport

To characterize the air quality of Paris during the MEGAPOLI winter campaign, the variability of specific compounds showing different sources and atmospheric lifetimes were analyzed. The three panels of the Fig. 6 present selected compounds and a discussion on their variability, sources and their relation to the air mass origin is given afterwards.

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In the upper panel of Fig. 6, nitrogen oxide and the sum of o, m and p xylenes are presented. The baseline levels of these two anthropogenic and short life-time (a few hours) compounds do not seem to be significantly influenced by the different origins of the air masses. Secondly, as previously presented, the NO represents a typical marker of the traffic exhaust emissions. The covariation of the xylenes with the NO suggests that the xylenes' variability (but also that of other aromatic compounds, like toluene) in Paris was mainly controlled by the traffic source. This finding is in agreement with the conclusions from Gaimoz et al. (2011) (see Sect. 3.2.1).

The second panel presents two anthropogenic compounds with longer lifetimes: CO and benzene. CO represents a combustion tracer, having fossil fuel and biomass burning as primary sources. Its atmospheric life-time is relatively long, ~2 month (Crutzen, 1994). Benzene has a lifetime of 12 days (for an OH concentration of 1.5×10^6 molecules cm^{-3} , Seinfeld and Pandis, 2006) and its emissions in Paris are, like other aromatic compounds, linked to traffic activities. Both compounds show higher baseline levels during periods I and III. As the life-time of these two compounds exceeds the day, additional imported sources are considered to be responsible for the higher levels registered during the periods where polluted air masses were originating from the continent.

In the third panel, two oxygenated compounds with different lifetimes are presented: acetone (66 days for an OH concentration of 1.5×10^6 molecules cm^{-3} , Seinfeld and Pandis, 2006) and acetaldehyde (11 h for an OH concentration of 1.5×10^6 molecules cm^{-3} , Seinfeld and Pandis, 2006). These two compounds have primary sources mainly of biogenic origin and an important secondary source: the oxidation of non-methane hydrocarbons (NMHC). As shown in Fig. 6, both compounds show relatively lower baseline levels during the oceanic episode (period II) (except for the two maximums observed for the acetone) and higher baseline levels monitored during the continental air masses episodes (periods I and III). The fact that this increase is observed also for acetaldehyde, an oxygenated compound with a relatively short life-time, suggests that during the continental imported air masses, two type of sources are

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



responsible of the high pollution levels: the direct pollution and the imported from over the continent and already processed air masses.

In summary, the examination of the variability of the compounds that have different lifetimes and sources shows that the measurement campaign was impacted during Period II only by local and regional (Paris and its surroundings, so-called “Ile de France”) emissions, whereas during the other periods (periods I and III), a significant contribution to the pollution levels in Paris was due to direct import of primary and secondary pollutants emissions that had undergone photochemical aging and mixing during transport to Paris.

The diurnal variation of the main gaseous pollutants in Paris is presented in Appendix B in Supplement. We have observed that the primary species – CO, NMHC and NO – show the same daily profile related to their common source, the traffic emissions. They show a minimum value during the night, a first peak in the early morning, a slight decrease in the afternoon and a second peak formed in the evening. This double peak profile of the primary pollutants has also been remarked in Paris during spring 2007 (Gros et al., 2011) but also in other city boundaries (de Gouw et al., 2009; Velasco et al., 2007). The main primary emission source in Paris is considered to be the traffic activity.

3.3 OH reactivity in Paris

3.3.1 OH reactivity variability during the MEGAPOLI campaign

The OH reactivity measurements were made from 23 January until 13 February. Figure 7 presents the OH reactivity levels measured: the 2 min and the 30 min averaged data series.

OH reactivity ranged from below the detection limit (3.5 s^{-1}) up to 130 s^{-1} . The median value over the entire campaign was of 33 s^{-1} . Missing points in the data set correspond to periods when instrument check and calibrations were run.

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Total OH reactivity
measurements in
Paris**C. Dolgorouky et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Generally we observed higher OH reactivity levels measured during the night time compared to daytime levels. This variability of the OH reactivity is explained by the evolution of the height of the boundary layer, which is lower during the night, and therefore at this time the OH reactivity is less diluted. This result is in good agreement with the higher level of the OH reactivity during nighttime than during daytime (except for the double peak due to traffic emissions) measured in other megacities around the world: Nashville, USA (1999) (Kovacs et al., 2003), Houston, USA (2000 and 2006) (Mao et al., 2010), New York (2001) (Ren et al., 2003) and Mexico City (2003) (Shirley et al., 2006).

These same studies showed also that, under the influence of local pollution, the OH reactivity usually followed the diurnal cycle of the pollutants emitted by traffic emissions: a first peak more or less accentuated, in the early morning (06:30), followed by a slight decrease in the late morning (10:30) and a second peak in the late after-noon and early evening. Nevertheless, we noticed that very few days during this campaign presented the typical diurnal (diel) cycle of a city impacted by traffic emissions. The main reason for this was that the diurnal variability was most of the time overtaken by the impact of long range transport as discussed below.

Regarding the entire MEGAPOLI winter campaign, three contrasting periods have been identified. The origins of the air masses corresponding to these OH reactivity periods have been previously described (see Sect. 3.1).

The “Period I” started with 26 January, mid-day, and ends on 27, at 21:00 h. The period is characterized by high OH reactivity levels with a maximum value of 120 s^{-1} reached on 27 January. The air masses originated from the eastern part of the continent and passed over Germany, before reaching Paris. They were therefore charged in pollutants (see discussion from Sect. 3.2.2), and can explain the high OH reactivity levels registered during “period I”.

“Period II” lasted from 2 to 4 February. During this period, low levels of OH reactivity were registered (20 s^{-1}). This result is consistent with the marine origin of the air masses during this period (see Fig. 5).

“Period III” started with 8 February, 15:00, and lasted until the end of the campaign, 13 February. The OH reactivity levels reached again very high values of 110 s^{-1} . As for “period I”, the air masses mainly originated from the north-eastern part of the continent and the Benelux area. Therefore an additional pollutant source due to the long range transport is considered to be responsible for the higher levels of the measured OH reactivity.

When considering the other time periods not-discussed here (time periods between period I, II and III), the air masses characterized intermediate regimes. The air masses originated either from the central part of France, either from intermediate regimes between the continental and oceanic (see Appendix A in Supplement). These periods correspond to intermediate OH reactivity levels.

Table 3 synthetises the mean values of OH reactivity and the main gaseous pollutants for the three periods previously mentioned.

Generally all the compounds showed lower levels during the marine air mass period (II), than for the continental import periods (I and III), as already described in the previous section. The primary compounds exclusively emitted by local sources (considering their short life-time), here presented by the NO and the xylenes, show levels almost double during “period I” compared to “period III”, both continental air masses periods. These high concentrations denote higher local pollution during period I, and implicitly, higher “local” OH reactivity. The other primary compounds presented (CO, benzene and NMHC) showed also the same difference between the two continental import periods, but because of their longer life-time, we can conclude that an additional imported source of primary compounds contributes to the total OH reactivity. Keeping in mind that the OH reactivity data are associated with an uncertainty of about 24 %, some differences can still be seen between these two different periods.

Surprisingly, the sum of the oxygenated VOCs shows higher values during the oceanic air masses period. This result is due to ethanol (which represents 50 % of the OVOCs) and which shows higher level during period II. Biogenic emissions are considered to be the main source of ethanol (Naik et al., 2010) and we note that Lewis

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

et al. (2005) observed also higher OVOCs compared to other species in North Atlantic marine air too.

We conclude that the lowest OH reactivity levels corresponded to the marine period and were than only impacted by the local and regional pollution. The higher OH reactivity registered during periods I and III was mainly due to long distance transport.

3.3.2 OH reactivity levels registered over a local pollution episode

The previous section suggested that the Paris OH reactivity larger variability is mainly driven by long range transport when comes to continental air masses. A discussion concerning the diel variation of the OH reactivity can therefore be done on periods influenced by oceanic air masses, and when only the local pollution is observed. The day of 4 February (Thursday) was chosen because of its mainly oceanic air mass origins (pure oceanic air masses before 09:00, and oceanic and central France after 09:00; see Fig. A1). We specifically use this day as it was the only one available with marine air mass origins and enough OH reactivity data. The other days had either larger blank period in the data set due to daily instrumental check, or either did not represent the typical diurnal cycle governed by the morning and evening pollution peaks because they were weekend days. Still, the 4 February presented a blank of 2 h in the data set, from 09:30 to 11:30, which was corresponding to the base line measurements.

Figure 8 presents in the top panel the variation of the OH reactivity with NO and in the bottom panel the OH reactivity plotted with acetone.

For this local pollution episode, NO showed a modified diurnal cycle. Indeed, instead of a double peak normally observed during the day and corresponding to the traffic rush hours, on 4 February, NO showed only the morning peak which was doubled. A small peak was observed in the early morning (06:30), followed by a larger one in the late morning, around 11:00. We note that 4 February was a Thursday, when the local public transportation had a strike, probably leading to enhanced use of private vehicle and unusual rush hour time. This conclusion is also sustained by the double

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



peak observed for the same day on other compounds related to traffic emissions: CO and xylenes.

The first maximum in the NO level corresponding to the morning rush hour is clearly seen on the OH reactivity as well as the decrease observed around 08:00 and the beginning of the increase observed at 09:00. Unfortunately the second maximum is not captured by the OH reactivity measurement because of two hours gap in the data set. The good agreement between the OH reactivity and the NO denotes that Paris local OH reactivity is well impacted by traffic related emissions. Nevertheless, in the second part of the day, an increase compared to the morning level (06:00) of the OH reactivity was observed, increase which was not showed by the NO variability. The second plot of Fig. 8 shows that acetone presents the same change in the concentration level, at the same time of the day. As acetaldehyde shows the same increase of the level measured as acetone, we concluded that for this episode, the in-situ photochemical formation of oxygenated compounds may be responsible of the increase of the OH reactivity in the afternoon.

3.4 Missing OH reactivity in Paris

The calculated OH reactivity represents the sum of the product of OH reactants and their reaction rate coefficient. As previously shown (Kovacs et al., 2003; Di Carlo et al., 2004), comparison between the measured and the calculated OH reactivity denotes the so-called “missing OH reactivity” and tests the assumption that all significant OH reactants have been measured and can be used to infer missing OH reactants.

Figure 9 presents the total measured OH reactivity plotted with the calculated OH reactivity due to the simultaneously measured OH sink compounds. The VOC species considered for these calculations are presented in Table 2. In addition, results of CO, NO and NO₂ were also taken in consideration. Because of the winter season, no biogenic emissions were identified (isoprene signal under the detection limit most of the time), and therefore no biogenic compounds were considered. No formaldehyde measurements were made at the LHVP site during the MEGAPOLI winter campaign.

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nevertheless, calculations considering high values of formaldehyde were conducted but the equivalent OH reactivity was not significant (only 2 s^{-1} for 10 ppbv of formaldehyde, $k_{\text{CH}_2\text{O}+\text{OH}} = 8.5 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, Atkinson et al., 2006). No formaldehyde reactivity was therefore considered. The CH_4 concentration of 1950 ppbv was assumed. Due to the temporal resolution of the instruments, calculated reactivity was performed every 1 h and 30 min.

When comparing the measured and the calculated OH reactivity, the calculated level is systematically smaller than the measured one, with the exception of 5 and 8 February, when instrumental interferences were noted. The median (mean) value of the calculated OH reactivity over the entire campaign is 16 s^{-1} (17.5 s^{-1}), while the measured OH reactivity is double, 33 s^{-1} (40 s^{-1}).

The breakdown of the total OH reactivity measured during the entire campaign is presented in Fig. 10. The mean values were taken in consideration for the calculations. The missing OH reactivity, given by the difference between the measured and the calculated reactivity, is quite high: 54 %, showing that when averaged over the entire campaign, more than half of the measured OH reactivity is not explained. Pie chart b presents the contribution of the different compounds to the calculated OH reactivity. The main compounds explaining the calculated part of the OH reactivity are NO_2 (55 %), followed by the sum of NMHCs (15 %), NO and CO (accounting each for 10 %) and finally the sum of OVOCs (8 %).

Examining now the missing reactivity according to the air mass origins, we note significant difference if the air mass was of oceanic or of continental origin. Figure 11 presents the enclosed piecharts of the breakdown total OH reactivity for periods II and III. During the period associated with oceanic air masses and therefore characteristic of the local pollution in Paris, the measured and the calculated OH reactivity agreed within 12 % (see piechart a), which is within the uncertainty range ($\pm 24 \%$) of the measured and calculated reactivity. This good agreement points out that during period II, all the main OH reactive species were measured. As Paris is impacted by traffic emissions, high levels of NO_x are emitted into the atmosphere. NO_2 and NO explain about 60 %

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

of the total measured OH reactivity. This large contribution is expected as in urban atmosphere, the main sink of OH being the formation of the nitric acid (HNO₃) (Delmas et al., 2005). The OH reactivity due to the NMHC accounts for ~14 %, while the OVOC and the CO contribute with 7 and respectively 8 % to the total OH reactivity. When considering only the measured VOCs presented in Table 2, among the NMHC, the species that contributed the most to the OH reactivity were ethene with 13 % and propene with 7.5 %. Among the oxygenated VOCs, the acetaldehyde and ethanol contributed with 16 % and respectively 14 % to the OH reactivity.

We note a very good agreement between the contribution of the different compounds to the total OH reactivity on the entire campaign (see piechart of Fig. 10) and during the local pollution period (pie chart a) from Fig. 11). This clearly points out that when not under continental pollution plumes, the species which were measured during the MEGAPOLI campaign (CO, NO_x and list of VOCs from Table 2) characterize correctly the Paris OH reactivity levels.

Part b of Fig. 11 indicates that during continental import, the measured OH reactivity was much higher than the calculated value. The missing OH reactivity was now of 75 %. This is a huge value compared to missing OH reactivities registered in other urban areas (for more details, see Sect. 3.5.3). Calculations indicate that the reactions of OH with NO₂ are the greatest contributors to the total OH reactivity, and followed by NMHC, CO, OVOCs and NO.

For a better understanding of the variability of the not-measured OH reactivity nature, Fig. 12 presents on the top panel the missing OH reactivity series.

As already predicted, the highest values were registered during the continental air mass imports, while the lowest level was registered under the marine air mass influence. We can conclude that during the local pollution period all of the main OH reactive species were measured and therefore the OH reactivity budget can be closed simply by using the already measured compounds. On the contrary, during continental import periods, the non-measured OH reactive species are considered to be at the origin of the high level of the missing OH reactivity. Therefore these non-measured species could

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

be: (a) either primary OH reactive compounds, or (b) secondary species, characteristic of reprocessed air masses. The first suggestion is less probable, if we consider that the most important classes of OH reactive species, already measured in other studies, were also measured during the MEGAPOLI winter campaign. Furthermore, the good agreement between the measured and calculated OH reactivity found for “period II” reinforces this conclusion.

The temporal variability of submicron secondary inorganic salts (ammonium nitrate and ammonium sulphate) measured by AMS is reported in Fig. 12 (second panel). At many locations ammonium nitrate has shown to be correlated with freshly (less-) photochemically aged organic aerosols (OA) whereas ammonium sulphate was more related to regional heavily aged OA (Lanz et al., 2007; Jimenez et al., 2009; Ulbrich et al., 2009). The sum of these two secondary ion components may be then used to infer the role of photochemical processes involved in the formation of secondary OA from ambient VOC. Low concentrations of these ions were observed during periods with oceanic air masses whereas significantly higher concentrations were observed during periods I and III which are influenced by continental air masses. These observations are consistent with those reported in Paris during springtime by Sciare et al. (2010) and clearly point to the major influence of photochemically processed continental EU anthropogenic emissions. The good agreement observed in Fig. 12 between the temporal variability of these ions and the missing OH reactivity strengthens the previous idea that during continental import, the OH reactivity in Paris is impacted by the non-measured oxidized compounds issues of the reprocessing of the air masses (products issued from multi-oxidation of primary compounds).

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.5 Comparison with OH reactivity in other urban areas

3.5.1 Comparison of OH reactivity levels with levels measured in other cities around the world

Table 4 synthesises the median and the maximum OH reactivity levels previously measured in other urban areas around the world.

As expected, the lowest median values were noticed in small cities. The median values registered in Nashville (United States), Houston (United States) and Mainz (Germany) hardly exceeded 10 s^{-1} . The highest OH reactivity value registered corresponded to summer measurements in Tokyo (40 s^{-1}), followed by Mexico City with 33 s^{-1} . Intermediate values were observed for New York City, when for both the winter and the summer campaign, the OH reactivity was about 20 s^{-1} . When considering the median value of the entire winter campaign, the Paris OH reactivity was pretty high, of about the same order as for Mexico City, 33 s^{-1} . Nevertheless, this discussion must take into consideration the two contrasting periods identified for the MEGAPOLI winter campaign and which were correlated with different air mass origins (see Sect. 3.3.1). During the local pollution period characterized by the oceanic air masses, the OH reactivity levels registered were quite low, about 20 s^{-1} , value comparable with values registered in New York for the same period of the year, but also comparable to the values registered in Tokyo during winter. When considering the continental long range transport, the median of the OH reactivity levels was much higher: 60 s^{-1} .

The maximum OH reactivity observed during the MEGAPOLI winter campaign was of the same order of magnitude as the maximum measured in Tokyo and New York, but lower than the maximum OH reactivity measured in Mexico City.

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.5.2 Analysis of the contributions of different atmospheric constituents to OH reactivity in different megacities

Figure 13 presents the contributions from different atmospheric constituents to the calculated OH reactivity for 5 different urban atmospheres and different periods of the year. For Tokyo values, the data was estimated from Fig. 9 (page 7879) of the Yoshino et al. (2006) reference. The Mexico City, New York and Houston data were obtained from Mao et al. (2010).

All of the studies presented in Fig. 13 take into consideration measurements of inorganic compounds (CO, NO_x), hydrocarbons (including isoprene), formaldehyde, oxygenated VOCs, but also biogenic emissions, with the exception of Paris, where no formaldehyde and biogenics (because of the winter season) were available. For more details see column “measured species” from Table 2. We can therefore conclude on the pertinence of the further discussion.

The main classes of the compounds actually identified as the main OH reactivity sinks were taken into consideration in Fig. 13. These are: CO, NO_x (sum of NO and NO₂), the sum of non-methane hydrocarbons (NMHC), sum of oxygenated VOCs (OVOCs) and the sum of biogenic VOC (BVOCs). Function of the season, the BVOC can be negligible. That is the reason why in Paris, the BVOCs contribution was not quantified. For the same reason, the BVOC contribution in Tokyo was small (<1 %). As the New York study was performed during summer (see Table 4), the contribution from the BVOC counted for 4 %.

The contribution from CO is very much alike in the five studies (12–15 %), although different seasons are considered. The contribution from the NO_x exceeds 50 % for Paris, Tokyo and New York cities, showing the large influence of the traffic-related emissions to the calculated OH reactivity. Still, the Paris NO_x contribution is the highest registered among the five studies (65 %), strengthening the idea that the local pollution in Paris is mainly impacted by traffic emissions. Mexico City and Houston showed low values of NO_x (only 23 % and respectively 15 %), but the contribution from the

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



NMHCs reaches 50 % in both cases. Although Mexico City and New York (comprising their Metropolitan Areas) are comparable in terms of population and size, the NMHC level registered in New York City is rather low because of the more advanced pollution abatement efforts deployed here. Previous study (de Gouw et al., 2006) showed that in Mexico City the high NMHC is due to higher biomass fuel being burnt and therefore explaining the high contribution from the NMHC to the OH reactivity. In the case of Houston, Leuchner and Rappenglück (2010) announced high industrial solvent emissions. Higher contributions from aromatics (15 %) were therefore reported for Houston, compared to only 3.6 % in Paris, 4 % in New York City and 7 % in Mexico City. The contribution to the total OH reactivity from behalf of the oxygenated VOCs is respectively low, comprised between 4 %, for Tokyo, 8 % for Paris and 14 % announced for New York and Houston.

3.5.3 Missing OH reactivity in other urban areas

Table 4 synthesises the percentages of the missing OH reactivity as previously measured and reported into the literature. Paris results present the highest missing OH reactivity determined in urban areas (up to 75 %). This is followed by New York and Tokyo with the values of the missing OH reactivity of about 30 %. A high missing OH reactivity has been registered also in Nashville, but in this case, the greater measured-than-calculated OH reactivity was due to the presence of the non-measured species (no biogenics were measured, although measurements were realized in summer time) during this campaign (Lewis et al., 2000; Kovacs et al., 2003).

OH reactivity measurements were performed for New York during summer time but also during winter and the results showed that the missing OH reactivity was higher in winter time than in summer. Opposite results have been obtained in Tokyo, when there was almost no missing OH reactivity during winter. As the suite of simultaneous VOC measurements was inadequate, no missing OH reactivity levels are available for the Mexico City (Shirley et al., 2006).

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The value of the missing OH reactivity registered in Paris during period II was in good agreement with the winter results in Tokyo and autumn in Houston.

4 Summary and conclusions

OH reactivity measurements using the CRM method were realized in Paris between 23 January and until 13 February, during the MEGAPOLI winter campaign 2010. As the measurements have been realized in strong NO field (10 to 160 ppbv), correction factors were applied to the data set to overcome the NO interference into the setup.

OH reactivity ranged from below the detection limit (3.5 s^{-1}) up to 130 s^{-1} . The median value registered for the entire campaign was of 33 s^{-1} . As Paris and its surrounding region (Ile de France) are surrounded by a rural area, it was possible to separate continental background influence from oceanic background influence, when discussing the OH reactivity levels. Considering these two main influences, we could conclude that during the marine air mass origins, the pollution in Paris was mainly due to traffic-related emissions and the measured OH reactivity level was of 20 s^{-1} , value comparable to values measured in other urban areas (New York and Tokyo). When considering the contribution of different atmospheric constituents to the OH reactivity in different megacities, two main profiles of urban areas have been identified: one related to traffic emissions, the other one to industrial emissions. Therefore, Paris has the same traffic-related emission profile as Tokyo and New York, the main compounds that contribute to the calculated OH reactivity being the NO_x , with more than 50 %, followed by NMHCs (15 to 28 %), CO (around 14 %) and OVOCs (less than 14 %). Mexico City and Houston were identified as cities with industrial profiles, the main contributor to the calculated OH reactivity being the NMHCs (around 50 %), followed by NO_x (15 to 28 %), OVOCs (11 to 14 %) and CO (14 %).

During the continental import air masses, an additional imported pollution added to the local one and very high OH reactivity levels were announced (120 s^{-1}).

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Total OH reactivity
measurements in
Paris**C. Dolgorouky et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Additional measurements and analysis of the variability of a large panel of compounds simultaneously measured within the MEGAPOLI framework (NO_x , CO , O_3 , NMHCs and OVOCs) pointed out a large part of the missing OH reactivity. When considering the entire MEGAPOLI campaign, the total missing OH reactivity was of 54 %, showing that when averaged over the entire campaign, more than half of the measured OH reactivity is not explained. The found value is higher than the values reported in other urban areas and when the maximum missing OH reactivity was of 33 % (Sadanaga et al., 2004; Ren et al., 2006). Nevertheless, during the marine air mass influences, the missing OH reactivity is of only 12 %, value comprised within the measurement uncertainties and which has shown that during the local pollution periods, the OH reactivity budget can be closed only by using the already measured compounds. Still, surprisingly, during the continental imported air masses, 75 % of the measured OH reactivity could not be explained by the measured species. This percentage was higher than the missing OH reactivity values registered in other urban areas. The missing OH reactivity showed a similar variation as longer life-time compounds and photochemically produced aerosol species suggesting that air masses corresponding to continental import were probably loaded in highly oxidized products, probably of anthropogenic origin.

We therefore concluded that levels of missing OH reactivity of the same magnitude as in forested areas were registered for the first time in an urban area. Still, if for the forested area, previous studies (Di Carlo et al., 2004; Sinha et al., 2010; Kim et al., 2011) identified the biogenic VOCs as mainly responsible of these high levels, in the case of Paris, the non-measured compounds were likely (multi)oxidized compounds issued from processed anthropogenic emissions.

Recently, Kato et al. (2011) proposed a relative rate method using GC-FID detectors to estimate the contribution of the OH reactivity of unidentified VOCs in the atmosphere of Tokyo (study conducted in spring 2004). They found that the contribution of the unidentified species was about 22 % of that for all the VOCs, and that when taken into consideration they improve the percentage of the missing OH reactivity that can be

explained. Still, the contribution to the total OH reactivity of the unidentified species using this method is underestimated because species with high polarity or heavy species are being lost during the concentration process.

Considering the vast complexity that the identification and quantification of the unknown OH reactive species in an urban environment pose for air quality, further studies involving more measurements of secondary compounds are required.

Supplementary material related to this article is available online at:
**[http://www.atmos-chem-phys-discuss.net/12/10937/2012/
acpd-12-10937-2012-supplement.pdf](http://www.atmos-chem-phys-discuss.net/12/10937/2012/acpd-12-10937-2012-supplement.pdf)**

Acknowledgements. We would like to thank T. Leonardis and N. Locoge (EMD) for support with GC-MS measurements, B. Temime (LCP) for the PTR-MS measurements and C. Gaimoz (LSCE and now at LISA) for the GC-FID measurements. We would like to thank also the MEGAPOLI field campaign participants and especially M. Beekmann for coordinating the field campaigns and A. Stohl (NILU) for providing Flexpart results. The research leading to these results has received funding from the European Union's Seventh Framework Programme FP/2007–2011 within the project MEGAPOLI grant agreement no. 212520. Support from CEA, CNRS, ADEME, ANR-MEGAPOLI are acknowledged.



The publication of this article is financed by CNRS-INSU.

ACPD

12, 10937–10994, 2012

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

- Ammann, C., Brunner, A., Spirig, C., and Neftel, A.: Technical note: Water vapour concentration and flux measurements with PTR-MS, *Atmos. Chem. Phys.*, 6, 4643–4651, doi:10.5194/acp-6-4643-2006, 2006.
- 5 Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., and Troe, J.: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume I – gas phase reactions of O_x, HO_x, NO_x and SO_x species, *Atmos. Chem. Phys.*, 4, 1461–1738, doi:10.5194/acp-4-1461-2004, 2004.
- 10 Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., Jenkin, M. E., Rossi, M. J., Troe, J., and IUPAC Subcommittee: Evaluated kinetic and photochemical data for atmospheric chemistry: Volume II – gas phase reactions of organic species, *Atmos. Chem. Phys.*, 6, 3625–4055, doi:10.5194/acp-6-3625-2006, 2006.
- 15 Canagaratna, M. R., Jayne, J. T., Jimenez, J. L., Allan, J. D., Alfarra, M. R., Zhang, Q., Onasch, T. B., Drewnick, F., Coe, H., Middlebrook, A., Delia, A., Williams, L. R., Trimborn, A. M., Northway, M. J., DeCarlo, P. F., Kolb, C. E., Davidovits, P., and Worsnop, D. R.: Chemical and microphysical characterization of ambient aerosols with the aerodyne aerosol mass spectrometer, *Mass Spectrom. Rev.*, 26, 185–222, doi:10.1002/mas.20115, 2007.
- Crutzen, P. J.: Global budgets for non-co₂ greenhouse gases, *Environ. Monit. Assess.*, 31, 1–15, doi:10.1007/bf00547177, 1994.
- 20 Day, B. M., Rappengluck, B., Clements, C. B., Tucker, S. C., and Brewer, W. A.: Nocturnal boundary layer characteristics and land breeze development in houston, texas during texaqs ii, *Atmos. Environ.*, 44, 4014–4023, doi:10.1016/j.atmosenv.2009.01.031, 2010.
- de Gouw, J. A., Warneke, C., Stohl, A., Wollny, A. G., Brock, C. A., Cooper, O. R., Holloway, J. S., Trainer, M., Fehsenfeld, F. C., Atlas, E. L., Donnelly, S. G., Stroud, V., and Lueb, A.: Volatile organic compounds composition of merged and aged forest fire plumes from alaska and western canada, *J. Geophys. Res.-Atmos.*, 111, D10303, doi:10.1029/2005jd006175, 2006.
- 25 de Gouw, J. A., Welsh-Bon, D., Warneke, C., Kuster, W. C., Alexander, L., Baker, A. K., Beyersdorf, A. J., Blake, D. R., Canagaratna, M., Celada, A. T., Huey, L. G., Junkermann, W., Onasch, T. B., Salcido, A., Sjostedt, S. J., Sullivan, A. P., Tanner, D. J., Vargas, O., Weber, R. J., Worsnop, D. R., Yu, X. Y., and Zaveri, R.: Emission and chemistry of organic carbon
- 30

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Total OH reactivity
measurements in
Paris**

C. Dolgorouky et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

in the gas and aerosol phase at a sub-urban site near Mexico City in March 2006 during the MILAGRO study, *Atmos. Chem. Phys.*, 9, 3425–3442, doi:10.5194/acp-9-3425-2009, 2009.

DeCarlo, P. F., Kimmel, J. R., Trimborn, A., Northway, M. J., Jayne, J. T., Aiken, A. C., Gonin, M., Fuhrer, K., Horvath, T., Docherty, K. S., Worsnop, D. R., and Jimenez, J. L.: Field-deployable, high-resolution, time-of-flight aerosol mass spectrometer, *Anal. Chem.*, 78, 8281–8289, doi:10.1021/ac061249n, 2006.

Delmas, R., Mégie, G., and Peuch, V.-H.: *Physique et chimie de l'atmosphère*, Ed. Belin, 2005.

Di Carlo, P., Brune, W. H., Martinez, M., Harder, H., Leshner, R., Ren, X. R., Thornberry, T., Carroll, M. A., Young, V., Shepson, P. B., Riemer, D., Apel, E., and Campbell, C.: Missing oh reactivity in a forest: Evidence for unknown reactive biogenic vocs, *Science*, 304, 722–725, doi:10.1126/science.1094392, 2004.

Gaimoz, C., Sauvage, S., Gros, V., Herrmann, F., Williams, J., Locoge, N., Perrussel, O., Bonsang, B., d'Argouges, O., Sarda-Esteve, R., and Sciare, J.: Volatile organic compounds sources in paris in spring 2007. Part ii: Source apportionment using positive matrix factorisation, *Environ. Chem.*, 8, 91–103, doi:10.1071/en10067, 2011.

Goldstein, A. H. and Galbally, I. E.: Known and unexplored organic constituents in the earth's atmosphere, *Environ. Sci. Technol.*, 41, 1514–1521, doi:10.1021/es072476p, 2007.

Gros, V., Sciare, J., and Yu, T.: Air-quality measurements in megacities: Focus on gaseous organic and particulate pollutants and comparison between two contrasted cities, paris and beijing, *Comptes Rendus Geoscience*, 339, 764–774, 2007.

Gros, V., Gaimoz, C., Herrmann, F., Custer, T., Williams, J., Bonsang, B., Sauvage, S., Locoge, N., d'Argouges, O., Sarda-Esteve, R., and Sciare, J.: Volatile organic compounds sources in paris in spring 2007. Part i: Qualitative analysis, *Environ. Chem.*, 8, 74–90, doi:10.1071/en10068, 2011.

Ingham, T., Goddard, A., Whalley, L. K., Furneaux, K. L., Edwards, P. M., Seal, C. P., Self, D. E., Johnson, G. P., Read, K. A., Lee, J. D., and Heard, D. E.: A flow-tube based laser-induced fluorescence instrument to measure OH reactivity in the troposphere, *Atmos. Meas. Tech.*, 2, 465–477, doi:10.5194/amt-2-465-2009, 2009.

Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, A. S. H., Zhang, Q., Kroll, J. H., DeCarlo, P. F., Allan, J. D., Coe, H., Ng, N. L., Aiken, A. C., Docherty, K. S., Ulbrich, I. M., Grieshop, A. P., Robinson, A. L., Duplissy, J., Smith, J. D., Wilson, K. R., Lanz, V. A., Hueglin, C., Sun, Y. L., Tian, J., Laaksonen, A., Raatikainen, T., Rautiainen, J., Vaattovaara, P., Ehn, M., Kulmala, M., Tomlinson, J. M., Collins, D. R., Cubison, M. J., Dunlea, E. J., Huffman, J.

**Total OH reactivity
measurements in
Paris**

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A., Onasch, T. B., Alfarra, M. R., Williams, P. I., Bower, K., Kondo, Y., Schneider, J., Drewnick, F., Borrmann, S., Weimer, S., Demerjian, K., Salcedo, D., Cottrell, L., Griffin, R., Takami, A., Miyoshi, T., Hatakeyama, S., Shimono, A., Sun, J. Y., Zhang, Y. M., Dzepina, K., Kimmel, J. R., Sueper, D., Jayne, J. T., Herndon, S. C., Trimborn, A. M., Williams, L. R., Wood, E. C., Middlebrook, A. M., Kolb, C. E., Baltensperger, U., and Worsnop, D. R.: Evolution of Organic Aerosols in the Atmosphere, *Science*, 326, 1525–1529, doi:10.1126/science.1180353, 2009.

Kato, S., Sato, T., and Kajii, Y.: A method to estimate the contribution of unidentified vocs to oh reactivity, *Atmos. Environ.*, 45, 5531–5539, doi:10.1016/j.atmosenv.2011.05.074, 2011.

Kim, S., Guenther, A., Karl, T., and Greenberg, J.: Contributions of primary and secondary biogenic VOC total OH reactivity during the CABINEX (Community Atmosphere-Biosphere Interactions Experiments)-09 field campaign, *Atmos. Chem. Phys.*, 11, 8613–8623, doi:10.5194/acp-11-8613-2011, 2011.

Kovacs, T. A. and Brune, W. H.: Total oh loss rate measurement, *J. Atmos. Chem.*, 39, 105–122, doi:10.1023/a:1010614113786, 2001.

Kovacs, T. A., Brune, W. H., Harder, H., Martinez, M., Simpas, J. B., Frost, G. J., Williams, E., Jobson, T., Stroud, C., Young, V., Fried, A., and Wert, B.: Direct measurements of urban oh reactivity during nashville sos in summer 1999, *J. Environ. Monit.*, 5, 68–74, doi:10.1039/b204339d, 2003.

Lanz, V. A., Alfarra, M. R., Baltensperger, U., Buchmann, B., Hueglin, C., and Prévôt, A. S. H.: Source apportionment of submicron organic aerosols at an urban site by factor analytical modelling of aerosol mass spectra, *Atmos. Chem. Phys.*, 7, 1503–1522, doi:10.5194/acp-7-1503-2007, 2007.

Leuchner, M. and Rappenglueck, B.: Voc source-receptor relationships in houston during texaqs-ii, *Atmos. Environ.*, 44, 4056–4067, doi:10.1016/j.atmosenv.2009.02.029, 2010.

Lewis, A. C., Carslaw, N., Marriott, P. J., Kinghorn, R. M., Morrison, P., Lee, A. L., Bartle, K. D., and Pilling, M. J.: A larger pool of ozone-forming carbon compounds in urban atmospheres, *Nature*, 405, 778–781, doi:10.1038/35015540, 2000.

Lewis, A. C., Hopkins, J. R., Carpenter, L. J., Stanton, J., Read, K. A., and Pilling, M. J.: Sources and sinks of acetone, methanol, and acetaldehyde in North Atlantic marine air, *Atmos. Chem. Phys.*, 5, 1963–1974, doi:10.5194/acp-5-1963-2005, 2005.

Lindinger, W., Hansel, A., and Jordan, A.: On-line monitoring of volatile organic compounds at pptv levels by means of proton-transfer-reaction mass spectrometry (ptr-ms) – medical

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



applications, food control and environmental research, *Int. J. Mass Spectrom.*, 173, 191–241, 1998.

Lou, S., Holland, F., Rohrer, F., Lu, K., Bohn, B., Brauers, T., Chang, C. C., Fuchs, H., Häsel, R., Kita, K., Kondo, Y., Li, X., Shao, M., Zeng, L., Wahner, A., Zhang, Y., Wang, W., and Hofzumahaus, A.: Atmospheric OH reactivities in the Pearl River Delta – China in summer 2006: measurement and model results, *Atmos. Chem. Phys.*, 10, 11243–11260, doi:10.5194/acp-10-11243-2010, 2010.

Mao, J., Ren, X., Brune, W. H., Olson, J. R., Crawford, J. H., Fried, A., Huey, L. G., Cohen, R. C., Heikes, B., Singh, H. B., Blake, D. R., Sachse, G. W., Diskin, G. S., Hall, S. R., and Shetter, R. E.: Airborne measurement of OH reactivity during INTEX-B, *Atmos. Chem. Phys.*, 9, 163–173, doi:10.5194/acp-9-163-2009, 2009.

Mao, J., Ren, X., Chen, S., Brune, W. H., Chen, Z., Martinez, M., Harder, H., Lefer, B., Rappenglueck, B., Flynn, J., and Leuchner, M.: Atmospheric oxidation capacity in the summer of houston 2006: Comparison with summer measurements in other metropolitan studies, *Atmos. Environ.*, 44, 4107–4115, doi:10.1016/j.atmosenv.2009.01.013, 2010.

Naik, V., Fiore, A. M., Horowitz, L. W., Singh, H. B., Wiedinmyer, C., Guenther, A., de Gouw, J. A., Millet, D. B., Goldan, P. D., Kuster, W. C., and Goldstein, A.: Observational constraints on the global atmospheric budget of ethanol, *Atmos. Chem. Phys.*, 10, 5361–5370, doi:10.5194/acp-10-5361-2010, 2010.

Ren, X., Brune, W. H., Mao, J., Mitchell, M. J., Leshner, R. L., Simpas, J. B., Metcalf, A. R., Schwab, J. J., Cai, C., Li, Y., Demerjian, K. L., Felton, H. D., Boynton, G., Adams, A., Perry, J., He, Y., Zhou, X., and Hou, J.: Behavior of OH and HO₂ in the winter atmosphere in new york city, *Atmos. Environ.*, 40, S252–S263, doi:10.1016/j.atmosenv.2005.11.073, 2006a.

Ren, X., Brune, W. H., Olliger, A., Metcalf, A. R., Simpas, J. B., Shirley, T., Schwab, J. J., Bai, C., Roychowdhury, U., Li, Y., Cai, C., Demerjian, K. L., He, Y., Zhou, X., Gao, H., and Hou, J.: Oh, ho(2), and oh reactivity during the pmtacs-ny whiteface mountain 2002 campaign: Observations and model comparison, *J. Geophys. Res.-Atmos.*, 111, D10s03, doi:10.1029/2005jd006126, 2006b.

Ren, X. R., Harder, H., Martinez, M., Leshner, R. L., Olliger, A., Shirley, T., Adams, J., Simpas, J. B., and Brune, W. H.: Hox concentrations and oh reactivity observations in new york city during pmtacs-ny2001, *Atmos. Environ.*, 37, 3627–3637, doi:10.1016/s1352-2310(03)00460-6, 2003.

**Total OH reactivity
measurements in
Paris**

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Roukos, J., Plaisance, H., Leonardis, T., Bates, M., and Locoge, N.: Development and validation of an automated monitoring system for oxygenated volatile organic compounds and nitrile compounds in ambient air, *Journal of Chromatography A*, 1216, 8642–8651, doi:10.1016/j.chroma.2009.10.018, 2009.
- 5 Sadanaga, Y., Yoshino, A., Kato, S., Yoshioka, A., Watanabe, K., Miyakawa, Y., Hayashi, I., Ichikawa, M., Matsumoto, J., Nishiyama, A., Akiyama, N., Kanaya, Y., and Kajii, Y.: The importance of NO₂ and volatile organic compounds in the urban air from the viewpoint of the oh reactivity, *Geophys. Res. Lett.*, 31, L08102, doi:10.1029/2004gl019661, 2004.
- Sadanaga, Y., Yoshino, A., Kato, S., and Kajii, Y.: Measurements of oh reactivity and photochemical ozone production in the urban atmosphere, *Environ. Sci. Technol.*, 39, 8847–8852, doi:10.1021/es049457p, 2005.
- 10 Sadanaga, Y., Kondo, S., Hashimoto, K., and Kajii, Y.: Measurement of the rate coefficient for the oh+no(2) reaction under the atmospheric pressure: Its humidity dependence, *Chem. Phys. Lett.*, 419, 474–478, doi:10.1016/j.cplett.2005.12.026, 2006.
- 15 Sciare, J., d'Argouges, O., Zhang, Q. J., Sarda-Estève, R., Gaimoz, C., Gros, V., Beekmann, M., and Sanchez, O.: Comparison between simulated and observed chemical composition of fine aerosols in Paris (France) during springtime: contribution of regional versus continental emissions, *Atmos. Chem. Phys.*, 10, 11987–12004, doi:10.5194/acp-10-11987-2010, 2010.
- Seinfeld, J. H. and Pandis, S. N.: *Atmospheric chemistry and physics: From air pollution to climate change*, Ed. Wiley-Interscience, 2006.
- 20 Shirley, T. R., Brune, W. H., Ren, X., Mao, J., Leshner, R., Cardenas, B., Volkamer, R., Molina, L. T., Molina, M. J., Lamb, B., Velasco, E., Jobson, T., and Alexander, M.: Atmospheric oxidation in the Mexico City Metropolitan Area (MCMA) during April 2003, *Atmos. Chem. Phys.*, 6, 2753–2765, doi:10.5194/acp-6-2753-2006, 2006.
- 25 Sinha, V., Williams, J., Crowley, J. N., and Lelieveld, J.: The Comparative Reactivity Method – a new tool to measure total OH Reactivity in ambient air, *Atmos. Chem. Phys.*, 8, 2213–2227, doi:10.5194/acp-8-2213-2008, 2008.
- Sinha, V., Custer, T. G., Kluepfel, T., and Williams, J.: The effect of relative humidity on the detection of pyrrole by ptr-ms for oh reactivity measurements, *Int. J. Mass Spectrom.*, 282, 108–111, doi:10.1016/j.ijms.2009.02.019, 2009.
- 30 Sinha, V., Williams, J., Lelieveld, J., Ruuskanen, T. M., Kajos, M. K., Patokoski, J., Hellen, H., Hakola, H., Mogensen, D., Boy, M., Rinne, J., and Kulmala, M.: OH reactivity measurements

**Total OH reactivity
measurements in
Paris**

C. Dolgorouky et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

within a boreal forest: Evidence for unknown reactive emissions, *Environ. Sci. Technol.*, 44, 6614–6620, doi:10.1021/es101780b, 2010.

Sinha, V., Williams, J., Diesch, J. M., Drewnick, F., Martinez, M., Harder, H., Regelin, E., Kubistin, D., Bozem, H., Hosaynali-Beygi, Z., Fischer, H., Andrés-Hernández, M. D., Kartal, D., Adame, J. A., and Lelieveld, J.: OH reactivity measurements in a coastal location in Southwestern Spain during DOMINO, *Atmos. Chem. Phys. Discuss.*, 12, 4979–5014, doi:10.5194/acpd-12-4979-2012, 2012.

Stohl, A., Forster, C., Frank, A., Seibert, P., and Wotawa, G.: Technical note: The Lagrangian particle dispersion model FLEXPART version 6.2, *Atmos. Chem. Phys.*, 5, 2461–2474, doi:10.5194/acp-5-2461-2005, 2005.

Tuch, T. M., Haudek, A., Müller, T., Nowak, A., Wex, H., and Wiedensohler, A.: Design and performance of an automatic regenerating adsorption aerosol dryer for continuous operation at monitoring sites, *Atmos. Meas. Tech.*, 2, 417–422, doi:10.5194/amt-2-417-2009, 2009.

Ulbrich, I. M., Canagaratna, M. R., Zhang, Q., Worsnop, D. R., and Jimenez, J. L.: Interpretation of organic components from Positive Matrix Factorization of aerosol mass spectrometric data, *Atmos. Chem. Phys.*, 9, 2891–2918, doi:10.5194/acp-9-2891-2009, 2009.

Vardoulakis, S., Gonzalez-Flesca, N., and Fisher, B. E. A.: Assessment of traffic-related air pollution in two street canyons in paris: Implications for exposure studies, *Atmos. Environ.*, 36, 1025–1039, 2002.

Velasco, E., Lamb, B., Westberg, H., Allwine, E., Sosa, G., Arriaga-Colina, J. L., Jobson, B. T., Alexander, M. L., Prazeller, P., Knighton, W. B., Rogers, T. M., Grutter, M., Herndon, S. C., Kolb, C. E., Zavala, M., de Foy, B., Volkamer, R., Molina, L. T., and Molina, M. J.: Distribution, magnitudes, reactivities, ratios and diurnal patterns of volatile organic compounds in the Valley of Mexico during the MCMA 2002 & 2003 field campaigns, *Atmos. Chem. Phys.*, 7, 329–353, doi:10.5194/acp-7-329-2007, 2007.

Xu, X., Stee, L. L. P., Williams, J., Beens, J., Adahchour, M., Vreuls, R. J. J., Brinkman, U. A., and Lelieveld, J.: Comprehensive two-dimensional gas chromatography (GC × GC) measurements of volatile organic compounds in the atmosphere, *Atmos. Chem. Phys.*, 3, 665–682, doi:10.5194/acp-3-665-2003, 2003.

Yoshino, A., Sadanaga, Y., Watanabe, K., Kato, S., Miyakawa, Y., Matsumoto, J., and Kajii, Y.: Measurement of total oh reactivity by laser-induced pump and probe technique – comprehensive observations in the urban atmosphere of tokyo, *Atmos. Environ.*, 40, 7869–7881, doi:10.1016/j.atmosenv.2006.07.023, 2006.

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Table 1. Gaseous pollutants levels registered in different urban atmospheres.

	Population ^a in 2010 (10 ⁶)	Period of the year	CO BGND level (ppbv)	CO max (ppbv)	NO _x BGND levels (ppbv)	NO _x max (ppbv)	Ozone BGND level (ppbv)	Ozone max (ppbv)	NMHC BGND (ppbv)	NMHC max (ppbv)
Houston	6	Aug–Sep 2000	200	250	5	22	25	70	100 ^c	200 ^c
New York City	22	Jun–Aug 2001	400	450	25	50	20	50	100 ^c	150 ^c
Mexico City	22	Apr 2003	100	3000	20	110	25	125	500 ^c	1000 ^c
Houston	6	Aug–Sep 2006	200	250	8	25	25	70	400 ^c	750 ^c
Tokyo	36 ^b	Jan–Feb 2004	400 ^d	1200 ^d	25 ^d	105 ^d	10 ^d	40 ^d	NA	NA
Tokyo	36 ^b	May 2004	300 ^d	550 ^d	13 ^d	50 ^d	50 ^d	80 ^d	NA	NA
Tokyo	36 ^b	November 2004	400 ^d	1000 ^d	20 ^d	65 ^d	10 ^d	60 ^d	NA	NA
Paris	12	Jan–Feb 2010	250	350	30	60	15	17	12 50 ^c	17 100 ^c

^a Cities and Metropolitan Areas

^b 2007

^c Values in ppbC (ppbCarbon)

^d Daily mean values were taken into consideration when median diurnal cycles were not available

NA = Non Available Data

BGND = Corresponds to background levels measured at midnight, local time

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 2. Mean values of the main volatile-organic-compounds (VOC) volume mixing ratio measured during the campaign (TMB, trimethylbenzene; MVK, methyl vinyl ketone).

Compound	Measurement technique	Mean (ppbv)	Min (ppbv)	Max (ppbv)	OH reactivity (s^{-1})
Ethane	GC-FID	2.45	1.14	7.12	0.015
Ethene	GC-FID	2.61	0.33	8.36	0.547
Propane	GC-FID	2.12	0.76	5.04	0.057
Propene	GC-FID	0.44	0.07	1.67	0.323
i-Butane	GC-FID	0.91	0.15	2.84	0.052
n-Butane	GC-FID	1.54	0.34	5.03	0.087
Acetylene	GC-FID	1.77	0.62	5.04	0.044
1-Butene	GC-FID	0.09	0.02	0.26	0.093
i-Butene	GC-FID	0.19	0.02	0.55	0.306
c-2-Butene	GC-FID	0.05	0.02	0.53	0.080
trans-2-Butene	GC-FID	0.07	0.01	0.23	0.126
Isoprene	GC-FID	0.17	0.05	1.43	0.423
n-Hexane	GC-FID	0.14	0.01	1.34	0.019
n-Pentane	GC-FID	0.40	0.07	1.39	0.040
Benzene	PTR-MS	0.33	0.08	1.65	0.010
Toluene	PTR-MS	0.54	0.09	2.36	0.079
Xylenes	GC-FID	0.67	0.13	2.88	0.265
TMB	GC-FID	0.29	0.08	0.76	0.289
Acetonitrile	PTR-MS	0.55	0.09	3.69	0.000
i-Propanol	GC-MS	0.24	0.04	1.06	0.031
Ethanol	GC-MS	7.68	2.22	23.57	0.605
MVK	GC-MS	0.09	0.02	0.44	0.042
Acetone	GC-MS	1.05	0.58	2.97	0.005
2-Butanone	GC-MS	0.29	0.06	0.94	0.009
Hexanal	GC-MS	0.09	0.01	0.22	0.000
Benzaldehyde	GC-MS	0.07	0.02	0.18	0.025
i-Butanol	GC-MS	0.08	0.02	0.20	0.014
Methanol	PTR-MS	2.35	0.89	6.18	0.052
Acetaldehyde	PTR-MS	1.87	0.92	4.49	0.691

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 3. Mean values of the main pollution compounds measured during period I, II and III.

	OH reactivity (s ⁻¹)	CO (ppbv)	NO (ppbv)	NO ₂ (ppbv)	Benzene (ppbv)	Xylenes (ppbv)	Acetone (ppbv)	NMHC (ppbv)	OVOC (ppbv)	Air masses
Period I 26/01–27/01	130	536	24	39	0.7	1.0	2.1	20.6	18.7	Continental
Period II 02/02–04/02	20	249	4	28	0.2	0.5	0.7	10.1	13.1	Oceanic
Period III 09/02–13/02	110	311	7	27	0.3	0.6	1.0	13.2	12.4	Continental

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 4. Synthesis of OH reactivity levels measured in other urban atmospheres.

Urban site	Season, Year	Employed technique	Average/ Median ^a (s ⁻¹)	Maximum (s ⁻¹)	Missing OH reactivity (%)	Measured species ^h (after Lou et al., 2010)	Reference
Nashville, USA	Summer 1999	LIF-H2O	11.3±4.8	25	31	SFO	Kovacs et al. (2003)
Houston, USA	Autumn 2000	LIF-H2O	10	12 ^c	~0 ^g	SFOB	Mao et al. (2010)
New York, USA	Summer 2001	LIF-FAGE	19±3	50	10	SFOB	Ren et al. (2003)
Mexico City	Spring 2003	LIF-FAGE	33	200	–	–	Shirley et al. (2006)
Tokyo, Japan	Summer 2003	LIF P&P	40	85	15 to 30	SFOB	Sadanaga et al. (2005)
New York, USA	Winter 2004	LIF-FAGE	20	100	33	SF	Ren et al. (2006a)
Tokyo, Japan	Winter 2004	LIF P&P	20*	100	5	SFOB	Yoshino et al. (2006)
Mainz, Germany	Summer 2008	CRM	10.4	18	–	–	Sinha et al. (2008)
Houston, USA	Autumn 2006	LIF-H2O	15 ^b	25	~0 ^g	SFOB	Mao et al. (2010b)
Paris, France	Winter 2010	CRM	33 ^d (19.6 ^e /60 ^f)	130	12 to 75	SO	This work

LIF- H₂O Laser Induced Fluorescence Technique - OH produced by H₂O photolysis

LIF- P&P Laser Induced Fluorescence – Pump and Probe Technique

LIF-FAGE Laser Induced Fluorescence – Fluorescence Assay by Gas Expansion

^a average was used when median not available

^b approximate values from graphs

^c large part of missing data

^d median on the entire campaign

^e median on the local pollution period


^f median on the continental import period

^g within the measurement and calculated uncertainty

^h Measured species that have been used for the calculation of the OH reactivity: *S* = inorganic compounds (O₃, CO, NO_x, etc) plus hydrocarbons (including isoprene); *F* = formaldehyde; *O* = OVOCs other than formaldehyde; *B* = biogenic VOCs, other than isoprene

Total OH reactivity measurements in Paris

C. Dolgorouky et al.

**Fig. 1.** OH reactivity system during MEGAPOLI winter 2010.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Total OH reactivity
measurements in
Paris**

C. Dolgorouky et al.




Fig. 2. Calibration results using two standards bottles: propane and acetone, methanol and isoprene standard.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Total OH reactivity
measurements in
Paris**

C. Dolgorouky et al.




Fig. 3. Change in the pyrrole signal (m/z 68) for different percentage of relative humidity into the setup. The temperature of the setup is maintained constant at 24 °C.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Total OH reactivity
measurements in
Paris**

C. Dolgorouky et al.




Fig. 4. NO test for OH reactivity of 50 s⁻¹: the red and the blue curves represent the 2 different slopes obtained for values of NO smaller or bigger than 20 ppbv.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Total OH reactivity
measurements in
Paris**

C. Dolgorouky et al.




Fig. 5. Backward trajectories characterizing the air masses of the three contrasted OH reactivity periods **(a)** continental air masses on 27 January, **(b)** oceanic air masses on 3 February and **(c)** continental air masses on 9 February Source: FLEXPART model.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Total OH reactivity
measurements in
Paris**

C. Dolgorouky et al.




Fig. 6. Campaign time series: top panel: NO in light-blue solid line and the sum of xylenes in dark-blue solid line; 2nd panel: CO in light-green solid line and benzene in dark-green solid line; 3rd panel: acetone in light-orange solid line and acetaldehyde in dark-orange solid line. The three air mass periods are colored as follows: in yellow the period I, in green, period II, and in violet, period III.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Total OH reactivity measurements in Paris

C. Dolgorouky et al.




Fig. 7. OH reactivity time series (2 min blue dots; 30 min black diamonds).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Total OH reactivity
measurements in
Paris**

C. Dolgorouky et al.




Fig. 8. Daily variability of the OH reactivity during a local pollution episode: top panel: OH reactivity time series (2 min blue dots and 10 min black diamonds) and NO time series (5 min green triangles). Bottom panel: OH reactivity time series (2 min blue dots and 10 min black diamonds); acetone time series (30 min red squares).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Total OH reactivity
measurements in
Paris**

C. Dolgorouky et al.




Fig. 9. Calculated (90 min time resolution red curve) versus the measured OH reactivity (2 min time resolution blue dots, 30 min average, black diamonds).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Total OH reactivity measurements in Paris

C. Dolgorouky et al.




Fig. 10. Breakdown of the (a) total OH reactivity and (b) calculated OH reactivity for the entire MEGAPOLI winter campaign.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Total OH reactivity
measurements in
Paris**

C. Dolgorouky et al.




Fig. 11. The Breakdown of the total OH reactivity for **(a)** period II and **(b)** period III.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Total OH reactivity
measurements in
Paris**

C. Dolgorouky et al.




Fig. 12. Top panel: missing OH reactivity; bottom panel: sum of SO₄, NO₃ and NH₄.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Total OH reactivity measurements in Paris

C. Dolgorouky et al.




Fig. 13. Contribution of different atmospheric compounds to OH reactivity in Paris (winter 2010), Tokyo (winter 2004), Mexico City (spring 2003), New York (summer 2001) and Houston (autumn 2006).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)
