

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

Estimating marine aerosol particle volume and number from Maritime Aerosol Network data

A. M. Sayer^{1,2}, A. Smirnov^{2,3}, N. C. Hsu², L. A. Munchak^{2,4}, and B. N. Holben²

¹Goddard Earth Sciences Technology And Research (GESTAR), Universities Space Research Association (USRA), Columbia, Maryland, USA

²NASA Goddard Space Flight Center, Greenbelt, Maryland, USA

³Sigma Space Corporation, Lanham, Maryland, USA

⁴Science Systems Applications Inc., Lanham, Maryland, USA

Received: 21 May 2012 – Accepted: 31 May 2012 – Published: 12 June 2012

Correspondence to: A. M. Sayer (andrew.sayer@nasa.gov)

Published by Copernicus Publications on behalf of the European Geosciences Union.

ACPD

12, 14959–15007, 2012

Marine aerosol concentration from MAN

A. M. Sayer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

As well as spectral aerosol optical depth (AOD), aerosol composition and concentration (number, volume, or mass) are of interest for a variety of applications. However, remote sensing of these quantities is more difficult than for AOD, as it is more sensitive to assumptions relating to aerosol composition. This study uses spectral AOD measured on Maritime Aerosol Network (MAN) cruises, with the additional constraint of a microphysical model for unpolluted maritime aerosol based on analysis of Aerosol Robotic Network (AERONET) inversions, to estimate these quantities over open ocean. When the MAN data are subset to those likely to be comprised of maritime aerosol, number and volume concentrations obtained are physically reasonable. Attempts to estimate surface concentration from columnar abundance, however, are shown to be limited by uncertainties in vertical distribution. Columnar AOD at 550 nm and aerosol number for unpolluted maritime cases are also compared with Moderate Resolution Imaging Spectroradiometer (MODIS) data, for both the present Collection 5.1 and forthcoming Collection 6. MODIS provides a best-fitting retrieval solution, as well as the average for several different solutions, with different aerosol microphysical models. The “average solution” MODIS dataset agrees more closely with MAN than the “best solution” dataset. Terra tends to retrieve lower aerosol number than MAN, and Aqua higher, linked with differences in the aerosol models commonly chosen. Collection 6 AOD is likely to agree more closely with MAN over open ocean than Collection 5.1. In situations where spectral AOD is measured accurately, and aerosol microphysical properties are reasonably well-constrained, estimates of aerosol number and volume using MAN or similar data would provide for a greater variety of potential comparisons with aerosol properties derived from satellite or chemistry transport model data.

ACPD

12, 14959–15007, 2012

Marine aerosol concentration from MAN

A. M. Sayer et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



1 Introduction

Columnar aerosol optical depth (AOD) has been mapped on a near-global basis for several decades from satellite measurements with varying degrees of accuracy (e.g., Husar et al., 1997; Hsu et al., 1999; Torres et al., 2002; Mishchenko et al., 2007; Remer et al., 2008; Thomas et al., 2009; Kahn et al., 2010; Sayer et al., 2012a). There is a similar wealth of ground-based aerosol observation from techniques such as sun photometry, lidar, or multifilter rotating shadowband radiometers, with records approaching two decades at some locations (e.g., Holben et al., 1998; Michalsky et al., 2001; Campbell et al., 2002). The AOD represents the vertically-integrated extinction of light by aerosol particles,

$$\tau_\lambda = \int_0^\infty \beta_\lambda(z) dz, \quad (1)$$

where τ_λ is the AOD at wavelength λ , and β_λ the aerosol extinction (sum of scattering and absorption) at that wavelength and altitude z , and as such is related to the aerosol mass loading of the atmosphere.

The spectral behaviour of AOD, frequently referred to in the context of the Ångström parameter α , where

$$\alpha = -\frac{d \ln(\tau_\lambda)}{d \ln(\lambda)}, \quad (2)$$

is often evaluated across the visible region of the solar spectrum and used as a first-order indication of aerosol type (Eck et al., 1999; Dubovik et al., 2002). However, α is not a unique identifier of a particular aerosol composition, so additional constraints such as microphysical aerosol particle models are necessary to infer physical aerosol amount from AOD. Further, particularly in low-AOD regimes, satellite and ground-based estimates of α can suffer from significant uncertainty (Wagner and Silva, 2008).

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



These factors mean that estimating aerosol number or volume from remotely-sensed AOD is, at present, not straightforward. Remote determination of aerosol number or volume/mass rather than solely AOD is of interest to estimate, for example, the deposition flux of mineral dust aerosols (Kaufman et al., 2005), near-surface particulate matter concentration in urban areas (Hoff and Christopher, 2009), aerosol radiative forcing (Quaas et al., 2008; Yu et al., 2012), or the available number of particles which can act as cloud condensation nuclei (CCN) and influence cloud development (Jefferson, 2010).

Recently, Sayer et al. (2012b) analysed size distribution inversions from a selection of Aerosol Robotic Network (AERONET) coastal and island sites (Holben et al., 1998; Dubovik and King, 2000), and arrived at a microphysical model of pure maritime aerosol as a combination of two lognormally-distributed components. The properties of maritime aerosol were found to be similar over the range of sites, such that the observed AERONET AOD record at these sites could be well-reproduced by varying only two free parameters, namely the fine-mode and coarse-mode aerosol volume. The Maritime Aerosol Network (MAN) provides ship-based observations of spectral AOD over the ocean (Smirnov et al., 2009, 2011), with typically four (sometimes five) channels in any individual measurement. Thus, the model can be used to infer columnar aerosol number and volume of maritime aerosol from the MAN AOD measurements (size distribution inversions as performed at the AERONET land sites are not possible from the hand-held measurements collected on MAN cruises). The aim of this study is to perform such an exercise.

This is first a test of whether the model of Sayer et al. (2012b) is able to produce physically-reasonable values of aerosol concentration. It also allows a comparison of derived aerosol number with the (unvalidated) aerosol columnar number concentration estimates provided in the Moderate Resolution Imaging Spectroradiometer (MODIS) satellite aerosol product (Remer et al., 2005, 2008). These open up further scope for evaluation of the aerosol parametrisation of chemistry transport models (CTMs) through comparisons with aerosol number or volume as well as AOD. This is important

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀◀	▶▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



because CTM aerosol fields are diverse, and it is possible for the CTM to produce the right AOD but with the wrong aerosol composition (e.g., Kinne et al., 2006; Textor et al., 2006). Uncertainties in the MAN-derived estimates are also discussed.

The notation adopted in this work follows that of Sayer et al. (2012b), in which various identities and derivations are presented. A summary of relevant notation is presented here. The aerosol number size distribution $dN(r)/d\ln(r)$ describes the number of aerosol particles with radius in the infinitesimal size range $r \pm d\ln(r)$; for spherical particles, this is related to the volume size distribution $dV(r)/d\ln(r)$ by

$$\frac{dV(r)}{d\ln(r)} = \frac{4\pi r^3}{3} \frac{dN(r)}{d\ln(r)}. \quad (3)$$

In this work and many others (Sayer et al., 2012b, and references therein) aerosol particle size distributions are represented as a sum of n_c lognormally-distributed components,

$$\frac{dN(r)}{d\ln(r)} = \sum_{i=1}^{n_c} \frac{C_{n,i}}{\sqrt{2\pi}\sigma_i} e^{-\frac{1}{2} \left(\frac{\ln(r) - \ln(r_{n,i})}{\sigma_i} \right)^2}, \quad (4)$$

where r_n and σ are the mode's modal (also median and geometric mean) radius and geometric standard deviation, respectively, and C_n the total number of aerosol particles in the mode. The equivalent aerosol volume distribution is given by the same expression, except substituting r_n with the volume median radius r_v , and C_n with the total aerosol volume C_v . The relationships between these quantities for a lognormal component (Sayer et al., 2012b) are

$$r_v = r_n e^{3\sigma^2} \quad (5)$$

and

$$C_v = \frac{4\pi}{3} r_n^3 e^{4.5\sigma^2} C_n, \quad (6)$$

enabling the conversion between number and volume radii, and calculation of aerosol number-to-volume ratio. In this work bimodal aerosol distributions are used (i.e., $n_c = 2$), with the smaller mode denoted “fine” with a subscripted f, and larger mode “coarse” with a subscripted c.

The remainder of the manuscript is organised as follows. Section 2 describes the MAN data and the method whereby aerosol number and volume can be estimated, along with derived maps of these data. Aerosol mass is not explicitly discussed (for a given density, it is a simple scaling of aerosol volume). Section 3 provides a comparison with MODIS estimates, and Sect. 4 looks at the relationship between total columnar and surface concentrations. Finally, Sect. 5 provides a summary and outlook.

2 Estimating columnar volume and number

2.1 MAN data

The AOD measurements on MAN cruises (Smirnov et al., 2009, 2011) are made with hand-held Microtops II Sun-photometers, which allow measurement of AOD with a total (one standard deviation) uncertainty of order ± 0.015 for typical oceanic conditions (Porter et al., 2001; Knobelispiese et al., 2003, 2004). The instruments have five filters which can be adjusted to observe the Sun at different wavelengths; typically on MAN cruises one is used to retrieve columnar water vapour, leaving four for AOD, in the spectral range 340 nm–1020 nm. The overwhelming majority of measurements consist of the combination τ_{440} , τ_{500} , τ_{675} , and τ_{870} (subscripted wavelengths are in nanometres throughout). The Angström parameter α is calculated in MAN from a least-squares fit (in logarithmic space) of AOD and wavelength over the spectral range 440 nm–870 nm.

Two MAN datasets are used in this work. The first is the “series average” product, where one measurement series is defined as the set of AOD measurements taken with a gap of no more than 2 min between an individual pair. The second is the “daily average” product, which is the average of all measurement series on a given day. Frequently

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

multiple series are obtained on a given day in identical or very close locations, so visual interpretation is typically clearer using daily data, while statistical analysis benefits from the larger sample size of the series average data. In practice the results change negligibly if only one or the other data product is used, suggesting that most of the observations collected on a MAN cruise over the course of a single day sample similar aerosol regimes. In all cases, only level 2.0 (cloud-screened and quality assured) data are used.

ACPD

12, 14959–15007, 2012

Marine aerosol concentration from MAN

A. M. Sayer et al.

2.2 Calculation and uncertainty

Size distribution parameters and refractive indices for the bimodal model of Sayer et al. (2012b) are given in Table 1. The free parameters, $C_{v,f}$ and $C_{v,c}$, are determined from a least-squares fit of each set of MAN spectral AOD to the spectral extinction per unit volume modelled using Mie theory (values at the common reference wavelength of 550 nm are also given in Table 1), with the constraint that the volumes cannot be negative. Aerosol number can then be calculated using the ratio C_n/C_v , from Eq. (6).

This process is shown conceptually in Fig. 1. In this case, the observed spectral AOD (black asterisks) is best reproduced by the combination $C_{v,f} = 0.005 \mu\text{m}^3 \mu\text{m}^{-2}$ and $C_{v,c} = 0.04 \mu\text{m}^3 \mu\text{m}^{-2}$ (total AOD given by the red curve). Note that as previously mentioned a real MAN observation would have five or fewer spectral AOD measurements, rather than the seven shown here, but this serves to illustrate the spectral dependence of the fine and coarse mode extinction across the Microtops bands.

Table 2 provides statistics on the quality with which the model is able to reproduce the spectral MAN AOD. At all wavelengths, the bias and scatter are small (<0.01), particularly over the wavelength range (440 nm–870 nm) which was used during development of the model. Note that some of the MAN measurements contain an interpolated rather than measured 500 nm AOD: in these cases, this interpolated AOD was used for the statistics in Table 2 but not when performing the least-squares fit (as it did not correspond to a real measurement).

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Clearly, not all of the observations in the MAN database will represent unpolluted maritime aerosol, and therefore derived aerosol volume and number may be significantly biased when this is not the case. For this reason, the MAN data have been stratified in a simple attempt to discriminate according to aerosol type, based on the 5 AOD and α . Three broad classes have been defined based on typical values for these parameters (e.g., Eck et al., 1999; Smirnov et al., 2004; Sayer et al., 2012b): pure maritime ($\tau_{500} \leq 0.2$, $\alpha \leq 1$), dust-influenced ($\tau_{500} > 0.2$, $\alpha \leq 0.6$), and fine continental (e.g., pollution/smoke or land organic)-influenced (all other points, referred to hereafter 10 as “continental” for brevity). Additionally, as in Smirnov et al. (2012), to minimise the likelihood of continental influence in the “maritime” subset it was further required that such points be at least 200 km from land, using a coarse (1°) land mask as a basis (to eliminate large land masses, but not small remote islands).

Maps of these classes and fit volumes from the daily MAN data are shown in Fig. 2; the points cluster in generally expected regions, suggesting that as a first-order attempt 15 this classification is reasonable, although there is inherently a degree of ambiguity in this type of classification. Altering these thresholds within sensible ranges does not significantly affect the spatial distribution or interpretation of results. Additionally, if the more conservative set of “pure maritime” MAN points used by Smirnov et al. (2012) is used, the results do not significantly change for this subset (which is the main focus of 20 the analysis). Information on the sampling of these subsets is given in Table 3.

The latitudinal distribution of the number of measurement series in these three classes is shown in Fig. 3. The large abundance of the “continental-influenced” class is not suggesting that the majority of the open ocean is influenced significantly by continental outflow. Rather, this happens because of the fact that the MAN cruises begin and 25 end on the coast, and often spend much of their time in coastal regions (Fig. 2). For example, the spike in this class at high southern latitudes comes from data collected near Antarctica. It is important to note that this “continental-influenced” classification is not purely an indicator of urban or smoke aerosol mixed with maritime aerosol (although it may contain these), but rather a catch-all for aerosol which, due to its proximity to the

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Marine aerosol concentration from MAN

A. M. Sayer et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



coast, has the potential to be influenced by land masses, and is additionally unlikely to be dominated by dust. The intent of this classification is to “protect” the “pure maritime” subset as much as possible.

Table 1 also contains information on microphysical models developed by Sayer et al. (2012a) to represent mixed maritime/continental and maritime/dust aerosols. The aerosol extinction per unit volume is generally within 20 % of that for the pure maritime model across the visible part of the spectrum (only extinction at 550 nm is shown in Table 1), suggesting a bias of order 20 % in aerosol volume if there is a non-maritime component to the aerosol and the maritime model is used. The per-particle extinction, however, shows differences in excess of 100 %, indicating that using the maritime model to predict aerosol number for a non-maritime aerosol would be significantly in error. For this reason, only MAN data from the maritime subset are used when aerosol number concentrations are analysed hereafter. Note that all three classes here were fit with the maritime model; the parameters from Sayer et al. (2012a) are presented here only to illustrate the potential magnitude in error of aerosol number/volume from an incorrect determination of aerosol type.

The reduced χ^2 statistic is used to measure the goodness of fit,

$$\chi^2 = \frac{1}{n_\lambda - 2} \sum_0^{n_\lambda} \left(\frac{\tau_{\lambda,\text{pred}} - \tau_{\lambda,\text{MAN}}}{\sigma_{\lambda,\text{MAN}}} \right)^2, \quad (7)$$

where $\tau_{\lambda,\text{MAN}}$ indicates the MAN-observed AOD, $\sigma_{\lambda,\text{MAN}}$ its uncertainty (assumed to be 0.015 and uncorrelated spectrally), $\tau_{\lambda,\text{pred}}$ the AOD predicted by the model, and n_λ the number of wavelengths in the measurement series (or daily average). The factor $n_\lambda - 2$ arises as two free parameters are fit; typically $n_\lambda = 4$, leading to 2 degrees of freedom. The expectation of the reduced χ^2 over a large number of samples is 1. Figure 2 also shows χ^2 for each of the three aerosol type subsets: this is almost always less than 1, even for those cases where the aerosol is likely not pure maritime in origin. There are two important implications of this. First, this highlights limitations in inferring aerosol type from spectral AOD (it is possible to obtain an acceptable quality of fit even

if the microphysical model is incorrect). Second, this suggests that $\sigma_{\lambda,MAN} = 0.015$ is not necessarily a good metric for the random error of the MAN AOD, i.e. systematic error is likely a significant component of the total uncertainty. This is also consistent with the results in Table 2, that the maritime model is able to reproduce spectral AOD with greater accuracy and precision than 0.015. Porter et al. (2001) and Knobelspiesse et al. (2004) estimate that among the largest uncertainty on Microtops AOD is the calibration gain coefficient. For a set of measurements taken on a single cruise with a single Microtops sun photometer, this is likely a systematic error, although over the whole MAN dataset (multiple instruments and calibration tests) biases may cancel out such that the errors are random.

The least-squares fit provides estimates of the uncertainty on C_v and hence C_n , under the assumption that $\sigma_{\lambda,MAN}$ represents the random error on the MAN AOD. Scaling these uncertainty estimates by $\sqrt{\chi^2}$ provides a lower bound on the estimated uncertainty, which is under the assumption that the true value of $\sigma_{\lambda,MAN}$ is unknown and the uncertainty is therefore related to the residuals on the fit AOD. The true error on derived C_v or C_n is therefore likely in between these two estimates.

Relative uncertainty on C_v is shown as a function of C_v for the maritime subset of daily-average MAN AOD for both these methods in Fig. 4. As, for a given microphysical aerosol model, the aerosol number is a simple scaling to the aerosol volume (rightmost column of Table 1), these relative uncertainties also apply to the equivalent aerosol numbers. The “unscaled” points (i.e., taking $\sigma_{\lambda,MAN} = 0.015$) typically fall into one of several curves, dependent on the selection of bands available for a particular MAN data point. The absolute uncertainty is fairly constant as a consequence of the fact that, for a given set of microphysical model parameters, AOD is linearly proportional to aerosol amount. If instead $\sigma_{\lambda,MAN}$ is assumed to be 0.01 or 0.02, the uncertainty estimates decrease and increase by approximately 50 %, respectively. The “scaled” uncertainties are much more diverse but generally lower by a factor of 2–5. For the unscaled uncertainty estimates, uncertainty on volume is around 100 % for low aerosol

Marine aerosol concentration from MAN

A. M. Sayer et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀◀	▶▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



loadings. However, the “scaled” uncertainty estimates are much lower, suggesting better sensitivity.

2.3 Derived volume and number concentration

For all three classes, coarse-mode volume tends to be larger than fine-mode volume by approximately a factor of 5–10 (Fig. 2); the maritime subset typically has 80 % or more of total aerosol volume in the coarse mode. Relatively higher fractional coarse-mode volumes are found for the dust subset, and relatively higher fractional fine mode volumes for the continental subset, which is expected from the α -based classification. Histograms of aerosol volume for these three cases appear to follow approximate log-normal distributions, illustrated in Fig. 5 (using the series average MAN dataset). This is not surprising, as AOD distributions have also been observed to be lognormal (e.g., O’Neill et al., 2000 and reference therein), and was also noted by Heintzenberg et al. (2000). The modes and spreads (geometric mean and geometric standard deviation, analogous to r_n and σ in Eq. (4), respectively) are given in Table 4, and are reasonably robust to small changes in histogram bin size. The median value of all points is also shown. As mentioned previously, volumes for the dusty and continental classes are likely to be qualitatively reliable but absolute values may have a bias of order 20 %.

The fine mode distribution for the continental-influenced class shows two peaks (or a very long tail); the lower-volume segment is similar to the maritime class’s peak, and likely corresponds to maritime aerosol which narrowly missed the τ_{500} , α , and/or land distance thresholds for inclusion in the maritime class. Both the maritime and dust-influenced distributions, however, are more distinct.

Median fine and coarse volumes for the maritime subset are $0.0056 \mu\text{m}^3 \mu\text{m}^{-2}$ and $0.062 \mu\text{m}^3 \mu\text{m}^{-2}$, respectively. These are similar to the average volumes found for island and coastal AERONET sites by Sayer et al. (2012b) of $0.0058 \mu\text{m}^3 \mu\text{m}^{-2}$ and $0.036 \mu\text{m}^3 \mu\text{m}^{-2}$ for fine and coarse modes, respectively; the higher median coarse-mode volume could be linked with increased wind-driven coarse particle emission over the open ocean. The majority of points where τ_{340} and τ_{380} are available were collected

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

in the tropical Atlantic and fall into the “dusty” subset, and so it is possible that the small negative AOD fitting bias at these wavelengths (Table 2) and elevated fine mode volume for this subset are related to the real aerosol being more absorbing than the maritime model.

Maps of derived aerosol number (for the maritime subset only) are shown in Fig. 6, and histograms in Fig. 7. The histogram are again approximated well by a lognormal distribution (parameters in Table 4). The aerosol number:volume ratio, $C_n/C_v = 190 \mu\text{m}^{-3}$ for the fine mode (Table 1), is in good agreement with previous studies in the range $167\text{--}225 \mu\text{m}^{-3}$ for marine aerosol (Hegg and Kaufman, 1998; van Dingenen et al., 1999; Hegg and Jonsson, 2000; Kaufman et al., 2001; Dusek et al., 2004). Dusek et al. (2004) note that in previous studies this ratio has been defined in different ways, often the ratio of total number of particles above a certain size to total volume of particles below a certain cutoff size, dependent often on the available instrumentation. As the coarse mode total aerosol number is generally only 1 % or so of the fine mode number, and the coarse mode volume below the typical cutoff size used in these studies (around $1 \mu\text{m}$) is small, it makes little difference to C_n/C_v in this case (although C_n/C_v would become slightly smaller and depend on the weighting between fine and coarse modes for each situation).

3 Comparison with MODIS data

20 3.1 Data description and methodology

The MODIS aerosol algorithm retrieves spectral AOD and α over ocean by mixing two aerosol components (one fine mode and one coarse mode, each from Table 1) to find the combination which matches the observed top-of-atmosphere (TOA) reflectance in six bands between 470 nm and $2.1 \mu\text{m}$ (Tanré et al., 1997; Remer et al., 2005, 2008). Two solutions are reported in the product: namely, the combination of fine and coarse modes which most closely fit the observed TOA reflectance (hereafter the

Marine aerosol concentration from MAN

A. M. Sayer et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



“best solution”), and the average (hereafter the “average solution”) of either all solutions with a root mean square fitting error of less than 3 %, or the three solutions with the smallest fitting error if none are less than 3 %. From this, the algorithm also derives an estimate of the total columnar number of aerosol particles of radius $0.03\text{ }\mu\text{m}$ or larger (Remer et al., 2005). The effect of this minimum size on the comparison is minor, as the maritime model applied to the MAN data results in fewer than 4 % of the fine mode particles having a radius smaller than $0.03\text{ }\mu\text{m}$.

The MODIS aerosol number estimate is to the authors’ best knowledge an unvalidated quantity, although it has been used, examined, or compared with other data in several studies (Gassó and Hegg, 2003; Li et al., 2010; Kaskaoutis et al., 2011; Krüger and Graßl, 2011). In contrast, the AOD over ocean has been compared more thoroughly with other satellite and ground-based datasets and its strengths and limitations are fairly well-understood (e.g., Zhang and Reid, 2006; Mishchenko et al., 2007, 2010; Remer et al., 2008; Kahn et al., 2009, 2011; Shi et al., 2011; Smirnov et al., 2011; Kleidman et al., 2012; Sayer et al., 2012a, and others). The MAN-derived data here are also subject to some uncertainties as discussed previously, so while a comparison between aerosol number from these two datasets cannot be considered a validation against ground truth, it does allow for an examination of their consistency and, hopefully, an understanding of their differences.

Matchups between MODIS and the MAN data are performed by averaging MODIS retrievals within 25 km of a MAN measurement separated in time by 30 min or less. In almost all cases, variability in MODIS AOD in this 25 km circle was small, and the same aerosol model was chosen as the “best solution” for each retrieval. Only MAN data defined previously as belonging to the maritime subset are considered, as C_n from others is expected to be less reliable. The AOD is compared at the standard reference wavelength of 550 nm, as provided in the MODIS product; interpolation of the MAN data to this wavelength using the Ångström power law (Eq. 2) introduces negligible uncertainty.

Marine aerosol concentration from MAN

A. M. Sayer et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



As well as the present Collection 5.1 (hereafter C5.1), results are presented using the forthcoming Collection 6 (C6) algorithm. C6 products should become available around the end of 2012. The over-ocean algorithmic concept is the same in C6 as C5.1, but there are changes related to gaseous transmittance corrections, improved masking of 5 cirrus clouds, accounting for wind-speed dependence of Sun glint and oceanic white-caps (which will decrease retrieved AOD in high-wind environments; Sayer et al., 2010; Shi et al., 2011; Kleidman et al., 2012), and definitions of quality flags. Sensor calibration will also be updated for C6 but final coefficients are not available at the time of writing (i.e., these results use C5 radiometric calibration coefficients), and the impact 10 over open ocean (on retrieved AOD at least) is expected to be small.

3.2 Results

Some statistics of the comparison of AOD and aerosol number are presented in Table 5, considering both the MODIS sensors on board the Terra and Aqua platforms, and for either the highest quality assurance (QA) flag ($QA = 3$) or the looser criteria of 15 $QA = 1, 2$, or 3 (almost no retrievals were assigned $QA = 2$). Scatter plots of the $QA = 3$ data are shown in Fig. 8, and maps for the “best” solution in Fig. 9, both for C5.1 data. Table 6 shows the fraction of MODIS-MAN matchups with an AOD at 550 nm within the expected MODIS absolute error (Remer et al., 2008) of $0.03 + 5\%$ of the MAN value. Looking first at τ_{550} , there is a tendency for MODIS to overestimate as compared to 20 the MAN data, and Terra to be higher than Aqua by around 0.01, as noted in previous studies (e.g., Remer et al., 2008; Shi et al., 2011; Smirnov et al., 2011). The bias is reduced by typically 30 %–50 % in C6 as compared to C5.1, and the fraction of matchups within the expected error increases.

A second set of comparisons of derived C_n have been performed, where the MODIS values have been scaled by the ratio of MAN to MODIS AOD at 550 nm (i.e., if MODIS overestimated MAN AOD by 20 %, the MODIS C_n would be decreased by a corresponding amount). This “scaled” C_n allows a first-order separation between the effects 25 of differences in AOD and differences in aerosol microphysical model assumptions on

Marine aerosol concentration from MAN

A. M. Sayer et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



the comparison. Tables 5 and 6 also show that restricting the data to QA = 3 (the retrievals with highest confidence) results in a poorer agreement of AOD (and also C_n) between the datasets. However, these results should not necessarily be expected to apply to the MODIS dataset as a whole, as the QA = 3 subset is comparatively small (about a factor of 3 fewer points than the QA = 1,2,3 case for C5.1), and these comparisons only consider MAN data from the maritime subset, which is a small proportion of the total dataset. The MODIS over-ocean data usage recommendation is that retrievals of QA 1, 2, or 3 are likely of similar quality and all suitable for analysis, which is consistent with these comparisons (Remer et al., 2005, 2008). The total number of C6 points is smaller than C5.1, likely due to stricter cloud screening, and QA = 3 becomes more common. Irrespective of QA threshold, the differences between best and average solutions, and scaled and unscaled QA, are broadly similar.

Relative to MAN-derived data, MODIS Terra tends to underestimate and Aqua to overestimate C_n (Table 5). The regional sampling of the two datasets is similar (Fig. 9), suggesting these differences are more related to the retrieval than spatial sampling. Performing the AOD-based scaling tends to reduce the absolute and root mean square differences between MAN and MODIS data; however, significant differences remain in this case, indicating microphysical model assumptions play a role. Further, large differences of either sign between the datasets are not confined to continental outflow areas, suggesting that errors in MAN C_n from non-maritime influences are not the primary cause for difference.

Table 1 shows that, as is the case for the maritime aerosol model of Sayer et al. (2012b), the fine mode per-particle extinction is around two orders of magnitude larger than the coarse mode, and so the total C_n will be determined largely by fine mode abundance. MODIS fine mode #2 has a similar per-particle extinction and C_n/C_v to the model of Sayer et al. (2012b), so if the two datasets report the same AOD and this fine mode is picked in the MODIS dataset, the two estimates of C_n should be close (although the partition between fine and coarse modes to the total AOD in both datasets will remain a factor).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

However, Fig. 9 shows that MODIS fine mode #2 is rarely picked as the “best” solution by either Terra or Aqua. Instead, Terra has a tendency to more frequently pick #3 or #4, and Aqua #1, which have lower (for Terra) or higher (for Aqua) C_n/C_v (and the reverse for per-particle extinction) than the maritime model applied to MAN data.

5 This will be responsible for the relative overestimate of Aqua and underestimate of Terra as compared to MAN. As the same algorithm is applied to both MODIS sensors, it seems reasonable to suspect that the differences in typical model choice between the two sensors could be linked with small systematic differences in their radiometric calibration; as the AOD is low for these cases, determination of size-related aerosol information is an inherently difficult task which is more sensitive than total AOD to un-
10 certainties and errors. These conclusions hold for both C5.1 and C6 (although it is likely that aerosol model selection will be more sensitive than total AOD to radiometric calibration changes which may be applied in the final C6 data).

The “average” MODIS solution matches MAN data closer than the “best” solution, for both AOD and C_n (Table 5), with higher correlations and smaller biases/absolute differences. This is likely because the averaging of several solutions reduces both retrieval noise and the effect of aerosol microphysical model assumptions on the retrieval. Although the MAN-derived C_n are not a “ground truth” for the satellite retrievals, these results suggest that the MODIS “average” solution may provide a better estimate of AOD and C_n for unpolluted maritime aerosol than the “best” solution. The differences
20 between these two solutions are generally smaller for C6 than C5.1.

4 Estimating surface concentration

The previous sections have dealt with columnar aerosol number and volume, as the MAN AOD measurements represent the column extinction. Determination of surface concentration requires knowledge of the vertical profile. Marine aerosol is often observed (e.g., Table 7) to show an exponentially decreasing number profile with height,
25

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[|◀](#)[▶|](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

$$n(z) = n_0 e^{-z/h}, \quad (8)$$

where $n(z)$ is the number concentration at altitude z , n_0 the surface concentration, and h the characteristic scale height. As by definition $\int_0^\infty n(z)dz = C_n$, simple integration of

- Eq. (8) yields $n_0 = C_n/h$. Table 7 suggests $h \approx 1.5$ km as a reasonable default assumption, although the typical range in Table 7 (1–2 km, a few around 0.5 km) will lead to around a factor of 2 variability in derived n_0 , highlighting the uncertainty in estimating surface concentration. Additionally, Table 8 shows that profiles other than exponentially-decreasing with height are also observed for marine aerosol. Finally, the assumption is required that the aerosol composition is invariant through the column, which may not always be true. Still, taking $h = 1.5$ km can provide a first-order estimate to examine average behaviour.

Unfortunately, direct comparison of MAN-derived estimates with in situ data is difficult due to a paucity of directly colocated data. However, general tendencies can be examined. Figure 10 shows a comparison between boundary layer aerosol number concentrations calculated from the maritime subset of the MAN data with other measurements and retrievals. These are intended to provide additional insight into typical values and spatial/temporal variability in different regions.

- The first of these is a collation of various field campaigns by Heintzenberg et al. (2000); the scale height used to convert the MAN column amounts to surface concentrations here ($h = 1.5$ km) corresponds to the typical value reported by Yu et al. (2010) for the geographical areas in which the data of Heintzenberg et al. (2000) were collected. MAN-derived latitudinal profiles for $h = 1$ km and $h = 2$ km are also given, to illustrate the magnitude of systematic uncertainty possible from scale height changes in this range. Field campaign measurements identified as belonging to maritime air masses by Bates et al. (2002) are also shown.

The World Meteorological Organisation (WMO) Global Atmosphere Watch (GAW) provide multiannual statistics of aerosol surface number concentration for multiple monitoring stations at <http://ebas.nilu.no>; shown here are multiannual median values and

Marine aerosol concentration from MAN

A. M. Sayer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



standard deviations for three low-lying island sites (Samoa, 14.5° S; Cape San Juan, 18.3° N; Sable Island, 43.9° N). Medians are used as it is likely these sites sometimes sample non-maritime air masses (so the median is likely a better estimate of the baseline maritime than the mean).

- 5 Finally, bimodal lognormal fits to median AERONET size distribution inversions representing maritime conditions at eleven sites were used by Sayer et al. (2012b) to constrain the maritime aerosol microphysical model applied in this work. These distributions have been used to calculate total columnar and then surface number concentration (again assuming $h = 1.5$ km), and also shown in Fig. 10. The error bars for
10 these points are taken using the relative standard deviation of τ_{440} at each site scaled to the total aerosol particle number, the rationale being that this wavelength is most sensitive to fine-mode particles. One site (Graciosa, 39.1° N) falls outside the scale at an estimated surface aerosol particle concentration of 3000 cm^{-3} ; Sayer et al. (2012b)
15 noted a higher fine-mode abundance here than at other sites, and speculated some contribution from a local aerosol source.

Poleward of 20° S, there is good agreement between the datasets. However, through the tropics and Northern Hemisphere the MAN-derived estimates are generally higher than the in situ data. There are multiple reasons why this could be the case. For the points poleward of 20° S, it could be that the scale height chosen is appropriate and
20 the aerosol is maritime in nature. Further north of this, land means there may be an increased influence of continental air masses and so the maritime model is less appropriate. Table 1 shows that discrepancies of this magnitude are possible due to uncertainty in aerosol microphysical properties. Non-coincidence of sampling (both spatial and temporal) is likely an important factor, as variabilities within each latitude range,
25 and at individual sites averaged over time, are large. A further possibility in the tropics is contamination by thin cirrus, which can be widespread and not always detected by ground-based instrumentation, leading to a positive bias in the MAN AOD (Chew et al., 2011).

Marine aerosol concentration from MAN

A. M. Sayer et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
	
	
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

Earlier reviews of field campaign data by Podzimek (1980) and Fitzgerald (1991) suggested typical concentrations in the range of several hundred cm^{-3} in open oceans but several thousand cm^{-3} in continentally-influenced air masses. Figure 2 indicates these could still be present in the data, as many of the “maritime” subset are still in continental outflow zones, and the remoter points tend to have lower number concentrations. On separate cruises through the Eastern Atlantic, Koponen et al. (2002) and Williams et al. (2007) reported similar high aerosol number concentrations along the coast of Europe and Northern Africa, but generally less than 1000 cm^{-3} in marine air masses in the Southern Atlantic at mid- and high latitudes. These results are also similar to recent CTM simulations, although such models are sensitive to e.g. emission and nucleation schemes (Spracklen et al., 2010).

Tables 7 and 8 show significant variability in vertical profile shape. If the bulk of the profile follows an exponential decrease with height but the near-surface layer is well-mixed and uniform, this would mean that assuming a relationship of the type of Eq. (8) overestimated surface concentration, as is observed. Instrumental artefacts such as incomplete sampling of the aerosol size distribution can also be a factor leading to underestimates of aerosol number in the in situ data (e.g., Reid et al., 2006), although Heintzenberg et al. (2000) performed additional filtering on their input data to minimise the likelihood of this.

The main conclusion from this exercise is therefore to illustrate the difficulties inherent in inferring near-surface quantities from columnar ones. This difficulty is also present when, for example, trying to estimate ground-level particulate matter concentrations from satellite measurements of AOD for air quality assessment (e.g., Hoff and Christopher, 2009, and references therein).

5 Conclusions

The remote sensing of spectral AOD from space is not a solved problem. Ground-based measurements by techniques such as sun photometry are able to make more

Marine aerosol concentration from MAN

A. M. Sayer et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



direct inferences about AOD, but with poorer spatial coverage. Remote sensing of aerosol number, volume, and mass would provide useful and important information about the Earth system, but is more complicated than retrieval of AOD, as it is more sensitive to assumptions relating to aerosol composition.

- 5 Using a microphysical model derived from AERONET inversions as a constraint, this study has attempted to determine columnar aerosol number and volume from ship-borne measurements of spectral AOD of fairly low uncertainty (~ 0.015) for cases where this microphysical model can be reasonably assumed to be appropriate (unpolluted maritime aerosol). Even with these constraints, the estimated uncertainty on
10 the derived quantities can be significant (10%–100 %), which probably precludes use of the technique with less accurate AOD data. Despite these uncertainties, the estimated concentrations are physically sensible. It is suggested that, in conditions where
15 a microphysical model for the dominant aerosol type can be prescribed with some confidence, accurate and precise spectral AOD measurements, such as from sun photometers deployed by the AERONET and MAN programs, could be used to estimate aerosol number or volume.

Potential applications of this method include an additional tool for comparison with CTM aerosol fields, and examining the fine/coarse partitions retrieved or assumed in satellite AOD retrieval algorithms. Currently, AERONET size distribution inversions are
20 sparse at some locations due to the requirement for clear skies and homogeneity over a period of one hour while almucantar scans necessary for the inversion algorithm are collected, plus a low Sun angle for high air mass factor (Dubovik and King, 2000). Although probably less accurate than these full inversions, estimates based on spectral AOD with the constraint of a microphysical model would expand the potential data
25 volume for comparison. In estimating volume/number from spectral AOD, this is complementary to the AERONET spectral deconvolution algorithm product (O'Neill et al., 2003), which uses a more generalised set of microphysical assumptions to estimate fine and coarse contributions to midvisible AOD (but not explicitly number or volume).

Marine aerosol concentration from MAN

A. M. Sayer et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



An attempt was made to convert MAN-derived columnar number concentrations into surface number concentrations, which were compared to typical values from in situ datasets and AERONET estimates. From around 20° S and poleward, similar values were obtained ($\sim 300\text{--}600\text{ cm}^{-3}$). However, in the tropics and Northern Hemisphere, the MAN-derived data tended to produce higher values than the in situ measurements. This poorer agreement is expected to be due to a combination of reasons including spatio-temporal variability of aerosol loading, uncertainties in aerosol microphysical properties, and, in particular, uncertainties in vertical profile shape.

- Columnar aerosol number and AOD at 550 nm were also compared with the current Collection 5.1 product from both MODIS sensors, and results using the forthcoming Collection 6 algorithm. Consistent with previous studies, MODIS was found to overestimate AOD as compared to the MAN data, with this overestimate being larger for MODIS Terra. The number to volume ratios and per-particle extinction of the different aerosol modes used in the MODIS retrieval over ocean can lead to significant differences in derived number concentration. It was found that Terra tended to estimate lower aerosol number than MAN, and Aqua higher, linked to differences between the sensors in the aerosol fine modes which are typically found to provide the best solution. The MODIS “average solution” dataset agreed more closely with the MAN data than the “best solution”, likely because some of the uncertainty associated with retrieval noise and microphysical model assumptions is averaged out. The results suggest that, at least for cases of pure maritime aerosol, quality assurance flags of 1, 2, and 3 are all of similar quality, and the “average solution” dataset is better than the “best solution” dataset. Collection 6 showed closer agreement with MAN AOD than Collection 5.1, but otherwise conclusions drawn were similar.
- Acknowledgements.** This work was supported by a grant from the NASA MEaSUREs program, managed by M. Maiden, and the EOS program, managed by H. Maring. The authors would like to acknowledge H. Maring for his support of the AERONET program. The authors are grateful to the AERONET, MAN, GAW, and field campaign PIs for the creation and stewardship of the ground-based data records used, and the Norwegian Institute for Air Research for hosting

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

the GAW data. L. A. Remer is thanked for useful discussions concerning the MODIS aerosol product. MODIS data were obtained from the NASA LAADS.

References

- Bates, T. S., Coffman, D. J., Covert, D. S., and Quinn, P. K.: Regional marine boundary layer aerosol size distributions in the Indian, Atlantic, and Pacific Oceans: a comparison of IN-DOEX measurements with ACE-1, ACE-2, and Aerosols99, *J. Geophys. Res.*, 107, 8026, doi:10.1029/2001JD001174, 2002. 14975
- Blanchard, D. C. and Woodcock, A. H.: The production, concentration, and vertical distribution of the sea-salt aerosol, in: *Annals of the New York Academy of Sciences*, Vol. 338, *Aerosols: Anthropogenic and Natural, Sources and Transport*, Wiley, New York, USA, doi:10.1111/j.1749-6632.1980.tb17130.x, 1980. 14996, 14997
- Campbell, J. R., Hlavka, D. L., Welton, E. J., Flynn, C. J., Turner, D. D., Spinhirne, J. . D., Scott, V. S., and Hawng, I. H.: Full-time, eye-safe cloud and aerosol lidar observation at atmospheric radiation measurement program sites: instrument and data processing, *J. Atmos. Ocean. Tech.*, 19, 431–442, doi:10.1175/1520-0426(2002)019j0431:FTESCA;2.0.CO;2, 2002. 14961
- Chew, B. N., Campbell, J. R., Reid, J. S., Giles, D. M., Welton, E. J., Salinas, S. V., and Liew, S. C.: Tropical cirrus cloud contamination in sun photometer data, *Atmos. Environ.*, 45, 6724–6731, doi:10.1016/j.atmosenv.2011.08.017, 2011. 14976
- Clarke, A. D. and Kapustin, V. N.: A Pacific aerosol survey. Part I: A decade of data on particle production, transport, evolution, and mixing in the troposphere, *J. Atmos. Sci.*, 59, 363–382, doi:10.1175/1520-0469(2002)059j0363:APASPI;2.0.CO;2, 2002. 14996
- Clarke, A. D., Uehara, T., and Porter, J. N.: Atmospheric nuclei and related aerosol fields over the Atlantic: clean subsiding air and continental pollution during ASTEX, *J. Geophys. Res.*, 102, 25281–25292, doi:10.1029/97JD01555, 1997. 14997
- Clarke, A. D., Collins, W. G., Rasch, P. J., Kapustin, V. N., Moore, K., Howell, S., and Fuelberg, H. E.: Dust and pollution transport on global scales: aerosol measurements and model predictions, *J. Geophys. Res.*, 106, 32555–32569, doi:10.1029/2000JD900842, 2001. 14996

12, 14959–15007, 2012

ACPD

Marine aerosol concentration from MAN

A. M. Sayer et al.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Marine aerosol concentration from MAN

A. M. Sayer et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- Heintzenberg, J., Covert, D. C., and van Dingenen, R.: Size distribution and chemical composition of marine aerosols: a compilation and review, *Tellus B*, 52, 1104–1122, doi:10.1034/j.1600-0889.2000.00136.x, 2000. 14969, 14975, 14977
- Hoff, R. and Christopher, S.: Remote sensing of particulate pollution from space: have we reached the promised land?, *J. Air Waste Manage.*, 59, 645–675, doi:10.3155/1047-3289.59.6.645, 2009. 14962, 14977
- Holben, B. N., Eck, T. F., Slutsker, I., Tanré, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A., Kaufman, Y. J., Nakajima, T., Lavenu, F., Jankowiak, I., and Smirnov, A.: AERONET: a federated instrument network and data archive for aerosol characterization, *Remote Sens. Environ.*, 66, 1–16, doi:10.1016/S0034-4257(98)00031-5, 1998. 14961, 14962
- Hsu, N. C., Herman, J. R., Torres, O., Holben, B. N., Tanré, D., Eck, T. F., Smirnov, A., Chatenet, B., and Lavenu, F.: Comparisons of the TOMS aerosol index with Sun-photometer aerosol optical thickness: results and applications, *J. Geophys. Res.*, 104, 6269–6279, doi:10.1029/1998JD200086, 1999. 14961
- Husar, R. B., Prospero, J. M., and Stowe, L. L.: Characterization of tropospheric aerosols over the oceans with the NOAA advanced very high resolution radiometer optical thickness operational product, *J. Geophys. Res.*, 102, 16889–16909, 1997. 14961
- Jefferson, A.: Empirical estimates of CCN from aerosol optical properties at four remote sites, *Atmos. Chem. Phys.*, 10, 6855–6861, doi:10.5194/acp-10-6855-2010, 2010. 14962
- Kahn, R. A., Nelson, D. L., Garay, M. J., Levy, R. C., Bull, M. A., Diner, D. J., Martonchik, J. V., Paradise, S. R., Hansen, E. G., and Remer, L. A.: MISR aerosol product attributes and statistical comparisons with MODIS, *IEEE T. Geosci. Remote*, 47, 4095–4114, doi:10.1109/TGRS.2009.2023115, 2009. 14971
- Kahn, R. A., Gaitley, B. J., Garay, M. J., Diner, D. J., Eck, T. F., Smirnov, A., and Holben, B. N.: Multiangle imaging spectroradiometer global aerosol product assessment by comparison with the Aerosol Robotic Network, *J. Geophys. Res.*, 115, D23209, doi:10.1029/2010JD014601, 2010. 14961
- Kahn, R. A., Garay, M. J., Nelson, D. L., Levy, R. C., Bull, M. A., Diner, D. J., Martonchik, J. V., Hansen, E. G., Remer, L. A., and Tanré, D.: Response to “Toward unified satellite climatology of aerosol properties: 3. MODIS versus MISR versus AERONET”, *J. Quant. Spectrosc. Ra.*, 112, 901–909, doi:10.1016/j.jqsrt.2010.11.001, 2011. 14971
- Kaskaoutis, D. G., Kharol, S. K., Sifakis, N., Nastos, P. T., Sharma, A. R., Badarinath, K. V. S., and Kambezidis, H. D.: Satellite monitoring of the biomass-burning aerosols during the

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

wildfires of August 2007 in Greece: climate implications, *Atmos. Environ.*, 45, 716–726, doi:10.1016/j.atmosenv.2010.09.043, 2011. 14971

Kaufman, Y. J., Smirnov, A., Holben, B. N., and Dubovik, O.: Baseline maritime aerosol: methodology to derive the optical thickness and scattering properties, *Geophys. Res. Lett.*, 28, 3251–3254, doi:10.1029/2001GL013312, 2001. 14970

Kaufman, Y. J., Koren, I., Remer, L. A., Tanré, D., Ginoux, P., and Fan, S.: Dust transport and deposition observed from the Terra-Moderate Resolution Imaging Spectroradiometer MODIS spacecraft over the Atlantic Ocean, *J. Geophys. Res.*, 110, D10S12, doi:10.1088/1748-9326/1/1/014005, 2005. 14962

Kinne, S., Schulz, M., Textor, C., Guibert, S., Balkanski, Y., Bauer, S. E., Berntsen, T., Berglen, T. F., Boucher, O., Chin, M., Collins, W., Dentener, F., Diehl, T., Easter, R., Feichter, J., Fillmore, D., Ghan, S., Ginoux, P., Gong, S., Grini, A., Hendricks, J., Herzog, M., Horowitz, L., Isaksen, I., Iversen, T., Kirkevåg, A., Kloster, S., Koch, D., Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Lesins, G., Liu, X., Lohmann, U., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S., Seland, O., Stier, P., Takemura, T., and Tie, X.: An AeroCom initial assessment – optical properties in aerosol component modules of global models, *Atmos. Chem. Phys.*, 6, 1815–1834, doi:10.5194/acp-6-1815-2006, 2006. 14963

Kleidman, R. G., Smirnov, A., Levy, R. C., Mattoo, S., and Tanré, D.: Evaluation and wind speed dependence of MODIS aerosol retrievals over open ocean, *IEEE T. Geosci. Remote*, 50, 429–435, doi:10.1109/TGRS.2011.2162073, 2012. 14971, 14972

Knobelispiesse, K. D., Pietras, C., and Fargion, G. S.: Sun-pointing-error correction for sea deployment of the MICROTOPS II handheld sun photometer, *J. Atmos. Ocean. Techn.*, 20, 767–771, 2003. 14964

Knobelispiesse, K. D., Pietras, C., Fargion, G. S., Wang, M., Frouin, R., Miller, M. A., Subramaniam, A., and Balch, W. M.: Maritime aerosol optical thickness measured by handheld Sun photometers, *Remote Sens. Environ.*, 93, 87–106, doi:10.1016/j.rse.2004.06.018, 2004. 14964, 14968

Koponen, I. K., Virkkula, A., Hillamo, R., Kerminen, V.-M., and Kulmala, M.: Number size distributions and concentrations of marine aerosols: observations during a cruise between the English Channel and the coast of Antarctica, *J. Geophys. Res.*, 107, 4753, doi:10.1029/2002JD002533, 2002. 14977

Kristament, I. S., Liley, J. B., and Harbey, M. J.: Aerosol variability in the vertical in the Southwest Pacific, *J. Geophys. Res.*, 98, 7129–7139, doi:10.1029/92JD02623, 1993. 14996

Marine aerosol concentration from MAN

A. M. Sayer et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Marine aerosol
concentration from
MAN

A. M. Sayer et al.

- Krüger, O. and Graßl, H.: Southern Ocean phytoplankton increases cloud albedo and reduces precipitation, *Geophys. Res. Lett.*, 38, L08809, doi:10.1029/2011GL047116, 2011. 14971
- Li, R., Min, Q.-L., and Harrison, L. C.: A case study: The indirect aerosol effects of mineral dust on warm clouds, *J. Atmos. Sci.*, 67, 805–816, doi:10.1175/2009JAS3235.1, 2010. 14971
- Livingston, J. M., Lapustin, V. N., Schmid, B., Russell, P. B., Quinn, P. K., Bates, T. S., Durkee, P. A., Smith, P. J., Freudenthaler, V., Wiegner, M., Covert, D. S., Gassó, S., Hegg, D., Collins, D. R., Flagan, R. C., Seinfeld, J. H., Vitale, V., and Tomasi, C.: Shipboard sunphotometer measurements of aerosol optical depth spectra and columnar water vapor during ACE-2, and comparison with selected land, ship, aircraft, and satellite measurements, *Tellus B*, 52, 594–619, doi:10.1034/j.1600-0889.2000.00045.x, 2000. 14996
- McClatchey, R. A., Fenn, R. W., Selby, J. E. A., Volz, F. E., and Garing, J. S.: Optical properties of the Atmosphere (3rd edn.), Tech. rep., report number AFCRL-TR-71-0279, environmental research paper 354, Air Force Cambridge Research Laboratories, Bedford, Mass, USA, 1972. 14996
- Michalsky, J. J., Schlemmer, J. A., Berkheiser, W. E., Berndt, J. L., Harrison, L. C., Laulainen, N. S., Larson, N. R., and Barnard, J. C.: Multiyear measurements of aerosol optical depth in the Atmospheric Radiation Measurement and Quantitative Links programs, *J. Geophys. Res.*, 106, 12099–12107, doi:10.1029/2001JD900096, 2001. 14961
- Mishchenko, M., Geogdzhayev, I. V., Cairns, B., Carlson, B. E., Chowdhary, J., Lacis, A. A., Liu, L., Rossow, W. B., and Travis, L. D.: Past, present, and future of global aerosol climatologies derived from satellite observations: a perspective, *J. Quant. Spectrosc. Ra.*, 106, 325–347, doi:10.1016/j.jqsrt.2007.01.007, 2007. 14961, 14971
- Mishchenko, M. I., Liu, L., Travis, L. D., Lacis, A. A., and Cairns, B.: Toward unified satellite climatology of aerosol properties: 3. MODIS versus MISR versus AERONET, *J. Quant. Spectrosc. Ra.*, 111, 540–552, 2010. 14971
- O'Neill, N. T., Ignatov, A., Holben, B. N., and Eck, T. F.: The lognormal distribution as a reference for reporting aerosol optical depth statistics; empirical tests using multi-year, multi-site AERONET sunphotometer data, *Geophys. Res. Lett.*, 27, 3333–3336, doi:10.1029/2000GL011581, 2000. 14969
- O'Neill, N. T., Eck, T. F., Smirnov, A., Holben, B. N., and Thulasiraman, S.: Spectral discrimination of coarse and fine mode optical depth, *J. Geophys. Res.*, 108, 4559–4573, doi:10.1029/2002JD002975, 2003. 14978

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
	
	
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



- Patterson, E. M., Kiang, C. S., Delany, A. C., Wartburg, A. F., Leslie, A. C. D., and Huebert, B. J.: Global measurements of aerosols in remote continental and marine regions: concentrations, size distributions, and optical properties, *J. Geophys. Res.*, 85, 7361–7376, doi:10.1029/JC085iC12p07361, 1980. 14996
- Podzimek, J.: Advances in marine aerosol research, *J. Rech. Atmos.*, 14, 35–61, 1980. 14977
- Porter, J. N., Miller, M., Pietras, C., and Motell, C.: Ship-based sun photometer measurements using Microtops sun photometers, *J. Atmos. Ocean. Techn.*, 18, 765–774, doi:10.1175/1520-0426(2001)018<0765:SBSPMU>2.0.CO;2, 2001. 14964, 14968
- Quaas, J., Boucher, O., Bellouin, N., and Kinne, S.: Satellite-based estimate of the direct and indirect aerosol climate forcing, *J. Geophys. Res.*, 113, D05204, doi:10.1029/2007JD008962, 2008. 14962
- Reid, J. S., Brooks, B., Crahan, K. K., Hegg, D. A., Eck, T. F., O'Neill, N., de Leeuw, G., Reid, E. A., and Anderson, K. D.: Reconciliation of coarse mode sea-salt aerosol particle size measurements and parameterizations at a subtropical ocean receptor site, *J. Geophys. Res.*, 111, D02202, doi:10.1029/2005JD006200, 2006. 14977
- Remer, L. A., Kaufman, Y. J., Tanré, D., Mattoo, S., Chu, D. A., Martins, J. V., Li, R.-R., Ichoku, C., Levy, R. C., Kleidman, R. G., Eck, T. F., Vermote, E., and Holben, B. N.: The MODIS aerosol algorithm, products, and validation, *J. Atmos. Sci.*, 62, 947–973, 2005. 14962, 14970, 14971, 14973
- Remer, L. A., Kleidman, R. G., Levy, R. C., Kaufman, Y. J., Tanré, D., Mattoo, S., Martins, J. V., Ichoku, C., Koren, I., Yu, H., and Holben, B. N.: Global aerosol climatology from the MODIS satellite sensors, *J. Geophys. Res.*, 113, D14S07, doi:10.1029/2007JD009661, 2008. 14961, 14962, 14970, 14971, 14972, 14973
- Remer, L. A., Tanré, D., and Kaufman, Y. J.: Algorithm for remote sensing of tropospheric aerosol from MODIS: collection 5, Tech. rep., ATBD-MOD-02, product ID MOD04/MYD04, available at: <http://modis.gsfc.nasa.gov/data/atbd/>, last access: May 2012, NASA Goddard Space Flight Center, 2009. 14990
- Russell, P. B., Hobbs, P. V., and Stowe, L. L.: Aerosol properties and radiative effects in the United States East Coast haze plume: an overview of the Tropospheric Aerosol Radiative Forcing Observational Experiment (TARFOX), *J. Geophys. Res.*, 104, 2213–2222, doi:10.1029/1998JD200028, 1999a. 14996
- Russell, P. B., Livingstone, J. M., Hignett, P., Kinne, S., Wong, J., Chien, A., Bergstrom, R., Durkee, P., and Hobbs, P. V.: Aerosol-induced radiative flux changes off the United

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Marine aerosol concentration from MAN

A. M. Sayer et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Smirnov, A., Holben, B. N., Giles, D. M., Slutsker, I., O'Neill, N. T., Eck, T. F., Macke, A., Croot, P., Courcoux, Y., Sakerin, S. M., Smyth, T. J., Zielinski, T., Zibordi, G., Goes, J. I., Harvey, M. J., Quinn, P. K., Nelson, N. B., Radionov, V. F., Duarte, C. M., Losno, R., Sciare, J., Voss, K. J., Kinne, S., Nalli, N. R., Joseph, E., Krishna Moorthy, K., Covert, D. S., Gulev, S. K., Milinevsky, G., Larouche, P., Belanger, S., Horne, E., Chin, M., Remer, L. A., Kahn, R. A., Reid, J. S., Schulz, M., Heald, C. L., Zhang, J., Lapina, K., Kleidman, R. G., Griesfeller, J., Gaitley, B. J., Tan, Q., and Diehl, T. L.: Maritime aerosol network as a component of AERONET – first results and comparison with global aerosol models and satellite retrievals, Atmos. Meas. Tech., 4, 583–597, doi:10.5194/amt-4-583-2011, 2011. 14962, 14964, 14971, 14972

Smirnov, A., Sayer, A. M., Holben, B. N., Hsu, N. C., Sakerin, S. M., Macke, A., Nelson, N. B., Courcoux, Y., Smyth, T. J., Croot, P., Quinn, P. K., Sciare, J., Gulev, S. K., Piketh, S., Losno, R., Kinne, S., and Radionov, V. F.: Effect of wind speed on aerosol optical depth over remote oceans, based on data from the Maritime Aerosol Network, Atmos. Meas. Tech., 5, 377–388, doi:10.5194/amt-5-377-2012, 2012. 14966

Spracklen, D. V., Carslaw, K. S., Merikanto, J., Mann, G. W., Reddington, C. L., Pickering, S., Ogren, J. A., Andrews, E., Baltensperger, U., Weingartner, E., Boy, M., Kulmala, M., Laakso, L., Lihavainen, H., Kivekäs, N., Komppula, M., Mihalopoulos, N., Kouvarakis, G., Jennings, S. G., O'Dowd, C., Birmili, W., Wiedensohler, A., Weller, R., Gras, J., Laj, P., Sellegrí, K., Bonn, B., Krejci, R., Laaksonen, A., Hamed, A., Minikin, A., Harrison, R. M., Talbot, R., and Sun, J.: Explaining global surface aerosol number concentrations in terms of primary emissions and particle formation, Atmos. Chem. Phys., 10, 4775–4793, doi:10.5194/acp-10-4775-2010, 2010. 14977

Sturm, B.: The atmospheric correction of remotely sensed data and the quantitative determination of suspended matter in marine water surface layers, in: Remote Sensing in Meteorology, Oceanography and Hydrology, edited by: Cracknell, A. P., Ellis Horwood, Chichester, UK, 1980. 14996

Tanré, D., Kaufman, Y. J., Herman, M., and Mattoo, S.: Remote sensing of aerosol properties over oceans using the MODIS/EOS spectral radiances, J. Geophys. Res., 102, 16971–16988, doi:10.1029/96JD03437, 1997. 14970

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



**Marine aerosol
concentration from
MAN**

A. M. Sayer et al.

5 Textor, C., Schulz, M., Guibert, S., Kinne, S., Balkanski, Y., Bauer, S., Berntsen, T., Berglen, T.,
Boucher, O., Chin, M., Dentener, F., Diehl, T., Easter, R., Feichter, H., Fillmore, D., Ghan, S.,
Ginoux, P., Gong, S., Grini, A., Hendricks, J., Horowitz, L., Huang, P., Isaksen, I., Iversen, I.,
Kloster, S., Koch, D., Kirkevåg, A., Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F.,
Liu, X., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S., Seland, Ø., Stier, P.,
Takemura, T., and Tie, X.: Analysis and quantification of the diversities of aerosol life cycles
within AeroCom, *Atmos. Chem. Phys.*, 6, 1777–1813, doi:10.5194/acp-6-1777-2006, 2006.
14963

10 Thomas, G. E., Poulsen, C. A., Sayer, A. M., Marsh, S. H., Dean, S. M., Carboni, E., Siddans, R.,
and Grainger, R. G.: Oxford-RAL Aerosol and Cloud (ORAC): aerosol retrievals from satellite
radiometers, in: *Aerosol Remote Sensing over Land*, edited by: Kokhanovsky, A. A. and de
Leeuw, G., Springer, Berlin, 2009. 14961

15 Torres, O., Bhartia, P. K., Herman, J. R., Sinyuk, A., Ginoux, P., and Holben, B.:
A long-term record of aerosol optical depth from TOMS observations and comparison
to AERONET measurements, *J. Atmos. Sci.*, 59, 398–413, doi:10.1175/1520-
0469(2002)059;0398:ALTROA;2.0.CO;2, 2002. 14961

20 van Dingenen, R., Raes, F., Putaud, J.-P., Virkkula, A., and Mangoni, M.: Processes determining
the relationship between aerosol number and non-sea-salt sulfate mass concentrations
in the clean and perturbed marine boundary layer, *J. Geophys. Res.*, 104, 8027–8038,
doi:10.1029/1998JD100059, 1999. 14970

25 Voss, K. J., Welton, E. J., Quinn, P. K., Johnson, J., Thompson, A. M., and Gordon, H. R.: Lidar measurements during Aerosols99, *J. Geophys. Res.*, 106, 20821–20831,
doi:10.1029/2001JD900217, 2001. 14997

30 Wagner, F. and Silva, A. M.: Some considerations about Ångström exponent distributions, *Atmos. Chem. Phys.*, 8, 481–489, doi:10.5194/acp-8-481-2008, 2008. 14961

Welton, E. J., Voss, K. J., Quinn, P. K., Flatau, P. J., Markowicz, K., Campbell, J. R., Spin-
hire, J. D., Gordon, H. R., and Johnson, J. E.: Measurements of aerosol vertical profiles
and optical properties during INDOEX 1999 using micropulse lidars, *J. Geophys. Res.*, 107,
8019, doi:10.1029/2000JD000038, 2002. 14997

Williams, P. I., McFiggans, G., and Gallagher, M. W.: Latitudinal aerosol size distribution variation
in the Eastern Atlantic Ocean measured aboard the FS-Polarstern, *Atmos. Chem. Phys.*,
7, 2563–2573, doi:10.5194/acp-7-2563-2007, 2007. 14977

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Yershov, O. A., Smirnov, A. V., and Shifrin, K. S.: The vertical profile of aerosol optical depths of maritime atmosphere (in Russian), Leningrad, in: Atmospheric and Marine Optics, edited by: Shifrin, K. S., Leningrad, Russia, 1988. 14996

5 Yu, F., Luo, G., Turco, R. P., Ogren, J. A., and Yantosca, R. M.: Decreasing particle number concentrations in a warming atmosphere and implications, *Atmos. Chem. Phys.*, 12, 2399–2408, doi:10.5194/acp-12-2399-2012, 2012. 14962

10 Yu, H., Chin, M., Winker, D. M., Omar, A. H., Liu, Z., Kittake, C., and Diehl, T.: Global view of aerosol vertical distributions from CALIPSO lidar measurements and GO-CART simulations: Regional and seasonal variations, *J. Geophys. Res.*, 115, D00H30, doi:10.1029/2009JD013364, 2010. 14975, 14996

Zhang, J. and Reid, J. S.: MODIS aerosol product analysis for data assimilation: assessment of over-ocean level 2 aerosol optical thickness retrievals, *J. Geophys. Res.*, 111, D22207, doi:10.1029/2005JD006898, 2006. 14971

Marine aerosol concentration from MAN

A. M. Sayer et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Marine aerosol concentration from MAN

A. M. Sayer et al.

Table 1. Aerosol microphysical model parameters. r_n is the modal radius, σ the geometric standard deviation, m the complex refractive index at 550 nm, and C_n/C_v the ratio of aerosol particle number to volume. The extinction coefficient at 550 nm is denoted β_{550} , given both per μm^3 and per particle.

Reference	Aerosol model	Parameters			β_{550} , μm^{-3}	β_{550} , per particle, $\times 10^{-14}$	C_n/C_v , μm^{-3}
		r_n , μm	σ	m			
Sayer et al. (2012b)	Marine, fine mode	0.0742	0.50	1.415–0.002 <i>i</i>	4.27	2.25	190
	Marine, coarse mode	0.547	0.72	1.363–3 $\times 10^{-9}$; 0.90	637		0.14
Sayer et al. (2012a)	Marine/continental, fine mode	0.106	0.44	1.43–0.0075 <i>i</i>	5.53	6.65	83
	Marine/continental, coarse mode	0.774	0.65	1.43–0.0075 <i>i</i>	0.78	1010	0.08
Remer et al. (2009)	Marine/dust, fine mode	0.0632	0.43	1.47–0.002 <i>i</i>	3.36	0.82	412
	Marine/dust, coarse mode	0.993	0.49	1.47–0.002 <i>i</i>	0.96	1060	0.08
MODIS #1 (fine mode)	0.07	0.40	1.45–0.0035 <i>i</i>	3.21	0.95	339	
	0.06	0.60	1.45–0.0035 <i>i</i>	5.17	2.36	219	
	0.08	0.60	1.40–0.002 <i>i</i>	5.09	5.51	92	
	0.1	0.60	1.40–0.002 <i>i</i>	5.36	11.4	47	
	0.4	0.60	1.35–0.001 <i>i</i>	2.06	278	0.74	
	0.6	0.60	1.35–0.001 <i>i</i>	1.26	576	0.22	
	0.8	0.60	1.35–0.001 <i>i</i>	0.90	973	0.09	
	0.6	0.60	1.53–0.001 <i>i</i>	1.22	557	0.22	
	0.5	0.80	1.53–0.001 <i>i</i>	0.71	658	0.11	

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



**Marine aerosol
concentration from
MAN**

A. M. Sayer et al.

Table 2. Statistics of errors on spectral AOD fit using the maritime model and the MAN series-average data. Biases are defined as the fit value minus the MAN value, i.e. positive indicates the model overpredicts AOD.

Wavelength	Number of points	Mean bias	Mean absolute bias	Standard deviation of bias
340 nm	3273	-0.004	0.006	0.007
380 nm	4200	-0.0008	0.003	0.005
440 nm	14 504	0.002	0.005	0.007
500 nm	15 804	-0.00007	0.005	0.008
675 nm	15 403	0.0003	0.004	0.006
870 nm	15 748	-0.0005	0.004	0.005
1020 nm	967	-0.002	0.004	0.007

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



**Marine aerosol
concentration from
MAN**

A. M. Sayer et al.

Table 3. Number of points in each aerosol class category, as described in the text, for series-average and daily-average MAN data. Figures in parentheses indicate the number which could not be fit to the maritime aerosol model with $\chi^2 < 1$.

Category	Series	Daily
Total	15 804 (1437)	2813 (177)
Maritime	4578 (253)	884 (50)
Dusty	1573 (267)	280 (33)
Continental	9653 (917)	1649 (94)



**Marine aerosol
concentration from
MAN**

A. M. Sayer et al.

Table 4. Lognormal mode and spread, and median values, of relative frequency histograms of aerosol volume (for all three classes) and number (for maritime only).

Category	Fine mode			Coarse mode		
	Mode	Spread	Median	Mode	Spread	Median
Maritime (volume, $\mu\text{m}^3 \mu\text{m}^{-2}$)	0.0036	0.70	0.0056	0.047	0.59	0.062
Maritime (number, μm^{-2})	0.71	0.66	1.07	0.0065	0.57	0.088
Dusty (volume, $\mu\text{m}^3 \mu\text{m}^{-2}$)	0.0160	0.41	0.0188	0.225	0.46	0.289
Continental (volume, $\mu\text{m}^3 \mu\text{m}^{-2}$)	0.0037	1.08	0.0136	0.013	1.09	0.037

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
◀	▶
◀	▶
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	



Marine aerosol concentration from MAN

A. M. Sayer et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Table 5. Statistics of comparison between colocated MAN and MODIS 550 nm AOD and aerosol columnar number concentration, for the MAN measurement series identified as marine aerosol. “Best” and “average” solutions are as defined in the text. “Scaled” refers to statistics when the MODIS data are scaled to match the MAN AOD at 550 nm. R is Pearson’s linear correlation coefficient and RMSD the root mean square difference. C6 statistics are given in parentheses, after C5.1 results in regular type.

Quantity/ solution	R	Median bias (MODIS – MAN)	Median absolute bias	RMSD
MODIS Terra, QA = 1,2,3 (306 C5.1 points, 260 C6 points)				
C_n , μm^{-2} , best	0.26 (0.39)	-0.17 (-0.20)	0.51 (0.63)	1.14 (1.17)
C_n , μm^{-2} , average	0.38 (0.47)	-0.016 (-0.16)	0.48 (0.60)	0.93 (0.96)
τ_{550} , best	0.74 (0.77)	0.027 (0.020)	0.028 (0.025)	0.046 (0.036)
τ_{550} , average	0.75 (0.77)	0.023 (0.016)	0.025 (0.021)	0.043 (0.034)
C_n , μm^{-2} , best, scaled	0.35 (0.43)	-0.32 (-0.03)	0.58 (0.64)	1.01 (1.13)
C_n , μm^{-2} , average, scaled	0.45 (0.57)	-0.23 (-0.24)	0.47 (0.50)	0.86 (0.85)
MODIS Aqua, QA = 1,2,3 (291 C5.1 points, 231 C6 points)				
C_n , μm^{-2} , best	0.34 (0.42)	0.65 (0.77)	1.02 (1.07)	1.61 (1.59)
C_n , μm^{-2} , average	0.48 (0.51)	0.42 (0.32)	0.71 (0.76)	1.13 (1.13)
τ_{550} , best	0.62 (0.73)	0.017 (0.009)	0.024 (0.021)	0.044 (0.031)
τ_{550} , average	0.63 (0.76)	0.016 (0.012)	0.023 (0.018)	0.044 (0.029)
C_n , μm^{-2} , best, scaled	0.46 (0.42)	0.29 (0.59)	0.87 (1.11)	1.62 (1.66)
C_n , μm^{-2} , average, scaled	0.57 (0.54)	0.10 (0.19)	0.51 (0.57)	1.06 (0.93)
MODIS Terra, QA = 3 (115 C5.1 points, 191 C6 points)				
C_n , μm^{-2} , best	0.53 (0.51)	-0.18 (-0.30)	0.41 (0.61)	0.91 (0.97)
C_n , μm^{-2} , average	0.58 (0.54)	-0.060 (-0.24)	0.43 (0.58)	0.79 (0.88)
τ_{550} , best	0.67 (0.82)	0.039 (0.021)	0.039 (0.023)	0.058 (0.036)
τ_{550} , average	0.69 (0.81)	0.038 (0.017)	0.038 (0.021)	0.052 (0.033)
C_n , μm^{-2} , best, scaled	0.64 (0.57)	-0.46 (-0.41)	0.53 (0.65)	0.95 (0.91)
C_n , μm^{-2} , average, scaled	0.67 (0.65)	-0.32 (-0.31)	0.46 (0.51)	0.82 (0.77)
MODIS Aqua, QA = 3 (97 C5.1 points, 96 C6 points)				
C_n , μm^{-2} , best	0.42 (0.50)	1.21 (0.77)	1.29 (1.00)	1.94 (1.73)
C_n , μm^{-2} , average	0.64 (0.61)	0.85 (0.44)	0.85 (0.68)	1.32 (1.16)
τ_{550} , best	0.49 (0.85)	0.037 (0.018)	0.039 (0.028)	0.059 (0.032)
τ_{550} , average	0.48 (0.86)	0.028 (0.017)	0.035 (0.021)	0.059 (0.028)
C_n , μm^{-2} , best, scaled	0.62 (0.47)	0.62 (0.67)	0.90 (1.08)	1.61 (1.84)
C_n , μm^{-2} , average, scaled	0.79 (0.62)	0.35 (0.24)	0.46 (0.51)	0.96 (0.98)

**Marine aerosol
concentration from
MAN**

A. M. Sayer et al.

Table 6. Fraction of MODIS-MAN matchups with an AOD at 550 nm within the expected MODIS absolute error of $0.03 + 5\%$ of the MAN value.

Sensor/QA threshold	Best solution		Average solution	
	C5.1	C6	C5.1	C6
Terra, QA = 1,2,3	0.60	0.71	0.62	0.74
Terra, QA = 3	0.41	0.72	0.46	0.77
Aqua, QA = 1,2,3	0.66	0.73	0.71	0.78
Aqua, QA = 3	0.45	0.71	0.52	0.76

Marine aerosol concentration from MAN

A. M. Sayer et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

◀

▶

◀

▶

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Table 7. Scale heights for marine aerosol, from various sources.

Reference	Scale height, km	Region	Note/origin in reference
Yu et al. (2010)	0.5–1.5	Global	Fig. 6, Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) profiles. Open ocean.
Clarke and Kapustin (2002)	1.5–2.5 ~ 2.2	Global Pacific	Continental outflow regions. Fig. 13; continentally-influenced air masses.
Clarke et al. (2001)	~ 1 ~ 1.5–2	Hawaii	Fig. 13; clean air masses. Figs. 1 and 3; polluted/dusty air masses, evidence of some distinct layers.
Schmid et al. (2000)	~ 0.8	Canary Islands	Fig. 2; also Figs. 3 and 6 of Gassó et al. (2000).
Livingston et al. (2000)	~ 0.9 ~ 0.9	Canary Islands Coastal N. Atlantic	Fig. 10. Fig. 4; little variability in lowest 0.4 km.
Russell et al. (1999a,b), Hegg et al. (1996)	~ 1–1.5 0.13–0.52	US coast (Virginia) US coast (California)	Plate 1 of Russell et al. (1999b). Table 5; some measurements sampled very limited altitude ranges.
Kristament et al. (1993) Fitzgerald (1989)	0.25–1.11 ~ 1.5 ~ 0.25–1.2	US coast (Washington) Southwest Pacific –	Fig. 1. Fig. 5; theoretical calculations for relative humidities between 52 % and 85 %. Helicopter measurements.
Yershov et al. (1988) Blanchard and Woodcock (1980)	1.7 ~ 1	Black Sea Atlantic, Pacific	Fig. 7; wind speeds of 8 ms ⁻¹ and 14 ms ⁻¹ . Coarse mode particles only.
Patterson et al. (1980)	~ 0.5	Pacific	Fig. 6; large particles.
Sturm (1980)	~ 1 1.1–2	–	Small particles. Eq. (11.22) in lower troposphere; range for 50 km visibility (1.1 km) to 10 km visibility (2 km). Based on McClatchey et al. (1972) profiles.
Sebacher et al. (1967)	~ 1.5	US coast (Virginia)	Fig. 3 (bottom).

Marine aerosol concentration from MAN

A. M. Sayer et al.

Table 8. Marine aerosol vertical profiles which are not well-represented with exponentially-decreasing vertical profiles, from various sources.

Reference	Region	Profile/origin in reference
Welton et al. (2002)	Indian Ocean	Figs. 9 and 13; peak ~ 0.5 km, decay above and below.
Voss et al. (2001)	Atlantic	Fig. 7; increasing from surface, capped at 1 km.
Clarke et al. (1997)	Atlantic	Fig. 9; distinct layer up to ~ 0.6 km.
Blanchard and Woodcock (1980)	Atlantic, Pacific	Fig. 7; wind speeds of 1 ms^{-1} and 3.5 ms^{-1} . Little vertical variation. Coarse mode particles only.
Sebacher et al. (1967)	US coast (Virginia)	Fig. 3 (top); peak around 1 km.

Title Page	Abstract	Introduction
Conclusions	References	
Tables	Figures	
◀	▶	
◀	▶	
Back	Close	
Full Screen / Esc		
Printer-friendly Version		
Interactive Discussion		



**Marine aerosol
concentration from
MAN**

A. M. Sayer et al.




Fig. 1. Conceptual overview of fitting spectral AOD to estimate aerosol volume as a combination of fine and coarse modes.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)

Marine aerosol concentration from MAN

A. M. Sayer et al.




Fig. 2. Aerosol optical depth and inferred columnar aerosol volume from daily average MAN data. From left-right, columns show plots for the maritime, dusty, and continental subsets, respectively. Rows show (a–c) 500 nm AOD, (d–f) the fine-mode aerosol volume, (g–i) the coarse-mode aerosol volume, (j–l) fraction of aerosol volume from the coarse mode, and (m–o) the fit χ^2 .

**Marine aerosol
concentration from
MAN**

A. M. Sayer et al.




Fig. 3. Latitudinal distribution of the number of MAN measurement series falling into the maritime, dusty, and continental-influenced classifications.

Marine aerosol concentration from MAN

A. M. Sayer et al.




Fig. 4. Columnar aerosol volume and relative uncertainty (i.e., 1 indicates 100 % uncertainty), for the marine subset of points from daily average MAN data. **(a)** shows the fine mode, and **(b)** the coarse mode. Black points show calculations assuming the MAN uncertainty is 0.015, and red points where the fit χ^2 is used to scale these uncertainty estimates.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Marine aerosol concentration from MAN

A. M. Sayer et al.




Fig. 5. Histograms of fit columnar aerosol volume for the three aerosol classes, for (a) fine and (b) coarse modes. The vertical axis indicates the number in each bin, as a fraction of the total number of points.

- [Title Page](#)
- [Abstract](#) [Introduction](#)
- [Conclusions](#) [References](#)
- [Tables](#) [Figures](#)
- [◀](#) [▶](#)
- [◀](#) [▶](#)
- [Back](#) [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)



Marine aerosol concentration from MAN

A. M. Sayer et al.




Fig. 6. Maps of fit columnar aerosol number for the maritime aerosol subset, for (a) fine and (b) coarse modes. From the daily-average MAN data.

Title Page	
Abstract	Introduction
Conclusions	References
Tables	Figures
	
	
Back	Close
Full Screen / Esc	
Printer-friendly Version	
Interactive Discussion	

Marine aerosol concentration from MAN

A. M. Sayer et al.




Fig. 7. Histograms of fit columnar aerosol number for the maritime aerosol subset, for **(a)** fine and **(b)** coarse modes. The vertical axis indicates the number in each bin, as a fraction of the total number of points. Black shows the binned data, and red the lognormal fit to it. From the series-average MAN data.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Marine aerosol concentration from MAN

A. M. Sayer et al.




Fig. 8. Comparison between MAN and MODIS-derived AOD and C_n , from MODIS Collection 5.1. The top row (**a–c**) shows the MODIS “best” solution, and the bottom row (**d–f**) the “average” solution. Panels show comparisons between (**a, d**) AOD at 550 nm (outer dashed lines show the MODIS expected error envelope of $0.03 + 5\%$); (**b, e**) C_n ; and (**c, f**) C_n scaled to the value which would be reported if MODIS and MAN AOD matched at 550 nm.

Title Page	Abstract	Introduction
Conclusions	References	
Tables	Figures	
◀	▶	
◀	▶	
Back	Close	
Full Screen / Esc		
Printer-friendly Version		
Interactive Discussion		



Marine aerosol concentration from MAN

A. M. Sayer et al.




Fig. 9. Locations and statistics of MODIS and MAN-derived C_n , for QA = 3 MODIS retrievals and the “best” MODIS solution. The top row (**a–c**) shows data for MODIS Terra, and the bottom row (**d–f**) MODIS Aqua, from MODIS Collection 5.1. Panels show (**a, d**) the MAN-derived C_n ; (**b, e**) the MODIS-MAN C_n difference; and (**c, f**) the index of the fine aerosol mode (Table 1) MODIS reported as providing the “best” solution.

- [Title Page](#)
- [Abstract](#) | [Introduction](#)
- [Conclusions](#) | [References](#)
- [Tables](#) | [Figures](#)
- [◀](#) | [▶](#)
- [◀](#) | [▶](#)
- [Back](#) | [Close](#)
- [Full Screen / Esc](#)
- [Printer-friendly Version](#)
- [Interactive Discussion](#)



Marine aerosol concentration from MAN

A. M. Sayer et al.




Fig. 10. Zonal estimates of aerosol surface number concentration from MAN data, from the maritime subset of the daily average dataset. The mean is shown in red and median in black, and the shaded grey area indicates the central 68 % of the data (for latitude ranges with more than one point), assuming an aerosol scale height of 1.5 km. Dashed and dot-dashed red lines indicate the mean profile if scale heights of 1 km or 2 km are used instead. Coloured open symbols show different comparative datasets, as described in the text, with error bars indicating the variability on the data (typically standard deviation) as quoted in the relevant references. Numbers below AERONET (Sayer et al., 2012b) and GAW points indicate stations as follows: 1, Crozet Island; 2, Amsterdam Island; 3, Tahiti; 4, Ascension Island; 5, Nauru; 6, Kaashidhoo; 7, Guam; 8, Lanai; 9, Midway Island; 10, Bermuda; 11, Samoa; 12, Cape San Juan; 13, Sable Island.

Title Page	Abstract	Introduction
Conclusions	References	
Tables	Figures	
◀	▶	
◀	▶	
Back	Close	
Full Screen / Esc		
Printer-friendly Version		
Interactive Discussion		