

**OMI, TES, and CTM  
tropospheric ozone  
columns**

Q. Tang and M. J. Prather

# Tropospheric column ozone: matching individual profiles from Aura OMI and TES with a chemistry-transport model

Q. Tang<sup>1,2</sup> and M. J. Prather<sup>1</sup>

<sup>1</sup>Department of Earth System Science, University of California, Irvine, California, 92697, USA

<sup>2</sup>Department of Biological and Environmental Engineering, Cornell University, Ithaca, New York, 14853, USA

Received: 6 June 2012 – Accepted: 10 June 2012 – Published: 29 June 2012

Correspondence to: Q. Tang (qtang@cornell.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Of all satellite measurements of ozone, only two instruments have coincident, spatially overlapping measurements to allow direct comparison of tropospheric column ozone (TCO): the Ozone Monitoring Instrument (OMI) and Tropospheric Emission Spectrometer (TES) on the NASA Aura spacecraft. For two years (2005–2006), we collect all observations between 60° S and 60° N from nadir (~ 65 000 from OMI and TES) and cross-track swaths (~ 30 000 000 from OMI) and compare with a chemistry-transport model (CTM) simulating each observation with corresponding spatial and temporal coincidence. High-frequency TCO variations are indicative of stratospheric intrusions of ozone-rich air, and the individual, level 2 data provide access to these short-lived phenomena. Although we can identify some seasonal and large-scale biases in the model, the CTM as a transfer standard identifies weaknesses in the observations and further helps quantify the measurement noise of individual profiles. The relatively noise-free CTM bridges these two satellite measurements and improves their cross-validation to better precision than a simple direct comparison. Previous validation studies of TES TCO versus ozonesondes found a bias of about +4 Dobson Units (DU) for large regions. The three-way comparison, with a far greater number of coincidences, indicates that monthly mean OMI-TES TCO biases fall within a few percent, and thus quantifies the OMI TCO bias at a few DU. For small regions (i.e.,  $5 \times 5^\circ$ ), however, the monthly mean OMI-TES differences can exceed  $\pm 10$  DU at many places (e.g., tropics) due to different tropospheric sensitivities of the two instruments at these locations. In addition, the CTM-TES comparison split into day versus night observations shows no apparent bias in TES at very low levels,  $\pm 1$  DU. These OMI-TES-CTM comparisons highlighted the importance of the a priori ozone profiles that went into each satellite retrieval, including a false agreement due to CTM-a priori similarity. This study highlights the advantages of overlapping measurements in terms of cross-validation and the application of a model as a transfer standard.

## OMI, TES, and CTM tropospheric ozone columns

Q. Tang and M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 1 Introduction

Among the four Earth Observing System (EOS) Aura instruments – High Resolution Dynamics Limb Sounder (HIRDLs), Microwave Limb Sounder (MLS), Ozone Monitoring Instrument (OMI), and Tropospheric Emission Spectrometer (TES) – only OMI and TES have coincident ozone ( $O_3$ ) measurements through the troposphere, providing the information content and the opportunity to compare tropospheric column ozone (TCO) with these two datasets directly on a single, overlapping measurement basis as is presented here. Applying a chemistry-transport model (CTM) as the transfer standard to compare different satellite measurements provides a unique opportunity for both model and measurement validation, but it brings into the comparison model errors, which are not always well understood (e.g., Eyring et al., 2010; Holmes et al., 2011). Other approaches to validate satellite data use sonde, lidar, or aircraft measurements, but these involve few direct comparisons, often with measurements that are not closely geo-located in space or time and are not globally or seasonally representative. Direct comparison of satellite measurements provides far more samples, pairs of observations that truly overlap, and coverage that is global and continuous. A geo-located, comprehensive and representative cross-validation of satellite measurements of ozone is especially important for studies focusing on processes with high-frequency variations and short-lived features, such as tropopause folds (TFs).

It remains difficult still to compare directly the profiles of seemingly coincident satellite measurements. For example, (Tang and Prather, 2012, Figs. 3, 4, A1–A4) compared the four Aura instruments for several swaths in which a major TF was observed by the sonde and predicted by the CTM, but found that differing horizontal and vertical resolution, sensitivities, as well as offset from the nadir track made it difficult to pinpoint the TFs. OMI observes a wide cross-track swath, while TES observes a sequence of nadir spots that fall in one of the OMI pixels (see below). Even these two overlapping nadir sounders have different sensitivities to  $O_3$  at different altitudes, and their retrieved profiles of  $O_3$  abundance have fundamentally different averaging kernels (AK) and initial

### OMI, TES, and CTM tropospheric ozone columns

Q. Tang and M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



guess for the O<sub>3</sub> profile (a priori). At altitudes where the observed spectrum is insensitive to the O<sub>3</sub> abundance, the retrieved profile is predominantly the a priori. Previous studies (Rodgers and Connor, 2003; Luo et al., 2007a; Ho et al., 2009; Zhang et al., 2010) have shown that if one is to exclude the impacts of different a priori constraints and smoothing errors due to different AK, then one must use the same a priori for both retrievals and to use equivalently coarse vertical resolution in the AK for each.

In this study, we examine the mean biases, covariances, and monthly spatial patterns of the coincident TCO fields from Aura OMI and TES level 2 (L2) swath data as well as the University of California, Irvine (UCI) CTM. Descriptions of the model and measurements are given in Sect. 2. Section 3 describes the main results, while Sect. 4 examines day-night differences in the TES measurements. The conclusions in Sect. 5 discuss the magnitude of model bias error and measurement noise for TCO measurements.

## 2 Measurements and model

The EOS Aura satellite was launched on 15 July 2004 to a sun-synchronous polar orbit 705 km above the sea surface with a 98° inclination. Aura is part of the A-Train and crosses the equator about 13:45 local time. Two nadir-viewing, passive sounding instruments, OMI and TES, are on Aura (Schoeberl et al., 2006). OMI measures at ultraviolet and visible wavelengths (UV-1: 264–311 nm, UV-2: 307–383 nm, VIS: 349–504 nm) (OMI Team, 2009), whereas TES measures thermal emissions in the infrared (650–3250 cm<sup>-1</sup>) (Osterman et al., 2009). OMI uses a 2-dimensional Charge-Coupled Device to scan a wide area (2600 km cross-orbit swath with 13 km×48 km resolution), while TES stares at a small spot (5 km×8 km nadir footprint) to measure the Fourier transform of the spectrum. As to the a priori O<sub>3</sub> profiles used in the retrievals, OMI uses a latitude-by-month climatology based upon observations (McPeters et al., 2007; Kroon et al., 2011), whereas TES adopts a climatology that varies with latitude and longitude derived from the MOZART CTM (Brasseur et al., 1998; Park et al., 2004). Level

### OMI, TES, and CTM tropospheric ozone columns

Q. Tang and M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## OMI, TES, and CTM tropospheric ozone columns

Q. Tang and M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2 (L2) orbit/swath data are used in this study: the OMI OMO3PR V003 data (de Haan and Veefkind, 2009) and the TES version 4 (V004, F05\_07) nadir global survey product. Although both datasets report ozone profiles, they contain only about 1–2 degrees of freedom for the signal (DOFS) in the troposphere (Nassar et al., 2008; Zhang et al., 5 2010), and thus provide limited profile resolution but reasonable TCO information. The tropopause heights simulated by the UCI CTM with a tropospheric-age tracer (e90) (Prather et al., 2011; Tang et al., 2011) are chosen for each satellite coincidence, and these pressure levels are applied to spline-interpolated OMI and TES profiles to calculate each TCO. TES data whose “SpeciesRetrievalQuality” or “O3\_Ccurve\_QA” is not equal to 1 are removed based on Osterman et al. (2009).

The UCI CTM has  $1 \times 1^\circ \times 40$ -layer  $\times 0.5$  hr resolution for years 2005–2006 and is driven by the pieced-forecast meteorology from the European Centre for Medium-Range Weather Forecasts (ECMWF) Integrated Forecast System developed by the University of Oslo (Kraabøl et al., 2002; Isaksen et al., 2005). The CTM includes tropospheric chemistry (Carver et al., 1997) as well as the linearized O<sub>3</sub> chemistry (Linoz version 2) (Hsu and Prather, 2009). For further details see Prather et al. (2008); Prather and Hsu (2010); Tang and Prather (2010). Satellite tracks are stored from the simulation to match the exact timing and location of OMI and TES observations (see Tang and Prather, 2012, for details). OMI and TES data are assigned to the  $1 \times 1^\circ$  model grid boxes that include the center point of observations. All comparisons here are made with matching pairs on the model grids.

Raw CTM profiles need to be convolved with the satellite operator (i.e., a priori and AK) before comparing with satellite data following the equation (e.g. Luo et al., 2007b; Worden et al., 2007):

$$25 \quad \hat{\mathbf{x}}_m = \mathbf{x}_a + \mathbf{A}(\mathbf{x}_m - \mathbf{x}_a) \quad (1)$$

where  $\mathbf{x}_a$  and  $\mathbf{A}$  are the a priori and AK, respectively.  $\mathbf{x}_m$  represents the CTM profiles interpolated on satellite levels. The convolved CTM profile  $\hat{\mathbf{x}}_m$  then contains the contributions from the a priori and the vertical smoothing of satellite retrievals and is thus

appropriate for comparison with satellite data. The CTM TCO in this study is calculated from  $\hat{x}_m$  with spline interpolation and the modeled tropopause.

OMI and TES use different a priori profiles, thus introducing artificial differences in their retrieved profiles. To remove the a priori influence in direct OMI-TES comparisons, the TES a priori ( $x_a^{\text{TES}}$ ) is replaced by the OMI a priori ( $x_a^{\text{OMI}}$ , interpolated to the TES pressure grid) by the equation (Rodgers and Connor, 2003; Luo et al., 2007a):

$$\hat{x}^{\text{TES}} = x^{\text{TES}} + (\mathbf{I} - \mathbf{A}^{\text{TES}})(x_a^{\text{OMI}} - x_a^{\text{TES}}) \quad (2)$$

where  $x^{\text{TES}}$  is the original TES profile and  $\hat{x}^{\text{TES}}$  is the TES retrieval with OMI a priori.  $\mathbf{I}$  is a unit matrix and  $\mathbf{A}^{\text{TES}}$  is the TES AK. Given the limited DOFS in the troposphere, their integrated TCO is expected to be less dependent on the AK, and we do not adjust the retrievals due to AK differences. Some TES profiles have no AK and are dropped from this analysis.

### 3 Comparisons in TCO for CTM, OMI and TES

The comparisons of TCO between coincident CTM, OMI and TES data are shown as 2-D probability density functions (PDFs) in Fig. 1 for Northern Hemisphere (NH) middle latitudes (25–60° N) July 2005 (left column) and January 2006 (right column). The number of comparisons ( $N$ ), mean bias ( $\mu$ ), standard deviation (STD,  $\sigma$ ) of the differences, and coefficient of determination ( $R^2$ ) are shown for each panel. The mean bias and STD are defined for the vertical-axis variable less the horizontal-axis variable. The correlation coefficient ( $R$ ) is calculated by removing only the means and without slope correction. There are ~400 000 CTM-OMI comparisons each month (Fig. 1a, b), while only ~8 000 CTM-TES pairs (Fig. 1c, d), because of the wide cross-orbit OMI swath and the sparse TES footprint. The high density areas (red and yellow pixels) are generally close to the 1:1 line (black line), reflecting small monthly mean biases between the model and measurements. The largest absolute monthly mean bias (–4.0

## OMI, TES, and CTM tropospheric ozone columns

Q. Tang and M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



---

**OMI, TES, and CTM  
tropospheric ozone  
columns**Q. Tang and M. J. Prather

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Dobson Unit (DU)) is found between CTM and OMI in July 2005 (Fig. 1a). The CTM is biased low relative to both measurements in July 2005, while it is high in January 2006, pointing to a CTM deficiency. The STDs are at least 1.5 times larger than the mean biases. They are larger in July than in January for both model-measurement comparisons, likely due to larger TCO variations over NH mid-latitudes in summer than in winter. The model-measurement biases and STDs are similar for both OMI and TES across all months (see Tables S1 and S2 in the Supplement). By contrast, the variations in individual measurements ( $R^2$ ) are better simulated by the model in July than in January for both OMI and TES. It is not clear why the CTM-OMI correlation is so much higher than that of the CTM-TES on a profile-by-profile basis ( $R^2 = 0.87$  vs.  $0.63$  in July 2005), since STD is similar in both comparisons. In spite of having better resolution in the troposphere, the individual TES TCO have greater measurement noise.

We selected large regions (e.g.,  $25\text{--}60^\circ\text{N}$ , all longitudes) because if  $5 \times 5^\circ$  regions were chosen, the number of CTM-TES matches drops to 16 (see later discussion of Fig. 3). Within these large regions, a high  $R^2$  reflects not only matching the local daily variability, but also the standing latitude-by-longitude patterns in TCO for that month.

Direct comparisons between coincident OMI and TES TCO for July 2005 and January 2006 are shown in Fig. 1e and f. Since OMI measures during the day time, there are only  $\sim 4000$  TES-OMI matching pairs per month. In general the TES-OMI biases are consistent with the difference between the two model-observation biases within the STD. Overall, this comparison is less satisfactory than either model-observation comparison as the PDFs are more dispersed; the STD larger; and the  $R^2$  much smaller (see Table S3 for the results of all months). With  $R^2 < 0.4$ , there is little skill in using one instrument to predict the variance of the other. We conclude that the noise in these two measurements is uncorrelated and larger than the model error.

The differences between TES and OMI TCO are partly attributable to their different retrieval a priori and AK. The influence of different a priori and AK for TCO is expected to be small, since the DOFS in the troposphere are 1–2 for TES (Nassar et al., 2008; Zhang et al., 2010) and about 1 for OMI (de Haan and Veefkind, 2009). To quantify

## OMI, TES, and CTM tropospheric ozone columns

Q. Tang and M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the dependence of TCO on a priori, TES profiles are processed with the OMI a priori by Eq. 2 (denoted by TES<sup>\*</sup>) and the results are shown in Fig. 1g, h. This TES<sup>\*</sup>-OMI comparison is improved compared with the TES-OMI one. The PDF patterns become slightly narrower and denser (Fig. 1g, h vs. Fig. 1e, f), the STD decrease, and the  $R^2$  increase. Further, the TES<sup>\*</sup>-OMI mean biases (−3.6 DU in July 2005 and −1.2 DU in January 2006) are now more predictable by the differences between CTM-OMI and CTM-TES (−3.4 DU and −0.5 DU, respectively).

Parallel results for tropics (25° S–25° N) are illustrated in Fig. 2. The mean CTM TCO bias ranges from 0.0 to +2.3 DU for both measurements and both months (see Fig. 2a–d). The STD are still much greater than the biases, but are generally smaller than in the NH middle latitudes. The model-measurement  $R^2$  is now better for TES than for OMI. The low TCO (<20 DU, found in Western Pacific) is well matched in CTM-TES, but not CTM-OMI because OMI entirely misses the low TCO. Direct TES-OMI comparisons in the tropics (Fig. 2e, f) show much less correlation than for those in the NH middle latitudes (Fig. 1e, f), except that the absolute bias and STD decrease by ~4.5 DU and 0.5 DU in July 2005. After substituting with the OMI a priori, the TES<sup>\*</sup>-OMI comparisons (Fig. 2g, h) are generally improved, but the low TCO values have disappeared. We conclude that the low tropical TCO reported by TES is probably just the a priori from the MOZART CTM calculations, and thus the apparently better agreement for CTM-TES in the tropics is driven by the close agreement between our CTM and the a priori in terms of tropical TCO patterns. In the tropics, direct TES<sup>\*</sup>-OMI comparisons look more like scatter plots ( $R^2 < 0.3$ ); whereas the model-observation comparisons indicate some skill in prediction ( $R^2 = 0.4 \sim 0.7$ ), only part of which can be due to the a priori.

Figure 3 shows the comparisons in monthly mean TCO among CTM, OMI and TES as a function of latitude and longitude on  $5 \times 5^\circ$  grids for July 2005 (first and second columns) and January 2006 (third and fourth columns). Inadvertently, the OMI data are excluded where the surface pressure is less than 700 hPa, causing white spaces at Tibet and Antarctic Plateau. The OMI TCO field (Fig. 3a) is much smoother compared to that of TES (Fig. 3b) in part because of the 800 vs. 16 points per grid cell.

**OMI, TES, and CTM  
tropospheric ozone  
columns**

Q. Tang and M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The uniformly high OMI TCO bands at NH middle latitudes and 30° S (noted in Tang and Prather, 2010) are unrealistic and largely attributable to latitudinal jumps in the OMI a priori profiles. Low TCO values, less than 20 DU, are seen over the equatorial Pacific in previous observations (e.g., Davis et al., 1996; Crawford et al., 1997; Browell et al., 2003), are modeled in most CTMs, and are reported by TES but not OMI, nor by the UCI CTM processed with the OMI operator (Fig. 3e). Although OMI has some sensitivity down to the surface, the inability to report such low TCO likely results from OMI's retrieval fitting algorithm (X. Liu, personal communication, 2011). The CTM means are averaged from coincident simulations for OMI (Fig. 3e) and TES (Fig. 3f). The patterns of corresponding pairs are quite similar, but the modeled TCO show consistent directions of biases relative to both measurements for many regions (see Fig. 3i, j for CTM-OMI and CTM-TES differences), such as underestimations at NH middle latitudes and Southern Hemisphere equatorial Pacific Ocean and overestimations at Africa, Southern Asia, and equatorial Pacific Ocean and South America. These model-measurement biases are consistent across both OMI and TES and suggest model deficiencies in these regions, but may also be a consequence of common features of OMI and TES, such as low sensitivities in the lower troposphere, causing these measurements to miss high O<sub>3</sub> from biomass burning emerging out of South America and Africa.

The zonally averaged OMI-TES biases are small ( $|\mu| < 5$  DU see Figs. 1 and 2), but hide the larger, systematic OMI-TES differences at the continental scale that can exceed  $\pm 10$  DU, over Africa, Pacific Ocean, and tropical Atlantic Ocean (Fig. 3m). When using the same a priori (Fig. 3n), the OMI-TES\* differences are generally reduced by a few DU but show the same large regional biases. The dependence of retrieved TCO on the a priori profile varies spatially, for instance, the OMI-TES comparisons are greatly improved over North and Central Pacific, but remain almost the same at 30° S and South Atlantic. The differences that cannot be explained by a priori are mainly caused by different AK in the OMI and TES retrievals.

Parallel results for January 2006 are shown in the third and fourth columns of Fig. 3. Compared to July 2005, the CTM is still biased high in both cases over South America, but biased in the opposite direction for most NH middle latitudes, indicating a possible pattern of model error. As to the OMI-TES differences, the negative-positive-negative longitudinal pattern in the tropics occurs for both July 2005 and January 2006. On the other hand, the OMI-TES differences change sign for NH middle latitudes and SH middle-high latitudes, reflecting the seasonality of a priori information. Removing the influence from different a priori (see Fig. 3o and p), differences are greatly reduced in extra-tropics of both hemispheres, while discrepancies remain in the tropics.

#### 4 Day-night differences in TES

TES provides both day and night observations of tropospheric  $O_3$  for each orbit. We do not expect, nor model, a significant diurnal cycle in TCO. Nevertheless, there is a possibility that day-night TES observations differ systematically because of the interference of scattered sunlight in the instrument or because of the different ambient conditions on the spacecraft such as temperature on the sunlit part of the orbit. Individual TES level 2 profiles can be separated by day vs. night and averaged into a gridded, monthly mean level 3 product, but different sampling times (e.g., relative to a TF) and low sampling densities (e.g., 8 profiles per  $5 \times 5^\circ$  daytime grid cell) can produce spurious day-night differences that are much larger than the true day-night bias. Using the CTM as a transfer standard between day and night observations provides a unique capability for removing this sampling error and quantifying the day-night bias to a better precision. The day-night comparison on small gridded regions (not shown) proved very noisy because of the limited number of monthly TES observations. Thus in Table 1 we examine the day-night biases and standard deviation in TCO (DU) for CTM minus TES using monthly zonally averaged regions, each containing about  $4\text{--}5 \times 10^3$  TES observations each for day and night. We use the OMI tracks as masks to separate TES profiles into day (with OMI data) vs. night (without OMI data).

## OMI, TES, and CTM tropospheric ozone columns

Q. Tang and M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## OMI, TES, and CTM tropospheric ozone columns

Q. Tang and M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



TES data have large day-night differences in some locations and some months, but these are predicted by the CTM and are due to sampling bias. For example, at NH mid-latitudes in July 2005, the daytime TCO, 47 DU, drops to 43 DU for night. The CTM-TES bias remains essentially unchanged ( $-0.7$  to  $-0.3$  DU) indicating that the 4 DU drop is predicted. The NH mid-latitude day-night difference is +2 DU in January 2006 and is also mostly predicted by the CTM. In the tropics there is no day-night difference in July 2005, but a +2 DU in January 2006. SH mid-latitudes have no day-night difference in both months, probably due to the zonal symmetry of the jet and pollution sources. Using the model, whose bias relative to TES does not change from day to night, we can confidently predict that an systematic day-night bias in TES TCO is less than 0.4 DU (1%) for the cases here. Part of the predictability of the day-night differences here involves different latitudinal sampling of day and night near  $60^\circ$  latitude at the solstices, but the tropical differences in January 2006 are not an obvious sampling issues. Without the CTM as a transfer, some day-night differences approach 10% and it would be difficult to reduce this by going to smaller regions with fewer observations. Another positive feature here is that the standard deviation of the CTM minus TES differences remains virtually unchanged between day and night. We conclude that the noise in individual TCO retrievals is unchanged from day to night. While the noise level of individual TES TCO is modest ( $\sim 10\%$ ) and not unexpected, the stability of the instrument and the day-night biases are remarkable ( $\sim 1\%$ ).

It is possible that the consistency in the TES day-night TCO only reflects a priori information based on the MOZART CTM and would thus be free of day-night bias. Table 2 shows the parallel statistics for the TES a priori day-night TCO using the same sampling as in Table 1. The corrections that TES retrieval adds to its a priori (i.e., Table 1 minus Table 2) vary with locations and months and represent the TES measurement signals. Despite the great reductions in the TCO monthly means (up to 8 DU) and in the STD of CTM minus TES a priori (up to 9 DU), the differences in the day-night CTM-TES bias between TES a priori and TES TCO remain essentially the same ( $\pm 0.1$  DU) or become smaller (from 1.2 DU to 0.4 DU in the tropics in July 2005). In other words, TES

measurement signals do not have day-night bias and can even improve the day-night consistency in our sampling of the TES a priori.

## 5 Discussion and conclusions

In this paper we present a detailed analysis of the NASA Aura level 2 (L2) tropospheric column ozone (TCO) data from OMI and TES. We compare coincident pairs of OMI, TES, and CTM simulations for years 2005–2006 that are time-matched and geo-located OMI on the CTM grid, using all recommended observations between 60° S and 60° N from OMI (~30 000 000) and TES (~65 000). There is generally good agreement in the TCO between the satellite observations and the model as well as between OMI and TES and hence these two datasets can be used to define a TCO climatology with recognition of the weakness in detecting lower tropospheric ozone. As to the monthly zonal means over different latitude bands, either CTM-OMI or CTM-TES biases are less than 6 DU (20%) but usually smaller. A consistent bias identified in the CTM relative to both observations suggests model errors, for example, the CTM is low in July 2005, but high in January 2006 at NH mid-latitudes. Direct comparisons of coincident OMI-TES TCO observations avoids these model uncertainties, yet it yields higher standard deviations implying large single profile noise in each observation. We conclude that the single-profile noise in these two TCO measurements is largely uncorrelated and is larger than the model uncertainties. Thus, there is better predictability between the almost noise-free model and the measurement than between the two measurements. This level of noise, typically 2–4 DU for small  $5 \times 5^\circ$  regions (see Tables S4, S8, S12, S16, S20, S24 for OMI-TES\* STD) makes it difficult to match single measurements involving tropopause folds or stratospheric intrusions of high-O<sub>3</sub> air. In some cases the event is large enough, or sampled multiply (as in the OMI swath data) that it can be detected (Pan et al., 2009). Indeed, we also find regions where a large fraction of the variability in OMI TCO, presumably due to TFs, is matched by the CTM (Tang and Prather, 2010).

## OMI, TES, and CTM tropospheric ozone columns

Q. Tang and M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## OMI, TES, and CTM tropospheric ozone columns

Q. Tang and M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



A common feature of the three comparisons here is that the standard deviations (STD) of the differences (or equivalently the root mean square errors) are much larger than the mean biases for large zonal regions in most months (see Tables S1–S24), and thus there must be some type of noise in the individual coincident comparisons.

This feature is consistent with the OMI and TES validation against ozone sondes (e.g., Nassar et al., 2008; Osterman et al., 2008; Zhang et al., 2010; Kroon et al., 2011). However, there are several exceptions, for example at SH mid-latitudes the CTM-OMI comparisons in May–December 2006 (Table S21) and CTM-TES comparison in November 2006 (Table S22). In these cases the relatively larger mean biases of model-measurement comparisons clearly identify model deficiencies, as for most months the OMI and TES TCO are consistent in terms of the much smaller mean biases than STD of OMI minus TES\* (Table S24).

Comparison of exactly overlapping OMI and TES measurements shows great advantages for cross-validation. Kroon et al. (2011) validated the OMI and TES ozone profiles for cloud-free scenes. We chose not to exclude the cloudy pixels to retain enough comparisons on  $5 \times 5^\circ$  grids. In spite of the fact that a priori information dominates the ozone profiles below clouds, especially for the infrared sounder TES, the monthly mean OMI-TES\* biases in TCO identified in this study are in good agreement with the persistent tropospheric structures illustrated by Kroon et al. (2011), which highlights the importance of correcting the a priori differences when comparing different satellite datasets.

A high bias of about +4 DU for TES TCO has been shown from the validation with sondes for large regions (Osterman et al., 2008), while no equivalent validation for the OMI TCO product is yet available. Our results show that the OMI and TES TCO agree within a few percent and imply that OMI TCO bias is at most a few Dobson Units. A bias of +4 DU is at the upper limit (depending on location) of those which might be inferred from the three-way comparison here of many times more profiles than possible with sonde comparisons. Nevertheless, when restricting the comparisons over smaller regions (i.e.,  $5 \times 5^\circ$  grids), even with corrections for a priori the local OMI-TES differences

are greater than  $\pm 10$  DU at many areas, for example tropics. These large local differences reflect different sensitivities (i.e., averaging kernels) of the two instruments at different locations.

Our unique approach to quantifying the possible day-night differences in the TES observations using all L2 data and coincident CTM profiles has demonstrated no biases and no change in individual measurement noise. This double result is astounding given that there are much larger latitudinal, seasonal, and even day-night differences, but these are generally predicted by the CTM.

**Supplementary material related to this article is available online at:**

**<http://www.atmos-chem-phys-discuss.net/12/16061/2012/acpd-12-16061-2012-supplement.pdf>**

*Acknowledgements.* We thank the Aura science team, particularly the OMI and TES researcher, for insight and assistance in this analysis. This research was supported by NASA's atmospheric chemistry and modeling, analysis, and prediction programs (grants NNG06GB84G and NNX09AJ47G to UCI) and NSF grant (AGS-1042787 to Cornell University).

## References

- Brasseur, G. P., Hauglustaine, D. A., Walters, S., Rasch, R. J., Müller, J., Granier, C., and Tie, X. X.: MOZART, a global chemical transport model for ozone and related chemical tracers. 1. Model description, *J. Geophys. Res.*, 103, 28265–28289, doi:10.1029/98JD02397, 1998. 16064
- Browell, E. V., Fenn, M. A., Butler, C. F., Grant, W. B., Brackett, V. G., Hair, J. W., Avery, M. A., Newell, R. E., Hu, Y., Fuelberg, H. E., Jacob, D. J., Anderson, B. E., Atlas, E. L., Blake, D. R., Brune, W. H., Dibb, J. E., Fried, A., Heikes, B. G., Sachse, G. W., Sandholm, S. T., Singh, H. B., Talbot, R. W., Vay, S. A., Weber, R. J., and Bartlett, K. B.: Large-scale ozone and aerosol distributions, air mass characteristics, and ozone fluxes over the Western Pacific

## OMI, TES, and CTM tropospheric ozone columns

Q. Tang and M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## OMI, TES, and CTM tropospheric ozone columns

Q. Tang and M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ocean in late winter/early spring, *J. Geophys. Res.*, 108, 8805, doi:10.1029/2002JD003290, 2003. 16069

Carver, G., Brown, P., and Wild, O.: The ASAD atmospheric chemistry integration package and chemical reaction database, *Comput. Phys. Commun.*, 105, 197–215, 1997. 16065

5 Crawford, J., Davis, D., Chen, G., Bradshaw, J., Sandholm, S., Kondo, Y., Liu, S., Browell, E., Gregory, G., Anderson, B., Sachse, G., Collins, J., Barrick, J., Blake, D., Talbot, R., and Singh, H.: An assessment of ozone photochemistry in the extratropical Western North Pacific: impact of continental outflow during the late winter/early spring, *J. Geophys. Res.*, 102, 28469–28487, 1997. 16069

10 Davis, D. D., Crawford, J., Chen, G., Chameides, W., Liu, S., Bradshaw, J., Sandholm, S., Sachse, G., Gregory, G., Anderson, B., Barrick, J., Bachmeier, A., Collins, J., Browell, E., Blake, D., Rowland, S., Kondo, Y., Singh, H., Talbot, R., Heikes, B., Merrill, J., Rodriguez, J., and Newell, R. E.: Assessment of ozone photochemistry in the Western North Pacific as inferred from PEM-West A observations during the fall 1991, *J. Geophys. Res.*, 101, 2111–2134, 1996. 16069

15 de Haan, J. F. and Veefkind, J. P.: OMO3PR readme, available at: [http://disc.sci.gsfc.nasa.gov/Aura/data-holdings/OMI/documents/v003/OMO3PRO\\_README.html](http://disc.sci.gsfc.nasa.gov/Aura/data-holdings/OMI/documents/v003/OMO3PRO_README.html), last access: 18 June 2012, 2009. 16065, 16067

20 Eyring, V., Shepherd, T. G., and Waugh, D. W. (Eds.): SPARC Report on the Evaluation of Chemistry-Climate Models, SPARC Report No. 5, WCRP-132, WMO/TD-No. 1526, 2010. 16063

25 Ho, S., Edwards, D. P., Gille, J. C., Luo, M., Osterman, G. B., Kulawik, S. S., and Worden, H.: A global comparison of carbon monoxide profiles and column amounts from tropospheric emission spectrometer (TES) and measurements of pollution in the troposphere (MOPITT), *J. Geophys. Res.*, 114, D21307, doi:10.1029/2009JD012242, 2009. 16064

Holmes, C. D., Tang, Q., and Prather, M. J.: Uncertainties in climate assessment for the case of aviation NO<sub>x</sub>, *P. Natl. Acad. Sci. USA*, 108, 10997–11002, doi:10.1073/pnas.1101458108, 2011. 16063

30 Hsu, J. and Prather, M. J.: Stratospheric variability and tropospheric ozone, *J. Geophys. Res.*, 114, D06102, doi:10.1029/2008JD010942, 2009. 16065

Isaksen, I. S. A., Zerefos, C., Kourtidis, K., Meleti, C., Dalsoren, S. B., Sundet, J. K., Grini, A., Zanis, P., and Balis, D.: Tropospheric ozone changes at unpolluted and

## OMI, TES, and CTM tropospheric ozone columns

Q. Tang and M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



semipolluted regions induced by stratospheric ozone changes, *J. Geophys. Res.*, 110, D02302, doi:10.1029/2004JD004618, 2005. 16065

Kraabøl, A. G., Berntsen, T. K., Sundet, J. K., and Stordal, F.: Impacts of NO<sub>x</sub> emissions from subsonic aircraft in a global three-dimensional chemistry transport model including plume processes, *J. Geophys. Res.*, 107, 4655, doi:10.1029/2001JD001019, 2002. 16065

Kroon, M., de Haan, J. F., Veeffkind, J. P., Froidevaux, L., Wang, R., Kivi, R., and Hakkarainen, J. J.: Validation of operational ozone profiles from the Ozone Monitoring Instrument, *J. Geophys. Res.*, 116, D18305, doi:10.1029/2010JD015100, 2011. 16064, 16073

Luo, M., Rinsland, C., Fisher, B., Sachse, G., Diskin, G., Logan, J., Worden, H., Kulawik, S., Osterman, G., Eldering, A., Herman, R., and Shephard, M.: Comparison of carbon monoxide measurements by TES and MOPITT: influence of a priori data and instrument characteristics on nadir atmospheric species retrievals, *J. Geophys. Res.*, 112, D09303, doi:10.1029/2006JD007663, 2007a. 16064, 16066

Luo, M., Rinsland, C. P., Rodgers, C. D., Logan, J. A., Worden, H., Kulawik, S., Eldering, A., Goldman, A., Shephard, M. W., Gunson, M., and Lampel, M.: TES carbon monoxide validation with DACOM aircraft measurements during INTEX-B 2006, *J. Geophys. Res.*, 112, D24S48, doi:10.1029/2007JD008803, 2007b. 16065

McPeters, R. D., Labow, G. J., and Logan, J. A.: Ozone climatological profiles for satellite retrieval algorithms, *J. Geophys. Res.*, 112, D05308, doi:10.1029/2005JD006823, 2007. 16064

Nassar, R., Logan, J. A., Worden, H. M., Megretskaia, I. A., Bowman, K. W., Osterman, G. B., Thompson, A. M., Tarasick, D. W., Austin, S., Claude, H., Dubey, M. K., Hocking, W. K., Johnson, B. J., Joseph, E., Merrill, J., Morris, G. A., Newchurch, M., Oltmans, S. J., Posny, F., Schmidlin, F. J., Vömel, H., Whiteman, D. N., and Witte, J. C.: Validation of tropospheric emission spectrometer (TES) nadir ozone profiles using ozonesonde measurements, *J. Geophys. Res.*, 113, D15S17, doi:10.1029/2007JD008819, 2008. 16065, 16067, 16073

OMI Team: Ozone monitoring instrument (OMI) data user's guide, OMI-DUG-3.0, available at: [http://disc.sci.gsfc.nasa.gov/Aura/additional/documentation/README.OMI\\_DUG.pdf](http://disc.sci.gsfc.nasa.gov/Aura/additional/documentation/README.OMI_DUG.pdf), last access: 4 August 2010, 2009. 16064

Osterman, G., Bowman, K., Eldering, A., Fisher, B., Herman, R., Jacob, D., Jourdain, L., Kulawik, S., Luo, M., Monarrez, R., Paradise, S., Payne, V., Poosti, S., Richards, N., Rider, D., Shepard, D., Shephard, M., Vilnrotter, F., Worden, H., Worden, J., Yun, H., and Zhang, L.: Earth Observing System (EOS) Tropospheric Emission Spectrometer (TES) Level 2 (L2) Data User's Guide (Up to & including Version 4 data), Tech. Rep. D-38042, Version 4.0, JPL,

## OMI, TES, and CTM tropospheric ozone columns

Q. Tang and M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Jet Propulsion Laboratory California Institute of Technology Pasadena, CA, 91109–8099,  
2009. 16064, 16065

Osterman, G. B., Kulawik, S. S., Worden, H. M., Richards, N. A. D., Fisher, B. M., Eldering, A.,  
Shephard, M. W., Froidevaux, L., Labow, G., Luo, M., Herman, R. L., Bowman, K. W., and  
5 Thompson, A. M.: Validation of tropospheric emission spectrometer (TES) measurements  
of the total, stratospheric, and tropospheric column abundance of ozone, *J. Geophys. Res.*,  
113, D15S16, doi:10.1029/2007JD008801, 2008. 16073

Pan, L. L., Randel, W. J., Gille, J. C., Hall, W. D., Nardi, B., Massie, S., Yudin, V., Khosravi,  
R., Konopka, P., and Tarasick, D.: Tropospheric intrusions associated with the secondary  
10 tropopause, *J. Geophys. Res.*, 114, D10302, doi:10.1029/2008JD011374, 2009. 16072

Park, M., Randel, W. J., Kinnison, D. E., Garcia, R. R., and Choi, W.: Seasonal variation  
of methane, water vapor, and nitrogen oxides near the tropopause: satellite observations  
and model simulations, *J. Geophys. Res.*, 109, D03302, doi:10.1029/2003JD003706, 2004.  
16064

15 Prather, M. J. and Hsu, J.: Coupling of nitrous oxide and methane by global atmospheric chem-  
istry, *Science*, 330, 952–954, doi:10.1126/science.1196285, 2010. 16065

Prather, M. J., Zhu, X., Strahan, S. E., Steenrod, S. D., and Rodriguez, J. M.: Quantifying  
errors in trace species transport modeling, *P. Natl. Acad. Sci. USA*, 105, 19617–19621,  
doi:10.1073/pnas.0806541106, 2008. 16065

20 Prather, M. J., Zhu, X., Tang, Q., Hsu, J., and Neu, J. L.: An atmospheric chemist in search of  
the tropopause, *J. Geophys. Res.*, 116, D04306, doi:10.1029/2010JD014939, 2011. 16065

Rodgers, C. D. and Connor, B. J.: Intercomparison of remote sounding instruments, *J. Geophys.*  
*Res.*, 108, 4116, doi:10.1029/2002JD002299, 2003. 16064, 16066

Schoeberl, M. R., Douglass, A. R., Hilsenrath, E., Bhartia, P. K., Beer, R., Waters, J. W.,  
25 Gunson, M. R., Froidevaux, L., Gille, J. C., Barnett, J. J., Levelt, P. F., and DeCola,  
P.: Overview of the EOS Aura Mission, *IEEE T. Geosci. Remote*, 44, 1066–1074,  
doi:10.1109/TGRS.2005.861950, 2006. 16064

Tang, Q. and Prather, M. J.: Correlating tropospheric column ozone with tropopause folds:  
the Aura-OMI satellite data, *Atmos. Chem. Phys.*, 10, 9681–9688, doi:10.5194/acp-10-9681-  
2010, 2010. 16065, 16069, 16072

30 Tang, Q. and Prather, M. J.: Five blind men and the elephant: what can the NASA Aura ozone  
measurements tell us about stratosphere-troposphere exchange?, *Atmos. Chem. Phys.*, 12,  
2357–2380, doi:10.5194/acp-12-2357-2012, 2012. 16063, 16065

Tang, Q., Prather, M. J., and Hsu, J.: Stratosphere-troposphere exchange ozone flux related to deep convection, *Geophys. Res. Lett.*, 38, L03806, doi:10.1029/2010GL046039, 2011. 16065

5 Worden, H. M., Logan, J. A., Worden, J. R., Beer, R., Bowman, K., Clough, S. A., Eldering, A., Fisher, B. M., Gunson, M. R., Herman, R. L., Kulawik, S. S., Lampel, M. C., Luo, M., Megretskaia, I. A., Osterman, G. B., and Shephard, M. W.: Comparisons of tropospheric emission spectrometer (TES) ozone profiles to ozonesondes: methods and initial results, *J. Geophys. Res.*, 112, D03309, doi:10.1029/2006JD007258, 2007. 16065

10 Zhang, L., Jacob, D. J., Liu, X., Logan, J. A., Chance, K., Eldering, A., and Bojkov, B. R.: Intercomparison methods for satellite measurements of atmospheric composition: application to tropospheric ozone from TES and OMI, *Atmos. Chem. Phys.*, 10, 4725–4739, doi:10.5194/acp-10-4725-2010, 2010. 16064, 16065, 16067, 16073

**OMI, TES, and CTM  
tropospheric ozone  
columns**

Q. Tang and M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## OMI, TES, and CTM tropospheric ozone columns

Q. Tang and M. J. Prather

**Table 1.** Monthly zonal mean TCO (unit: DU) from TES separated by day and night and compared with coincident CTM simulations\*.

Regions	TES	July 2005		January 2006		TES	TES	TES	TES
		Day CTM–TES	Night CTM–TES	Day CTM–TES	Night CTM–TES				
NH mid	47.0	$-0.7 \pm 5.9$	42.9	$-0.3 \pm 5.2$	34.6	$+2.6 \pm 4.5$	32.5	$+3.0 \pm 4.7$	
Tropics	32.6	$+1.8 \pm 4.4$	32.7	$+1.4 \pm 4.2$	32.6	$+2.4 \pm 4.6$	30.6	$+2.7 \pm 4.8$	
SH mid	28.5	$+1.1 \pm 4.2$	27.8	$+1.0 \pm 4.2$	24.6	$+1.2 \pm 3.5$	24.8	$+1.3 \pm 3.3$	

\* Latitude ranges are NH mid (25–60° N), tropics (25° S–25° N), SH mid (60–25° S). See Figs. 1 and 2 for number of points in the sample, about  $4\text{--}5 \times 10^3$  for both day and night in each region. Single numbers are the TES monthly means, while the remaining shows bias  $\pm$  STD.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


## OMI, TES, and CTM tropospheric ozone columns

Q. Tang and M. J. Prather

**Table 2.** Same as Table 1 for TES a priori.

Regions	TES	July 2005		January 2006		TES	TES	TES	TES
		Day CTM–TES	Night CTM–TES	Day CTM–TES	Night CTM–TES				
NH mid	54.7	$-8.3 \pm 13.3$	51.4	$-8.8 \pm 14.1$	40.1	$-3.0 \pm 7.9$	38.2	$-2.7 \pm 7.5$	
Tropics	31.4	$+3.0 \pm 8.4$	32.3	$+1.8 \pm 7.8$	35.6	$-0.5 \pm 5.9$	33.5	$-0.2 \pm 5.6$	
SH mid	29.0	$+0.6 \pm 5.9$	28.6	$+0.2 \pm 5.6$	27.6	$-1.8 \pm 6.2$	28.1	$-1.9 \pm 6.3$	

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## OMI, TES, and CTM tropospheric ozone columns

Q. Tang and M. J. Prather



**Fig. 1.** 2-D TCO PDFs (unit:  $\text{DU}^{-2}$ ) of coincident CTM, OMI, and TES data for NH middle latitudes ( $25\text{--}60^\circ\text{N}$ ) July 2005 (left column) and January 2006 (right column). Comparisons are shown for each matching pair: **(a), (b)** CTM vs. OMI; **(c), (d)** CTM vs. TES; **(e), (f)** TES vs. OMI; **(g), (h)** TES\* vs. OMI. TES\* denotes the TES data processed with OMI a priori. The number of comparisons ( $N$ ), mean bias ( $\mu$ , unit: DU), standard deviation ( $\sigma$ , unit: DU), and coefficient of determination ( $R^2$ ) are shown for each panel.  $R$  is calculated by removing only the means and no slope correction. The black line represents the 1:1 line.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


**OMI, TES, and CTM tropospheric ozone columns**

Q. Tang and M. J. Prather




Fig. 2. Same as Fig. 1 for tropics (25° S–25° N).

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶


Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 3.** Latitude-by-longitude comparisons of monthly mean CTM, OMI, and TES TCO (unit: DU) on  $5 \times 5^\circ$  grid boxes. Results for July 2005 are presented in the first two columns: **(a)** OMI; **(b)** TES; **(e)** CTM coincident with OMI; **(f)** CTM coincident with TES; **(i)** CTM-OMI; **(j)** CTM-TES; **(m)** OMI-TES; **(n)** OMI-TES\*. All the differences are calculated from exact matching pairs on  $5 \times 5^\circ$  grids. Parallel results for January 2006 are shown in the third and fourth columns.

OMI, TES, and CTM  
tropospheric ozone  
columns

Q. Tang and M. J. Prather

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

