

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

# Age of stratospheric air in the ERA-Interim

M. Diallo<sup>1</sup>, B. Legras<sup>1</sup>, and A. Chedin<sup>2</sup>

<sup>1</sup>Laboratoire de Météorologie Dynamique, UMR8539, IPSL, UPMC/ENS/CNRS/Ecole Polytechnique, Paris, France

<sup>2</sup>Laboratoire de Météorologie Dynamique, UMR8539, IPSL, CNRS/Ecole Polytechnique/ENS/UPMC, Palaiseau, France

Received: 14 June 2012 – Accepted: 16 June 2012 – Published: 11 July 2012

Correspondence to: M. Diallo (mdiallo@lmd.ens.fr)

Published by Copernicus Publications on behalf of the European Geosciences Union.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[I◀](#)

[▶I](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



## Abstract

The age of stratospheric air is calculated over 22 yr of the ERA-Interim reanalysis using an off-line Lagrangian transport model and heating rates.

At low and mid-latitudes, the mean age of air is in good agreement with observed ages from aircraft flights, high altitude balloons and satellite observations of CO<sub>2</sub> and SF<sub>6</sub>. The mid-latitude age spectrum in the lower stratosphere exhibits a long tail with a peak at 0.5 yr, which is maximum at the end of the winter, and a secondary flat maximum between 4 and 5 yr due to the combination of fast and slow branches of the Brewer-Dobson circulation and the reinforced barrier effect of the jet. At higher altitudes, the age spectrum exhibits the footprint of the annual modulation of the deep Brewer-Dobson circulation.

The variability of the mean age is analysed through a decomposition in terms of annual cycle, QBO, ENSO and trend. The annual modulation is the dominating signal in the lower stratosphere and in the tropical pipe with amplitude up to one year. The phase of the oscillation is opposite in both hemisphere beyond 20° and is also reversed below and above 25 km with maximum arising in mid-March in the Northern Hemisphere and in mid-September in the Southern Hemisphere. The tropical pipe signal is in phase with the lower southern stratosphere and the mid northern stratosphere. The maximum amplitude of the QBO modulation is of about 0.5 yr and is mostly concentrated within the tropics between 25 and 35 km. It lags the QBO wind at 30 hPa by about 8 months. The ENSO signal is small and limited to the lower northern stratosphere.

The trend is significant and negative, of the order of  $-0.3$  to  $-0.5$  yr dec<sup>-1</sup>, within the lower stratosphere in the Southern Hemisphere and under 40° N in the Northern Hemisphere below 25 km. It is positive (of the order of 0.3 yr dec<sup>-1</sup>) in the mid stratosphere but there is no region of consistent significance. This suggests that the shallow and deep Brewer-Dobson circulations may evolve in opposite directions. It is however difficult to estimate a reliable long-term trend from only 22 yr of data. For instance, a positive trend is found in the lower stratosphere if only the second half of the period

ACPD

12, 17087–17134, 2012

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



is considered in agreement with MIPAS SF<sub>6</sub> data excepted in the northern polar region and at high altitude.

Finally, it is found that the long lasting influence of the Pinatubo eruption can be seen on the age of air from June 1991 until the end of 1993 and can bias the statistics encompassing this period. In our analysis, this eruption shifts the trend towards negative values by about 0.2 to 0.3 yr dec<sup>-1</sup>.

## 1 Introduction

Over the last twenty years, the Brewer-Dobson circulation has been recognized as a major component of the climate system (Andrews et al., 1987; Holton et al., 1995; Salby and Callaghan, 2005, 2006) which affects radiative budgets and atmospheric circulation.

Reanalysed winds from operational weather centres are used to drive Chemistry Transport Models (CTM). It is thus essential that they represent properly the Brewer-Dobson circulation to account for the dependence of the distribution of chemical species in the stratosphere onto the transport properties. It is also important per se to assess the ability of the combined system of a numerical weather forecast model and the associated assimilation system to reproduce the observed behaviour of the stratospheric circulation.

A commonly used metric of the Brewer-Dobson circulation is the age of air defined as the time spent by a particle in the stratosphere since its entry across the tropopause (Li and Waugh, 1999; Waugh and Hall, 2002). As each air parcel is a mixture of particles with different histories and ages, the age of the parcel is an average over these particles (Kida, 1983; Hall and Plumb, 1994). The age can be further averaged over time and space to define a *mean age* over this ensemble or can be described as a distribution denoted as the *age spectrum* (Waugh and Hall, 2002). A main advantage of the age of air is that it can be directly measured from observations of long-lived species (Andrews et al., 2001b; Waugh and Hall, 2002; Stiller et al., 2008; Garcia et al., 2011). The age

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



of air is also used as a mean to compare models (Eyring et al., 2006). The distribution of age in latitude and altitude is convenient to visualize the Brewer-Dobson circulation, its strength and its variability (Li and Waugh, 1999; Waugh and Hall, 2002; Austin and Li, 2006).

Another metric of the Brewer-Dobson circulation is based on the calculation of the residual vertical and meridional velocities (Andrews et al., 1987) which are a representation of the mean zonally averaged mass transport in the stratosphere. This *residual circulation* is used to calculate transit times from the tropopause crossing (see, e.g. Birner and Bonisch, 2011). These transit times, however, are generally not identical to the age of air as this latter is also influenced by the fast stirring and mixing induced by horizontal quasi-isentropic motion in the stratosphere (Waugh and Hall, 2002; Birner and Bonisch, 2011).

The Brewer-Dobson circulation undergoes an annual cycle and changes from year to year. A major mode of variability is the quasi-biennial oscillation (QBO) (Baldwin et al., 2001) which triggers a modulation of vertical transport in the stratosphere by affecting temperature and thus heating rates (Niwano et al., 2003; Punge et al., 2009). An other important factor are volcanic eruptions: in 1991, the Pinatubo has injected massive amount of dust in the stratosphere which have affected its circulation for several years (Thompson and Solomon, 2009). ENSO (Shu et al., 2011) and solar variations are two other sources of Brewer-Dobson variability.

A major source of concern is the existence of a trend in the Brewer-Dobson circulation. Changes in wave propagation and dissipation as well as possible increases in tropospheric wave activity are thought to be the primary driver of a strengthened Brewer-Dobson circulation observed in many models (Butchart and Scaife, 2001; Sigmond et al., 2004; Butchart et al., 2006; Li et al., 2008; Garcia and Randel, 2008). Thompson and Solomon (2005) have observed a cooling of the tropical stratosphere in radiosonde records over the last decades, which is consistent with increased upwelling in the tropical stratosphere.

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Age of stratospheric air in the ERA-Interim**

M. Diallo et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Analysis of tracer data, however, do not provide evidence that such trend is already observed. Engel et al. (2009) and Stiller et al. (2012) even suggest that the age of air might be increasing in some parts of the stratosphere. A possible reason for this discrepancy is that short term trends over one decade are not representative of the trend over one century (Waugh, 2009). Another possibility, however, is that the models or the diagnostics do not fully account of the stratospheric processes. According to Ray et al. (2010) who analyse the observed trends in mean age and ozone under the light of a simple tropical pipe model, “the best quantitative agreement with the observed mean age and ozone trends over the past three decades is found assuming a small strengthening of the mean circulation in the lower stratosphere, a moderate weakening of the mean circulation in the middle and upper stratosphere, and a moderate increase in the horizontal mixing into the tropics”. Similarly, Bonisch et al. (2011) found an increase of the Brewer-Dobson circulation in the lower stratosphere but no change at upper levels.

In this paper, we present the age of stratospheric air over 22 yr from Lagrangian transport calculations based on most recent reanalysed winds and heating rates from the ERA-Interim reanalysis of the European Center for Medium Range Weather Forecast (ECMWF). The age of stratospheric air is investigated using backward deterministic trajectories which are integrated over 10 yr in time to evaluate the residence time in the stratosphere. The method and data used in this study are described in Sect. 2. The mean climatology of the age of air is discussed and compared with observations in Sect. 3. The age variability, the impact of annual cycle and QBO, and the age trend are discussed in Sect. 4. Section 5 offers further discussions and conclusions.

**2 Method and data****2.1 Backward trajectories**

Backward deterministic trajectories are calculated using the Lagrangian model TRACZILLA (Legras et al., 2005) which is a modified version of FLEXPART (Stohl et al.,

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2005) performing direct interpolation from fields provided on hybrid levels. TRACZILLA uses analysed winds to move particles in the horizontal direction. In the vertical direction, it uses either pressure coordinate and Lagrangian pressure tendencies, or potential temperature coordinate and heating rates. In the first case, we denote the trajectories as *kinematic* and in the second case as *diabatic* following a convention established by Eluszkiewicz et al. (2000). Particles are initialised on each level on a longitude-latitude grid with  $2^\circ$  resolution in latitude and an almost uniform spacing in longitude of  $2^\circ/\cos(\phi)$ , where  $\phi$  is the latitude, generating 10 255 particles on each level. For convenience the vertical levels of the initial grid are chosen to be the hybrid levels of the ECMWF models. In order to encompass the whole stratosphere at any latitude, the 30 levels from about 400 hPa (varying according to the surface pressure) to 2 hPa are selected. Trajectories starting below the tropopause are immediately stopped and therefore do not induce any computational cost. Particles starting in the stratosphere are integrated backward in time until they cross the tropopause (see Sect. 2.3). Ensembles of particles have been launched at the end of every months over the 22-yr period 1989–2010 and they are integrated backward for 10 yr.

## 2.2 Data

The wind data and heating rates used in this study have been produced by the ERA-Interim reanalysis of ECMWF (Dee et al., 2011). This reanalysis uses a 12 h 4D-Var assimilation cycle with a T255 partially desaliased horizontal truncature in spherical harmonics and 60 hybrid levels in the vertical from the surface to 0.1 hPa or 66 km. The model has, on the average, 8 levels between 300 and 100 hPa which encompass the extra-tropical lower stratosphere and the tropical tropopause layer and 25 levels above 100 hPa. Wind data are a standard product of the analysis which are available at 6 h interval (00:00 UT, 06:00 UT, 12:00 UT and 18:00 UT). They are completed by wind fields from 3 h and 9 h forecasts at 03:00 UT, 09:00 UT, 15:00 UT and 21:00 UT. Heating rates are obtained as temperature tendencies at 3 h intervals from the twice-daily

assimilation cycles starting at 00:00 UT and 12:00 UT. Hence they are available at 01:30 UT, 04:30 UT, 07:30 UT, 10:30 UT, 13:30 UT, 16:30 UT, 19:30 UT and 22:30 UT.

Several studies have shown that winds from analysis or reanalysis are noisy and induce unrealistic diffusive transport and too fast apparent Brewer-Dobson circulation in the stratosphere (Schoeberl et al., 2003; Meijer et al., 2004; Scheele et al., 2005). This effect is mostly noticed in the vertical direction where velocities are naturally very small. There are two main reasons for this behaviour. The first is the gravity wave noise induced by the assimilation system. Such noise is transient and damped during subsequent evolution and medium-range forecasts exhibit less diffusion than the analysis (Stohl et al., 2005; Legras et al., 2005). This effect is more pronounced in assimilation systems using 3-D-Var assimilation, like in the ERA-40, and is significantly reduced with 4D-Var assimilation, like in the ERA-Interim. The second reason lies in the fact that archived analysis used for off-line transport studies are instantaneous winds sampled typically at 6 h interval. As a result, fast perturbations with time-scale smaller than 6 h are under-sampled. In the limit of very fast uncorrelated perturbations, a sampling with interval  $\tau$  induces a spurious diffusion which is proportional to  $\tau$ . Other reasons might be found in the parameterisations of gravity-wave drag, the representation of convection and the radiative calculations.

The undersampling effect can be reduced by using higher sampling rates at 3 h resolution (Stohl et al., 2005; Legras et al., 2005) or averaging the wind field (Schoeberl et al., 2003; Schoeberl and Dessler, 2011) which both tend to reduce the level of noise. There are indications (Pisso and Legras, 2008) that increasing the sampling rate to 1 h does not improve the noise in the stratosphere with current generation of reanalysis.

There are several reasons for which the vertical motion as represented by heating rates in isentropic coordinates is expected to be less noisy than that represented by vertical velocities in pressure coordinates. Using isentropic coordinates in the vertical separates the fast isentropic motion from slower vertical cross-isentropic motion in the stratosphere and avoids spurious numerical transport effects when particles move with respect to oscillating isobaric surfaces. Another reason is that the heating rates are

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



usually archived as accumulations over finite periods and not as instantaneous values like the velocities. Consequently, the heating rates integrate the noisy fluctuations and are much smoother in time and space than the kinematic velocities.

Data from the ERA-Interim have been used from 1979 to 2010. Since backward trajectory calculations are performed over a duration of 10 yr, the age of air has been estimated monthly over a 22-yr period between 1989 and 2010. For the sake of comparison with the ERA-40 reanalysis and the calculations of Monge-Sanz et al. (2007), some integrations with a perpetual 2000 yr have also been performed.

### 2.3 Metric of Brewer-Dobson circulation

The residence time of a particle in the stratosphere since it has crossed the tropopause is defined as the *age of air* (Waugh and Hall, 2002) and is a common metric of the Brewer-Dobson circulation. As each air parcel results from the mixing of a large number of particles with different trajectories within the stratosphere, the age is actually distributed over a range of values for all the particles contributing to a given parcel. This distribution is denoted as the *age spectrum* which can be mathematically defined (Kida, 1983; Hall and Plumb, 1994; Waugh and Hall, 2002) as generated by a Green function describing the probability that a particle located at the tropopause at time  $t - \tau$  is found within the considered parcel at time  $t$ . The average of  $\tau$  over the distribution is the *mean age*.

The age of air can be measured from trace chemical species, such as sulfur hexafluoride SF<sub>6</sub> and carbon dioxide CO<sub>2</sub> which are well-mixed in the troposphere with a known trend and are nearly passive tracers in the stratosphere (SF<sub>6</sub> is only oxidized in the mesosphere and CO<sub>2</sub> has limited sources through the oxydation of CH<sub>4</sub>) (Andrews et al., 2001a; Waugh and Hall, 2002; Stiller et al., 2008; Garcia et al., 2011). Such quantities have been measured from aircraft and balloons for several decades (Andrews et al., 2001a) and more recently from satellites (Stiller et al., 2008, 2012; Foucher et al., 2011). These data are used in this study in order to compare modeled ages to observations. The comparison to more recent observations of CO<sub>2</sub>

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





stratospheric profiles from ACE-FTS (Foucher et al., 2009, 2011) is postponed to another study. It is in principle possible to derive the full age spectrum if a large density of observations is available (Andrews et al., 2001b) but in practice only the mean age can be retrieved without ad hoc assumptions.

5 In our backward calculations, the age along a given backward trajectory is obtained as the time of first crossing of the tropopause defined by the lower envelop of the surfaces  $\theta = 380\text{K}$  and  $|P| = 2 \times 10^{-6} \text{Kkg}^{-1} \text{m}^2 \text{s}^{-1}$  where  $P$  is the Ertel potential vorticity. The mean age for a given box in latitude and altitude (typically  $2^\circ \times$  model level spacing) and for a given month is calculated as the average in longitude over all particles  
10 falling within this box. Owing to the quasi-uniform spread of the discrete trajectories at the initialisation stage, the average is made over 180 particles at the equator and over 67 particles at  $68^\circ \text{N}$  or  $\text{S}$ . Latitudes closer to the pole are grouped into enlarged latitude bins ( $69^\circ\text{--}73^\circ$ ,  $73^\circ\text{--}77^\circ$ ,  $77^\circ\text{--}81^\circ$ ,  $81^\circ\text{--}90^\circ$ ) to avoid large fluctuations due to the reduced number of particles on a latitude circle near the pole. Further averaging over  
15 time is performed in the sake of improving statistics and reducing noise. These averaging procedures are a simple way to account for mixing in the stratosphere and gather within each box a distribution of particles with different histories.

As observed by Scheele et al. (2005), the number of backward trajectories launched at a given date and remaining within the stratosphere after some delay  $\tau$  decreases exponentially with  $\tau$ . Figure 1 shows that this law is indeed very well satisfied for  $\tau > 3\text{yr}$  with an exponential decrement  $b = 0.2038 \text{yr}^{-1}$  for the mean decay and that the standard deviation from the mean (when each month is considered separately) decays at the same rate. After 10 yr, 88 % of the particles launched within the stratosphere have  
20 left it. We follow Scheele et al. (2005) in using this property to correct the estimated ages for the truncature of trajectory lengths at 10 yr. If we define  $F(\tau)$  as the probability density of the age  $\tau$ , the mean age is  
25

$$\bar{\tau} = \int_0^{\infty} F(\tau) d\tau. \quad (1)$$

**Age of stratospheric air in the ERA-Interim**

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The truncated version of this integral up to  $t_f = 10\text{yr}$  can be calculated explicitly from the trajectory calculations. Assuming that  $F(\tau) = F(t_f)\exp(-b(\tau - t_f))$  beyond this bound, the mean age can be estimated as

$$\bar{\tau} = \int_0^{t_f} F(\tau)d\tau + \frac{F(t_f)}{b}. \quad (2)$$

5 The decrement coefficient  $b$ , shown in Fig. 1 as a time average, has been calculated over the whole stratosphere for each month. It has also been calculated as a 22-yr mean for each latitude and altitude box. The two estimates differ by a standard deviation of 0.008yr. The resulting correction to the mean age varies from zero to about one year at high altitude and latitude. The impact of choosing one definition of the decrement or the other does not change the estimated age by more than 3%. Hence, the correction which is not negligible per se is quite insensitive to the arbitrary details of the calculations.

### 3 Mean climatology of the age of air

#### 3.1 Global distribution of the mean age

15 The mean diabatic age of air, obtained with diabatic trajectories, is calculated as a function of latitude and altitude after averaging over the 22-yr dataset between 1989 and 2010. The left panel of Fig. 2 shows that mean age contours follow the tropopause except in the tropics where they bend up as a result of the tropical upwelling. Gradients of the age of air are concentrated within the extra-tropical lowermost stratosphere with approximately 0.5 yr per km. In the mid extra-tropical stratosphere, the mean age of air varies between 6 and 7.5 yr, with maximum values near the poles, and is older in the Northern Hemisphere above 25 km than in the Southern Hemisphere. The *tropical pipe*

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(Neu and Plumb, 1999), which is the ascending branch of the Brewer-Dobson circulation, is revealed by young air moving upward in the tropics. This tropical pipe is slightly shifted from the equator with a maximum near 5° S. Its relative isolation is visualised by the horizontal gradients on its northern and southern edges.

5 The right panel of Fig. 2 shows the standard deviation of the mean diabatic age calculated using the equivalent sampling size (Bence, 1995; von Storch and Zwiers, 1999) that accounts for the time correlation of monthly ages. This standard deviation shows that the mean diabatic age is estimated with fairly good accuracy within the framework of the ERA-Interim, with patterns that clearly offset the level of fluctuations.  
10 It means also that the variability of the mean age is bound to a limited range of altitudes between 20 and 30 km and, as we shall see below, that the maximum of variance is correlated with the maximum of QBO wind modulation.

In order to understand better the relation between air parcel origins and ages, Fig. 3 shows the distributions of maximum vertical excursion and of altitudes of tropopause crossing. The distributions are shown separately for particles launched below 113 hPa in the extra-tropical lowermost stratosphere (lower row) and those launched at this level and above in the tropics and the extra-tropics (upper row), in the region of the stratosphere denoted as the *overworld* (Holton et al., 1995).

For overworld parcels, Fig. 3 shows that most of the entries to the stratosphere occur through the tropical tropopause between the isentropic levels 370–380 K. The histogram of maximum vertical excursion shows three maxima. The first one below 500 K corresponds to the fast branch of the Brewer-Dobson circulation which is bound to the lower stratosphere. The two other maxima are associated with the tropical pipe. The plume of air rising through the pipe is progressively stripped by detrainment to the mid-latitudes. Most particles reach a maximum value under 1500 K. Above this level, there is very little leakage from the tropical pipe between 1800 K and 2500 K and the third maximum near 2800 K is associated with particles reaching the top of the stratosphere and the mesosphere in the model.  
25

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



For particles initialised in the extra-tropical lowermost stratosphere, Fig. 3 shows that the majority of tropopause crossings still occur between 370 and 380 K in the tropics but a significant proportion of particles enter the stratosphere through the subtropical and extra-tropical tropopause at lower potential temperatures down to 300 K. The maximum vertical excursion is mainly contained within the 300–450 K range with a peak at 380 K. Only a small portion of the particles (about 6%, not shown), have maximum vertical excursion exceeding 700 K.

Hence, the low value and the strong gradient of the mean age above the extra-tropical tropopause are due to the combined effect of isentropic mixing of tropical and extra-tropical across the subtropical tropopause and the fast shallow branch of the Brewer-Dobson circulation (Hoor et al., 2004; Bonisch et al., 2009). Although the deep branch of the Brewer-Dobson is important for the distribution of ages in the stratosphere and for stratospheric chemistry, it processes only a small portion of the air which is eventually found within the lowermost extra-tropical stratosphere (except within the winter polar vortex).

### 3.2 Comparison with observations and models

As a basis for comparison, we use the age of air obtained from in situ aircraft data prior to 1998 reported in Andrews et al. (2001a) and the ages derived from MIPAS retrieval of SF<sub>6</sub> in 2002–2004 (Stiller et al., 2008). Figure 4 shows that these two estimates overlap in the mid-latitudes but the SF<sub>6</sub> ages are older at high latitudes. This is consistent with the impact of photochemical dissociation of SF<sub>6</sub> in the mesosphere which contaminates the stratospheric air within the winter polar vortex (Waugh and Hall, 2002).

Figure 4 shows that the ERA-Interim mean diabatic ages for the period 1989–2010 (red curve) are in good agreement with the aircraft observations except at high latitude. They are generally smaller than the MIPAS SF<sub>6</sub> ages by about 1 yr except in the southern mid-latitudes where the agreement between observations and simulation is the best. This latter region is also, consistently, where the observations are less dispersed. In the tropics the SF<sub>6</sub> MIPAS mean ages is about twice that of the ERA-Interim and

in situ observations of SF<sub>6</sub> and CO<sub>2</sub>. This comparison should be appreciated with the reservation that observed and simulated ages are obtained over periods which overlap but are not identical.

It has been noticed that kinematic and diabatic trajectories produce fairly similar statistics in the ERA-Interim (Liu et al., 2010). It is however visible here that kinematic trajectories tend to produce significantly older ages in the Southern Hemisphere (black curve). These kinematic trajectories have been calculated for two years only, 2007 and 2008, but the diabatic age of air averaged over the same years (blue curve) does not depart significantly from the 22-yr mean. Schoeberl and Dessler (2011) found a similar result with the MERRA reanalysis and attributed the effect to the stronger mixing of tropical and extra-tropical air that favours recirculation of particles within the stratosphere and longer residence times. It is not obvious, however, why mixing should not instead lower the mean age of stratospheric air (Schoeberl et al., 2003). This is indeed observed in a comparison of age of air for kinematic and diabatic trajectories using the ERA-40 reanalysis (not shown). Hence the respective bias introduced in the age of air by kinematic versus diabatic trajectories are variable and not yet fully understood.

We stress that our calculations are all based on full historical records of velocity fields and heating rates over the length of the integration. In a number of previous studies, simulations using perpetual repetition of a given year have been used. This choice leads to considerable fluctuations of the age of air. For instance, the kinematic ages obtained for ERA-Interim based on a 2000 perpetual are significantly older than the 22-yr average (see Fig. 4) while its ERA-40 counterpart provides much too young ages. Large fluctuations, positive or negative, are also observed for diabatic trajectories calculated over perpetual years for 2000 and other years.

When compared with Chemistry-Climate-Model (CCM) estimates (Butchart et al., 2010), our ages based on Lagrangian trajectories are usually older, by about 1 yr in the tropics above 30 km and often 2 yr at mid and high latitudes. The horizontal gradient between the tropical pipe and the mid-latitudes is also stronger. Age is often estimated

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



from an age tracer in a CCM and is subject to a large numerical diffusion which may explain the discrepancy with our Lagrangian calculations.

### 3.3 Vertical profiles of the mean age

A detailed comparison of the vertical profiles of the calculated mean ages with those derived from observations of middle stratosphere balloon flights (Andrews et al., 2001b; Ray et al., 1999) and from SF<sub>6</sub> MIPAS profiles is shown in Fig. 5. In the tropics, the ages from SF<sub>6</sub> MIPAS are higher than those from in situ measurements at all altitudes. The diabatic ages are in good agreement with the in situ measurements and the seasonal dispersion is much smaller than the discrepancy between in situ and satellite data. The mean diabatic age increases almost uniformly in  $z$  from 18 to 34 km at 0.35 yr km<sup>-1</sup>.

In the mid-latitudes, the agreement between in situ and satellite observations is much better except for the SF<sub>6</sub> profiles of Harnisch et al. (1996) above 25 km. The diabatic ages follow the main group of observations, being slightly on the younger side below 20 km, perhaps a sign of excessive mixing across the mid-latitude tropopause. The age increases from 16 to 28 km at a rate of 0.4 yr km<sup>-1</sup> and exhibits only a weak vertical gradient above 28 km, consistently with the findings of Waugh and Hall (2002).

At high latitude during winter, the descent of mesospheric air within the vortex and on its edge induces strong contrast between in vortex and out of vortex air. Because of the limited representation of the mesospheric circulation in the ERA-Interim, it is expected to find a young bias at high altitude and high latitude. This is what is observed above 30 km where mean diabatic ages stay close to out of vortex ages. At lower levels the mean diabatic ages stay within the range of observations. The SF<sub>6</sub> ages are themselves biased within the polar vortex due to the partial photo-chemical dissociation of SF<sub>6</sub> in the mesosphere (Stiller et al., 2012).

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



### 3.4 Age spectrum

As a significant portion of particles remain in the stratosphere with old ages (see Fig. 1), it is important to consider not only the mean age but the age spectrum. Figure 6 shows that there is a clear distinction between the tropical and the extra-tropical spectra at 20 and 25 km. In the tropics at 20 km, the distribution of ages is mono-disperse and compact and decays rapidly to zero for ages above 1 yr indicating that very few particles with old ages return to the tropics from mid-latitudes. In the extra-tropics, the peak is at 0.5 yr, which is small compared to the mean value of about 3.5 yr. The ages exhibit a long tail which extends well to large ages and the curves display in both hemisphere a secondary flat maximum at about 4–5 yr. This distribution of ages corroborates the existence of fast and slow branches of the Brewer circulation (Bonisch et al., 2009, 2011). The fast branch is associated with the particles which have travelled directly and rapidly from the tropics to the mid-latitudes through quasi-isentropic motion (Haynes and Shuckburgh, 2000a; Hoor et al., 2005; Shuckburgh et al., 2009), staying at levels below 450 K. The slow branch corresponds to the deep Brewer-Dobson circulation in which the particles enter the tropical pipe and circulate to high altitudes in the stratosphere. A strong seasonal modulation of the fast branch is observed in the Northern Hemisphere with a younger peak during summer than during winter. The modulation is much smaller in the Southern Hemisphere. This variation is associated with the seasonal modulation of the subtropical jet and the meridional exchanges which is larger in the Northern Hemisphere than in the Southern Hemisphere (Hoor et al., 2005; Shuckburgh et al., 2009).

At higher altitudes (see Fig. 6), the age distribution in both the tropics and the extra-tropics shifts to larger values. The tail of the tropical distribution gets thicker with increasing altitude but remains much less developed than the extra-tropical tail up to 30 km, in agreement with the relative isolation of the tropical pipe (Neu and Plumb, 1999). The peak of the tropical distribution is always about one year younger than the broader main maximum of the extra-tropical distribution. A striking feature is the

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





presence of oscillations in the distributions with an interval of about one year between maxima. Moreover, it is visible in the extra-tropical spectra that these oscillations propagate towards old ages with the seasonal cycle. These oscillations and their propagation are the result of the modulation of the deep Brewer-Dobson circulation by the annual cycle and their relative amplitude depends of the distribution of pathways among stratospheric particles.

#### 4 Variability and trends

The age of air contains an integrated footprint of the variability of the Brewer-Dobson circulation. The first hint on the variability is provided by considering the temporal variation of the number of remaining particles with respect to the mean decay shown in Fig. 1. It is visible (upper left panel of Fig. 7) that the variations are dominated by the annual cycle. The mean annual cycle (lower left panel) exhibits a negative deviation during winter and a positive deviation during summer, implying that more particles cross the tropopause during winter than during summer corresponding to the winter time intensification of the tropical upwelling. This annual cycle propagates through ages and it is useful to notice that the amplitude of the cycle is largest for ages of 2 to 3 yr which indicates the duration over which the exiting particles strongly feel the annual cycle. The amplitude decays for larger values of the ages as remaining particles get uniformly distributed within the stratosphere.

The right panel of Fig. 7 shows the percentage of remaining particles after removal of the mean and annual cycle. Special events that impact transfers at the tropopause are seen as discontinuities in the vertical and variations of the Brewer-Dobson circulation are seen as the oblique patterns.

The most prominent feature is clearly associated with the Pinatubo eruption in June 1991 which induces a reduction of the tropopause crossings and a slowing of the Brewer-Dobson circulation. The impact is strongest on particles with ages of 2 to 3 yr but extends to much older ages and over most of the following decade.

### Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





The second smaller pattern after 2008 is possibly due to the cumulative effects of the eruptions of the Soufrières Hills in May 2006 and Tavurvur in October 2006. As observed by Vernier et al. (2011), the remains of the plumes of these two eruptions have taken about 2 yr to be transported through the depth of the stratosphere (while the Pinatubo plume reached instantly 35 km), which might explain the delay in the stratospheric response.

#### 4.1 Regression method

The temporal evolution of monthly mean diabatic age-of-air at specific altitudes and latitudes (bined as described in Sect. 2.3) has been analysed using a linear response model over the 22 yr of data available from TRACZILLA integrations. This model yields

$$\text{age}(t) = a \cdot t + C(t) + b_1 \cdot \text{qbo}(t - \tau_{\text{qbo}}) + b_2 \cdot \text{enso}(t - \tau_{\text{enso}}) + \varepsilon(t) \quad (3)$$

where qbo is a normalised quasi-biennial oscillation index from CDAS/Reanalysis zonally averaged winds at 30 hPa and enso is the normalised Multivariate El Niño Southern Oscillation Index (MEI) (Wolter and Timlin, 1993, 1998), both provided by NOAA website. The coefficients are a linear trend  $a$ , the annual cycle  $C(t)$  (12 coefficients), the amplitude  $b_1$  and the delay  $\tau_{\text{qbo}}$  associated to QBO and the amplitude  $b_2$  and the delay  $\tau_{\text{enso}}$  associated to ENSO. The constraint applied to determine the 17 parameters  $a$ ,  $b_1$ ,  $b_2$ ,  $\tau_{\text{qbo}}$ ,  $\tau_{\text{enso}}$  and  $C$  is to minimise the residual  $\varepsilon(t)$  in the least square sense. As the combination of amplitude and delay introduces a non linear dependency, there are multiple minima which are sorted out as described below in Sect. 4.3.

Notice that we discard here the influence of Pinatubo as there is no simple index suitable to describe the effect of this single event on the Brewer-Dobson circulation and we do not consider solar forcing as well because our dataset covers only two solar periods.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 4.2 Annual cycle

The annual cycles calculated by the minimising procedure (which takes in account all the factors of variability together) or by a simple monthly composite over the 22 yr turn out to be almost identical.

Figure 8 shows the amplitude and the phase calculated by fitting a pure annual cosine variation to the annual cycle. The phase of the cosine is fixed to be zero in mid-January. The amplitude is maximum in the extra-tropical lowermost stratosphere with peak values of about one year at all latitudes in the Northern Hemisphere and of half a year except at high latitudes in the Southern Hemisphere. The phase is in opposition between the two hemisphere, the maximum being in mid-March for the Northern Hemisphere and mid-September in the Southern Hemisphere. This maximum signal at the end of the winter is consistent with the reinforced barrier effect of the jet and a stronger descent of old air due to the intensification of the deep Brewer-Dobson circulation during winter (Holton et al., 1995; Waugh and Hall, 2002). In turn, younger ages are observed during summer and autumn. Above 25 km, where the amplitude modulation is less than 0.5 yr, the phase is in opposition with the extra-tropical lowermost stratosphere. At these altitudes, the stronger descent during winter favours the replacement of old air by younger air detrained from the tropical pipe. The maximum modulation in the Southern Hemisphere is found at 60° S and might be associated with the strong descent occurring on the polar vortex edge during late winter and spring (Mariotti et al., 2000).

The enhanced penetration of subtropical and tropical air masses in the extra-tropics during summer and autumn, due to a decreased barrier against quasi-horizontal transport and mixing at the subtropical jet, is consistent with previous works on stratosphere-troposphere exchanges using observations of tracer (Hoor et al., 2005; Krebsbach et al., 2006; Sawa et al., 2008) or based on model simulations (Chen, 1995; Haynes and Shuckburgh, 2000b; Sprenger and Wernli, 2003). It is also consistent with our findings on the age spectrum and those of Bonisch et al. (2009).

### Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In the tropics, the annual modulation above the tropopause and in the tropical pipe exceeds rarely 0.5 yr. In the Northern Hemisphere, it is confined below 30 km with a maximum during summer. In the Southern Hemisphere, it is confined between 30 and 40 km with a maximum at the end of the winter.

The seasonal modulation is shown with more details at 55 hPa (about 20 km) in Fig. 9 where the diabatic ages are compared with ages derived from MIPAS SF<sub>6</sub> data (Stiller et al., 2012). It is visible that the Northern Hemisphere modulation is larger in the tropics and in the extra-tropics while the Southern Hemisphere modulation is larger in the subtropics. At high latitudes, both hemispheres exhibit the same amplitude. It is also visible that if SF<sub>6</sub> ages and diabatic ages are similar at mid and high latitudes, their seasonal variation are not related.

### 4.3 Quasi-Biennial-Oscillation and ENSO

Because of the presence of lags in the QBO and ENSO terms in Eq. (3), the problem is non linear and the residual may have multiple minima as a function of the parameters. In order to determine the optimal values of  $\tau_{qbo}$  and  $\tau_{enso}$ , the residual is first minimized at fixed lag and then over the range of lags. This is done in sequence for QBO and ENSO. Figure 10 shows the variations of the QBO amplitude coefficient and the residual amplitude as a function of latitude and lag at several levels in the vertical, roughly corresponding to 20, 25 and 30 km height. In most cases, the minimum residual corresponds also to a maximum of the QBO amplitude coefficient in absolute value and the QBO correlates with the age over more than one period. The optimal lag depends strongly on latitude varying, e.g. by more than a year between 0 and 20° S at 25 km.

Figure 11 shows the amplitudes and lags of the QBO and ENSO contributions to the variability of the age of air. The QBO modulation reaches 0.6 yr with a lag of about 8 months in the tropics near 30 km, with a stronger component in the Northern Hemisphere. This region has already been identified as displaying the largest variability in the age of air in Fig. 2. The influence of the QBO extends upward in the tropical pipe

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper | Discussion Paper

and towards the extratropical stratosphere in both hemispheres with amplitudes of the order of 3 months. The phase lag with respect to the wind at 30 hPa is fairly symmetric with respect of the equator and varies most rapidly with latitude near 25 km. The dependence on ENSO is much less pronounced (less than 0.2 yr) and bound mostly to the lower stratosphere in the Northern Hemisphere.

The standard deviation of the residual in Eq. (3), shown in the right panel of Fig. 11, is larger than the signal explained by QBO and ENSO and peaks in the same regions as the QBO. Hence, the variability not linked to QBO and ENSO dominates the age of air at any location in the stratosphere.

#### 4.4 Trends

As for the annual cycle, the trends calculated by the minimising procedure (which takes in account all the factors of variability together) or by a simple linear fit turn out to be almost identical. The trend as a function of latitude and altitude is shown in Fig. 12 (left panel). The trend is negative within the lower stratosphere with a larger magnitude in the tropics and the Southern Hemisphere than in the Northern Hemisphere. The trend is positive in the extra-tropics above 25 km. In the tropics, the situation is contrasted between the Southern Hemisphere where the trend is negative up to 33 km and the Northern Hemisphere where it is positive above 28 km. The maxima of the trend are located where the amplitude of the QBO modulation is also the largest.

The significance of the trend has been assessed, following von Storch and Zwiers (1999), by performing a Student's t-test among the 264 months of our record using an equivalent number of degrees of freedom calculated as in Zwiers and von Storch (1995) and Bence (1995). This equivalent number ranges between 35 and 80 in the region of maximum negative trend. The right panel of Fig. 12 shows the one-sided  $p$ -value of the Student's test for the hypothesis of a null trend. It is visible that the whole region of large negative trend is highly significant. Although such a simple test is known to overestimate the significance in many cases (Zwiers and von Storch, 1995) and we neglect sub-monthly contribution to the mean age variance, this is an indication that

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the negative trend in the lower stratosphere is a robust feature. On the opposite, we do not find any area of consistent significance above 25 km except perhaps near 60° N and 30 km. In the tropics near 30 km, the  $p$ -value is 0.30.

The observed negative trend in the lower stratosphere is consistent with the finding of an acceleration of the shallow branch of the Brewer-Dobson circulation by Bonisch et al. (2011).

In order to assess better the contributions to the age of air, Fig. 13 shows three illustrative cases. The first one at 20 km and 60° N is chosen where the amplitude of the annual cycle is the largest and indeed dominates the variability of the age of air, the QBO component being much smaller. The second case at 28 km and 4° N is chosen at a location of large amplitude of the QBO component dominating over the annual cycle. The ENSO component makes a negligible contribution in both cases. The trend is negative of  $-0.29 \text{ yr dec}^{-1}$  in the first case and slightly positive of  $0.11 \text{ yr dec}^{-1}$  in the second case. The fluctuations of the age are large in the second case, so that the trend is hardly significant. In the first case, it is visible that the last section of the record, after 2004, would suggest a positive trend while the negative trend over the whole period has a  $p$ -value of 0.02. The third case at 19.7 km and 40° S is located where the trend is the largest of  $-0.56 \text{ yr dec}^{-1}$ . This is obtained in spite of a flattening or a very slight increase over the last period but it is also visible that the period following the Pinatubo eruption exhibits increased ages which contribute significantly to the trend. The amplitude of this effect is tested by recalculating the trend after removing the period mid-1991 to the end of 1994, leading to a reduced value of  $-0.33 \text{ yr dec}^{-1}$ . It is thus clear that the age of air undergoes decadal scale variability, induced by volcanoes and other less known processes, which may affect trends calculated over durations of a few decades. Our own estimate, based on 22 yr of estimated ages is prone to such effect and our statistical test is not made against such decadal variability.

We compare now our calculations to the age of air obtained by Stiller et al. (2012) using SF<sub>6</sub> data from the MIPAS instrument on board ENVISAT over the period (2002–2010) (see Fig. 14). The comparisons are performed at three altitudes 16.5, 20 and

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



25 km and four bands of latitudes (30° S–20° S, 20° S–10° S, 40° N–50° N, 70° N–80° N). There is an old bias of SF<sub>6</sub> ages at 16.5 km at all latitudes and at 20 km in the tropics with respect to our calculations. The tropical bias, which was already noticed on Fig. 4 appears as “robust” in the sense that the dispersion of SF<sub>6</sub> ages is much smaller than the bias. The other panels show a good agreement between our estimations and SF<sub>6</sub> ages. A striking feature is that the age of air tends to flatten or to rise in most locations over the 2002–2010 period, even when a clear negative trend is found over the 22-yr period. This is also in agreement with Stiller et al. (2012) who found a well-spread positive trend over 2002–2010 although we disagree in some locations like at 25 km and 70° N–80° N but this is also where the statistics are noisy and the trends unreliable.

## 5 Conclusions

We have performed direct calculations of the stratospheric age of air using Lagrangian trajectories guided by reanalysed winds and heating rates from the ERA-Interim. This study is based on 32 yr of data and provides estimates of the age over the last 22 yr of the dataset (1989–2010). This study complements previous works on the Brewer-Dobson circulation in the ERA-Interim (Iwasaki et al., 2009; Garny et al., 2011; Seviour et al., 2011) based on residual velocities which did not take into account stirring and mixing.

Our analysis corroborates the significant improvement of the stratospheric circulation in the ERA-Interim compared to the older ERA-40 which has already been noted in other recent works (Monge-Sanz et al., 2007; Fueglistaler et al., 2009b; Dee et al., 2011; Seviour et al., 2011). In contrast with the ERA-40, diabatic heating rates provide younger ages than the kinematic velocity, a feature shared with the MERRA reanalysis (Schoeberl and Dessler, 2011) but which is not yet properly understood.

On the overall, the agreement between ages calculated with the heating rates and observations is very good but for the SF<sub>6</sub> ages in the tropics and near 20 km where our estimates are much younger by almost a factor 2 and fit in situ observations.

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



This discrepancy was already noticed by Stiller et al. (2012). The younger ages and smoother gradients observed in CCMs (Butchart et al., 2010) might be due to limited resolution and excessive numerical diffusion in these models.

The age spectrum corroborates that the lower extra-tropical stratosphere contains young air coming from the mass residual circulation (Birner and Bonisch, 2011) and horizontal mixing, both combined in the lower branch of the Brewer-obson circulation. At higher altitude, the age spectrum reveals peaks spaced by annual intervals that propagate on the spectrum with the seasonal cycle and are associated with the annual modulation of the Brewer-Dobson circulation. These modulations do not necessarily show up or weakly when the sole mean age is considered. While the age spectrum is hardly accessible from observations without strong hypothesis (Andrews et al., 2001b), we suggest that it would be useful to expand its usage within the framework of model inter-comparisons as it contains much more informations on the stratospheric circulation than mean age only.

The variability of mean age as a function of latitude and altitude had been analysed as a linear combination of several contributions: annual cycle, QBO, ENSO and trend. The annual variability dominates in the extra-tropical lower stratosphere below 25 km with higher amplitude, reaching 1.5 yr, in the extra-tropical Northern Hemisphere than in the Southern Hemisphere. The age reaches its maximum during January–February in the Northern Hemisphere and during August–September in the Southern Hemisphere. This is in agreement with an intensified Brewer-Dobson circulation during the winter in each hemisphere which has been described in many studies (e.g. Iwasaki et al., 2009; Garcia et al., 2011; Seviour et al., 2011) but it has little relation with the annual modulation as derived by Stiller et al. (2012). In the tropics, the amplitude of the annual modulation does not exceed six months and is basically in opposition with the modulation of the tropical ascent which reaches its maximum during winter.

The QBO modulation is mostly pronounced within the tropics between 25 and 35 km where its maximum reaches 6 months. The ENSO signal is found to be small and noisy, with a maximum value of about 3 months, and limited to the lower northern

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



stratosphere. This does not contradict the recent finding that warm ENSO events accelerate the upwelling in the lowermost tropical stratosphere (Calvo et al., 2010).

A negative trend of the order of  $-0.2$  to  $-0.4$  yr dec<sup>-1</sup> is found in the lower stratosphere of the Southern Hemisphere and under 40° N in the Northern Hemisphere below 25 km. A positive trend of 0.2 to 0.3 yr dec<sup>-1</sup> is found in the mid extra-tropical stratosphere above 25 km. The negative trend is significant with respect to a test that ignores the decadal variations of the stratospheric circulation. It is, however, influenced by the Pinatubo eruption and reduced when the Pinatubo years are removed. The positive trend is not significant but is consistent with studies based on in situ and satellite observations (Engel et al., 2009; Stiller et al., 2012). The positive trend in our calculation is increased if only the decade 2000–2010 is considered and this is consistent with the strong positive trend found by Stiller et al. (2012) over the same period.

The whole pattern suggests that the shallow and deep branches of the Brewer-Dobson circulation have not evolved recently in the same direction. The trends are consistent with the recent conclusions of Ray et al. (2010), that the increased in the tropical upwelling documented by Randel et al. (2006) is associated with an acceleration of the shallow branch of the Brewer-Dobson circulation (Bonisch et al., 2011) while the deep branch does not change or slightly weakens.

CCMs and studies based on residual circulation all predict an increased Brewer-Dobson circulation in the whole stratosphere of 2 to 3% per decade (McLandress and Shepherd, 2009; Iwasaki et al., 2009; Butchart et al., 2010; Garny et al., 2011; Garcia et al., 2011; Seviour et al., 2011). The discrepancy with our results and those of Engel et al. (2009) and Stiller et al. (2012) might arise from the fact that the age of air does not only depend on the residual circulation but also on the stirring and mixing properties of the stratospheric motion. Owing to their limited resolution, CCMs may not necessarily represent well the mixing properties. A separate diagnostic of the isentropic mixing was performed by Ray et al. (2010) who found a large dispersion between reanalysis but concluded that mixing properties may have varied recently. Such study remains to be done for the ERA-Interim.

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Age of stratospheric air in the ERA-Interim**

M. Diallo et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

It is difficult to estimate a reliable long-term trend from only 22 yr of data. There are evidences that volcanic eruptions, ENSO and other badly represented processes like solar variations may affect the stratospheric circulation over decadal scale. An other factor is ozone depletion to which a significant part of the recent evolution of the Brewer-Dobson circulation has been attributed by Li et al. (2008) while long term changes this century will more likely be caused by global warming.

There are also limitations due to the fact that a reanalysis dataset like the ERA-Interim is based on a single model but on a constantly changing observation systems. For instance it is known that the introduction of AMSU-A data in 1998 (Dee and Upala, 2009) or the introduction of radio-occultation data at the end of 2006 (Poli et al., 2010) had significant impact on the stratospheric temperature. To which extend these changing biases may affect the age of air calculations remains to be determined. It is also worth mentioning that the budget of a reanalysis system, both for heat and momentum, is not closed but is biased by the assimilation increment. Such bias may affect trajectory calculations and residual velocities. It is known, however, that the biases are significantly reduced in the ERA-Interim with respect to previous reanalysis (?).

Finally the understanding of stratospheric circulation is limited by the poor accuracy and sparseness of available measurements of the mean age of air from tracers. More observations and in situ measurements combined with new processing of available data (see, e.g. Foucher et al., 2011) will improve our understanding.

*Acknowledgements.* We thank G. Stiller for providing SF<sub>6</sub> age measurements from MIPAS.



The publication of this article is financed by CNRS-INSU.

## References

- Andrews, A. E., Boering, K. A., Daube, B. C., Wofsy, S. C., Loewenstein, M., Jost, H., Podolske, J. R., Webster, C. R., Herman, R. L., Scott, D. C., Flesch, G. J., Moyer, E. J., Elkins, J. W., Dutton, G. S., Hurst, D. F., Moore, F. L., Ray, E. A., Romashkin, P. A., and Strahan, S. E.: Mean age of stratospheric air derived from in situ observations of CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O, *J. Geophys. Res.*, 106, 32295–32314, doi:10.1029/2001JD000465, 2001a. 17094, 17098, 17124, 17125
- Andrews, A. E., Boering, K. A., Wofsy, S. C., Daube, B. C., Jones, D. B., Alex, S., Loewenstein, M., Podolske, J. R., and Strahan, S. E.: Empirical age spectra for the lower tropical stratosphere from in situ observations of CO<sub>2</sub>: quantitative evidence for a sub-tropical barrier to horizontal transport, *J. Geophys. Res.*, 106, 32295–32314, doi:10.1029/2001JD000465, 2001b. 17089, 17095, 17100, 17109
- Andrews, D. G., Holton, J. R., and Leovy, C. B.: *Middle Atmosphere Dynamics*, vol. 40 of *International Geophysics Series*, Academic Press, San Diego, USA, 1987. 17089, 17090
- Austin, J. and Li, F.: On the relationship between the strength of the Brewer-Dobson circulation and the age of stratospheric air, *Geophys. Res. Lett.*, 33, L17807, doi:10.1029/2006GL026867, 2006. 17090
- Baldwin, M. P., Gray, L. J., Dunkerton, T. J., Hamilton, K., Haynes, P. H., Randel, W. J., Holton, J. R., Alexander, M. J., Hirota, I., Horinouchi, T., Jones, D. B. A., Kinnerson, J. S., Marquardt, C., Sato, K., and Takahashi, M.: The quasi-biennial oscillation, *Rev. Geophys.*, 39, 179–229, doi:10.1029/1999RG000073, 2001. 17090
- Bence, J. R.: Analysis of short time series: correcting for autocorrelation, *Ecology*, 76, 628–639, 1995. 17097, 17106
- Birner, T. and Bönisch, H.: Residual circulation trajectories and transit times into the extratropical lowermost stratosphere, *Atmos. Chem. Phys.*, 11, 817–827, doi:10.5194/acp-11-817-2011, 2011. 17090, 17109
- Boering, K. A., Wofsy, S. C., Daube, B. C., Schneider, H. R., Loewenstein, M., Podolske, J. R., and Conway, T. J.: Stratospheric mean ages and transport rates from observations of carbon dioxide and nitrous oxide, *Science*, 274, 1340–1343, doi:10.1126/science.274.5291.1340, 1996. 17125

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Bönisch, H., Engel, A., Curtius, J., Birner, Th., and Hoor, P.: Quantifying transport into the lowermost stratosphere using simultaneous in-situ measurements of SF<sub>6</sub> and CO<sub>2</sub>, *Atmos. Chem. Phys.*, 9, 5905–5919, doi:10.5194/acp-9-5905-2009, 2009. 17098, 17101, 17104
- Bönisch, H., Engel, A., Birner, Th., Hoor, P., Tarasick, D. W., and Ray, E. A.: On the structural changes in the Brewer-Dobson circulation after 2000, *Atmos. Chem. Phys.*, 11, 3937–3948, doi:10.5194/acp-11-3937-2011, 2011. 17091, 17101, 17107, 17110
- Butchart, N. and Scaife, A. A.: Removal of chlorofluorocarbons by increased mass exchange between the stratosphere and troposphere in a changing climate, *Nature*, 410, 799–802, doi:10.1038/35071047, 2001. 17090
- Butchart, N., Scaife, A. A., Bourqui, M., De Grandpré, J., Hare, S. H. E., Kettleborough, J., Langematz, U., Manzini, E., Sassi, F., Shibata, K., Shindell, D., and Sigmond, M.: Simulation of anthropogenic change in the strength of Brewer-Dobson circulation, *Clim. Dynam.*, 27, 727–741, doi:10.1007/s00382-006-0162-4, 2006. 17090
- Butchart, N., Cionni, I., Eyring, V., Shepherd, T. G., Waugh, D. W., Akiyoshi, H., Austin, J., Brühl, C., Chipperfield, M. P., Cordero, E., Dameris, M., Deckert, R., Dhomse, S., Frith, S. M., Garcia, R. R., Gettelman, A., Giorgetta, M. A., Kinnison, D. E., Li, F., Mancini, E., McLandress, C., Pawson, S., Pitari, G., Plummer, D. A., Rozanov, E., Sassi, F., Scinocca, J. F., Shibata, K., Steil, B., and Tian, W.: Chemistry-Climate Model simulations of twenty-first century stratospheric climate and circulation changes, *J. Climate*, 23, 5349–5374, doi:10.1175/2010JCLI3404.1, 2010. 17099, 17109, 17110
- Calvo, N., Garcia, R. R., Randel, W. J., and Marsh, D. R.: Dynamical mechanism for the increase in tropical upwelling in the lowermost tropical stratosphere during warm ENSO events, *J. Atmos. Sci.*, 67, 2331–2340, doi:10.1175/2010JAS3433.1, 2010. 17110
- Chen, P.: Isentropic cross-tropopause mass exchange in the extratropics, *J. Geophys. Res.*, 100, 16661–16674, doi:10.1029/95JD01264, 1995. 17104
- Dee, D. and Uppala, S.: Variational bias correction of satellite radiance data in the ERA-Interim reanalysis, *Q. J. Roy. Meteorol. Soc.*, 135, 1830–1841, doi:10.1002/qj.493, 2009. 17111
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and performance of the data assimilation system, *Q. J. Roy. Meteor. Soc.*, 137, 553–597, doi:10.1002/qj.828, 2011. 17092, 17108

Elkins, J. W., Gilligan, J. M., Dutton, G. S., Baring, T. J., Volk, C. M., Dunn, R. E., Myers, R. C., Montzka, S. A., Wamsley, P. R., Hayden, A., H., Butler, J. H., Thompson, T. M., Swanson, T. H., Dlugokencky, E. J., Novelli, P. C., Hurst, D. F., Lobert, J. M., Ciciora, S. J., McLaughlin, R. J., Thompson, T. L., Winkler, R. H., Fasser, P. J., Steele, L. P., and Lucarelli, M. P.: Air-borne gas chromatograph for in-situ measurements of long-lived species in upper troposphere and lower stratosphere, *Geophys. Res. Lett.*, 23, 347–350, doi:10.1029/96GL00244, 1996. 17124

Eluszkiewicz, J., Hemler, R. S., Mahlman, J. D., Bruhwiler, L., and Takacs, L. L.: Sensitivity of age-of-air calculations to the choice of advection scheme, *J. Atmos. Sci.*, 57, 3185–3201, doi:10.1175/1520-0469(2000)057<3185:SOAOAC>2.0.CO;2, 2000. 17092

Engel, A., Möbius, T., Bönisch, H., Schmidt, U., Heinz, R., Levin, I., Atlas, E., Aoki, S., Nakazawa, T., Sugawara, S., Moore, F., Hurst, D., Elkins, J., Schauffler, S., Andrews, A., and Boering, K.: Age of stratospheric air unchanged within uncertainties over the past 30 years, *Nature Geosci.*, 2, 28–31, doi:10.1038/ngeo388, 2009. 17091, 17110

Eyring, V., Butchart, N., Waugh, D. W., Akiyoshi, H., Austin, J., Bekki, S., Bodeker, G. E., Boville, B. A., Brühl, C., Chipperfield, M. P., Cordero, E., Dameris, M., Deushi, M., Fioletov, V. E., Frith, S. M., Garcia, R. R., Gettelman, A., Giorgetta, M. A., Grewe, V., Jourdain, L., Kinnison, D. E., Mancini, E., Manzini, E., Marchand, M., Marsh, D. R., Nagashima, T., Newman, P. A., Nielsen, J. E., Pawson, S., Pitari, G., Plummer, D. A., Rozanov, E., Schraner, M., Shepherd, T. G., Shibata, K., Stolarski, R. S., Struthers, H., Tian, W., and Yoshiki, M.: Assessment of temperature, trace species, and ozone in chemistry-climate model simulations of the recent past, *J. Geophys. Res.*, 111, D22308, doi:10.1029/2006JD007327, 2006. 17090

Foucher, P. Y., Chédin, A., Dufour, G., Capelle, V., Boone, C. D., and Bernath, P.: Technical Note: Feasibility of CO<sub>2</sub> profile retrieval from limb viewing solar occultation made by the ACE-FTS instrument, *Atmos. Chem. Phys.*, 9, 2873–2890, doi:10.5194/acp-9-2873-2009, 2009. 17095

Foucher, P. Y., Chédin, A., Armante, R., Boone, C., Crevoisier, C., and Bernath, P.: Carbon dioxide atmospheric vertical profiles retrieved from space observation using ACE-FTS solar occultation instrument, *Atmos. Chem. Phys.*, 11, 2455–2470, doi:10.5194/acp-11-2455-2011, 2011. 17094, 17095, 17111

Fueglistaler, S., Dessler, A. E., Dunkerton, T. J., Folkins, I., Fu, Q., and Mote, P. W.: Tropical tropopause layer, *Rev. Geophys.*, 47, RG1004, doi:10.1029/2008RG000267, 2009a.

**Age of stratospheric air in the ERA-Interim**

M. Diallo et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Fueglistaler, S., Legras, B., Beljaars, A., Morcrette, J.-J., Simmons, A., Tompkins, A. M., and Uppala, S.: The diabatic heat budget of the upper troposphere and lower/mid stratosphere in ECMWF reanalyses, *Q. J. Roy. Meteor. Soc.*, 135, 21–37, doi:10.1002/qj.361, 2009b. 17108
- 5 Garcia, R. R. and Randel, W. J.: Acceleration of Brewer-Dobson circulation due to the increases in greenhouse gases, *J. Atmos. Sci.*, 65, 2731–2739, 2008. 17090
- Garcia, R. R., Randel, W. J., and Kinnison, D. E.: On the determination of age of air trends from atmospheric trace species, *J. Atmos. Sci.*, 68, 139–154, doi:10.1175/2010JAS3527.1, 2011. 17089, 17094, 17109, 17110
- 10 Garny, H., Dameris, M., Randel, W., Bodeker, G. E., and Deckert, R.: Dynamically forced increase of tropical upwelling in the lower stratosphere, *J. Atmos. Sci.*, 68, 1214–1233, doi:10.1175/2011JAS3701.1, 2011. 17108, 17110
- Hall, T. and Plumb, R. A.: Age as a diagnostic of stratospheric transport, *J. Geophys. Res.*, 99, 1059–1070, doi:10.1029/93JD03192, 1994. 17089, 17094
- 15 Harnisch, J., Borchers, R., Fabian, P., and Maiss, M.: Tropospheric trends for CF<sub>4</sub> and C<sub>2</sub>F<sub>6</sub> since 1982 derived from SF<sub>6</sub> dated stratospheric air, *Geophys. Res. Lett.*, 23, 1099–1102, doi:10.1029/96GL01198, 1996. 17100, 17125
- Haynes, P. H. and Shuckburgh, E.: Effective diffusivity as a diagnostic of atmospheric transport 1. Stratosphere, *J. Geophys. Res.*, 105, 22777–22794, doi:10.1029/2000JD900093, 2000a. 17101
- 20 Haynes, P. H. and Shuckburgh, E.: Effective diffusivity as a diagnostic of atmospheric transport 2. Troposphere and lower stratosphere, *J. Geophys. Res.*, 105, 22795–22810, doi:10.1029/2000JD900092, 2000b. 17104
- Holton, J. R., Haynes, P. H., McIntyre, M. E., Douglass, A. R., Rood, R. B., and Pfister, L.: Stratosphere-troposphere exchange, *Rev. Geophys.*, 33, 403–440, doi:10.1029/95RG02097, 1995. 17089, 17097, 17104
- 25 Hoor, P., Gurk, C., Brunner, D., Hegglin, M. I., Wernli, H., and Fischer, H.: Seasonality and extent of extratropical TST derived from in-situ CO measurements during SPURT, *Atmos. Chem. Phys.*, 4, 1427–1442, doi:10.5194/acp-4-1427-2004, 2004. 17098
- Hoor, P., Fischer, H., and Lelieveld, J.: Tropical and extratropical tropospheric air in the lowest stratosphere over Europe: a CO-based budget, *Geophys. Res. Lett.*, 32, L13812, doi:10.1029/2005GL022495, 2005. 17101, 17104
- 30

**Age of stratospheric air in the ERA-Interim**

M. Diallo et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Iwasaki, T., Hamada, H., and Miyazaki, K.: Comparisons of Brewer-Dobson circulations diagnosed from reanalyses, *J Meteorol. Soc. Jpn.*, 87, 997–1006, doi:10.2151/jmsj.87.997, 2009. 17108, 17109, 17110
- Kida, H.: General circulation of air parcels and transport characteristics derived from a hemispheric GCM, Part 2, Very long-term motions of air parcels in the troposphere and stratosphere, *J. Meteorol. Soc. Jpn.*, 61, 510–522, 1983. 17089, 17094
- Krebsbach, M., Schiller, C., Brunner, D., Günther, G., Hegglin, M. I., Mottaghy, D., Riese, M., Spelten, N., and Wernli, H.: Seasonal cycles and variability of O<sub>3</sub> and H<sub>2</sub>O in the UT/LMS during SPURT, *Atmos. Chem. Phys.*, 6, 109–125, doi:10.5194/acp-6-109-2006, 2006. 17104
- Legras, B., Pisso, I., Berthet, G., and Lefèvre, F.: Variability of the Lagrangian turbulent diffusion in the lower stratosphere, *Atmos. Chem. Phys.*, 5, 1605–1622, doi:10.5194/acp-5-1605-2005, 2005. 17091, 17093
- Li, F., Austin, J., and Wilson, J.: The strength of the Brewer Dobson circulation in a changing climate: coupled chemistry climate model simulations, *J. Climate*, 21, 40, doi:10.1175/2007JCLI1663.1, 2008. 17090, 17111
- Li, S. and Waugh, D. W.: Sensitivity of mean age and long-lived tracers to transport parameters in a two-dimensional model, *J. Geophys. Res.*, 104, 30559–30569, doi:10.1029/1999JD900913, 1999. 17089, 17090
- Liu, S., Fueglistaler, S., and Haynes, P. H.: The advection-condensation paradigm for stratospheric water vapour, *J. Geophys. Res.*, 115, D24307, doi:10.1029/2010JD014352, 2010. 17099
- Mariotti, A., Mechoso, C., Legras, B., and Chi, Y.: The evolution of the ozone “collar” in the Antarctic lower stratosphere during early August 1994, *J. Atmos. Sci.*, 57, 402–414, doi:10.1175/1520-0469(2000)057<0402:TEOTOC>2.0.CO;2, 2000. 17104
- McLandress, C. and Shepherd, T. G.: Simulated anthropogenic changes in the Brewer-Dobson circulation, including its extension to high latitudes, *J. Climate*, 22, 1516, doi:10.1175/2008JCLI2679.1, 2009. 17110
- Meijer, E. W., Bregman, B., Segers, A., and van Velthoven, P. F. J.: The influence of data assimilation on the age of air calculated with a global chemistry-transport model using ECMWF wind fields, *Geophys. Res. Lett.*, 31, L23114, doi:10.1029/2004GL021158, 2004. 17093
- Monge-Sanz, B. M., Chipperfield, M. P., Simmons, A. J., and Uppala, S. M.: Mean age of air and transport in a CTM: comparison of different ECMWF analyses, *Geophys. Res. Lett.*, 340, L04801, doi:10.1029/2006GL028515, 2007. 17094, 17108

**Age of stratospheric air in the ERA-Interim**

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Neu, J. L. and Plumb, R. A.: Age of air in a “leaky pipe” model of stratospheric transport, *J. Geophys. Res.*, 104, 19243–19255, doi:10.1029/1999JD900251, 1999. 17097, 17101
- Niwano, M., Yamazaki, K., and Shiotani, M.: Seasonal and QBO variations of ascent rate in the tropical lower stratosphere as inferred from UARS HALOE trace gas data, *J. Geophys. Res.*, 108, 4794, doi:10.1029/2003JD003871, 2003. 17090
- Pisso, I. and Legras, B.: Turbulent vertical diffusivity in the sub-tropical stratosphere, *Atmos. Chem. Phys.*, 8, 697–707, doi:10.5194/acp-8-697-2008, 2008. 17093
- Poli, P., Healy, S. B., and Dee, D. P.: Assimilation of global positioning system radio occultation data in the ECMWF ERA-Interim reanalysis, *Q. J. Roy. Meteor. Soc.*, 136, 1972–1990, doi:10.1002/qj.722, 2010. 17111
- Punge, H. J., Konopka, P., Giorgetta, M. A., and Müller, R.: Effect of the quasi-biennial oscillation on low-latitude transport in the stratosphere derived from trajectory calculations, *J. Geophys. Res.*, 114, D03102, doi:10.1029/2008JD010518, 2009. 17090
- Randel, W. J., Wu, F., Vömel, H., Nedoluha, G. E., and Forster, P.: Decreases in stratospheric water vapor after 2001: links to changes in the tropical tropopause and the Brewer-Dobson circulation, *J. Geophys. Res.*, 111, D12312, doi:10.1029/2005JD006744, 2006. 17110
- Ray, E. A., Moore, F. L., Elkins, J. W., Dutton, G. S., Fahey, D. W., Vomel, H., Oltmans, S. J., and Rosenlof, K. H.: Transport into the Northern Hemisphere lowermost stratosphere revealed by in-situ tracer measurements, *J. Geophys. Res.*, 104, D21304, doi:10.1029/1999JD900323, 1999. 17100, 17125
- Ray, E. A., Moore, F. L., Rosenlof, K. H., Davis, S. M., Boenisch, H., Morgenstern, O., Smale, D., Rozanov, E., Hegglin, M., Pitari, G., Mancini, E., Braesicke, P., Butchart, N., Hardiman, S., Li, F., Shibata, K., and Plummer, D. A.: Evidence for changes in stratospheric transport and mixing over the past three decades based on multiple data sets and tropical leaky pipe analysis, *J. Geophys. Res.*, 115, D21304, doi:10.1029/2010JD014206, 2010. 17091, 17110
- Salby, M. L. and Callaghan, P. F.: Interaction between the Brewer Dobson circulation and the Hadley circulation, *J. Climate*, 18, 4303–4316, doi:10.1175/JCLI3509.1, 2005. 17089
- Salby, M. L. and Callaghan, P. F.: Influence of the Brewer-Dobson circulation on stratosphere-troposphere exchange, *J. Geophys. Res.*, 111, D21106, doi:10.1029/2006JD007051, 2006. 17089
- Sawa, Y., Machida, T., and H., M.: Seasonal variations of CO<sub>2</sub> near the tropopause observation by commercial aircraft, *J. Geophys. Res.*, 113, D23301, doi:10.1029/2008JD010568, 2008. 17104



- Scheele, M. P., Siegmund, P. C., and Velthoven, P. F. J.: Stratospheric age of air computed with trajectories based on various 3D-Var and 4D-Var data sets, *Atmos. Chem. Phys.*, 5, 1–7, doi:10.5194/acp-5-1-2005, 2005. 17093, 17095
- Schoeberl, M. R. and Dessler, A. E.: Dehydration of the stratosphere, *Atmos. Chem. Phys.*, 11, 8433–8446, doi:10.5194/acp-11-8433-2011, 2011. 17093, 17099, 17108
- Schoeberl, M. R., Douglass, A. R., Zhu, Z., and Pawson, S.: A comparison of the lower stratospheric age spectra derived from a general circulation model and two data assimilation systems, *J. Geophys. Res.*, 108, 4113, doi:10.1029/2002JD002652, 2003. 17093, 17099
- Seviour, W. J. M., Butchart, N., and Hardiman, S. C.: The Brewer-Dobson circulation inferred from ERA-Interim, *Q. J. Roy. Meteor. Soc.*, 138, 878–888, doi:10.1002/qj.966, 2011. 17108, 17109, 17110
- Shu, J., Tian, W., Austin, J., Chipperfield, M. P., Xie, F., and Wang, W.: Effects of sea surface temperature and greenhouse gas changes on the transport between the stratosphere and troposphere, *J. Geophys. Res.*, 116, 0148–0227, doi:10.1029/2010JD014520, 2011. 17090
- Shuckburgh, E., D'Ovidio, F., and Legras, B.: Local mixing events in the upper troposphere and lower stratosphere. Part II: Seasonal and interannual variability, *J. Atmos. Sci.*, 66, 3695–3706, doi:10.1175/2009JAS2983.1, 2009. 17101
- Sigmond, M. P., Siegmund, P. C., Manzini, E., and Kelder, H.: A simulation of the separate climat effects of middle atmospheric and tropospheric CO<sub>2</sub> doubling, *J. Climate*, 17, 2352–2367, doi:10.1175/1520-0442(2004)017<2352:ASOTSC>2.0.CO;2, 2004. 17090
- Sprenger, M. and Wernli, H.: A northern hemispheric climatology of cross-tropopause exchange for the ERA15 time period (1979–1993), *J. Geophys. Res.*, 108, 8521, doi:10.1029/2002JD002636, 2003. 17104
- Stiller, G. P., von Clarmann, T., Höpfner, M., Glatthor, N., Grabowski, U., Kellmann, S., Kleinert, A., Linden, A., Milz, M., Reddmann, T., Steck, T., Fischer, H., Funke, B., López-Puertas, M., and Engel, A.: Global distribution of mean age of stratospheric air from MIPAS SF<sub>6</sub> measurements, *Atmos. Chem. Phys.*, 8, 677–695, doi:10.5194/acp-8-677-2008, 2008. 17089, 17094, 17098, 17124
- Stiller, G. P., von Clarmann, T., Haenel, F., Funke, B., Glatthor, N., Grabowski, U., Kellmann, S., Kiefer, M., Linden, A., Lossow, S., and López-Puertas, M.: Observed temporal evolution of global mean age of stratospheric air for the 2002 to 2010 period, *Atmos. Chem. Phys.*, 12, 3311–3331, doi:10.5194/acp-12-3311-2012, 2012. 17091, 17094, 17100, 17105, 17107, 17108, 17109, 17110, 17125

**Age of stratospheric air in the ERA-Interim**

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





- Stohl, A., Forster, C., Frank, A., Seibert, P., and Wotawa, G.: Technical note: The Lagrangian particle dispersion model FLEXPART version 6.2, *Atmos. Chem. Phys.*, 5, 2461–2474, doi:10.5194/acp-5-2461-2005, 2005. 17091, 17093
- Thompson, D. W. J. and Solomon, S.: Recent stratospheric climat trends as evidence in radiosonde data: Global structure and tropospheric linkages., *J. Climate*, 18, 4785–4795, doi:10.1175/JCLI3585.1, 2005. 17090
- Thompson, D. W. J. and Solomon, S.: Understanding recent stratospheric climate change, *J. Climate*, 22, 1934–1943, doi:10.1175/2008JCLI2482.1, 2009. 17090
- Uppala, S., Kallberg, P., Simmons, A., Andrae, U., da Costa Bechtold, V., Fiorino, M., Gibson, J., Haseler, J., Hernandez, A., Kelly, G., Li, X., Onogi, K., Saarinen, S., Sokka, N., Allan, R., Andersson, E., Arpe, K., Balmaseda, M., Beljaars, A., van de Berg, L., Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M., Fisher, M., Fuentes, M., Hagemann, S., Holm, E., Hoskins, B., Isaksen, L., Janssen, P., Jenne, R., McNally, A., Mahfouf, J.-F., Morcrette, J.-J., Rayner, N., Saunders, R., Simon, P., Sterl, A., Trenberth, K., Untch, A., Vasiljevic, D., Viterbo, P., and Woollen, J.: The ERA-40 re-analysis, *Q. J. Roy. Meteor. Soc.*, 131, 2961–3012, doi:10.1256/qj.04.176, 2005.
- Vernier, J. P., Thomason, L. W., Pommereau, J. P., Bourassa, A., Pelon, J., Garnier, A., Hauchecorne, A., Trepte, C., Degenstein, D., and Vargas, F.: Major influence of tropical volcanic eruptions on the stratospheric aerosol layer during the last decade, *Geophys. Res. Lett.*, 38, L12807, doi:10.1029/2011GL047563, 2011. 17103
- von Storch, H. and Zwiers, F. W.: *Statistical Analysis in Climate Research*, Cambridge Univ. Press, 1999. 17097, 17106
- Waugh, D.: Atmospheric dynamics: the age of stratospheric air, *Nature Geosci.*, 2, 14–16, doi:10.1038/ngeo397, 2009. 17091
- Waugh, D. and Hall, T.: Age of stratospheric air: theory, observations, and models, *Rev. Geophys.*, 40, 1010, doi:10.1029/2000RG000101, 2002. 17089, 17090, 17094, 17098, 17100, 17104, 17124
- Wolter, K. and Timlin, M. S.: Monitoring ENSO in COADS with a seasonally adjusted principal component index, in: *Proc. of the 17th Climate Diagnostics Workshop*, Oklahoma Clim. Survey, NOAA/NMC/CAC, NSSL, CIMMS and the School of Meteor., Univ. of Oklahoma, Norman, OK, 52–57, 1993. 17103
- Wolter, K. and Timlin, M. S.: Measuring the strength of ENSO – how does 1997/98 rank?, *Weather*, 53, 315–324, 1998. 17103

**Age of stratospheric air in the ERA-Interim**

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Age of stratospheric air in the ERA-Interim

M. Diallo et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





**Fig. 1.** Mean percentage of remaining air parcels in the stratosphere as a function of elapsed time  $\tau$  (age). Blue: mean over the 264 months of the 22-yr range. Green and red: mean  $\pm$  one standard deviation. Basis 100 at time 0. Dotted: fit of the mean by the exponential decay law  $93.48 \exp(-0.2038\tau)$  where  $\tau$  is in yr.

**Age of stratospheric air in the ERA-Interim**

M. Diallo et al.

Title Page

Abstract	Introduction
Conclusions	References
Tables	Figures

◀
▶

◀
▶

Back	Close
------	-------

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Age of stratospheric air in the ERA-Interim**

M. Diallo et al.




**Fig. 2.** (Left): distribution of mean age of stratospheric air (in yr) averaged over 22 yr (264 months) from 10 km to 42 km as a function of latitude and altitude. (Right): variance of the mean age.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.



**Fig. 3.** Distribution (in potential temperature) of maximum vertical excursion (left) and altitude of tropopause crossings (center) for the populations of particles launched in the extratropical lowermost stratosphere (bottom) and in the overworld (top). The separation is made at the pressure level 113 hPa (level 26 of the ECMWF grid). The right panels show the joint histograms of vertical excursions and tropopause crossings. In the bottom panels the particles with vertical excursion larger than 700 K (6 % of the total) are not shown. The histograms are calculated over the ensemble of particles launched during the 264 months that were located in the stratosphere at day 1 of the backward trajectory.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Age of stratospheric air in the ERA-Interim

M. Diallo et al.



**Fig. 4.** Mean age of stratospheric air at 55 hPa ( $\sim 20$  km) as a function of latitude, using ERA-Interim data unless specified. (Red): ages from diabatic trajectories averaged over 22 yr (1989–2010). (Blue): ages from kinematic trajectories averaged over 2 yr (2007–2008). (Black): ages from diabatic trajectories averaged over 2 yr (2007–2008). (Green): ages from kinematic trajectories and a perpetual run based on the year 2000 only. (Cyan): ages from kinematic trajectories and a perpetual run based on the year 2000 only with ERA-40 winds. (Yellow shaded area): envelop of ages from SF<sub>6</sub> MIPAS observations during November 2002–February 2003 and November 2003–February 2004 from Stiller et al. (2008). (Triangles): ages from airborne observations of SF<sub>6</sub> (Elkins et al., 1996; Waugh and Hall, 2002). (Dots and error bars): compilation of ages from airborne observations of CO<sub>2</sub> until 1998 by Andrews et al. (2001a). Notice that the statistical uncertainty is shown for the mean age (red curve) as  $\pm$  one standard deviation (cyan shaded area) but this uncertainty is small enough to be hidden by the thickness of the curve.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Age of stratospheric air in the ERA-Interim

M. Diallo et al.



**Fig. 5.** (Dashed curves): calculated vertical profiles of mean age (red: all year, blue: winter, cyan: summer). (Curves with circles): mean age profiles from SF<sub>6</sub> MIPAS (Stiller et al., 2012) (magenta: all year, black: winter, green: summer). (Symbols): in situ measurements of CO<sub>2</sub> (diamonds) (Boering et al., 1996; Andrews et al., 2001a), SF<sub>6</sub> (triangle) (Ray et al., 1999) and whole air samples of SF<sub>6</sub> (square outside vortex and asterisk inside vortex) (Harnisch et al., 1996).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Age of stratospheric air in the ERA-Interim

M. Diallo et al.



**Fig. 6.** Age spectra calculated with diabatic trajectories (corresponding to red curve in Fig. 4) at three altitudes and over three latitude bands. The altitudes are 20 km (upper left), 25 km (upper right), 31 km (lower left) and 35 km (lower right). Each panels shows three latitude ranges: 35° N–65° N (top), 15° S–15° N (mid) and 65° S–35° S (bottom). (Red): winter (DJF) age spectrum averaged over 1989–2010. (Blue): spring (MAM) age spectrum. (Green): summer (JJA) age spectrum. (Black): autumn (SON) age spectrum.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion





## Age of stratospheric air in the ERA-Interim

M. Diallo et al.




**Fig. 7.** Upper left: percentage of remaining parcels as a function of time and age. Lower left: annual cycle in the percentage shown over two cycles. Right: percentage of remaining parcels after removal of the mean and annual cycle for each age.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Age of stratospheric air in the ERA-Interim**

M. Diallo et al.




**Fig. 8.** Altitude-latitude cross-sections of amplitudes (left) and phases (right) of the annual variation derived from least squares method applied on the mean age of air between 85° S and 85° N in latitude and within 10–42 km height.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.




**Fig. 9.** Amplitude of the seasonal cycle of the mean age over 22 yr at 20 km height in the inner tropics (upper left: 0–10° S and N), in the subtropics (lower left: 20–30° S and N), in the mid-latitudes (lower right: 40–50° N and S) and the polar regions (upper right: 60–70° N and S) compared with the age derived from SF<sub>6</sub> MIPAS data. (Red and blue): Southern Hemisphere for model and SF<sub>6</sub> ages, respectively. (Black and green): same for Northern Hemisphere

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


## Age of stratospheric air in the ERA-Interim

M. Diallo et al.



**Fig. 10.** Time-latitude cross-sections of the amplitude variation of the QBO component (upper) and residual (lower) at altitudes 20 km, 26 km and 33 km. The magenta curves show the calculated lag as a function of latitude. Each panel has its own scale.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Age of stratospheric air in the ERA-Interim

M. Diallo et al.



**Fig. 11.** (Left): altitude-latitude cross-sections of the amplitude of the age variations attributed to the QBO (upper) and its phase with respect to wind at 30 hPa (lower). (Mid): same for the amplitude and delay of the age variability correlated with ENSO. (Righ): standard deviation of the residual in Eq. (3).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Age of stratospheric air in the ERA-Interim**

M. Diallo et al.




**Fig. 12.** (Left): altitude-latitude cross-sections of the trend derived from linear regression over the 22 yr of available data. (Right):  $p$ -value of its associated statistical test of significance where the  $p$ -value is less than 0.05.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Age of stratospheric air in the ERA-Interim

M. Diallo et al.



**Fig. 13.** Temporal evolution of the different components of mean age at several heights and latitudes. (Upper): 20 km and 60° N, (mid): 28 km and 4° N, (lower): 20 km and 40° S. (Black): mean age over 22-yr period. (Green): residual of the least squares method. (Magenta): amplitude of the QBO component. (Blue): amplitude of the ENSO component. (Cyan): annual cycle of the mean age. (Red): trend of the mean age derived from the least square method. (Dash blue): trend of the mean age derived from the least squares method after removed the Pinatubo years.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Age of stratospheric air in the ERA-Interim

M. Diallo et al.



**Fig. 14.** Evolution of mean age compared with SF<sub>6</sub> ages at several heights and latitudes. (Three first rows of left column): southern subtropics (30° S to 20° S) at 16.5, 20 and 25.5 km. (Last two rows of left column): inner tropics (20° S to 10° S) at 20 and 25.5 km. (Two first rows of right column): northern mid-latitudes (40° N to 50° N) at 17 and 20 km. (Three last rows of right column): northern polar region (71° S to 79° N) at 16.5, 20 and 25.5 km. (Black): mean age. (Cyan): annual cycle of the mean age. (Red): trend of the mean age derived from the least squares method. (Green diamond): SF<sub>6</sub> MIPAS data. (Dash blue): trend of the mean age derived from the least squares method over 2000–2010.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

