

1 **Supplement for:**

2

3 **The mixing state of carbonaceous aerosol particles in northern and southern**
4 **California measured during CARES and CalNex 2010**

5

6 John F. Cahill¹, Kaitlyn Suski¹, John H. Seinfeld², Rahul A. Zaveri³, Kimberly A. Prather^{1,4}.

7 *¹Dept. of Chemistry and Biochemistry, University of California San Diego, CA*

8 *²Division of Chemistry and Chemical Engineering, California Institute of Technology, Pasadena,*
9 *CA*

10 *³Atmpsheric Sciences & Global Division, Pacific Northwest National Laboratory, Richland,*
11 *WA*

12 *⁴Scripps Institute of Oceanography, University of California San Diego, La Jolla, CA*

13

14

15

16 **Corresponding author: kprather@ucsd.edu, 858-822-5312. Fax: 858-534-7042*

17

18

19

20

21

22

23 *A-ATOFMS data processing*

24 High sensitivity of the A-ATOFMS detectors occasionally led to the acquisition of gas phase
25 species ionized by a laser pulse. These signals were occasionally counted as particles, and were
26 removed from analysis by retroactively raising the peak area threshold above the gas phase
27 baseline. During CalNex sampling inlet pressures were changed after 10 May 2010, hence the
28 transmission efficiency of the aerodynamic lens was altered. Therefore, a direct comparison of
29 particle number between early flights and later flights is not possible. However fractional
30 compositions of particles can still be compared as there was no noticeable change in size
31 distribution with the differing inlet pressure as indicated by a high correlation of size
32 distributions before and after inlet pressure change ($R^2 = 0.97$).

33

34 *Extended particle classifications*

35 Particle classifications were established based upon characteristic peaks identified in
36 previous lab studies. The dominant carbonaceous particle types are explained in the main text.
37 Vanadium mixed with OC (V-OC), high mass OC (HMOC), amine (AM), biological (BIO), dust
38 (D), and sea salt (SS) represented 2.78, 0.86, 0.56, 0.30, 0.50, 2.80% of total aerosol measured
39 by the A-ATOFMS, respectively. Vanadium mixed with OC (V-OC) emitted from the
40 combustion of ship fuels composed ~3% of particles measured by the A-ATOFMS ([Ault et al.,](#)
41 [2009](#)). This particle type has intense peaks at $^{51}\text{V}^+$ and $^{67}\text{VO}^+$ as well as OC peaks at
42 m/z $^{27}\text{C}_2\text{H}_3^+/\text{CHN}^+$, $^{29}\text{C}_2\text{H}_5^+$, $^{37}\text{C}_3\text{H}^+$, $^{39}\text{C}_3\text{H}_3^+/\text{K}^+$, and $^{43}\text{C}_2\text{H}_3\text{O}^+/\text{CHNO}^+$ ([Ault et al., 2009](#)).
43 HMOC consists of OC peaks at m/z $^{27}\text{C}_2\text{H}_3^+/\text{CHN}^+$, $^{37}\text{C}_3\text{H}^+$, $^{39}\text{C}_3\text{H}_3^+/\text{K}^+$ as well as many intense
44 peaks >100 m/z . These types likely represent polycyclic aromatic hydrocarbons or other
45 oligomers formed through cooking processes ([Silva and Prather, 2000](#)). Occasionally this

46 particle type contained peaks similar to OS that may lead to an overestimation of OS number
47 fractions, especially during CalNex where HMOC was more prevalent (1.42% compared to
48 0.16% for CalNex and CARES, respectively). However the number fractions of HMOC are
49 significantly smaller than the observed number fraction of OS (28 and 35% for CalNex and
50 CARES, respectively); hence overestimation of OS number fractions is likely small. Amines are
51 OC particles that contain an intense peak at m/z $^{56}\text{C}_2\text{HNO}^+$, $^{59}\text{C}_3\text{H}_9\text{N}^+$, $^{86}(\text{C}_2\text{H}_5)_2\text{NCH}_2^+$,
52 and/or $^{118}(\text{C}_2\text{H}_5)_3\text{NOH}^+$ and originate from agricultural processes, animal husbandry, or
53 photochemical processing ([Angelino et al., 2001](#); [Pratt and Prather, 2010](#); [Sorooshian et al., 2008](#)).
54 Biological particles contain an intense $^{40}\text{Ca}^+$, $^{56}\text{CaO}^+$, and $^{96}\text{Ca}_2\text{O}^+$ with OC
55 ($^{27}\text{C}_2\text{H}_3^+/\text{CHN}^+$, $^{37}\text{C}_3\text{H}^+$, $^{39}\text{C}_3\text{H}_3^+/\text{K}^+$), soot ($^{12}\text{C}^+$, $^{24}\text{C}_2^+$, $^{48}\text{C}_3^+$), and phosphate ($^{79}\text{PO}_3^-$) peaks
56 ([Fergenson et al., 2004](#); [Pratt and Prather, 2010](#); [Russell, 2009](#)). Dusts contained a wide variety of
57 metals (Na, K, Ti, Ca, and Fe), as well as phosphate ($^{79}\text{PO}_3^-$) and silicate ($^{44}\text{SiO}^-$, $^{60}\text{SiO}_2^-$, and
58 $^{103}\text{Si}_2\text{O}_3^-$). Sea salt is characterized by an intense sodium peak ($^{23}\text{Na}^+$) and chlorine peaks ($^{35}\text{Cl}^-$
59 and $^{37}\text{Cl}^-$) as well as clusters of the two ($^{81}\text{Na}_2\text{Cl}^+$) ([Gard et al., 1998](#); [Pratt and Prather,](#)
60 [2010](#); [Silva and Prather, 2000](#)). Often SS was aged significantly, containing significant nitrate,
61 sulfate, and OC peaks.

62

63 *Temporal changes within CARES*

64 A distinct change in chemistry, particle concentrations, and meteorological variables was
65 seen during the CARES study; hence the study was split into two periods, 2 June 2010 – 19 June
66 2010 (NoCal-1) and 21 June 2010 – 28 June 2010 (NoCal-2) as can be seen in SI Table 1 and SI
67 Figure 1. An increase in A-ATOFMS Soot-OC and BB fractions was seen during NoCal-2,
68 which coincided with a general increase in temperature and particulate matter > 2.5 μm in the

69 region (SI Figure 1). More detail on the differences in chemistry can be found in the main text.
70 It is hypothesized that higher SO₂ and NO_x concentrations during NoCal-2 lead to the growth of
71 soot. The broadening in A-ATOFMS size distributions for NoCal-2 signifies the growth of
72 particles, supporting this hypothesis (SI Figure 2). Unlike OC:soot ratios (Figure 9),
73 sulfate:nitrate ratio distributions remained relatively unchanged between NoCal-1 and NoCal-2
74 (SI Figure 3); hence sulfate was still the most common secondary species present on particles in
75 northern California.

76

77 *The absence of negative ion spectra*

78 Negative ion spectra were absent in 13% of particles in California. This has previously been
79 attributed to significant amounts of water present on the particle which inhibits the formation of
80 negative ions ([Neubauer et al., 1998, 1997](#)). However, due to the low average relative humidity
81 (RH) during the studies, 49±30% and 39±14%, for CalNex and CARES respectively, and typical
82 deliquescent RH thresholds of >60% ([Neubauer et al., 1998](#)), it is unlikely that there was
83 significant water present on the particles to justify the lack of negative spectra. Similar
84 conclusions were deduced from modeling of the CARES study ([Fast et al., 2012](#)). Further,
85 spectra with only positive ions were less frequent during CalNex (4%) than CARES (24%)
86 despite the higher RH during CalNex. Temporal comparisons of positive only spectra with RH
87 do not indicate any correlation between the two. Further, significantly higher fractions of
88 particles contain negative ion spectra during NoCal-1, 94%, compared to NoCal-2, 62%. This is
89 despite the higher RH of 41±15% compared to 36±12% for NoCal-1 and NoCal-2, respectively.
90 It is hypothesized that for these studies the acquisition of negative ion spectra was dependent on
91 the presence of secondary species, like sulfate or nitrate, rather than the amount of water present.

92

93

References

94

95 Angelino, S., Suess, D. T., and Prather, K. A.: Formation of aerosol particles from reactions of
96 secondary and tertiary alkylamines: Characterization by aerosol time-of-flight mass
97 spectrometry, *Environmental science & technology*, 35, 3130-3138, 2001.

98 Ault, A. P., Moore, M. J., Furutani, H., and Prather, K. A.: Impact of Emissions from the Los
99 Angeles Port Region on San Diego Air Quality during Regional Transport Events,

100 *Environmental science & technology*, 43, 3500-3506, 10.1021/es8018918, 2009.

101 Fast, J. D., Gustafson Jr., W. I., Berg, L. K., Shaw, W. J., Pekour, M., Shrivastava, M., Barnard,

102 J. C., Ferrare, R. A., Hostetler, C. A., Hair, J. A., Erickson, M., Jobson, B. T., Flowers, B.,

103 Dubey, M. K., Springston, S., Pierce, R. B., Dolislager, L., Pederson, J., and Zaveri, R. A.:

104 Transport and mixing patterns over Central California during the carbonaceous aerosol and
105 radiative effects study (CARES), *ACPD*, 2012.

106 Fergenson, D. P., Pitesky, M. E., Tobias, H. J., Steele, P. T., Czerwieniec, G. A., Russell, S. C.,

107 Lebrilla, C. B., Horn, J. M., Coffee, K. R., Srivastava, A., Pillai, S. P., Shih, M. T. P., Hall, H. L.,

108 Ramponi, A. J., Chang, J. T., Langlois, R. G., Estacio, P. L., Hadley, R. T., Frank, M., and Gard,

109 E. E.: Reagentless detection and classification of individual bioaerosol particles in seconds,

110 *Analytical chemistry*, 76, 373-378, Doi 10.1021/Ac034467e, 2004.

111 Gard, E. E., Kleeman, M. J., Gross, D. S., Hughes, L. S., Allen, J. O., Morrical, B. D.,

112 Fergenson, D. P., Dienes, T., Galli, M. E., Johnson, R. J., Cass, G. R., and Prather, K. A.: Direct

113 observation of heterogeneous chemistry in the atmosphere, *Science*, 279, 1184-1187, 1998.

114 Neubauer, K. R., Johnston, M. V., and Wexler, A. S.: On-line analysis of aqueous aerosols by
115 laser desorption ionization, *International Journal of Mass Spectrometry and Ion Processes*, 163,
116 29-37, 1997.

117 Neubauer, K. R., Johnston, M. V., and Wexler, A. S.: Humidity effects on the mass spectra of
118 single aerosol particles, *Atmospheric Environment*, 32, 2521-2529, 1998.

119 Pratt, K. A., and Prather, K. A.: Aircraft measurements of vertical profiles of aerosol mixing
120 states, *Journal of Geophysical Research*, 115, 10.1029/2009jd013150, 2010.

121 Russell, S. C.: Microorganism characterization by single particle mass spectrometry, *Mass*
122 *spectrometry reviews*, 28, 376-387, 10.1002/mas.20198, 2009.

123 Silva, P. J., and Prather, K. A.: Interpretation of Mass Spectra from Organic Compounds in
124 Aerosol Time-of-Flight Mass Spectrometry, *Analytical chemistry*, 72, 3553-3562,
125 10.1021/ac9910132, 2000.

126 Sorooshian, A., Murphy, S. N., Hersey, S., Gates, H., Padro, L. T., Nenes, A., Brechtel, F. J.,
127 Jonsson, H., Flagan, R. C., and Seinfeld, J. H.: Comprehensive airborne characterization of
128 aerosol from a major bovine source, *Atmos Chem Phys*, 8, 5489-5520, 2008.

129

130

131

Campaign	Flight Name ¹ (yyyymmdd)	Temperature (°C)	RH (%)	UF-CPC (#/ccm)*10 ⁴	CPC (#/ccm)*10 ³	PCASP/UHSAS ² (#/ccm)*10 ³
Calnex	20100504a	20.8 ± 2.6	39 ± 12	1.5 ± 1.0	11.6 ± 5.0	1.5 ± 0.4
	20100505a	18.8 ± 2.4	51 ± 33	1.0 ± 0.5	N/A	5.8 ± 5.1
	20100506a	17.0 ± 1.9	58 ± 54	1.2 ± 0.6	10.6 ± 4.5	1.4 ± 0.4
	20100507a	21.6 ± 2.3	35 ± 22	1.4 ± 0.6	11.4 ± 4.3	3.3 ± 3.9
	20100510a	13.7 ± 1.3	61 ± 37	1.3 ± 0.7	11.0 ± 4.9	0.7 ± 0.3
	20100512a	18.8 ± 3.2	35 ± 15	1.4 ± 0.8	11.2 ± 5.3	2.9 ± 3.8
	20100513a	21.3 ± 3.9	31 ± 14	1.1 ± 0.7	8.2 ± 4.1	1.0 ± 0.6
	20100514a	16.6 ± 2.3	64 ± 10	1.5 ± 0.8	12.1 ± 5.2	1.3 ± 0.4
	20100515a	19.3 ± 3.2	56 ± 29	1.2 ± 0.5	10.8 ± 4.0	1.6 ± 0.4
All Flights	18.6 ± 3.6	48 ± 31	1.3 ± 0.7	10.9 ± 4.8	2.2 ± 2.9	
CARES	20100603a	20.2 ± 4.0	60 ± 7	2.2 ± 1.9	18.1 ± 12.5	N/A
	20100606a	23.5 ± 2.1	55 ± 5	2.2 ± 1.8	18.0 ± 11.4	N/A
	20100606b	24.4 ± 6.7	41 ± 9	1.3 ± 1.0	10.8 ± 6.4	N/A
	20100608a	20.1 ± 2.0	56 ± 10	2.0 ± 2.0	1.0 ± 0.8	N/A
	20100608b	20.5 ± 4.6	40 ± 14	1.3 ± 1.3	0.9 ± 0.7	N/A
	20100610a	17.4 ± 3.8	38 ± 8	1.7 ± 1.0	1.0 ± 0.3	1.1 ± 0.6
	20100612a	20.9 ± 2.7	28 ± 2	1.2 ± 1.8	0.5 ± 0.7	0.9 ± 0.2
	20100612b	25.1 ± 2.5	25 ± 4	1.4 ± 1.0	0.7 ± 0.3	1.5 ± 0.8
	20100614a	23.8 ± 2.9	32 ± 9	2.8 ± 2.4	1.3 ± 1.0	2.1 ± 1.9
	20100615a	17.3 ± 1.9	55 ± 12	2.2 ± 1.9	1.1 ± 1.5	1.6 ± 1.9
	20100615b	20.7 ± 5.5	42 ± 9	1.5 ± 0.9	1.1 ± 0.6	2.6 ± 0.8
	20100618a	21.5 ± 5.7	25 ± 12	2.2 ± 1.6	1.1 ± 0.8	2.3 ± 1.9
	20100619a	18.5 ± 3.6	39 ± 9	2.0 ± 1.0	1.5 ± 0.7	1.9 ± 1.0
	20100621a	18.8 ± 1.7	43 ± 7	1.9 ± 2.2	1.1 ± 1.2	1.9 ± 1.1
	20100621b	25.2 ± 6.6	21 ± 7	1.1 ± 0.9	0.8 ± 0.5	1.8 ± 1.1
	20100623a	19.6 ± 4.0	40 ± 10	0.8 ± 1.2	0.6 ± 1.0	3.2 ± 1.0
	20100623b	25.0 ± 6.8	30 ± 8	1.3 ± 0.7	0.9 ± 0.5	3.9 ± 1.8
	20100624a	19.1 ± 2.1	44 ± 15	2.1 ± 2.1	1.0 ± 1.0	1.2 ± 0.8
	20100624b	22.1 ± 3.8	37 ± 8	2.1 ± 1.6	2.4 ± 3.1	2.3 ± 1.0
	20100627a	25.6 ± 2.1	41 ± 10	0.6 ± 0.9	0.5 ± 0.6	3.4 ± 3.1
	20100628a	28.2 ± 2.6	38 ± 7	1.0 ± 1.0	0.8 ± 0.8	3.9 ± 1.9
20100628b	35.5 ± 5.7	25 ± 4	0.8 ± 0.5	0.6 ± 0.3	3.5 ± 1.6	
All Flights	22.4 ± 5.7	39 ± 14	1.6 ± 1.6	2.7 ± 6.1	2.2 ± 1.7	
NoCal-1	21.0 ± 4.7	41 ± 14	1.9 ± 1.6	3.9 ± 7.6	1.8 ± 1.4	
NoCal-2	24.3 ± 6.5	36 ± 1	1.3 ± 1.5	1.0 ± 1.3	2.7 ± 1.9	

¹ Flight names labeled "a" occurred in the morning, while those labeled with a "b" were in the afternoon

² PCASP was used during CalNex while the UHSAS was used during CARES

132

133 SI Table 1: Mean (± std dev) meteorological data and particle concentrations over all of CalNex,

134


CARES, NoCal-1, and NoCal-2.

135

136

137

138



139

140

141

142

143

144

145

146

147

148

149

150

SI Figure 1: A-ATOFMS relative fractions of particle types and average PM2.5 mass


concentrations for each flight during the CARES study. Flight labels indicate the date of the

flight and if it was in the morning (a) or afternoon (b). A change in chemistry and a general

increase in PM2.5 mass were observed after 6/19/10.

151

152




153

154 SI Figure 2: A-ATOFMS measured size distributions for NoCal-1 (red) and NoCal-2 (blue).

155 The broadening of the size distribution indicates the growth of particles in NoCal-2.

156



158

159 SI Figure 3: Fraction of particles containing sulfate and nitrate with RPA > 0.5% in NoCal-1

160 (red) and NoCal-2 (blue, left panel). Sulfate:nitrate peak ratios are shown in (right panel).

161 Values < 0 indicate more soot than nitrate and values > 0 indicate more sulfate than soot. Ratios

162 representing 1:1, 2:1, and 10:1 are shown by solid, dotted, and dashed lines respectively.

163