

This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

# Relationship between level of neutral buoyancy and dual-Doppler observed mass detrainment levels in deep convection

G. L. Mullendore<sup>1</sup>, A. J. Homann<sup>2</sup>, S. T. Jorgenson<sup>1</sup>, and T. J. Lang<sup>3</sup>

<sup>1</sup>Department of Atmospheric Sciences, University of North Dakota, Grand Forks, North Dakota, USA

<sup>2</sup>National Weather Service, Indianapolis, IN, USA

<sup>3</sup>Department of Atmospheric Science, Colorado State University, Fort Collins, Colorado, USA

Received: 12 July 2012 – Accepted: 6 August 2012 – Published: 21 August 2012

Correspondence to: G. L. Mullendore (gretchen@atmos.und.edu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

ACPD

12, 21267–21290, 2012

**Relationship  
between LNB and  
LMD**

G. L. Mullendore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## Abstract

Although it is generally accepted that the level of neutral buoyancy (LNB) is only a coarse estimate of updraft depth, the LNB is still used to understand and predict storm structure in both observations and modeling. This study uses case studies to quantify the variability associated with using environmental soundings to predict detrainment levels. Nine dual-Doppler convective cases were used to determine the observed level of maximum detrainment (LMD) to compare with the LNB. The LNB for each case was calculated with a variety of methods and with a variety of sources (including both observed and simulated soundings). The most representative LNB was chosen as the proximity sounding from NARR using the most unstable parcel and including ice processes.

The observed cases were a mix of storm morphologies, including both supercell and multicell storms. As expected, the LMD was generally below the LNB, the mean offset for all cases being 2.2 km. However, there was a marked difference between the supercell and non-supercell cases. The two supercell cases had LMDs of 0.3 km and 0.0 km below the LNB. The remaining cases had LMDs that ranged from 4.0 km below to 1.6 km below the LNB, with a mean offset of 2.8 km below. Observations also showed that evolution of the LMD over the lifetime of the storm can be significant (e.g., >2 km altitude change in 30 min), and this time evolution is lacking from models with coarse time steps, missing significant changes in detrainment levels that may strongly impact the amount of boundary layer mass transported to the upper troposphere and lower stratosphere.

## 1 Introduction

The most efficient method for transporting heat, moisture, and chemical tracers from the boundary layer to the upper troposphere and lower stratosphere is through moist convection (e.g., Dickerson, 1987; Pickering et al., 1988; Mullendore et al., 2005). Deep

ACPD

12, 21267–21290, 2012

### Relationship between LNB and LMD

G. L. Mullendore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



convection allows constituents from the boundary layer to be transported within minutes into the upper troposphere and lower stratosphere, where longer chemical residence times can increase their impact on the radiative budget, and higher wind speeds can increase their influence region from local to regional or even global (e.g., Dickerson, 1987; Pickering et al., 1993; Thompson et al., 1997).

For atmospheric chemistry models to correctly simulate the upper troposphere and lower stratosphere and their interactions, the level at which mass is being detrained from convection must be correctly simulated. These models must be initialized with accurate data and methodologies that provide reliable detrainment level estimations. In large-scale models where convection is not explicitly resolved, parameterizations of convective transport are utilized. These convective transport parameterizations use the level of neutral buoyancy (LNB), as calculated by parcel theory, as an upper bound of significant transport, assuming entrainment of environmental air will cause most parcels to be detrained below the LNB. Parcel theory struggles to accurately predict detrainment levels given the number of upper-air observations that are available and the lack of detail about storm morphology in the estimations. The purpose of this study is to attempt to quantify the variability associated with predicting detrainment levels.

In parcel theory, an initial parcel is chosen to best represent the environment being studied. The parcel is then lifted to obtain estimates of various environmental indices, such as convective available potential energy (CAPE) and LNB. The LNB (also known as equilibrium level, EL) is calculated from an environmental sounding by lifting a representative parcel of air from the surface to the level at which a parcel is no longer positively buoyant. Although simple to use, parcel theory can provide significantly varying estimations based on the choice of the initial parcel and the fact that the parcel chosen to be lifted is unlikely to be completely independent from the environment (Manzato and Morgan, 2003).

More specifically, using parcel theory to determine detrainment heights is highly idealized, and several of the problems inherent in the idealization are listed here. One, using a single column to represent the environment of a mesoscale system ignores the

## Relationship between LNB and LMD

G. L. Mullendore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



high spatial and temporal variability at cloud-scale, and choosing and/or measuring the “best” column is difficult. Two, parcel theory assumes that there is no entrainment of environmental air into the updraft, which one would expect to reduce buoyancy and, thus, detrainment heights. Approximating the amount of entrained air in a storm is important to accurately determine the level of detrainment, which is dependent on the thermodynamic profile, storm classification, and storm size (Cohen, 2000; Mullendore et al., 2005). Three, parcel theory neglects the dynamically induced vertical perturbation pressure gradients (e.g., Doswell and Markowski, 2004), producing an inaccurate picture of parcel accelerations and therefore inaccurate detrainment height estimates. Four, parcel theory assumes that a representative convective updraft can be defined with a single parcel, when in fact parcels entering the storm have a range of starting locations and properties. Existing convective parameterizations attempt to address some of the above challenges (e.g., parcel variability and entrainment estimates), but do not yet account for variability in updraft characteristics due to storm classification, such as possible updraft enhancements due to vertical pressure gradients in supercell storms (e.g., Arakawa, 2004; Randall et al., 2003).

In summary, to understand the potential for deep convective transport in certain storm regimes, we need to understand the relationship between the LNB and the observed level of maximum detrainment (LMD, Mullendore et al., 2009). Environmental soundings, from both observations and simulations, are analyzed and compared to one another to understand the near-storm environment before and during nine case studies and to estimate the LNB for various parcel theory assumptions. Vertical velocities derived from dual-Doppler analysis of each case study are then used to calculate the LMD in order to investigate the relationship between LNB and LMD. An overview of the field campaigns and associated observational platforms utilized in this study is presented in Sect. 2. Sections 3 and 4 describe the methodology and results for calculations of LNB and LMD, respectively. Section 5 summarizes the LNB and LMD results, and conclusions are presented in Sect. 6.

## Relationship between LNB and LMD

G. L. Mullendore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2 Data

Dual-Doppler radar data from nine midlatitude storms, representing several storm types, were utilized to calculate observed detrainment heights. Six cases were observed during the Severe Thunderstorm Electrification and Precipitation Study (STEPS, Lang et al., 2004). Three additional cases were observed as they passed through the dual-Doppler lobes of the Colorado State University (CSU) CHILL-Pawnee radar network. In addition, estimated idealized detrainment levels were calculated by utilizing both simulated and observed soundings (STEPS) or simulated soundings only (CHILL-Pawnee).

### 2.1 Radar

The STEPS campaign took place from 17 May to 20 July 2000 in eastern Colorado and western Kansas. Among the main observational platforms used in the campaign, three radars were used to create a triple-Doppler network (Fig. 1): the Colorado State University dual-polarization Doppler radar (CSU-CHILL) located near Burlington, Colorado, the NCAR S-band dual-polarization Doppler radar (S-Pol) located near Idalia, Colorado, and the National Weather Service (NWS) Weather Surveillance Radar-1988 Doppler (WSR-88D) at Goodland, Kansas (KGLD). The CHILL/Pawnee cases utilized CSU-CHILL and a second S-band weather radar located at the CSU Pawnee site. Note that all the raw data quality control measures, interpolation, and dual-Doppler processing of radar data described herein were performed prior to this study; for more information on the processing done see Lang and Rutledge (2002, 2008); Tessendorf et al. (2005, 2007b,a); Wiens (2005). All altitudes for the CHILL/Pawnee cases were processed as above ground level (a.g.l.). The STEPS cases were converted to a.g.l. from mean sea level (MSL) by subtracting the altitude of the KGLD radar (1.1 km). All cases had both reflectivity and velocities mapped to a regular Cartesian grid with 0.5 km resolution in the vertical, and either 0.5 or 1.0 km resolution in the horizontal.

ACPD

12, 21267–21290, 2012

## Relationship between LNB and LMD

G. L. Mullendore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 2.2 Mobile soundings

NCAR mobile soundings systems (M-GLASS) were deployed during STEPS to characterize the storm environment (Lang et al., 2004). The times (UTC) and locations of the mobile soundings collected for the 29 June 2000 case are shown in Fig. 1 as triangles.

Some soundings from the campaign were not used due to missing upper-level data (e.g., the 29 June 2000, 23:55 UTC sounding recorded no data above 300 hPa). Bad data points, e.g., anomalous spike in the 29 June 2000, 20:22 UTC sounding, and superadiabatic layers near the surface in some mobile soundings, were removed from the sounding data before calculating parcel ascents. Mobile sounding altitudes are given as a.g.l.

## 2.3 NARR soundings

As previously discussed, traditional environmental soundings are limited in time and space and therefore do not always provide reliable information about pre-storm environments. One source of information that is useful for supplementing environmental soundings after a storm has occurred is the North American Regional Reanalysis (NARR) dataset (Mesinger et al., 2006). The reanalysis data have a horizontal grid spacing of 32-km, 29 layers in the vertical, and 3-hourly output. To obtain a sounding representative of the pre-storm environment using NARR data, criterion was followed for proximity soundings of using a point 30 min or 40 km upwind and in the path of the storm (Thompson et al., 2003). For this study, a point was chosen along the eastern or southern edge of the reflectivity region, depending on storm propagation direction, of a given storm at 00:00 UTC (except for the 21 July 1998 CHILL case, which used the time 21:00 UTC). The NARR grid point closest to this location was calculated, and then the proximity NARR sounding was chosen as the location one or two points further east or south (resulting in a distance from storm of 33 to 63 km). This proximity sounding was also labeled as “most representative”, and additional simulated soundings were read from the grid points surrounding the most representative location (Fig. 1, plus signs), to

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sample the local variability. All NARR soundings were converted to a.g.l. by subtracting out the surface altitude of the chosen grid point and interpolated to 250 m vertical grid spacing using cubic interpolation. Throughout the remainder of this article, all altitudes are given as a.g.l.

## 2.4 Case overview

Table 1 lists the nine cases that were used for this study. The cases studied included single cell (unicell) and multicell storms, supercells, and convective lines. Information on the STEPS cases can be found in the mission summaries from the STEPS campaign archive (UCAR, 2012a). In addition, many of these cases have been described in detail in previous studies (Lang and Rutledge, 2002; Lang et al., 2004; Tessendorf et al., 2005; Kuhlman et al., 2006; Lang and Rutledge, 2008). Classification of these storms was based on all sources cited herein; the “convective line” designation was chosen to be general, describing a system with significant areas of both convective and stratiform structures, with the convective elements being at least quasi-linear. Note that while the 11 June 2000 convective line case did not have an observed embedded supercell, a relatively large and persistent single cell was observed at the southern end of the line.

## 3 LNB calculations

It is generally accepted that level of neutral buoyancy (LNB) derived from environmental profiles is highly variable, and sensitive to several parameters including choice of initial parcel and method of calculation (e.g., irreversible or reversible ascent). With the goal of quantifying the variability, several calculations were performed for each observed and simulated sounding. Three different initial parcels were used in each case – the surface-based parcel (surface air and dewpoint temperatures), the most unstable parcel (air and dewpoint temperatures at the height where the equivalent potential

### Relationship between LNB and LMD

G. L. Mullendore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



temperature,  $\Theta_e$ , was the greatest in the lowest 500 hPa), and the mixed-layer parcel (average of the lowest 1500 m). Calculations were performed using all three initial parcels assuming both liquid water processes only and also incorporating both liquid and ice processes. Note that calculations were also done assuming all condensed water fell out of the parcel (irreversible, or pseudoadiabatic ascent) and again under the assumption that no condensed water fell out of the parcel (reversible ascent). However, the results incorporating reversible ascent were not significantly different from the irreversible calculations. Therefore, only the irreversible results are shown throughout the remainder of this study.

Table 2 shows the calculated LNB for multiple methodologies for the 29 June 2000 STEPS case. Parcels with no buoyancy, or insignificant buoyancy (LNB <5 km), were labeled as “NA” in the table, and were not used in calculation of LNB statistics (Table 1). For parcels with significant buoyancy, LNB ranged from 8.5 to 12.3 km, showing significant variability due to methodology and data sources. Focusing on differences due to methodology alone, choice of surface parcel caused differences up to 1.6 km for calculated LNB (Table 2, 17:57 UTC), and often was the difference between a buoyant ascent and no buoyancy at all (e.g., 17:27 UTC). For a given initial parcel type, inclusion of ice processes had a relatively small impact on the calculated LNB (<0.5 km). However, inclusion of ice is more physically accurate, so should be included in the absence of computational constraints. The differences between LNB estimations obtained using just one sounding emphasize the amount of variability in estimating the LNB from environmental soundings.

Variability in LNB is also seen due to spatial and temporal variability. As seen in results for a single methodology (Table 2; most unstable parcel with ice), calculated LNB ranges from 10.9 to 12.3 km, representing soundings taken over a 6.5 h time period and covering a distance of 236 km. Based on time and position, the most representative of the mobile soundings is the 20:22 UTC sounding (the 23:38 UTC sounding was likely contaminated by convection occurring to the north of the main storm). Comparing the most representative mobile sounding to the most representative NARR sounding

## Relationship between LNB and LMD

G. L. Mullendore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





shows that the LNB calculated from NARR is over 1 km higher than the LNB calculated from the mobile sounding. This positive bias in the NARR soundings was seen in all the STEPS cases, with the offset between the most representative NARR sounding and the average mobile soundings ranging from 0.2 km (SPEPS 6/19) to 2.1 km (SPEPS 6/11, 6/23).

Table 1 shows the variability in LNB for all cases analyzed. The most representative LNB (Table 1, column 3) is the most unstable parcel calculated with ice processes from the central NARR sounding. Columns 4 and 5 show the mean value and standard deviation for a particular case using all data methodologies and all soundings. As is expected, the mean LNB is lower in all cases than the most representative, as the most unstable parcel with ice processes is most likely to produce the highest LNB values. The variability seen in the 29 June 2000 case is typical for all cases, as seen in the standard deviation values.

In summary, determining a single LNB value for a given storm is difficult, as the LNB is highly variable, dependent on the choice of lifted parcel, method of calculation, source of sounding (observed versus simulated), local spatial and temporal variability, and storm-relative location. In any storm, the likelihood of having a single representative sounding valid for the lifecycle of the entire storm is low. However, this case study is a “best case scenario”, as the temporal and spatial resolution is significantly greater in a field campaign with special soundings than cases that rely only on regular weather station soundings. Although the LNB is widely recognized as only an approximation to the actual level at which parcels will detrain from the updraft, it remains an important index for understanding convective dynamics and predicting the depths of convective mixing. Therefore, systematic testing of the relationship between LNB and observed storm detrainment height is important.

## Relationship between LNB and LMD

G. L. Mullendore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## 4 LMD calculations

To compare with LNB calculations, an estimate of the actual detrainment heights is needed. Aircraft chemical retrievals are not available for the cases presented, and even when available, in situ measurements present the challenge of being only point measurements of the complex three-dimensional, time-variable storm. Chemical concentrations from satellite retrievals are not able to resolve both the horizontal and vertical cloud-scale transports (i.e. horizontal scales of order 1 km and vertical scales of order 100 m). Vertical velocities derived from dual-Doppler radar retrievals can be used to estimate the level at which mass was detrained in the observed storm (Mullendore et al., 2009). At the level where the vertical momentum decelerates most sharply, parcels will accumulate due to a maximum in vertical convergence. Assuming local changes in density are negligible *relative to the mass flux*, this means that the same amount of mass must be exiting the volume horizontally. Therefore, the level at which a maximum in vertical convergence occurs is expected to be the same level at which horizontal divergence is maximized. Throughout this study, the level of maximum horizontal divergence will be considered to be synonymous with the level of maximum detrainment (LMD), i.e. the location at which the transfer of air from an updraft is maximized.

The vertical divergence profile ( $\partial\rho w/\partial z$ , Fig. 2) was calculated for each vertical column and then horizontally integrated over the entire dual-Doppler domain at each time. Vertical divergence was calculated by taking a first order forward difference along each vertical column. This calculation included an estimate of the variability of density ( $\rho$ ) with height by incorporating the basic equation

$$\rho = \rho_0 \exp(z/H),$$

where  $\rho_0$  is  $1.22 \text{ kg m}^{-3}$ ,  $H$  is 7 km, and  $z$  is the altitude (km) of the calculation, as was used in the dual-Doppler calculations. These vertical divergence values were then horizontally-integrated to derive a net vertical divergence value at each height. Note that the horizontally-integrated vertical divergence values were calculated by including

### Relationship between LNB and LMD

G. L. Mullendore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



positive vertical velocities only. Negative velocities were removed because the main goal is to diagnosis the level of detrainment for deep transport from the boundary layer, and, as shown in Fig. 4c of Mullendore et al. (2009), vertical divergence in deep down-drafts can alter the signature of vertical convergence due to the deep updrafts alone.

Figure 2 shows the calculated horizontal detrainment for the 29 June 2000 supercell case at 23:57 UTC (dashed line). From this graph, we can define the LMD, 12.1 km, and the “detrainment envelopment”, or the range of values over which vertical divergence is negative, suggesting horizontal detrainment. For the time shown, the detrainment envelopment extends from 5.3 to 15.6 km. By averaging over all sampled times, we can obtain an average detrainment envelope for the storm (Fig. 2; solid line).

To investigate how the deep convective detrainment evolved with time, the detrainment envelope was plotted at each observation time and the altitudes of the LMD (Fig. 3; red line), and minimum (gray line) and maximum (blue line) extent, were recorded. The black dashed line shows the most representative LNB. As stated previously, the LNB is generally considered an upper bound on the LMD, but from observation times 23:57 UTC onward, the LMD is approximately at or above the most representative LNB. The magnitude of the LMD was also recorded at all times; the case-normalized LMD is shown on the right-hand y-axis of Fig. 3 (solid black line). The 29 June 2000 case was unique in that it showed a significant increase in the altitude (Fig. 3; red line) of the LMD over time, with a mean altitude of 11.0 km before the strengthening period (23:20 to 23:38 UTC; solid black line) and a mean altitude of 12.3 km after the strengthening period. This increase in LMD strength and altitude corresponds to the time of the strong right turn (“severe right mature phase” Tessendorf et al., 2005). The timing of this increase suggests that the enhancement in transport corresponds to the co-location of the supercell updraft with positive vertical pressure gradient forcing to the south of the original updraft (e.g., Rotunno and Klemp, 1985) and highlights the importance of storm morphology in understanding deep convective transport.

## Relationship between LNB and LMD

G. L. Mullendore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Most of the cases analyzed fit what was expected, i.e. the mean LMD was below the most representative LNB. Figure 4 shows the time evolution of detrainment envelope, identical to what is shown in Fig. 3 for the 29 June 2000 case, for the additional eight cases. The 2 July 1998 CHILL supercell case (panel a) had LMDs (red line) near the altitude of the most representative LNB (black dashed line). The 11 June 2000 STEPS convective line (which had a markedly large cell embedded in the line) had two observed times where the LMD was near the LNB. Figure 4 also demonstrates that all the cases studied had significant variability in time, in either the LMD, the depth of the detrainment envelope, or the magnitude of detrainment (shown as normalized detrainment magnitude in Fig. 4; black line). The time variability of detrainment is not captured in a signal snapshot, highlighting an additional challenge in estimating detrainment values from a single environmental sounding.

In addition, even when the LMD was below the most representative LNB, the upper edge of the detrainment envelope (blue lines) was at most times above the LNB, suggesting some transport occurred above the LNB, and in certain cases, extending up to 16 km. This depth suggests that at least in some cases the storms were transporting boundary layer mass directly into the lowermost stratosphere. The regions of the detrainment envelopes from the LMD to the upper edge of the detrainment envelope include transport occurring in the overshooting tops. As discussed in Mullendore et al. (2005), the majority of the air in overshooting tops is negatively buoyant, and therefore descends back to lower altitudes. However, due to vigorous mixing at the top of the storm (e.g., Fig. 9 of Mullendore et al., 2005; Lane et al., 2003), a portion of the overshooting air is mixed with stratospheric air, thereby attaining neutral buoyancy and irreversible transport. Although the percentage of the air irreversibly transported via overshooting tops (and, more generally, via mixing at cloud top) is small compared to the amount of transport occurring at the LMD, this amount can be very important because of transport of even small concentrations of reactive chemicals into the lowermost stratosphere. Additional analysis of the amount transported to the stratosphere requires careful treatment of the tropopause, as the tropopause region becomes poorly

## Relationship between LNB and LMD

G. L. Mullendore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



defined above active convection (e.g., Mullendore et al., 2005), therefore further quantification of the cross-tropopause transport is beyond the scope of the current analysis.

It should be noted the detrainment envelope calculations are sensitive to dual-Doppler processing, e.g., the case presented in Mullendore et al. (2009) from the LBA-TRMM campaign had spurious near-surface velocities away from main updrafts, which contaminated an analysis of the full domain and forced a more subjective selection of analysis areas. Even in the cases presented here, some lower level contamination existed; for times where significant lower level uncertainty existed, no lower bound on the detrainment envelope (Fig. 4a,b; dashed gray lines) was recorded. An objective methodology for choosing the analysis region, with focus on the anvil region of a given case, is currently being tested (e.g., Feng et al., 2011). Another interesting feature noted particularly in the convective line detrainment profiles was a secondary, lower magnitude detrainment maximum at a higher altitude than the main LMD. This feature is hypothesized to be due to dissipating convective cells transitioning into the stratiform region (e.g., Houze Jr., 1997) and is also under further investigation.

## 5 Relationship between LNB and LMD

The mean LMD calculated for each case is given in Table 1. The mean LMD was calculated by averaging the LMD values over all times observed, except in the case of the 29 June 2000 STEPS supercell. For the 29 June 2000 case, two LMDs were calculated, one before strengthening (to 23:43 UTC, 11.0 km) and one after strengthening (from 23:51 UTC, 12.3 km); the post-strengthening LMD is used in comparison figures. As follows from the discussion of the time-varying LMD in Sect. 4, the mean LMD was below the LNB in all cases except for the 29 June 2000 STEPS case, which had a mean LMD equal to the LNB. Figure 5 summarizes the comparison between the most representative LNB (x-axis) and mean LMD (y-axis). The one-to-one comparison is plotted as a gray solid line. The supercell cases (shown as diamonds) had markedly higher LMDs than the other storm types (x is unicell, plus signs are multicells, circles

### Relationship between LNB and LMD

G. L. Mullendore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



are convective lines). On average, all other storms had mean LMD values 2.8 km below the most representative LNB (Fig. 5, gray dashed line).

## 6 Conclusions

As stated in Sect. 1, the purpose of this study is to quantify the variability associated with predicting detrainment levels, specifically the prediction from environmental soundings. Although it is generally accepted that the level of neutral buoyancy (LNB) is only a coarse estimate of updraft depth (as detailed in Sect. 3), the LNB is still used to understand and predict storm structure. Using a method proposed in Mullendore et al. (2009), nine dual-Doppler convective cases were used to determine the observed level of maximum detrainment (LMD) to compare with the LNB. The LNB for each case was calculated with a variety of methods and with a variety of sources (including both observed and simulated soundings). The most representative LNB was chosen as the proximity sounding from NARR using the most unstable parcel and including ice processes.

The observed cases were a mix of storm morphologies, including both supercell and multicell storms. As expected, the LMD was generally below the LNB, the mean offset for all cases being 2.2 km. However, there was a marked difference between the supercell and non-supercell cases. The two supercell cases had LMDs just below (0.3 km; 2 July 1998) or equal to (29 June 2000) the LNB. The mean offset for all other cases was a LMD 2.8 km below the LNB. These findings are significant for two reasons. First, the LNB is considered an upper bound on significant convective mass transport, as calculations of LNB ignore parcel mixing, which would be expected to reduce updraft depth. The findings presented here suggest that, for time-averaged values, using LNB as an upper bound may be appropriate, but only with careful selection of environmental sounding. Additionally, the assumption often made that the LMD is likely below the LNB does not always hold true, and at certain observed times, the LMD is even above the LNB. Second, storm morphology has a significant impact on mass transport (e.g.,

### Relationship between LNB and LMD

G. L. Mullendore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Mullendore et al., 2005), suggesting that internal storm dynamics are important for transport calculations. However, with further case studies, a predictable relationship between LNB and LMD may yet be determined for a given storm regime. The Deep Convective Clouds and Chemistry (UCAR, 2012b) field campaign will provide several additional cases that can provide further insights into the relationships presented in this study.

This study has also demonstrated the evolution of the LMD over the lifetime of the storm can be significant (e.g., >2 km altitude change in 30 min for 29 June 2000 supercell), and this time evolution is lacking from models with coarse time steps, missing significant changes in detrainment levels that may strongly impact the amount of boundary layer mass transported to the lower stratosphere.

*Acknowledgements.* This work was supported by the National Science Foundation Division of Atmospheric Sciences, NSF grant ATM-0918010. The authors would like to thank George Bryan for making available his parcel ascent code to calculate CAPE, Aaron Kennedy for assistance in obtaining the NARR data, and Kurtis Pinkney for assistance with data analysis. The authors also thank Jeffrey Tilley, Matt Gilmore, and Mark Askelson for helpful comments and discussions.

## References

- Arakawa, A.: The cumulus parameterization problem: Past, present, and future, *J. Climate*, 17, 2493–2525, 2004. 21270
- Cohen, C.: A quantitative investigation of entrainment and detrainment in numerically simulated cumulonimbus clouds, *J. Atmos. Sci.*, 57, 1657–1674, 2000. 21270
- Dickerson, R. R.: Thunderstorms: An important mechanism in the transport of air pollutants, *Science*, 235, 460–465, 1987. 21268, 21269
- Doswell III, C. A. and Markowski, P. M.: Is Buoyancy a Relative Quantity?, *Mon. Weather Rev.*, 132, 853–863, 2004. 21270

### Relationship between LNB and LMD

G. L. Mullendore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





- Feng, Z., Dong, X., Xi, B., Schumacher, C., Minnis, P., and Khaiyer, M.: Top-of-atmosphere radiation budget of convective core/stratiform rain and anvil clouds from deep convective systems, *J. Geophys. Res. Atmos.*, 116, D23202, doi:10.1029/2011JD016451, 2011. 21279
- Houze Jr., R. A.: Stratiform Precipitation in Regions of Convection: A Meteorological Paradox?, *B. Am. Meteor. Soc.*, 78, 2179–2196, 1997. 21279
- Kuhlman, K. M., Ziegler, C. L., Mansell, E. R., MacGorman, D. R., and Straka, J. M.: Numerically simulated electrification and lightning of the 29 June 2000 STEPS supercell storm, *Mon. Weather Rev.*, 134, 2734–2757, 2006. 21273
- Lane, T. P., Sharman, R. D., Clark, T. L., and Hsu, H.-M.: An investigation of turbulence generation mechanisms above deep convection, *Journal of the Atmos. Sci.*, 60, 1297–1321, 2003. 21278
- Lang, T. J. and Rutledge, S. A.: Relationships between convective storm kinematics, precipitation, and lightning, *Mon. Weather Rev.*, 130, 2492–2506, 2002. 21271, 21273
- Lang, T. J. and Rutledge, S. A.: Kinematic, microphysical, and electrical aspects of an asymmetric bow-echo mesoscale convective system observed during STEPS 2000, *J. Geophys. Res. Atmos.*, 113, D08213, doi:10.1029/2006JD007709, 2008. 21271, 21273
- Lang, T. J., Miller, L. J., Weisman, M., Rutledge, S. A., Barker, III, L. J., Bringi, V. N., Chandrasekar, V., Detwiler, A., Doesken, N., Helsdon, J., Knight, C., Krehbiel, P., Lyons, W. A., MacGorman, D., Rasmussen, E., Rison, W., Rust, W. D., and Thomas, R. J.: The Severe Thunderstorm Electrification and Precipitation Study, *B. Am. Meteor. Soc.*, 1107–1125, 2004. 21271, 21272, 21273
- Manzato, A. and Morgan, G.: Evaluating the sounding instability with the Lifted Parcel Theory, *Atmos. Res.*, 67-68, 455–473, 12, 2003. 21269
- Mesinger, F., DiMego, G., Kalnay, E., Mitchell, K., Shafran, P. C., Ebisuzaki, W., Jović, D., Woollen, J., Rogers, E., Berbery, E. H., Ek, M. B., Fan, Y., Grumbine, R., Higgins, W., Li, H., Lin, Y., Manikin, G., Parrish, D., and Shi, W.: North American regional reanalysis, *B. Am. Meteorol. Soc.*, 87, 343–360, 2006. 21272
- Mullendore, G. L., Durran, D. R., and Holton, J. R.: Cross-tropopause tracer transport in midlatitude convection, *J. Geophys. Res.*, 110, D06113, doi:10.1029/2004JD005059, 2005. 21268, 21270, 21278, 21279, 21281
- Mullendore, G. L., Homann, A. J., Bevers, K., and Schumacher, C.: Radar reflectivity as a proxy for convective mass transport, *J. Geophys. Res. Atmos.*, 114, D16103, doi:10.1029/2008JD011431, 2009. 21270, 21276, 21277, 21279, 21280

## Relationship between LNB and LMD

G. L. Mullendore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





## Relationship between LNB and LMD

G. L. Mullendore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Pickering, K. E., Dickerson, R. R., Huffman, G. J., Boatman, J. F., and Schanot, A.: Trace Gas Transport in the Vicinity of Frontal Convective Clouds, *J. Geophys. Res.*, 93, 759–773, doi:10.1029/JD093iD01p00759, 1988. 21268

Pickering, K. E., Thompson, A. M., Tao, W.-K., and Kucsera, T. L.: Upper Tropospheric Ozone Production Following Mesoscale Convection During STEP/EMEX, *J. Geophys. Res.*, 98, 8737–8749, doi:10.1029/93JD00875, 1993. 21269

Randall, D., Khairoutdinov, M., Arakawa, A., and Grabowski, W.: Breaking the Cloud Parameterization Deadlock, *Bull. Amer. Meteorol. Soc.*, 1547–1564, 2003. 21270

Rotunno, R. and Klemp, J. B.: On the rotation and propagation of simulated supercell thunderstorms., *J. Atmos. Sci.*, 42, 271–292, 1985. 21277

Tessendorf, S. A., Miller, L. J., Wiens, K. C., and Rutledge, S. A.: The 29 June 2000 supercell observed during STEPS. Part I: Kinematics and microphysics, *J. Atmos. Sci.*, 62, 4127–4150, 2005. 21271, 21273, 21277

Tessendorf, S. A., Rutledge, S. A., and Wiens, K. C.: Radar and lightning observations of normal and inverted polarity multicellular storms from STEPS, *Mon. Weather Rev.*, 135, 3682–3706, 2007a. 21271

Tessendorf, S. A., Wiens, K. C., and Rutledge, S. A.: Radar and lightning observations of the 3 June 2000 electrically inverted storm from STEPS, *Mon. Weather Rev.*, 135, 3665–3681, 2007b. 21271

Thompson, A. M., Tao, W.-K., Pickering, K. E., Scala, J. R., and Simpson, J.: Tropical Deep Convection and Ozone Formation, *Bulletin of the American Meteorological Society*, 78, 1043–1054, doi:10.1175/1520-0477(1997)078<1043:TDCAOF>2.0.CO;2, 1997. 21269

Thompson, R. L., Edwards, R., Hart, J. A., Elmore, K. L., and Markowski, P.: Close proximity soundings within supercell environments obtained from the Rapid Update Cycle, *Weather Forecast.*, 18, 1243–1261, 2003. 21272

UCAR: STEPS Operations Journal, <http://www.mmm.ucar.edu/pdas/STEPS/ops-journal.html>, (last access: 27 June 2012), 2012a. 21273

UCAR: DC3 Home Page, <http://www.eol.ucar.edu/projects/dc3/index.html>, (last access: 1 July 2012), 2012b. 21281

Wiens, K. C.: Kinematic, microphysical, and electrical structure and evolution of thunderstorms during the Severe Thunderstorm Electrification and Precipitation Study (STEPS), Ph.D. thesis, Colorado State University, USA, 2005. 21271

## Relationship between LNB and LMD

G. L. Mullendore et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Table 1.** Summary of detrainment analysis for each case. Most representative level of neutral buoyancy (LNB) is calculated from the central NARR simulated sounding; mature level of maximum detrainment (LMD) is calculated by taking the mean LMD during the mature stage of the observed storm. The mean and standard deviation calculations for LNB include all non-zero methods from all collected soundings (e.g. see Table 2 for LNB calculations for STEPS 6/29 case). All STEPS cases are from 2000; all CHILL cases are from 1998. All altitudes are given as a.g.l.

Case	Storm Type	Most Representative LNB (km)	Mean LNB (km)	LNB Standard Deviation (km)	Mean LMD (km)
STEPS 6/3	unicell	11	10.1	0.8	8.2
STEPS 6/11	convective line	11.4	10.4	1	8.3
STEPS 6/19	convective line	10.7	10.4	1.4	6.7
STEPS 6/22	convective line	12.3	12	0.6	10.5
STEPS 6/23	multicell	12.7	11.3	1.3	8.8
STEPS 6/29	supercell	12.3	11.4	1	12.3
CHILL 7/2	supercell	11.1	10.7	0.3	10.8
CHILL 7/15	convective line	11.9	10.7	1.3	9.4
CHILL 7/21	multicell	12.6	12.2	0.2	10.9

## Relationship between LNB and LMD

G. L. Mullendore et al.

**Table 2.** Level of neutral buoyancy (LNB) calculated for the STEPS 6/29 case, including simulated (NARR) and observed soundings. LNB is calculated assuming irreversible (pseudoadiabatic) ascent using two methods (liquid only hydrometeors and liquid and frozen hydrometeors) and using three different initial parcel assumptions (surface-based, mixed layer, and most unstable).

STEPS 6/29		Liquid Only LNB (km)			With Ice LNB (km)		
Location	Time (UTC)	Surface	Mixed Layer	Most Unstable	Surface	Mixed Layer	Most Unstable
NARR 38.78° N, 101.30° W	6/30 00:00	12	11.8	12	12.3	12	12.3
NARR South	6/30 00:00	12.1	11.8	12.1	12.3	12.1	12.3
NARR West	6/30 00:00	11.6	11.4	11.6	11.6	11.6	11.9
NARR East	6/30 00:00	12.1	11.8	12.1	12.1	12.1	12.1
39.367° N, 101.703° W	6/29 17:27	NA	NA	8.9	NA	NA	NA
40.441° N, 102.306° W	6/29 17:57	10.8	9.4	11	10.6	9.3	10.9
39.367° N, 101.703° W	6/29 20:22	10.9	10.8	11.1	10.7	10.6	10.9
40.441° N, 102.306° W	6/29 23:38	NA	NA	8.5	NA	NA	NA

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



**Relationship  
between LNB and  
LMD**

G. L. Mullendore et al.



**Fig. 1.** Horizontal cross-section of reflectivity (dBZ) at 2 km a.g.l. of 29 June 2000 supercell case from the STEPS campaign at time 23:57 UTC. Solid dots show radar locations, triangles show locations of mobile soundings (with launch times labeled), and plus marks show locations of NARR simulated soundings.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## Relationship between LNB and LMD

G. L. Mullendore et al.




**Fig. 2.** Horizontally-integrated vertical divergence for the 29 June 2000 supercell case at 23:57 UTC (dashed line) and averaged over all times (solid line). The level of maximum de-trainment (LMD) is approximately 12.3 km for both divergence lines.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Relationship between LNB and LMD

G. L. Mullendore et al.




**Fig. 3.** Left axis: Time evolution of detrainment in 29 June 2000 supercell case including level of maximum detrainment (LMD; red line) and maximum (blue line) and minimum (gray line) extent of detrainment envelope. The altitude of the LNB from the most representative sounding is shown as a black dashed line. Right axis: Time evolution of normalized detrainment magnitude (black line) for same case.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

# Relationship between LNB and LMD

G. L. Mullendore et al.




**Fig. 4.** Same as shown in Fig. 3, but for additional case studies.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## Relationship between LNB and LMD

G. L. Mullendore et al.



**Fig. 5.** Most representative LNB height (km) plotted versus mean LMD height (km, see text for details). Each case is designated by different color, and the shapes designate storm type: x for unicell, plus sign for multicell, circle for convective line, and diamond for supercell. The LNB versus LMD one-to-one line is plotted for reference as a solid gray line; a line representing LNB 2.8 km above the LMD is plotted as dashed gray.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

