

# Auxiliary Material: Impact of Very Short-lived Halogens on Stratospheric Ozone Abundance and UV radiation in a Geo-engineered Atmosphere

S. Tilmes<sup>1</sup>, Doug E. Kinnison<sup>1</sup>, Rolando R. Garcia<sup>1</sup>, Ross Salawitch<sup>2</sup>, Tim Canty<sup>2</sup>, Julia Lee-Taylor<sup>1</sup>, Sasha Madronich<sup>1</sup>, and Kelly Chance<sup>3</sup>

<sup>1</sup>National Center for Atmospheric Research, Boulder, Colorado, USA

<sup>2</sup>University of Maryland College Park, MD, USA

<sup>3</sup>Harvard-Smithsonian Center for Astrophysics, Cambridge, MA, USA

*Correspondence to:* Simone Tilmes, National Center for Atmospheric Research, Boulder, Colorado, USA (tilmes@ucar.edu)

## 1 Meteorology of the Polar Vortex

As described in the main text, we select meteorological conditions of the year 2040 from Tilmes et al. (2009) for simulations presented in this study. Chemical ozone loss in high latitudes is strongly linked to temperatures in the polar vortex, since the activation of chlorine is strongly temperature dependent (Drdla and Müller, 2012, and references therein). The activation of chlorine containing reservoirs is dominated by reactions on cold binary aerosol particles. A formulation to calculate the threshold temperature for the activation of chlorine ( $T_{ACL}$ ) was derived by Drdla and Müller (2012).  $T_{ACL}$  is dependent on potential temperature, water vapor and the surface area density (SAD). Vortex temperatures are therefore not directly correlated to chlorine activation. Tilmes et al. (2007) have introduced a measure for the Potential for activated chlorine (PaCl). PaCl describes the fraction of the vortex where temperatures are below  $T_{ACL}$ . This measure can be directly linked to chlorine activation and therefore ozone loss in the polar vortex (WMO, 2010).

In Figure S1, we compare the meteorology of the polar vortex used in this study based on WACCM results to MERRA reanalysis (Rienecker et al., 2011) for recent winters (as shown in Figure S1, left panel). For this, average temperatures of the Arctic and Antarctic polar vortex are compared. The polar vortex edge is calculated by using the criterion by Nash et al. (1996). In addition, we compare PaCl derived using the WACCM meteorology and background SAD conditions to values derived using MERRA reanalysis (Figure S1, right panel).

Polar vortex temperatures over Antarctica in WACCM are lower compared to MERRA reanalysis,

20 resulting in a colder and longer-lasting winter. After reaching  $T_{ACL}$ , colder temperatures do not trigger more chlorine activation. PACI is therefore in good agreement with the values derived from MERRA between July and October. PACI in WACCM is larger in November compared to the reanalysis dataset, since temperatures are below  $T_{ACL}$  for a longer period. An overestimation of chemical ozone depletion in November compared to observations can be expected in this study.

25 Arctic average polar vortex temperatures tend to be warmer between January and March than recent cold Arctic winters. However, PACI is comparable to values derived from MERRA reanalysis for moderately cold Arctic winters. This is, because WACCM temperatures describe a temperature gradient in the polar vortex ranging from rather warm to very cold temperatures towards the vortex core. We therefore expect chlorine activation in the Arctic vortex to be comparable to a recently  
30 observed moderately cold Arctic winter. As for Antarctica, the Arctic vortex is more persistent and long-lasting compared to observed cold winters. In April, vortex temperatures are well below observed temperatures and PACI values are larger than what we expect from recent observations. However, the Arctic vortex is rather small compared to observations (not shown) and the expected overestimation in ozone loss only covers a small area north of  $75^{\circ}N$ .

## 35 **2 Contributions of VSL organics ( $Br_y^{VSL}$ ) to the stratospheric total bromine ( $Br_y$ )**

In the main paper, three cases for stratospheric  $Br_y^{VSL}$  are considered. Here we justify the enhancement of  $Br_y$  of 6 ppt as an average estimate derived from earlier studies, and the enhancement of  $Br_y$  of 10 ppt plus an increase of  $Cl_y$  by 100 ppt, as an upper limit estimate considering stratospheric  $Br_y^{VSL}$ .

40 The past three WMO/UNEP Scientific Assessment of Ozone Depletion reports have quantified the enhancement to stratospheric total inorganic bromine ( $Br_y$ ) due to VSL organics ( $Br_y^{VSL}$ ). Chapter 2 of WMO (2003) provides an overview of the chemical, biological, and physical processes that govern  $Br_y^{VSL}$ . The largest contributions are from dibromomethane ( $CH_2Br_2$ ) and bromoform ( $CHBr_3$ ), which are produced by biological processes in the ocean. Sea-salt and halide-containing aerosols  
45 supply bromine to the lower troposphere, but likely do not play a strong role in affecting stratospheric  $Br_y$  (Yang et al., 2005).

Table 1-9 of WMO (2010) gives a range for  $Br_y^{VSL}$  of 1 to 8 ppt based on tropospheric focused studies. Table 1-14 of WMO (2010) gives an ensemble mean estimate of 6 ppt for  $Br_y^{VSL}$  based on stratospheric focused studies. The tropospheric focused estimate is based on observed abundances  
50 of VSL bromocarbons near the cold point tropopause combined with model estimates of efficiency of transport, to the stratosphere, of product gases formed when VSL bromocarbons decompose in the troposphere. The stratospheric studies are all based on observations of BrO;  $Br_y^{VSL}$  is inferred from the difference between  $Br_y$  calculated from observed BrO (typically in the middle and upper stratosphere) and the amount of  $Br_y$  that is supplied by long-lived methyl bromine ( $CH_3Br$ ) and




Fig. 1: Left panels: WACCM averaged polar vortex temperature between 400-550 K for the Antarctic (top) and Arctic (bottom) compared to MERRA reanalysis for different years (different colors). For the Arctic, three cold winters (1997, 2000, and 2011) and one very warm winter (1997) are selected. Right panels: Potential for activated chlorine PACl (Tilmes et al., 2009) derived using WACCM results and MERRA reanalysis for background SAD conditions. See text for more details.

55 halons.

In the main text we consider three cases for  $\text{Br}_y^{\text{VSL}}$ : 0, 6, and 10 ppt. The 6 ppt value is based on the ensemble mean from Table 1-14 of WMO (2010). The upper limit of 10 ppt is chosen because 7 of the 12 studies represented in Table 1-14 have upper limits of 10 ppt or higher. The plausibility of the 10 ppt upper limit is further supported by an analysis of consistency between satellite observations of total column BrO versus the sum of tropospheric partial column BrO (based on in situ aircraft data) and stratospheric partial column BrO (based on a model), published after WMO (2010), that reported best agreement for  $\text{Br}_y^{\text{VSL}}$  of 9 ppt (Choi et al., 2012).

Chapter 1 and 2 of (WMO, 2007) and Chapter 1 of (WMO, 2010) quantified the enhancement to stratospheric total inorganic chlorine ( $\text{Cl}_y$ ) due to VSL chlorocarbons ( $\text{Cl}_y^{\text{VSL}}$ ). Two lines of evidence were pursued. Time series of HCl measured in the upper stratosphere by three instruments, ATMOS, MLS, and ACE, are most consistent with expectation if 100 ppt is added to the burden of stratospheric  $\text{Cl}_y$  supplied by long lived sources. In contrast, upper stratospheric HCl from HALOE suggests  $\text{Cl}_y^{\text{VSL}}$  is close to zero (Froidevaux, et al., 2006, Fig. 1-2 of WMO). Interpretation of

70 tropospheric data is complicated because: a) various VSL organics decompose to form phosgene  
(COCl<sub>2</sub>) that is likely transported into the stratosphere, but phosgene is also a byproduct of the  
decomposition of long-lived chlorocarbons (which undergo slight loss in the troposphere) (WMO,  
2007, Section 2.5.1.2); b) HCl has been observed in the sub-tropical upper troposphere but the source  
could be either local decomposition of VSL chlorocarbons or irreversible stratosphere to troposphere  
mixing of air (Marcy et al., 2004). In this study, we consider two cases for Cl<sub>y</sub><sup>VSL</sup>: 0 and 100 ppt.  
75 The 100 ppt value is consistent with upper stratospheric HCl from ATMOS, MLS, and ACE as well  
as the range for Cl<sub>y</sub><sup>VSL</sup> of 25 to 170 ppt given in Table 1-9 of WMO (2010).

## 2.1 Changes in ozone loss rates: Baseline – Geo-engineering

The impact of volcanic aerosols on ozone destroying cycles was discussed in various studies (e.g.,  
Brasseur et al., 1990; Solomon, 1999), as well as the sensitivity to ozone loss cycles on increases in  
80 bromine (e.g., Salawitch et al., 1994; Solomon et al., 1996; Solomon, 1999). Here, we investigate  
the impact of geo-engineering on ozone loss rates for projected 2040 chemical conditions and further  
quantify the impact of additional stratospheric bromine from VSL.

The increase of aerosol particles and therefore heterogeneous reactions was shown to lead to a  
reduced NO<sub>x</sub>/NO<sub>y</sub> ratio between 15-25 km (e.g., Fahey et al., 1993). This results in decreasing  
85 ozone loss rates in mid- and low latitudes around 20 km. The reduction of NO<sub>x</sub> further induces  
changes in the gas phase chemistry (Solomon et al., 1996). For mid-latitudes, NO<sub>x</sub> ozone loss  
rates are strongly reduced for the geo-engineering simulation compared to the baseline simulation,  
whereas HO<sub>x</sub>, ClO<sub>x</sub>, and BrO<sub>x</sub> loss rates are increased at the same altitudes (Figure S2, 5th and  
6th row), second column. Changes in Br<sub>y</sub><sup>VSL</sup> and Cl<sub>y</sub><sup>VSL</sup> do not influence NO<sub>x</sub> ozone loss rates, as  
90 expected, see Figure S2, 3th and 4th column.

In high polar latitudes in winter and spring, increased heterogeneous reactions due to geo-engineering  
lead to a doubling of ClO<sub>x</sub> + BrO<sub>x</sub> loss rates in the lower stratosphere. At this time of the year, the  
HO<sub>x</sub> cycle is less important and geo-engineering induces only a small increase in HO<sub>x</sub> ozone loss  
rates. For the NH, enhanced bromine (VSL = 6) results in an increase of ClO<sub>x</sub> + BrO<sub>x</sub> ozone loss  
95 rates by another 50%. For Br<sub>y</sub><sup>VSL</sup> = 10 ppt, Cl<sub>y</sub><sup>VSL</sup> = 100 ppt, ClO<sub>x</sub> + BrO<sub>x</sub> loss rates double com-  
pared to the VSL = 0 case and reach 3 times the amount derived from the baseline simulation.

In the Austral spring and summer in the SH, Figure S2, third row, the increase of ClO<sub>x</sub> + BrO<sub>x</sub>  
rates due to geo-engineering is of similar magnitude as for the NH, but is relatively less important.  
Additional Br<sub>y</sub> and Cl<sub>y</sub> in the stratosphere results in a decrease of the ClO<sub>x</sub> + BrO<sub>x</sub> rates in a  
100 geo-engineered environment for some altitudes (Figure S2, third row). In contrast to the VSL = 0  
case, ozone reaches very small values at the end of spring and the reduction of loss rates for the  
geo-engineering simulation is limited.

During polar summer (MJJ for the NH and NDJ for the SH), temperatures are warming up and  
heterogeneous reaction are less important. Only a small increase of the ClO<sub>x</sub> + BrO<sub>x</sub> cycle is

105 simulated around 12 km due to geo-engineering. On the other hand, the HO<sub>x</sub> cycle is much more important at this time of the year, especially for the NH. The additional bromine approximately doubles the HO<sub>x</sub> cycle and strongly increases the ClO<sub>x</sub> + BrO<sub>x</sub> ozone loss rates in the lowermost stratosphere (around 12 km).

110 For summer mid-latitudes in the NH, increased heterogeneous reactions in spring and summer result in reduced NO<sub>x</sub> ozone loss rates and increased ClO<sub>x</sub> + BrO<sub>x</sub> and HO<sub>x</sub> rates around 25 km. The consideration of additional Br<sub>y</sub> from VSL halogen compounds results in a large increase in the ClO<sub>x</sub> + BrO<sub>x</sub> rates, which doubles the impact of geoengineering on column ozone (from about 2% to 4% for MJJ, as discussed in the main text).

## References

- 115 Brasseur, G., Granier, C., and Walters, S.: Future changes in stratospheric ozone and the role of heterogeneous chemistry, *Nature*, 348, 626–628, 1990.
- Choi, S., Wang, Y., Salawitch, R. J., Canty, T., Joiner, J., Zeng, T., Kurosu, T. P., Chance, K., Richter, A., Huey, L. G., Liao, J., Neuman, J. A., Nowak, J. B., Dibb, J. E., Weinheimer, A. J., Diskin, G., Ryerson, T. B., da Silva, A., Curry, J., Kinnison, D., Tilmes, S., and Levelt, P. F.: Analysis of satellite-  
120 derived Arctic tropospheric BrO columns in conjunction with aircraft measurements during ARCTAS and ARCPAC, *Atmospheric Chemistry and Physics*, 12, 1255–1285, doi:10.5194/acp-12-1255-2012, <http://www.atmos-chem-phys.net/12/1255/2012/>, 2012.
- Drdla, K. and Müller, R.: Temperature thresholds for chlorine activation and ozone loss in the polar stratosphere, *Annales Geophysicae*, 30, 1055–1073, doi:10.5194/angeo-30-1055-2012, [http://www.ann-geophys.net/30/](http://www.ann-geophys.net/30/1055/2012/)  
125 1055/2012/, 2012.
- Fahey, D. W., Kawa, S. R., Woodbridge, E. L., Tin, P., Wilson, J. C., Jonsson, H. H., Dye, J. E., Baumgardner, D., Borrmann, S., and Toohey, D. W.: In situ measurements constraining the role of sulphate aerosols in mid-latitude ozone depletion, *Nature*, 363, 509–514, 1993.
- Froidevaux, et al., L.: Temporal decrease in upper atmospheric chlorine, *Geophys. Res. Lett.*, 33, L23812, doi:10.1029/2006GL027600, 2006.  
130
- Marcy, T. P., Fahey, D. W., Gau, R. S., Popp, P. J., Richard, E. C., Thompson, T. L., Rosenlof, K. H., Ray, E. A., Salawitch, R. J., Atherton, C. S., Bergmann, D. J., Ridley, B. A., Winheimer, A. J., Weinstock, M. L. E. M., and Mahoney, M. J.: Quantifying Stratospheric Ozone in upper troposphere with in situ measurements of HCl, *Science*, 2004.
- 135 Nash, E. R., Newman, P. A., Rosenfield, J. E., and Schoeberl, M. R.: An objective determination of the polar vortex using Ertel’s potential vorticity, *J. Geophys. Res.*, 101, 9471–9478, 1996.
- Rienecker, M., Suarez, M., Gelaro, R., Todling, R., Bacmeister, J., Liu, E., Bosilovich, M., Schubert, S., Takacs, L., Kim, G.-K., Bloom, S., Chen, J., Collins, D., Conaty, A., and A. da Silva, e. a.: MERRA - NASA’s Modern-Era Retrospective Analysis for Research and Applications, *J. Climate*, 24, 3624–3648,  
140 doi:10.1175/JCLI-D-11-00015, 2011.
- Salawitch, R. J., Wofsy, S. C., Wennberg, P. O., Cohen, R. C., Anderson, J. G., Fahey, D. W., Gao, R. S., Keim, E. R., Woodbridge, E. L., Stimpfle, R. M., Koplow, P., Kohn, D. W., Webster, C. R., May, R. D., Pfister, L., Gottlieb, E. W., Michelsen, H. A., Yue, G. K., Prather, M. J., Wilson, J. C., Brock, C. A., Jonsson, H. H., Dye, J. E., Baumgardner, D., Proffitt, M. H., Loewenstein, M., Podolske, J. R., Elkins, J. W., Dutton, G. S.,  
145 Hints, E. J., Dessler, A. E., Weinstock, E. M., Kelly, K. K., Boering, K. A., Daube, B. C., Chan, K. R., and Bowen, S. W.: The diurnal variation of hydrogen, nitrogen, and chlorine radicals: implications for the heterogeneous production of HNO<sub>2</sub>, *Geophys. Res. Lett.*, 21, 2551–2554, doi:10.1029/94GL02781, 1994.
- Solomon, S.: Stratospheric ozone depletion: A review of concepts and history, *Rev. Geophys.*, 37, 275–316, doi:10.1029/1999RG900008, 1999.
- 150 Solomon, S., Portmann, R., Garcia, R. R., Thomason, L., Poole, L. R., and McCormick, M. P.: The role of aerosol variations in anthropogenic ozone depletion at northern midlatitudes, *J. Geophys. Res.*, 101, 6713–6727, 1996.
- Tilmes, S., Kinnison, D., Müller, R., Sassi, F., Marsh, D., Boville, B., and Garcia, R.: Evaluation of hetero-

- geneous processes in the polar lower stratosphere in the Whole Atmosphere Community Climate Model,  
155 J. Geophys. Res., 112, D24301, doi:10.1029/2006JD008334, 2007.
- Tilmes, S., Garcia, R. R., Kinnison, D. E., Gettelman, A., and Rasch, P. J.: Impact of geoengineered aerosols  
on the troposphere and stratosphere, J. Geophys. Res., 114, D12305, doi:10.1029/2008JD011420, 2009.
- WMO: Scientific assessment of ozone depletion: 2002, Global Ozone Research and Monitoring Project-Report  
No. 47, Geneva, Switzerland, 2003.
- 160 WMO: Scientific assessment of ozone depletion: 2006, Global Ozone Research and Monitoring Project-Report  
No. 50, Geneva, Switzerland, 2007.
- WMO: Scientific assessment of ozone depletion: 2010, Global Ozone Research and Monitoring Project-Report  
No. 51, Geneva, Switzerland, 2010.
- Yang, X., Cox, R. A., Warnwick, N. J., Pyle, J. A., Carver, G. D., O'Connor, F. M., , and Savage, N. H.:  
165 Tropospheric bromine chemistry and its impacts on ozone: A model study, J. Geophys. Res., 110, D23311,  
doi:10.1029/2005JD006244, 2005.




Fig. 2: First column: ozone loss rates for different ozone destroying cycles (different colors) and total ozone loss rates (black); baseline simulations (solid) and geo-engineering simulation (dashed) in 2040 for VSL = 0 (see main text for definition) considering different seasons and regions. Second column: differences between geo-engineering and baseline simulations as shown on the left. Third column: differences between VSL = 6 and VSL = 0 case for the geo-engineering simulations only. Fourth column: differences between VSL = 10 + 100 and VSL = 0 case for the geo-engineering simulation only.