

¹⁰Max Planck Institute for Meteorology, Hamburg, Germany

¹¹Department of Geosciences, University of Oslo, Norway

¹²NCAR Earth System Laboratory, National Center for Atmospheric Research, Boulder, CO, USA

¹³Department of Atmospheric, Oceanic, and Space Sciences, University of Michigan, Ann Arbor, Michigan, USA

¹⁴Atmospheric Sciences Research Center, State University of New York at Albany, New York, USA

¹⁵University of Oxford, Oxford, UK

¹⁶Research Institute for Applied Mechanics, Kyushu University, Japan

¹⁷Chinese Academy of Meteorological Sciences, Beijing 100081, China

¹⁸Scripps Institution of Oceanography, University of California San Diego, La Jolla, California, USA

¹⁹Laboratory for Climate Studies, National Climate Center, China Meteorological Administration, Beijing 100081, China

Received: 31 July 2012 – Accepted: 21 August 2012 – Published: 30 August 2012

Correspondence to: G. Myhre (gunnar.myhre@cicero.uio.no)

Published by Copernicus Publications on behalf of the European Geosciences Union.

ACPD

12, 22355–22413, 2012

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

We report on the AeroCom Phase II direct aerosol effect (DAE) experiment where 15 detailed global aerosol models have been used to simulate the changes in the aerosol distribution over the industrial era. All 15 models have estimated the radiative forcing (RF) of the anthropogenic DAE, and have taken into account anthropogenic sulphate, black carbon (BC) and organic aerosols (OA) from fossil fuel, biofuel, and biomass burning emissions. In addition several models have simulated the DAE of anthropogenic nitrate and anthropogenic influenced secondary organic aerosols (SOA). The model simulated all-sky RF of the DAE from total anthropogenic aerosols has a range from -0.58 to -0.02 W m^{-2} , with a mean of -0.30 W m^{-2} for the 15 models. Several models did not include nitrate or SOA and modifying the estimate by accounting for this with information from the other AeroCom models reduces the range and slightly strengthens the mean. Modifying the model estimates for missing aerosol components and for the time period 1750 to 2010 results in a mean RF for the DAE of -0.39 W m^{-2} . Compared to AeroCom Phase I (Schulz et al., 2006) we find very similar spreads in both total DAE and aerosol component RF. However, the RF of the total DAE is stronger negative and RF from BC from fossil fuel and biofuel emissions are stronger positive in the present study than in the previous AeroCom study. We find a tendency for models having a strong (positive) BC RF to also have strong (negative) sulphate or OA RF. This relationship leads to smaller uncertainty in the total RF of the DAE compared to the RF of the sum of the individual aerosol components. The spread in results for the individual aerosol components is substantial, and can be divided into diversities in burden, mass extinction coefficient (MEC), and normalized RF with respect to AOD. We find that these three factors give similar contributions to the spread in results.

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Global estimates of the direct aerosol effect (DAE) have improved tremendously over the last two decades, due to improvements in processes taken into account and handling of complexities in the aerosol schemes (Bauer et al., 2010; Chin et al., 2009; Seland et al., 2008; Stier et al., 2005; Takemura et al., 2002). The first study of radiative forcing (RF) of the DAE with a global aerosol model included only sulphate aerosols (Charlson et al., 1991). Now multi-model studies with multi aerosol components and including some complex aerosol microphysical schemes have been performed (Schulz et al., 2006).

In the estimate of the RF from the direct aerosol effect from AeroCom Phase I models (Schulz et al., 2006) anthropogenic sulphate, black carbon and organic carbon were considered. In the interim increased attention has been given to the direct aerosol effect of anthropogenic nitrate and secondary organic aerosols (SOA). The formation of ammonium nitrate is complex and is dependent on sufficient ammonia and thus competition with sulphate is important (Adams et al., 2001; Metzger et al., 2002). With the expected reduction in SO₂ emission in the future, nitrate may become more important as a climate forcer (Bauer et al., 2007; Bellouin et al., 2011). Over the last few years increasing consideration has also been paid to secondary organic aerosols (SOA), with advanced understanding of their anthropogenic influence (Hoyle et al., 2011; Robinson et al., 2007). Anthropogenic influence on SOA is now included with varying degrees of complexity in some global aerosol models (Hoyle et al., 2009; Liao and Seinfeld, 2005; Spracklen et al., 2011; Tsigaridis et al., 2006).

The mean RF of the total direct aerosol effect in Schulz et al. (2006) from 9 global aerosol models was -0.22 W m^{-2} . This is much weaker than several observational based estimates (Bellouin et al., 2005; Bellouin et al., 2008; Chung et al., 2005; Quaas et al., 2008). Observational based estimates have been made possible through the advancement of remote sensing from ground and space (Holben et al., 1998; Remer et al., 2008). Myhre (2009) showed that estimates of the direct aerosol effect must

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



rely on global aerosol model estimates, but consistency between models and observational based methods could be reached by accounting for anthropogenic changes in the aerosol optical properties in the latter method.

Various factors lie behind the large range in the global estimates of the DAE, which in AeroCom Phase I was from -0.41 to $+0.04 \text{ W m}^{-2}$ (Schulz et al., 2006). Schulz et al. (2006) showed that the variations in RF could be separated into diversities in aerosol residence time, aerosol mass extinction coefficients, and normalized forcing (RF divided by aerosol optical depth). Normalized forcing was found to be the single largest contributor to the variability. Differences in the aerosol vertical profile, known to be significant between the models, is among the factors that contribute to differences in the normalized RF (Heald et al., 2011; Schwarz et al., 2010; Textor et al., 2006; Zarzycki and Bond, 2010).

In this paper RF of the DAE from 15 recent state-of-the-art global aerosol models in the AeroCom Phase II are presented. Two of the 15 global aerosol models are from Asia, five from Europe and eight from North America. Additional models compared to Schulz et al. (2006) are included in Phase II, and the models have been further developed over the 7–8 yr since AeroCom Phase I. Results for anthropogenic nitrate and SOA are included in the present study. Evaluation of the global aerosol models is an ongoing activity within AeroCom, with comparisons of various measurements being performed (Huneeus et al., 2011; Koch et al., 2009; Koffi et al., 2012). Activities to understand the model differences are also an important part of AeroCom (Randles et al., 2012; Samset et al., 2012; Stier et al., 2012; Textor et al., 2007).

2 Methods

The RF of the total anthropogenic DAE is calculated as the difference between the reflected solar radiation at TOA for simulations with present (2000, or 2006 for a few models) and pre-industrial (1850) emissions of aerosols and their precursors, denoted respectively as CTRL and PRE simulations, but the same cloud and surface conditions.

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The RF has been simulated by 15 global aerosol models with various complexities with regard to number of aerosol species included, aerosol microphysical treatment and their aerosol optical properties. The host model is also of great importance since aerosol transport and removal is dependent on the transport scheme and distribution of precipitation, respectively. In the radiative transfer calculations cloud distribution and properties and surface albedo, among other factors, are important.

Tables 1 and 2 show a list of the models included in the AeroCom Phase II DAE experiment, together with the resolutions used, aerosols components treated and comments on their microphysics scheme. Meteorological data for year 2006 have been used for all the models, except for IMPACT with meteorological data for 1997 and for CAM4-Oslo, NCAR-CAM3.5, and CAM5.1 where nudging possibilities were not available. A protocol describing the data submitted from the global aerosol models are available from the AeroCom web site (<http://aerocom.met.no/aerocomhome.html>). Most of the simulations for this DAE experiment are performed with identical cloud cover and cloud optical properties in PRE and CTRL, i.e. no indirect aerosol effects are included, but one (CAM5.1) estimated the DAE from the CTRL-PRE difference (Δ) in the direct radiative forcing by all aerosols in each simulation (Ghan et al., 2012): $\Delta(S - S_{\text{clean}})$, where S is the net downward solar at TOA calculated with all aerosols present, and S_{clean} is the solar calculated neglecting scattering and absorption by all aerosols.

Aerosol optical properties are treated differently in the global aerosol models but in all models the optical properties are derived using Mie theory; see further description in model references. All models include treatment of sulphate, black carbon, primary organic carbon, sea salt, and mineral dust. Internal mixing between BC and scattering aerosols enhances the absorption from BC (Fuller et al., 1999; Haywood and Shine, 1995), and about half (7) of the 15 models includes some degree of internal mixture for BC. Most of the models have used emissions of aerosols and their precursors from Lamarque et al. (2010). CAM4-Oslo and SPRINTARS have used these IPCC AR5 emissions for PRE (year 1850), but for the CTRL simulations emissions for year 2006 from the AeroCom Phase II dataset has been used (HCA0 v1 or v2 by T. Diehl, see

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



http://aerocom.met.no/emissions.html). This dataset also includes emissions estimates of BC, SO₂ and POM from aviation. BCC used year 2000 emissions, but preindustrial emissions for 1750. GEOS-Chem uses emissions for 2006 which is close to AeroCom emission for 2000 and employs pre-industry emission corresponding to 1750 (Dentener et al., 2006).

If the diagnostics for the different components were difficult to establish, separate model simulations were performed to allow for an attribution to eg biomass burning (BB) or fossil fuel (FF) and bio fuel (BF) sources. For some of the models with internally-mixed aerosols the separation of AOD and RF by aerosol component is challenging. Several groups have solved this by additional radiation calls. Even with this method a few models have been unable to properly extract the component AOD. Where relevant, these results have been removed from the analysis. All simulations were performed such that the cloud fields were identical, e.g. by neglecting the influence of aerosols on clouds or keeping the influence constant.

Only anthropogenic aerosol effects will be treated in the present study, and reference to the DAE apply to the anthropogenic change in the aerosol distribution and its influence on the scattering and absorption of solar radiation. Results will be presented for sulphate, nitrate, BC from fossil fuel and biofuel, OA from fossil fuel and biofuel, BC and OA combined from biomass burning, and SOA.

3 Results

All results are for anthropogenic aerosols only, unless otherwise stated. Anthropogenic aerosols are defined as the difference between the CTRL and PRE simulations described above.

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.1 Total direct aerosol effect

The underlying albedo is crucial for RF of the direct aerosol effect (Haywood and Shine, 1995). Scattering aerosols are efficient at producing a RF over dark surfaces and absorbing aerosols are efficient over bright surfaces. Figure 1a shows the latitudinal variation of the top of the atmosphere (TOA) albedo of the 15 models and the satellite retrieval from CERES (Smith et al., 2011; Wielicki et al., 1996). Note that the albedo at the TOA consists of reflection at the surface and scattering in the atmosphere from clouds, aerosols, and Rayleigh scattering, and thus the scattering may occur above the anthropogenic aerosols. The higher albedo at high latitudes is mainly due to high surface albedo from snow and ice, but larger cloud fraction also contributes. For most of the models the albedo compares well to the CERES data, except between 30° N and 30° S where almost all overestimate the albedo. By inspection of the geographical distribution (not shown) of the albedo the overestimation is largest in regions with low cloud fraction, indicating problems with too large cloud fractions in these regions for the models. The GOCART model has a very low albedo compared to the rest of the models and the CERES data, which is identified to be caused by the meteorological data. The agreement between the simulated latitudinal variation in effective broadband surface albedo (computed from surface level down and up welling short wave radiation fluxes as diagnosed in the models) is in general excellent (see Fig. 1b), except at high latitudes due to differences in snow cover or albedo for snow and for the Northern Hemisphere sea ice coverage and albedo for sea ice.

Figure 2 shows the latitudinal variation in the RF of the total DAE for all sky (Fig. 2a) and clear sky (Fig. 2b) conditions. The negative RF is at maximum around 30° N for most of the models, since the anthropogenic change in aerosol burden (see Fig. 3) has its maximum in this region combined with a relatively high solar radiation flux. GEOS-Chem has the strongest all-sky RF in the Northern Hemisphere, but for clear-sky the OsloCTM2 and IMPACT models show stronger RF there than GEOS-Chem. The RF in the Southern Hemisphere is generally weak. How clouds affect the RF of the direct

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

aerosol effect depends on a variety of factors. Among them are the degree of aerosol absorption, vertical position of aerosols in relation to clouds, and the cloud distribution in general. The all-sky RF is around half in magnitude of the clear sky RF, except for the models with close to zero all sky RF. All models turn over into a positive net RF at high northern latitudes. This is caused by the higher surface and cloud albedo at high latitudes, as can be seen in Fig. 2b. Figure 3c shows that the single scattering albedo of anthropogenic aerosols is rather similar at 30° N and in the Arctic for most of the models, and therefore the surface and cloud albedo is the main cause of the positive RF. CAM4-Oslo has the strongest positive all-sky RF at high latitudes. As can be seen from Fig. 3b and 3c, this is due to having more anthropogenic absorbing aerosols at high latitudes than the other models. This is partly due to the different CTRL emission data sets, which are for year 2006 and have more biomass burning from forest fires than the 2000 emissions. Assumed emission height of biomass burning (up to 6 km) may be another reason. The treatment of convective mixing of aerosols and aerosol precursors probably also leads to somewhat overestimated aerosol burdens in CAM4-Oslo (see Kirkevåg et al., 2012).

Figure 3 shows the latitudinal variation in the anthropogenic AOD (550 nm) used in the radiative transfer calculations, anthropogenic absorbing AOD (550 nm), anthropogenic single scattering albedo (550 nm), and the change in single scattering albedo (550 nm) between pre-industrial (PRE) and current (CTRL). The anthropogenic AOD in Fig. 3a shows a strong hemispheric difference which is a consistent model characteristic. CAM4-Oslo has the highest and CAM5.1 and BCC the lowest anthropogenic AOD almost at all latitudes. The pattern for the anthropogenic absorbing AOD (Fig. 3b) is rather similar in the models, with two peaks, one around the equator where biomass burning emissions of BC are high, and the second around 30° N, where fossil fuel and biofuel emissions of BC are high. The range in the anthropogenic single scattering albedo (defined as the ratio of anthropogenic AOD-AAOD and AOD) (Fig. 3c) is large, with most values lying between 0.9 and 1.0 but even lower values than 0.9 seen at high latitudes for a few models. The lowest anthropogenic single scattering albedo is

simulated near the equator and can be seen in Fig. 3b to be caused by strong anthropogenic absorption AOD that is not associated with a similar strong maximum in the anthropogenic AOD in Fig. 3a. Most models simulate a reduction in the single scattering albedo during the industrial era at all latitudes, except GISS-MATRIX, GOCART and IMPACT. The reduction in the single scattering albedo in the majority of the models is due to stronger growth in absorbing aerosol (BC) than scattering aerosols relative to pre-industrial distribution of absorbing and scattering aerosols. GOCART has an increase in the single scattering albedo in the NH due to low single scattering albedo in the PRE simulation compared to most of the other models. IMPACT shows a weaker increase in single scattering albedo than GOCART, but still positive in the NH. As will be shown later this model has a much larger anthropogenic influenced SOA than the other models, which is likely the cause of the increase in single scattering albedo.

Table 3 summarizes the global and annual mean values for RF, anthropogenic AOD, anthropogenic absorption AOD (AAOD), atmospheric absorption, anthropogenic single scattering albedo, and cloud fraction. The anthropogenic AOD has a mean of 0.031, which is slightly higher than the Schulz et al. (2006) value of 0.029. A main reason for this small increase is likely the inclusion of additional aerosol components in this experiment. The anthropogenic AAOD of 0.0016 is lower than the value of 0.0019 for AAOD attributed to BC in Schulz et al. (2006). However, here it should be mentioned that some of the models in this experiment include absorption by OA which is non-negligible at 550 nm and thus the anthropogenic AAOD is not solely due to BC. A main reason for the lower AAOD compared to Schulz et al. (2006) is likely due to the much smaller anthropogenic BB emissions of BC in a majority of the models using the Lamarque et al. (2010) data. Figure 4 shows the model simulated global and annual mean total RF and the RF for the six aerosol components. The simulated mean RF for the DAE is -0.30 W m^{-2} which is stronger than the mean of -0.22 W m^{-2} found in Schulz et al. (2006). A stronger RF of the DAE in this experiment than in Schulz et al. (2006) is in accordance with the stronger AOD and weaker AAOD as noted above. It can also be noted that the models in this experiment have lower global and annual

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

mean cloud cover (mean of 60 %) than in the models in Schulz et al. (2006) (mean of 63 %). As shown in many previous studies of the DAE the total RF consists of aerosol components causing negative RF and positive RF. Figure 4 shows clearly that this balance varies substantially between the models. Further, it shows that some models have weak RF for the aerosol components (CAM5 and BCC) whereas other models have stronger aerosol component RF (CAM4-Oslo and OsloCTM2).

Not all models have all six aerosol components included. Figure 5 shows the total RF where modifications for missing aerosol components have been taken into account. We calculate the mean of each aerosol component from all models that treat it, then add that mean to the total RF for each component that an individual model does not treat. Since the missing aerosol components in the models are predominantly scattering aerosols (nitrate, SOA, and in some few cases BB aerosols), the modification leads to stronger negative RF for all models with modification. The mean RF of the DAE changes from -0.30 W m^{-2} to -0.36 W m^{-2} with the aerosol component modification.

Table 3 also shows the normalized clear sky RF (NRF, clear sky RF divided by AOD), which ranges from -17 W m^{-2} to -76 W m^{-2} with a mean of -27 W m^{-2} . The NRF depends on a variety of factors such as spatial distribution, optical properties, and the vertical profile. The weakening of the DAE RF due to clouds (Schulz et al., 2006) depends on vertical positioning of the aerosols and the clouds (Chand et al., 2009; Liao and Seinfeld, 1998; Samset and Myhre, 2011) as well as the degree of absorption of the aerosols (single scattering albedo) (Haywood and Shine, 1995).

Atmospheric absorption by the aerosols is calculated as the difference between the RF at TOA and radiative effect of aerosols at the surface. Absorbing aerosols such as BC are the main cause for the atmospheric absorption by the aerosols, but scattering aerosols enhance the absorption by gases in the atmosphere and make a small contribution (Randles et al., 2012; Stier et al., 2012). Figure 6 shows the atmospheric absorption by anthropogenic aerosols as a function of absorbing AOD. The mean atmospheric absorption (see Table 3) is 0.81 W m^{-2} , which is slightly weaker than the 0.82 W m^{-2} reported in Schulz et al. (2006). The normalized values (atmospheric absorption divided

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

et al. (2006). Likewise the range in RF is very similar in the two AeroCom DAE experiments. The global mean MEC is calculated as the ratio of global mean AOD to global mean burden. The MEC depends on various factors, where growth by water uptake for sulphate aerosols is a major cause of the difference between the models. To illustrate the importance of water uptake, the MEC in OsloCTM2 is $4.0 \text{ m}^2 \text{ g}^{-1}$ for low relative humidity (below 30 %) and is $11.2 \text{ m}^2 \text{ g}^{-1}$ as a global mean with ambient relative humidity. The mean of the MEC is $13 \text{ m}^2 \text{ g}^{-1}$, but also without the two models with very high MEC (NCAR-CAM3.5 and GISS modelE) the MEC is $10.3 \text{ m}^2 \text{ g}^{-1}$ and about 10 % higher than in Schulz et al. (2006). The weaker normalized RF with respect to AOD and stronger with respect to burden than in Schulz et al. (2006) could be a result of sulphate at generally lower altitudes, but other factors could contribute.

Figure 8 shows the latitudinal variation in RF, burden, AOD, NRF (with respect to AOD) for sulphate. All models have a maximum in the burden at around 30°N with a large difference in the magnitude. The difference in the magnitude of the burden is particularly large at high northern latitudes and is likely linked to transport and scavenging problems in the models (Rasch et al., 2000). The difference in RF of sulphate at high latitudes is not as evident as for the burden (and AOD), since NRF at high latitudes is weak due to the high surface albedo, and some of the models with strongest burden (and AOD) have weak NRF. The RF of sulphate is very weak in the Southern Hemisphere.

3.3 RF of BC

In Fig. 9 the latitudinal variation is shown for RF, burden, AOD, NRF (with respect to AOD) for BC from FF and BF emissions. Global numbers are shown in Table 5. The importance of surface and cloud albedo is evident in the NRF for BC, especially if it is compared to NRF for sulphate in Fig. 8d, which weakens (for a majority of the models) rather than strengthens over the poles as for BC. The much stronger NRF at high latitudes for BC than sulphate and the relatively large range in burden at high latitudes are also clearly evident for the RF. Schulz et al. (2006) showed the latitudinal

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Radiative forcing of
the direct aerosol
effect from AeroCom
Phase II simulations**

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

variation in burden of BC (total and not only FF and BF as in Fig. 9b). There is no indication of a reduced range in the BC burden among the AeroCom models from Phase I to Phase II. For the maximum in RF at around 30° N there is almost an order of magnitude difference between the model with strongest RF (OsloCTM2) and the model with the weakest RF (BCC). Table 4 shows global mean burden, MEC, AOD, RF and NRF. The model differences in the NRF are larger for BC than for sulphate, and main reason for this is likely a strong dependence on altitude of BC and variations in absorbing properties of BC (Samset and Myhre, 2011; Zarzycki and Bond, 2010). The model mean mass extinction coefficient is $10.5 \text{ m}^2 \text{ g}^{-1}$. Unfortunately the model mass absorption coefficient for BC from FF and BF emissions was not diagnosed. Based on observations, Bond et al. (2006) recommend a mass absorption coefficient (MAC) of around $7.5 \text{ m}^2 \text{ g}^{-1}$ for fresh BC particles and 50 % higher for aged BC. Jacobson (2012) and Chung et al. (2012) use higher absorption enhancement factors. As the single scattering albedo for BC often is in the range of 0.2 to 0.3 (Bond and Bergstrom, 2006), the BC MEC should be around 25 % higher than the MAC, implying the models underestimate the MEC and MAC. The low MEC for a few of the models is the cause of the generally low mean MEC; but half of the models have MEC according to the recommendations. Internal mixing of BC with other aerosol components increases the absorption, so simulating sufficiently high MEC requires representing internal mixing of BC and its effects on absorption. The difference in MEC for models using internal mixture ($12.2 \text{ m}^2 \text{ g}^{-1}$) is on average 45 % higher than for models assuming external mixture ($8.4 \text{ m}^2 \text{ g}^{-1}$).

The model mean global average burden of BC is 0.14 mgm^{-2} and a RF of 0.19 Wm^{-2} . The RF estimate is similar to the IPCC AR4 estimate, but much lower than Ramanathan and Carmichael (2008). However, the estimates here and in IPCC AR4 (Forster et al., 2007) are solely from FF and BF, whereas Ramanathan and Carmichael (2008) included also BC from BB emissions and estimate forcing for all BC, not just anthropogenic. The spread in NRF with respect to burden is larger than in NRF with respect to AOD due to variations in MEC included in the former. For the two NRF mea-

sures the model difference is shown to arise from differences in the vertical profile of BC (Samset et al., 2012), host model difference (Stier et al., 2012) and likely differences in optical properties, especially the single scattering albedo of BC.

3.4 RF of OA

5 The RF of primary OA from FF and BF is weak and has a mean of -0.04 Wm^{-2} . The RF together with burden, MEC, AOD and NRF are shown in Table 6. Primary OA from biomass burning and SOA are described in separate sections below. The burden of OA is on average about 20 % of the sulphate burden; however, this fraction varies between the models. RF from primary OA from FF and BF is around 10 % of the sulphate RF, and thus the NRF with respect to burden for OA is much weaker than for sulphate. One main reason for this is the much smaller water uptake for OA. This is also reflected in the MEC. Due to the smaller water uptake for OA compared to sulphate the particles are smaller, leading to a larger Ångström exponent and a stronger NRF with respect to AOD. Compared to BC from FF, OA has a lower burden in the Arctic (see Fig. 10).
10 Combined with the weak NRF in the Arctic for scattering aerosols, this causes the quite weak RF from OA in this region seen in Fig. 10.

Another cause for the differences in NRF between the models as well as between sulphate and OA are differences in the single scattering albedo for OA. Several models have single scattering albedo for OA at 550 nm at 0.96 with increasing values for higher relative humidity. Some OA components have absorption at short solar wavelength with a reduction in the absorption with wavelength stronger than for BC (Jacobson, 2012; Kanakidou et al., 2005). Uncertainties in the refractive indexes and thus absorption for OA are large. However, some of the major components of OA do not absorb (Kanakidou et al., 2005). OsloCTM2 has the strongest RF for OA from FF and BF emissions and uses pure scattering OA aerosols. One additional simulation was performed with OsloCTM2 using refractive indexes for OA as in CAM4-Oslo, and with a 25 % weakening in the RF of OA from FF and BF emissions. This reduced the NRF with respect to AOD from -28 Wm^{-2} to -21 Wm^{-2} , very close to the mean value. It can also be
25

22369

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



noted that using these refractive indexes for OA from FF and BF emissions as well as for SOA enhanced the total anthropogenic AAOD (at 550 nm) by 6 % compared to the standard OsloCTM2 simulations.

3.5 RF SOA

Five models have explicitly performed RF simulations for SOA; the two GISS-models have SOA included in the total DAE simulations and some information is available for GISS-modelE. The uncertainties in the modelling of SOA are large (Spracklen et al., 2011), and thus it is not surprising that we find a huge range in the burden of anthropogenic SOA from 0.09 to 0.97 mg m⁻² (see Table 7). The burden in IMPACT-Umich is substantially higher than the other models. Figure 11 shows that burden of SOA has a larger fraction around the equator than for the other models. IMPACT-Umich and CAM5.1 have a clearer maximum over latitudes where industrialization is strong, whereas ECHAM5-HAM and OsloCTM2 have the maximum in the RF shifted slightly towards the equator. The zonal pattern of the NRF is rather similar for the models, but the magnitude differs substantially, and one major cause here is assumptions regarding the optical properties and in particular the degree of absorption at short wavelength similar to primary OA discussed above. OsloCTM2 has the strongest NRF and uses pure scattering SOA. CAM5.1 has some positive NRF at very low SOA abundances and this model includes weak absorption of SOA at short wavelengths.

3.6 RF nitrate

Nine models have simulated RF of nitrate aerosols. The resulting range of nitrate burdens has a larger span among the aerosol components than for sulphate and primary BC and OA, ranging from 0.16 to 0.82 mg m⁻² (see Table 8 and Fig. 12). The mean RF is -0.09 Wm⁻² but, like the burden, the RF also varies widely among the AeroCom Phase II models with a range from -0.21 to -0.03 Wm⁻². The pattern of the zonal mean burden of nitrate resembles the sulphate pattern. There is a tendency for models

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

with weak sulphate burden to have a strong nitrate burden and vice versa and to some extent this can be expected since ammonium nitrate can only be formed if there is an excess of ammonia and thus it competes with sulphate (Adams et al., 2001). Although, there are some exceptions to this relationship with some models having both low burden for sulphate and nitrate. The hygroscopic growth for nitrate is generally stronger than for sulphate (Fitzgerald, 1975) so one might expect stronger or at least similar MEC for nitrate compared to sulphate, but this is not the case for GMI and NCAR-CAM3.5. However, regional variation in the aerosols and thus water uptake may influence the results. The NRF with respect to burden and with respect to AOD are rather similar for nitrate and sulphate.

The mean RF of the DAE of nitrate is substantial and with its potentially larger role in the future it is important to investigate the model differences and compare with observations. We encourage studies comparing nitrate surface concentrations and in the atmosphere from aircraft measurements, which over Europe indicate a large role of nitrate (Crosier et al., 2007; Putaud et al., 2010), as well as over ocean where observations show small fractions fine mode nitrate (Quinn and Bates, 2005).

3.7 RF BB

The RF for simulated BB aerosol varies between positive and negative RF, highly dependent on SSA and likely also the altitude of the BB aerosols. The mean RF from the models is close to zero. The BB RF is weak in magnitude, but it is important to note that the RF of BB consists of a positive RF from BC and a negative RF from OA of much larger magnitude than for the total RF from BB aerosols. The MECs for BB aerosols given in Table 9 are in the upper range of what is measured during aerosol campaigns in Africa (Haywood et al., 2003; Johnson et al., 2008). In Schulz et al. (2006) the burden of primary carbonaceous was given separately for BC and OA, but not differentiated by sources. Since RF of OAFF and BCFF are at least not weaker and the total primary carbonaceous burden is lower than in Schulz et al. (2006) this indicates lower BB burden in Phase II compared to Phase I. The model range in burden of BB is large

as illustrated by the large difference between the mean and the median values. Zonal mean RF, burden, AOD and NRF with respect to AOD is shown for BB aerosols in Fig. S1.

4 Discussion

The geographical distributions of the RF of the total DAE and its uncertainty (given as one standard deviation) are shown in Fig. 13. As shown in earlier studies the RF of the DAE is strongly negative in industrialized regions of Asia, Europe and Northern America and reaches positive RF at high latitudes and in other regions with high albedo (either high surface albedo or high cloud cover).

The model mean RF and standard deviation of DAE for sulphate, BC, and OA are shown in Fig. S2. Maximum strength in RF for sulphate, BC, and OA is in Southeast Asia. For sulphate a quite strong RF is also simulated in Europe and Eastern US Over regions with high surface albedo such as Sahara and Arctic the RF for sulphate and OA are quite weak, but RF from BC is relatively strong. The standard deviations are highest in the regions of high RF.

The geographical distribution of RF by nitrate and SOA (Fig. S3) show similar pattern as for the other aerosol components (Fig. S2). For BB aerosols the geographical pattern of RF is very different from the other aerosol components. In some areas (Southeast US) this has been caused by reduction in BB emissions, but in other regions differing signs may be caused by the underlying albedo such as Southern Africa (Chand et al., 2009; Keil and Haywood, 2003), where low albedo over land causes negative RF and large coverage of low lying clouds with BB aerosols above cause positive RF over ocean to the west of the coast. The high surface albedo at high latitudes is the major cause of the positive RF at high latitudes.

Schulz et al. (2006) introduced a way to investigate the contribution to the uncertainty from various key components in estimates of the RF of the direct aerosol effect. The same approach is shown in Fig. 14 for burden, MEC, NRF (with respect to AOD) and

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



RF. For sulphate the uncertainties in RF are slightly stronger than for the burden, MEC, NRF. However, the range for burden, MEC, NRF seems quite similar. CAM5, IMPACT, and BCC have the weakest RF. Weak MEC is shown to be a main reason for this for CAM5 and BCC, whereas for IMPACT burden and NRF are the main factors. OsloCTM2 has the strongest sulphate RF, and has among the largest burden and strongest MEC and NRF, although not the strongest for any of these three.

Whereas Schulz et al. (2006) found that diversity in NRF clearly dominated the diversity in RF of BC, Fig. 14 shows that diversities in burden and MEC are equally important in this study. Illustrative of how differences in burden, MEC, and NRF may compensate each other, the HadGEM2 model has the highest BC burden, but a RF of BC close to the mean, which can be seen to be caused by a rather low MEC. For OA the dominant cause for the spread in RF is differences in MEC, overwhelming the importance of differences in burden and NRF. ECHAM5-HAM has the weakest MEC of $1.6 \text{ m}^2 \text{ g}^{-1}$ and IMPACT the strongest of $14.1 \text{ m}^2 \text{ g}^{-1}$. The weak MEC for ECHAM5-HAM explains why it shows the weakest RF among the global aerosol models, since its burden and NRF are close to mean values. OsloCTM2 has the strongest RF of OA and as for sulphate and BC this is due to strong burden, MEC, and NRF, although not the strongest for any of these three essential variables for the RF.

Global aerosol models with strong RF for sulphate have a tendency to also have strong RF for BC and similar for OA in relation to BC. The RF sulphate and RF BC have an anti-correlation of 0.73 and RF OA and BC an anti-correlation of 0.47. Figure 15 shows the relationship between burden, RF and NRF for sulphate, BC and OA. Aerosol burden is dependent on dry and wet deposition and the transport schemes in the models, and the simulated burdens of various aerosol compounds are therefore correlated. Similarly, there is a tendency also for relationship between the NRF (with respect to burden) for various aerosol compounds. The total DAE which is the combined effect of all aerosol components with both positive RF (BC, and in some cases OA) and negative RF (rest of components) thus also consists of RF contributions that are correlated. Altogether, this leads to smaller range in the total DAE than combining

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the range through error propagation rules for the individual aerosol components. This is illustrated in Fig. 16, which shows the PDFs for the individual aerosol components and their sum, and the modelled total RF. A narrower PDF for the model total DAE than aerosol component sum is evident. Further, Fig. 16 illustrates the smaller model diversity when the aerosol missing component modification is performed compared to the total DAE directly from the various models.

The emission data used in most of the simulations (Lamarque et al., 2010) is generated for the period 1850 to 2000. Anthropogenic emissions of aerosols and their precursors had started in 1850 and emission changes have occurred after 2000. Skeie et al. (2011) performed RF simulations for the different aerosol species for several time intervals for the period 1750 to 2000 based on Lamarque et al. (2010), as well as for 2010 based on the RCP4.5 scenario. GISS model simulations in Shindell et al. (2012) were also performed under the RCP4.5 scenario for years 2000 and 2010. Figure 17 shows a scaling of the RF AeroCom simulation for 1850–2000 using results from Skeie et al. (2011) combined with the GISS values for 2000 to 2010. The scaling is done for all aerosol components, except for BB due to large uncertainties in BB activity between 1750 and 1850 as well as the inhomogeneous RF (see Fig. S3). The strengthening in the RF for 1750–2010 relative to 1850–2000 is particularly strong for OA from FF and BF and nitrate (40–45%), quite important for BC from FF and BF and SOA (25–30%), but rather small for sulphate (5%). For OA from FF and BF, nitrate and SOA the largest strengthening occurs for the period 1750–1850 in comparison to the 2000–2010 period. For BC and sulphate the strengthening is rather similar in the 1750–1850 and 2000–2010 periods. The increase in BC emission from 2000 to 2010 is of similar magnitude in other emission data sets as we have applied in for the scaling (Granier et al., 2011). In Skeie et al. (2011) the increase in RF from BC (from FF and BF emissions) for the time period 2000 to 2010 was slightly larger than the associated increase in emissions due to also changes in the spatial pattern of the emissions. The increase in the RF of the primary carbonaceous aerosols from 2000 to 2010 were very similar in OsloCTM2 and GISS, but somewhat weaker for the secondary components in the

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



GISS model compared to OsloCTM2. This indicates that uncertainties in the scaling are largest for nitrate and SOA which had a quite large change from 1750 to 1850. Taking into account the changes between 1750–1850 and 2000–2010 for the aerosol components changes the total DAE from -0.36 W m^{-2} (modified for missing aerosol components) for 1850–2000 to -0.39 W m^{-2} for the 1750–2010 period.

5 Conclusions

We have documented the RF of the DAE for anthropogenic aerosol emissions from 15 global aerosol models, combined and for sulphate, BC, OA, BB, nitrate and SOA separately. This AeroCom Phase II DAE experiment shows many similarities with the Phase I (Schulz et al., 2006), even though more models are included here and model development has taken place over the intervening years. We summarize here the most important findings:

- The total RF of DAE is stronger negative in the current simulations than in AeroCom Phase I. The main reason for this is addition of new species such as nitrate and SOA.
- Sulphate and BC from FF and BF emissions are the two aerosol components with strongest absolute RF. The RF of the DAE of sulphate is very similar in the simulations presented here and in Schulz et al. (2006). On the other hand BC RF from FF and BF emissions is almost twice as strong compared to in Schulz et al. (2006).
- The global aerosol models have undergone substantial development over the last 7–8 yr, and while there are some changes relative to AeroCom Phase I (Schulz et al., 2006), the present analysis shows that the main estimates in AeroCom Phase I and II are quite robust.

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- 5
- The mean from the 15 models of the RF of the total DAE is -0.30 W m^{-2} . Several of the AeroCom models have not included either nitrate aerosols or SOA or both. Modifying these models with information from AeroCom models altogether results in a mean RF of -0.36 W m^{-2} . The model simulations are to a large extent performed for the period 1850–2000 and using available information to scale to 1750–2010 results in a mean RF of the total DAE of -0.39 W m^{-2} .
- 10
- The AeroCom Phase II simulated 1850–2000 (four models use slightly different time period) mean RF of the DAE for sulphate is -0.35 W m^{-2} , BC from FF and BF emissions is $+0.19 \text{ W m}^{-2}$, OA from FF and BF emissions is -0.04 W m^{-2} , BB is -0.01 W m^{-2} , nitrate is -0.09 W m^{-2} and SOA is -0.07 W m^{-2} . When these numbers were scaled to 1750–2010 numbers the RF for sulphate is -0.37 W m^{-2} , BC from FF and BF emissions is $+0.24 \text{ W m}^{-2}$, OA from FF and BF emissions is -0.05 W m^{-2} , BB is -0.01 W m^{-2} (no change included), nitrate is -0.13 W m^{-2} and SOA is -0.08 W m^{-2} .
- 15
- The differences between models are large for the DAE of the aerosol components, as illustrated by the relative standard deviation of more than 40% for RF of BC and a range from 0.05 to 0.38 W m^{-2} . A further understanding of the causes of the uncertainties is attempted by decomposing the results into diversities due to burden, MEC, and NRF. In general these three factors contribute equally to the uncertainty in the DAE.
- 20
- We find a correlation in the magnitude of RF of BC (positive) and RF of sulphate or OA (negative) among the models, due to transport, deposition, and optical properties being treated similarly in each of the models. We therefore find a smaller uncertainty in the total DAE than the sum of the DAE by aerosol components.

Supplementary material related to this article is available online at:
[http://www.atmos-chem-phys-discuss.net/12/22355/2012/
acpd-12-22355-2012-supplement.pdf](http://www.atmos-chem-phys-discuss.net/12/22355/2012/acpd-12-22355-2012-supplement.pdf).

Acknowledgements. S. Ghan, X. Liu, R. Easter, P. Rasch and J.-H. Yoon were funded by the US Department of Energy, Office of Science, Scientific Discovery through Advanced Computing (SciDAC) Program and by the Office of Science Earth System Modeling Program. Computing resources were provided by the Climate Simulation Laboratory at NCAR's Computational and Information Systems Laboratory (CISL), sponsored by the National Science Foundation and other agencies. The Pacific Northwest National Laboratory is operated for DOE by Battelle Memorial Institute under contract DE-AC06-76RLO 1830. Simulations of the ECHAM5-HAM, INCA, CAM4-Oslo and HadGEM2 models have been supported with funds from the FP6 project EUCAARI (Contract 34684). A. Kirkevåg, T. Iversen and Ø. Seland (CAM4-Oslo) were supported by the Research Council of Norway through the EarthClim (207711/E10) and NO-TUR/NorStore projects, by the Norwegian Space Centre through PM-VRAE, and through the EU projects PEGASOS and ACCESS. G. Myhre and B. Samset were funded by the Research Council of Norway through the EarthClim and SLAC projects.

References

- Adams, P. J., Seinfeld, J. H., Koch, D., Mickley, L., and Jacob, D.: General circulation model assessment of direct radiative forcing by the sulfate-nitrate-ammonium-water inorganic aerosol system, *J. Geophys. Res.-Atmos.*, 106, 1097–1111, 2001.
- Balkanski, Y.: L'Influence des Aérosols sur le Climat, Thèse d'Habilitation à Diriger des 871 Recherches Thesis, Université Versailles Saint Quentin, 2011.
- Balkanski, Y., Schulz, M., Moulin, C., and Ginoux, P.: The formulation of dust emissions on global scale: formulation and validation using satellite retrievals. In: *Emissions of Atmospheric Trace Compounds*, edited by: Granier, C., Artaxo, P., and Reeves, C., Kluwer Academic Publishers, Dordrecht, The Netherlands, 239–267, 2004.

ACPD

12, 22355–22413, 2012

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Bauer, S. E., Koch, D., Unger, N., Metzger, S. M., Shindell, D. T., and Streets, D. G.: Nitrate aerosols today and in 2030: a global simulation including aerosols and tropospheric ozone, *Atmos. Chem. Phys.*, 7, 5043–5059, doi:10.5194/acp-7-5043-2007, 2007.
- 5 Bauer, S. E., Wright, D. L., Koch, D., Lewis, E. R., McGraw, R., Chang, L.-S., Schwartz, S. E., and Ruedy, R.: MATRIX (Multiconfiguration Aerosol TRacker of mIXing state): an aerosol microphysical module for global atmospheric models, *Atmos. Chem. Phys.*, 8, 6003–6035, doi:10.5194/acp-8-6003-2008, 2008.
- Bauer, S. E., Menon, S., Koch, D., Bond, T. C., and Tsigaridis, K.: A global modeling study on carbonaceous aerosol microphysical characteristics and radiative effects, *Atmos. Chem. Phys.*, 10, 7439–7456, doi:10.5194/acp-10-7439-2010, 2010.
- 10 Bellouin, N., Boucher, O., Haywood, J., and Reddy, M. S.: Global estimate of aerosol direct radiative forcing from satellite measurements, *Nature*, 438, 1138–1141, 2005.
- Bellouin, N., Jones, A., Haywood, J., and Christopher, S. A.: Updated estimate of aerosol direct radiative forcing from satellite observations and comparison against the Hadley Centre climate model, *J. Geophys. Res.-Atmos.*, 113, D10205, doi:10.1029/2007JD009385, 2008.
- 15 Bellouin, N., Rae, J., Jones, A., Johnson, C., Haywood, J., and Boucher, O.: Aerosol forcing in the Climate Model Intercomparison Project (CMIP5) simulations by HadGEM2-ES and the role of ammonium nitrate, *J. Geophysical Res.-Atmos.*, 116, D20206, doi:10.1029/2007JD009385, 2011.
- 20 Bian, H., Chin, M., Rodriguez, J. M., Yu, H., Penner, J. E., and Strahan, S.: Sensitivity of aerosol optical thickness and aerosol direct radiative effect to relative humidity, *Atmos. Chem. Phys.*, 9, 2375–2386, doi:10.5194/acp-9-2375-2009, 2009.
- Bond, T. C. and Bergstrom, R. W.: Light absorption by carbonaceous particles: an investigative review, *Aerosol Sci. Tech.*, 40, 27–67, 2006.
- 25 Bond, T. C., Habib, G., and Bergstrom, R. W.: Limitations in the enhancement of visible light absorption due to mixing state, *J. Geophys. Res.-Atmos.*, 111, D20211, doi:10.1029/2006JD007315, 2006.
- Chand, D., Wood, R., Anderson, T. L., Satheesh, S. K., and Charlson, R. J.: Satellite-derived direct radiative effect of aerosols dependent on cloud cover, *Nat. Geosci.*, 2, 181–184, 2009.
- 30 Charlson, R. J., Langner, J., Rodhe, H., Leovy, C. B., and Warren, S. G.: Perturbation of the Northern-Hemisphere radiative balance by backscattering from anthropogenic sulfate aerosols, *Tellus A*, 43, 152–163, 1991.

**Radiative forcing of
the direct aerosol
effect from AeroCom
Phase II simulations**

G. Myhre et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Chin, M., Rood, R. B., Lin, S. J., Muller, J. F., and Thompson, A. M.: Atmospheric sulfur cycle simulated in the global model GOCART: model description and global properties, *J. Geophys. Res.-Atmos.*, 105, 24671–24687, 2000.

Chin, M., Ginoux, P., Kinne, S., Torres, O., Holben, B. N., Duncan, B. N., Martin, R. V., Logan, J. A., Higurashi, A., and Nakajima, T.: Tropospheric aerosol optical thickness from the GOCART model and comparisons with satellite and Sun photometer measurements, *J. Atmos. Sci.*, 59, 461–483, 2002.

Mian Chin, Diehl, T., Dubovik, O., Eck, T. F., Holben, B. N., Sinyuk, A., and Streets, D. G.: Light absorption by pollution, dust, and biomass burning aerosols: a global model study and evaluation with AERONET measurements, *Ann. Geophys.*, 27, 3439–3464, doi:10.5194/angeo-27-3439-2009, 2009.

Chung, C. E., Ramanathan, V., Kim, D., and Podgorny, I. A.: Global anthropogenic aerosol direct forcing derived from satellite and ground-based observations, *J. Geophys. Res.-Atmos.*, 110, D24207, doi:10.1029/2005JD006356, 2005.

Chung, C. E., Lee, K., and Muller, D.: Effect of internal mixture on black carbon radiative forcing, *Tellus B*, 64, 1–13, 2012.

Crosier, J., Allan, J. D., Coe, H., Bower, K. N., Formenti, P., and Williams, P. I.: Chemical composition of summertime aerosol in the Po Valley (Italy), Northern Adriatic and Black Sea, *Q. J. Roy. Meteor. Soc.*, 133, 61–75, 2007.

Feng, Y. and Penner, J. E.: Global modeling of nitrate and ammonium: Interaction of aerosols and tropospheric chemistry, *J. Geophys. Res.-Atmos.*, 112, D01304, doi:10.1029/2005JD006404, 2007.

Fitzgerald, J. W.: Approximation formulas for equilibrium size of an aerosol particle as a function of its dry size and composition and ambient relative humidity, *J. Appl. Meteorol.*, 14., 1044–1049, 1975.

Forster, P., Ramaswamy, V., Artaxo, P., Bernsten, T., Betts, R., Fahey, D. W., Haywood, J., Lean, J., Lowe, D. C., Myhre, G., Nganga, J., Prinn, R., Raga, G., Schulz, M., and Van Dorland, R.: Changes in atmospheric constituents and in radiative forcing, in: *Climate Change 2007: The Physical Science Basis, Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, Cambridge, UK and New York, NY, USA, 2007.

Fuller, K. A., Malm, W. C., and Kreidenweis, S. M.: Effects of mixing on extinction by carbonaceous particles, *J. Geophys. Res.-Atmos.*, 104, 15941–15954, 1999.

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ghan, S. J., Liu, X., Easter, R. C., Zaveri, R., Rasch, P. J. et al.: Toward a minimal representation of aerosols in climate models: Comparative decomposition of aerosol direct, semi-direct and indirect radiative forcing, *J. Climate*, doi:10.1175/JCLI-D-11-00650.1, in press, 2012.

Ginoux, P., Chin, M., Tegen, I., Prospero, J. M., Holben, B., Dubovik, O., and Lin, S. J.: Sources and distributions of dust aerosols simulated with the GOCART model, *J. Geophys. Res.-Atmos.*, 106, 20255–20273, 2001.

Granier, C., Bessagnet, B., Bond, T., D'Angiola, A., van der Gon, H. D., Frost, G. J., Heil, A., Kaiser, J. W., Kinne, S., Klimont, Z., Kloster, S., Lamarque, J. F., Liousse, C., Masui, T., Meleux, F., Mieville, A., Ohara, T., Raut, J. C., Riahi, K., Schultz, M. G., Smith, S. J., Thompson, A., van Aardenne, J., van der Werf, G. R., and van Vuuren, D. P.: Evolution of anthropogenic and biomass burning emissions of air pollutants at global and regional scales during the 1980–2010 period, *Climatic Change*, 109, 163–190, 2011.

Haywood, J. M. and Shine, K. P.: The effect of anthropogenic sulfate and soot aerosol on the clear-sky planetary radiation budget, *Geophys. Res. Lett.*, 22, 603–606, 1995.

Haywood, J. M., Osborne, S. R., Francis, P. N., Keil, A., Formenti, P., Andreae, M. O., and Kaye, P. H.: The mean physical and optical properties of regional haze dominated by biomass burning aerosol measured from the C-130 aircraft during SAFARI 2000, *J. Geophys. Res.-Atmos.*, 108, 8473, doi:10.1029/2002/JD002226, 2003.

Heald, C. L., Coe, H., Jimenez, J. L., Weber, R. J., Bahreini, R., Middlebrook, A. M., Russell, L. M., Jolleys, M., Fu, T.-M., Allan, J. D., Bower, K. N., Capes, G., Crosier, J., Morgan, W. T., Robinson, N. H., Williams, P. I., Cubison, M. J., DeCarlo, P. F., and Dunlea, E. J.: Exploring the vertical profile of atmospheric organic aerosol: comparing 17 aircraft field campaigns with a global model, *Atmos. Chem. Phys.*, 11, 12673–12696, doi:10.5194/acp-11-12673-2011, 2011.

Holben, B. N., Eck, T. F., Slutsker, I., Tanre, D., Buis, J. P., Setzer, A., Vermote, E., Reagan, J. A., Kaufman, Y. J., Nakajima, T., Lavenu, F., Jankowiak, I., and Smirnov, A.: AERONET – a federated instrument network and data archive for aerosol characterization, *Remote Sens. Environ.*, 66, 1–16, 1998.

Hoyle, C. R., Myhre, G., Berntsen, T. K., and Isaksen, I. S. A.: Anthropogenic influence on SOA and the resulting radiative forcing, *Atmos. Chem. Phys.*, 9, 2715–2728, doi:10.5194/acp-9-2715-2009, 2009.

Hoyle, C. R., Boy, M., Donahue, N. M., Fry, J. L., Glasius, M., Guenther, A., Hallar, A. G., Huff Hartz, K., Petters, M. D., Petäjä, T., Rosenoern, T., and Sullivan, A. P.: A review of the

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)




[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

anthropogenic influence on biogenic secondary organic aerosol, *Atmos. Chem. Phys.*, 11, 321–343, doi:10.5194/acp-11-321-2011, 2011.

Huneus, N., Schulz, M., Balkanski, Y., Griesfeller, J., Prospero, J., Kinne, S., Bauer, S., Boucher, O., Chin, M., Dentener, F., Diehl, T., Easter, R., Fillmore, D., Ghan, S., Ginoux, P., Grini, A., Horowitz, L., Koch, D., Krol, M. C., Landing, W., Liu, X., Mahowald, N., Miller, R., Morcrette, J.-J., Myhre, G., Penner, J., Perlwitz, J., Stier, P., Takemura, T., and Zender, C. S.: Global dust model intercomparison in AeroCom phase I, *Atmos. Chem. Phys.*, 11, 7781–7816, doi:10.5194/acp-11-7781-2011, 2011.

Jacobson, M. Z.: Investigating cloud absorption effects: global absorption properties of black carbon, tar balls, and soil dust in clouds and aerosols, *J. Geophys. Res.-Atmos.*, 117, D06205, doi:10.1029/2011JD017218, 2012.

Johnson, B. T., Osborne, S. R., Haywood, J. M., and Harrison, M. A. J.: Aircraft measurements of biomass burning aerosols over West Africa during DABEX, *J. Geophys. Res.-Atmos.*, 113, D00C06, doi:10.1029/2007JD009451, 2008.

Kanakidou, M., Seinfeld, J. H., Pandis, S. N., Barnes, I., Dentener, F. J., Facchini, M. C., Van Dingenen, R., Ervens, B., Nenes, A., Nielsen, C. J., Swietlicki, E., Putaud, J. P., Balkanski, Y., Fuzzi, S., Horth, J., Moortgat, G. K., Winterhalter, R., Myhre, C. E. L., Tsigaridis, K., Vignati, E., Stephanou, E. G., and Wilson, J.: Organic aerosol and global climate modelling: a review, *Atmos. Chem. Phys.*, 5, 1053–1123, doi:10.5194/acp-5-1053-2005, 2005.

Keil, A. and Haywood, J. M.: Solar radiative forcing by biomass burning aerosol particles during SAFARI 2000: A case study based on measured aerosol and cloud properties, *J. Geophys. Res.-Atmos.*, 108, 8467, doi:10.1029/2002JD002315, 2003.

Kirkevåg, A., Iversen, T., Seland, Ø., Hoose, C., Kristjansson, J. E., Struthers, H., Ekman, A. M. L., Ghan, S., Griesfeller, J., Nilsson, E. D., and Schulz, M.: Aerosol-climate interactions in the Norwegian Earth System model – Nor ESM, *Geosci. Model Dev.*, submitted, 2012.

Koch, D., Schmidt, G. A., and Field, C. V.: Sulfur, sea salt, and radionuclide aerosols in GISS ModelE, *J. Geophys. Res.-Atmos.*, 111, D06206, doi:10.1029/2004JD005550, 2006.

Koch, D., Bond, T. C., Streets, D., Unger, N., and van der Werf, G. R.: Global impacts of aerosols from particular source regions and sectors, *J. Geophys. Res.-Atmos.*, 112, D02205, doi:10.1029/2005JD007024, 2007.

Koch, D., Schulz, M., Kinne, S., McNaughton, C., Spackman, J. R., Balkanski, Y., Bauer, S., Berntsen, T., Bond, T. C., Boucher, O., Chin, M., Clarke, A., De Luca, N., Dentener, F., Diehl, T., Dubovik, O., Easter, R., Fahey, D. W., Feichter, J., Fillmore, D., Freitag, S., Ghan, S.,

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Ginoux, P., Gong, S., Horowitz, L., Iversen, T., Kirkevåg, A., Klimont, Z., Kondo, Y., Krol, M., Liu, X., Miller, R., Montanaro, V., Moteki, N., Myhre, G., Penner, J. E., Perlwitz, J., Pitari, G., Reddy, S., Sahu, L., Sakamoto, H., Schuster, G., Schwarz, J. P., Seland, Ø., Stier, P., Takegawa, N., Takemura, T., Textor, C., van Aardenne, J. A., and Zhao, Y.: Evaluation of black carbon estimations in global aerosol models, *Atmos. Chem. Phys.*, 9, 9001–9026, doi:10.5194/acp-9-9001-2009, 2009.

Koffi, B., Schulz, M., Breon, F. M., Griesfeller, J., Winker, D., Balkanski, Y., Bauer, S., Bernsten, T., Chin, M. A., Collins, W. D., Dentener, F., Diehl, T., Easter, R., Ghan, S., Ginoux, P., Gong, S. L., Horowitz, L. W., Iversen, T., Kirkevåg, A., Koch, D., Krol, M., Myhre, G., Stier, P., and Takemura, T.: Application of the CALIOP layer product to evaluate the vertical distribution of aerosols estimated by global models: AeroCom phase I results, *J. Geophys. Res.-Atmos.*, 117, D10201, doi:10.1029/2011JD016858, 2012.

Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, *Atmos. Chem. Phys.*, 10, 7017–7039, doi:10.5194/acp-10-7017-2010, 2010.

Lamarque, J.-F., Emmons, L. K., Hess, P. G., Kinnison, D. E., Tilmes, S., Vitt, F., Heald, C. L., Holland, E. A., Lauritzen, P. H., Neu, J., Orlando, J. J., Rasch, P. J., and Tyndall, G. K.: CAM-chem: description and evaluation of interactive atmospheric chemistry in the Community Earth System Model, *Geosci. Model Dev.*, 5, 369–411, doi:10.5194/gmd-5-369-2012, 2012.

Liao, H. and Seinfeld, J. H.: Effect of clouds on direct aerosol radiative forcing of climate, *J. Geophys. Res.-Atmos.*, 103, 3781–3788, 1998.

Liao, H. and Seinfeld, J. H.: Global impacts of gas-phase chemistry-aerosol interactions on direct radiative forcing by anthropogenic aerosols and ozone, *J. Geophys. Res.*, 110, D18208, doi:10.1029/2005JD005907, 2005.

Lin, G., Penner, J. E., Sillman, S., Taraborrelli, D., and Lelieveld, J.: Global modeling of SOA formation from dicarbonyls, epoxides, organic nitrates and peroxides, *Atmos. Chem. Phys.*, 12, 4743–4774, doi:10.5194/acp-12-4743-2012, 2012.

Liu, X., Easter, R. C., Ghan, S. J., Zaveri, R., Rasch, P., Shi, X., Lamarque, J.-F., Gettelman, A., Morrison, H., Vitt, F., Conley, A., Park, S., Neale, R., Hannay, C., Ekman, A. M. L., Hess, P., Mahowald, N., Collins, W., Iacono, M. J., Bretherton, C. S., Flanner, M. G., and

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Mitchell, D.: Toward a minimal representation of aerosols in climate models: description and evaluation in the Community Atmosphere Model CAM5, *Geosci. Model Dev.*, 5, 709–739, doi:10.5194/gmd-5-709-2012, 2012.

Ma, X., Yu, F., and Luo, G.: Aerosol direct radiative forcing based on GEOS-Chem-APM and uncertainties, *Atmos. Chem. Phys.*, 12, 5563–5581, doi:10.5194/acp-12-5563-2012, 2012.

Metzger, S., Dentener, F., Krol, M., Jeuken, A., and Lelieveld, J.: Gas/aerosol partitioning – 2. Global modeling results, *J. Geophys. Res.-Atmos.*, 107, 4313, doi:10.1029/2001JD001103, 2002.

Myhre, G.: Consistency between satellite-derived and modeled estimates of the direct aerosol effect, *Science*, 325, 187–190, 2009.

Myhre, G., Bellouin, N., Berglen, T. F., Berntsen, T. K., Boucher, O., Grini, A., Isaksen, I. S. A., Johnsrud, M., Mishchenko, M. I., Stordal, F., and Tanré, D.: Comparison of the radiative properties and direct radiative effect of aerosols from a global aerosol model and remote sensing data over ocean, *Tellus B*, 59, 115–129, 2007.

Myhre, G., Berglen, T. F., Johnsrud, M., Hoyle, C. R., Berntsen, T. K., Christopher, S. A., Fahey, D. W., Isaksen, I. S. A., Jones, T. A., Kahn, R. A., Loeb, N., Quinn, P., Remer, L., Schwarz, J. P., and Yttri, K. E.: Modelled radiative forcing of the direct aerosol effect with multi-observation evaluation, *Atmos. Chem. Phys.*, 9, 1365–1392, doi:10.5194/acp-9-1365-2009, 2009.

Putaud, J. P., Van Dingenen, R., Alastuey, A., Bauer, H., Birmili, W. et al.: A European aerosol phenomenology-3: Physical and chemical characteristics of particulate matter from 60 rural, urban, and kerbside sites across Europe, *Atmos. Environ.*, 44, 1308–1320, 2010.

Quaas, J., Boucher, O., Bellouin, N., and Kinne, S.: Satellite-based estimate of the direct and indirect aerosol climate forcing, *J. Geophys. Res.-Atmos.*, 113, D05204, doi:10.1029/2007jd008962, 2008.

Quinn, P. K. and Bates, T. S.: Regional aerosol properties: Comparisons of boundary layer measurements from ACE 1, ACE 2, aerosols99, INDOEX, ACE asia, TARFOX, and NEAQS, *J. Geophys. Res.-Atmos.*, 110, D14202, doi:10.1029/2004JD004755, 2005.

Ramanathan, V. and Carmichael, G.: Global and regional climate changes due to black carbon, *Nat. Geosci.*, 1, 221–227, 2008.

Randles, C. A., Kinne, S., Myhre, G., Schulz, M., Stier, P., Fischer, J., Doppler, L., Highwood, E., Ryder, C., Harris, B., Huttunen, J., Ma, Y., Pinker, R. T., Mayer, B., Neubauer, D., Hitzenberger, R., Oreopoulos, L., Lee, D., Pitari, G., Genova, G. D., Rose, F. G., Kato, S., Rumbold,

**Radiative forcing of
the direct aerosol
effect from AeroCom
Phase II simulations**

G. Myhre et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

S. T., Vardavas, I., Hatzianastassiou, N., Matsoukas, C., Yu, H., Zhang, F., Zhang, H., and Lu, P.: Intercomparison of shortwave radiative transfer schemes in global aerosol modeling: Results from the AeroCom Radiative Transfer Experiment, submitted to Atmos. Chem. Phys., 2012.

5 Remer, L. A., Kleidman, R. G., Levy, R. C., Kaufman, Y. J., Tanre, D., Mattoo, S., Martins, J. V., Ichoku, C., Koren, I., Yu, H. B., and Holben, B. N.: A comparison of scavenging and deposition processes in global models: results from the WCRP Cambridge Workshop of 1995, Tellus B, 52, 1025–1056, 2000.

10 Remer, L. A., Kleidman, R. G., Levy, R. C., Kaufman, Y. J., Tanré, D., Mattoo, S., Martins, J. V., Ichoku, C., Koren, I., Yu, H. B., and Holben, B. N.: Global aerosol Global aerosol climatology from the MODIS satellite sensors, J. Geophys. Res.-Atmos., 113, D14S07, doi:10.1029/2007JD009661, 2008.

15 Robinson, A. L., Donahue, N. M., Shrivastava, M. K., Weitkamp, E. A., Sage, A. M., Grieshop, A. P., Lane, T. E., Pierce, J. R., and Pandis, S. N.: Rethinking organic aerosols: Semivolatile emissions and photochemical aging, Science, 315, 1259–1262, 2007.

Samset, B. H. and Myhre, G.: Vertical dependence of black carbon, sulphate and biomass burning aerosol radiative forcing, Geophys. Res. Lett., 38, L24802, doi:10.1029/2011GL049697, 2011.

20 Samset, B. H., Myhre, G., Schulz, M., Balkanski, Y., Bauer, S., Bellouin, N., Berntsen, T. K., Chin, M., Diehl, T., Easter, R. E., Ghan, S. J., Iversen, T., Kirkevåg, A., Lamarque, J.-F., Lin, G., Penner, J., Seland, Ø., Skeie, R. B., Stier, P., Takemura, T., Tsigaridis, K., and Zhang, K.: Black carbon vertical profiles strongly affect its radiative forcing uncertainty, submitted, 2012.

Schulz, M.: Constraining model estimates of the aerosol radiative forcing, Habilitation Thesis Thesis, Université Pierre et Marie Curie, Paris VI, 2007.

25 Schulz, M., Textor, C., Kinne, S., Balkanski, Y., Bauer, S., Berntsen, T., Berglen, T., Boucher, O., Dentener, F., Guibert, S., Isaksen, I. S. A., Iversen, T., Koch, D., Kirkevåg, A., Liu, X., Montanaro, V., Myhre, G., Penner, J. E., Pitari, G., Reddy, S., Seland, Ø., Stier, P., and Takemura, T.: Radiative forcing by aerosols as derived from the AeroCom present-day and pre-industrial simulations, Atmos. Chem. Phys., 6, 5225–5246, doi:10.5194/acp-6-5225-2006, 2006.

30 Schwarz, J. P., Spackman, J. R., Gao, R. S., Watts, L. A., Stier, P. et al.: Global-scale black carbon profiles observed in the remote atmosphere and compared to models, Geophys. Res. Lett., 37, L18812, doi:10.1029/2010GL044372, 2010.

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)




[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Seland, Ø., Iversen, T., Kirkevåg, A. and Storelvmo, T.: Aerosol-climate interactions in the CAM-Oslo atmospheric GCM and investigation of associated basic shortcomings, *Tellus A*, 60, 459–491, 2008.

Shindell, D. T., Lamarque, J.-F., Schulz, M., Flanner, M., Jiao, C., Chin, M., Young, P., Lee, Y. H., Rotstayn, L., Milly, G., Faluvegi, G., Balkanski, Y., Collins, W. J., Conley, A. J., Dalsoren, S., Easter, R., Ghan, S., Horowitz, L., Liu, X., Myhre, G., Nagashima, T., Naik, V., Rumbold, S., Skeie, R., Sudo, K., Szopa, S., Takemura, T., Voulgarakis, A., and Yoon, J.-H.: Radiative forcing in the ACCMIP historical and future climate simulations, submitted to *Atmos. Chem. Phys.*, 2012.

Skeie, R. B., Berntsen, T. K., Myhre, G., Tanaka, K., Kvalevåg, M. M., and Hoyle, C. R.: Anthropogenic radiative forcing time series from pre-industrial times until 2010, *Atmos. Chem. Phys.*, 11, 11827–11857, doi:10.5194/acp-11-11827-2011, 2011.

Smith, G. L., Priestley, K. J., Loeb, N. G., Wielicki, B. A., Charlock, T. P., Minnis, P., Doelling, D. R., and Rutan, D. A.: Clouds and Earth Radiant Energy System (CERES), a review: past, present and future, *Adv. Space Res.*, 48, 254–263, 2011.

Spracklen, D. V., Jimenez, J. L., Carslaw, K. S., Worsnop, D. R., Evans, M. J., Mann, G. W., Zhang, Q., Canagaratna, M. R., Allan, J., Coe, H., McFiggans, G., Rap, A., and Forster, P.: Aerosol mass spectrometer constraint on the global secondary organic aerosol budget, *Atmos. Chem. Phys.*, 11, 12109–12136, doi:10.5194/acp-11-12109-2011, 2011.

Stier, P., Feichter, J., Kinne, S., Kloster, S., Vignati, E., Wilson, J., Ganzeveld, L., Tegen, I., Werner, M., Balkanski, Y., Schulz, M., Boucher, O., Minikin, A., and Petzold, A.: The aerosol-climate model ECHAM5-HAM, *Atmos. Chem. Phys.*, 5, 1125–1156, doi:10.5194/acp-5-1125-2005, 2005.

Stier, P., Schutgens, N. A. J., Bellouin, N., Bian, H., Boucher, O., Chin, M., Ghan, S., Huneus, N., Kinne, S., Lin, G., Myhre, G., Penner, J. E., Randles, C., Samset, B., Schulz, M., Yu, H., and Zhou, C.: Host model uncertainties in aerosol radiative forcing estimates: results from the AeroCom prescribed intercomparison study, submitted to *Atmos. Chem Phys.*, 2012.

Szopa, S., Balkanski, Y., Cozic, A., Déandreis, C., Dufresne, J.-L., Hauglustaine, D., Lathière, J., Schulz, M., Vuolo, R., Yan, N., Marchand, M., Bekki, S., Cugnet, D., Idelkadi, A., Denvil, S., Foujols, M.-A., Caubel, A., and Fortems-Cheiney, A.: Aerosol and ozone changes as forcing for climate evolution between 1850 and 2100, *Clim. Dynam.*, in press, 2012.

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Takemura, T., Nakajima, T., Dubovik, O., Holben, B. N., and Kinne, S.: Single-scattering albedo and radiative forcing of various aerosol species with a global three-dimensional model, *J. Climate*, 15, 333–352, 2002.

Takemura, T., Nozawa, T., Emori, S., Nakajima, T. Y. and Nakajima, T.: Simulation of climate response to aerosol direct and indirect effects with aerosol transport-radiation model, *J. Geophys. Res.-Atmos.*, 110, D02202, doi:10.1029/2004JD005029, 2005.

Takemura, T., Egashira, M., Matsuzawa, K., Ichijo, H., O'ishi, R., and Abe-Ouchi, A.: A simulation of the global distribution and radiative forcing of soil dust aerosols at the Last Glacial Maximum, *Atmos. Chem. Phys.*, 9, 3061–3073, doi:10.5194/acp-9-3061-2009, 2009.

Textor, C., Schulz, M., Guibert, S., Kinne, S., Balkanski, Y., Bauer, S., Berntsen, T., Berglen, T., Boucher, O., Chin, M., Dentener, F., Diehl, T., Easter, R., Feichter, H., Fillmore, D., Ghan, S., Ginoux, P., Gong, S., Grini, A., Hendricks, J., Horowitz, L., Huang, P., Isaksen, I., Iversen, I., Kloster, S., Koch, D., Kirkevåg, A., Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Liu, X., Montanaro, V., Myhre, G., Penner, J., Pitari, G., Reddy, S., Seland, Ø., Stier, P., Takemura, T., and Tie, X.: Analysis and quantification of the diversities of aerosol life cycles within AeroCom, *Atmos. Chem. Phys.*, 6, 1777–1813, doi:10.5194/acp-6-1777-2006, 2006.

Textor, C., Schulz, M., Guibert, S., Kinne, S., Balkanski, Y., Bauer, S., Berntsen, T., Berglen, T., Boucher, O., Chin, M., Dentener, F., Diehl, T., Feichter, J., Fillmore, D., Ginoux, P., Gong, S., Grini, A., Hendricks, J., Horowitz, L., Huang, P., Isaksen, I. S. A., Iversen, T., Kloster, S., Koch, D., Kirkevåg, A., Kristjansson, J. E., Krol, M., Lauer, A., Lamarque, J. F., Liu, X., Montanaro, V., Myhre, G., Penner, J. E., Pitari, G., Reddy, M. S., Seland, Ø., Stier, P., Takemura, T., and Tie, X.: The effect of harmonized emissions on aerosol properties in global models – an AeroCom experiment, *Atmos. Chem. Phys.*, 7, 4489–4501, doi:10.5194/acp-7-4489-2007, 2007.

Tsigaridis, K., Krol, M., Dentener, F. J., Balkanski, Y., Lathière, J., Metzger, S., Hauglustaine, D. A., and Kanakidou, M.: Change in global aerosol composition since preindustrial times, *Atmos. Chem. Phys.*, 6, 5143–5162, doi:10.5194/acp-6-5143-2006, 2006.

Vignati, E., Wilson, J., and Stier, P.: M7: an efficient size-resolved aerosol microphysics module for large-scale aerosol transport models, *J. Geophys. Res.-Atmos.*, 109, D22202, doi:10.1029/2003JD004485, 2004.

Wang, M. and Penner, J. E.: Aerosol indirect forcing in a global model with particle nucleation, *Atmos. Chem. Phys.*, 9, 239–260, doi:10.5194/acp-9-239-2009, 2009.

**Radiative forcing of
the direct aerosol
effect from AeroCom
Phase II simulations**

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Wielicki, B. A., Barkstrom, B. R., Harrison, E. F., Lee, R. B., Smith, G. L., and Cooper, J. E.: Clouds and the earth's radiant energy system (CERES): an earth observing system experiment, *B. Am. Meteorol. Soc.*, 77, 853–868, 1996.
- 5 Xu, L. and Penner, J. E.: Global simulations of nitrate and ammonium aerosols and their radiative effects, *Atmos. Chem. Phys. Discuss.*, 12, 10115–10179, doi:10.5194/acpd-12-10115-2012, 2012.
- Yu, F.: A secondary organic aerosol formation model considering successive oxidation aging and kinetic condensation of organic compounds: global scale implications, *Atmos. Chem. Phys.*, 11, 1083–1099, doi:10.5194/acp-11-1083-2011, 2011.
- 10 Yu, F. and Luo, G.: Simulation of particle size distribution with a global aerosol model: contribution of nucleation to aerosol and CCN number concentrations, *Atmos. Chem. Phys.*, 9, 7691–7710, doi:10.5194/acp-9-7691-2009, 2009.
- Zarzycki, C. M. and Bond, T. C.: How much can the vertical distribution of black carbon affect its global direct radiative forcing?, *Geophys. Res. Lett.*, 37, L20807, doi:10.1029/2010GL044555, 2010.
- 15 Zhang, H., Wang, Z. L., Wang, Z. Z., Liu, Q. X., Gong, S. L., Zhang, X. Y., Shen, Z. P., Lu, P., Wei, X. D., Che, H. Z., and Li, L.: Simulation of direct radiative forcing of aerosols and their effects on East Asian climate using an interactive AGCM-aerosol coupled system, *Clim. Dynam.*, 38, 1675–1693, 2012a.
- 20 Zhang, K., O'Donnell, D., Kazil, J., Stier, P., Kinne, S., Lohmann, U., Ferrachat, S., Croft, B., Quaas, J., Wan, H., Rast, S., and Feichter, J.: The global aerosol-climate model ECHAM-HAM, version 2: sensitivity to improvements in process representations, *Atmos. Chem. Phys. Discuss.*, 12, 7545–7615, doi:10.5194/acpd-12-7545-2012, 2012b.

Table 1. Model description, general information. If meteorology was nudged or driven by re-analysis fields, the year 2006 meteorology was used.

Model	Type	Resolution	Levels	Meteorology	Responsible
BCC	GCM	2.8° × 2.8°	26	NCEP/NCAR reanalysis	Hua Zhang Zhili Wang
CAM4-Oslo	GCM	2.5° × 1.8°	26	Produced by CAM4 atmospheric physics with CAM4-Oslo cloud tuning and boundary data from the data ocean and sea-ice models of CCSM4.	Alf Kirkevåg Trond Iversen Øyvind Seland
CAM5.1	GCM	2.5° × 1.8°	30	CAM5.1	X. Liu R. C. Easter Steve Ghan P. J. Rasch J.-H. Yoon
GEOS.Chem	CTM	5.0° × 4.0°	47	GEOS-5, reanalysis, nudged	Fangqun Yu Gan Luo Xiaoyan Ma
GISS-MATRIX	GCM	2.5° × 2.0°	40	Nudged to NCEP winds	Susanne Bauer Kostas Tsigaridis
GISS-modelE	GCM	2.5° × 2.0°	40	Nudged to NCEP winds	Kostas Tsigaridis Susanne Bauer
GMI	CTM	2.5° × 2.0°	72	GEOS-5 MERRA reanalysis for 2006, nudged	Huisheng Bian Hongbin Yu
GOCART	CTM	2.5° × 2.0°	30	GEOS-4 DAS (Goddard Earth Observing System version 4 Data Assimilation System), reanalysis for year 2006	Thomas Diehl Mian Chin
IMPACT	CTM	5.0° × 4.0°	46	DAO assimilation fields for 1997, reanalysis.	Guangxing Lin Joyce Penner Li Xu Cheng Zhou
INCA	GCM	3.8° × 1.9°	19	ECMWF reanalysis from the Integrated Forecast System (IFS) model for year 2006.	Yves Balkanski Michael Schulz Didier Hauglustaine
HadGEM2	GCM	1.8° × 1.2°	38	ERA Interim data for 2006, nudged	Nicolas Bellouin
ECHAM5-HAM	GCM	1.8° × 1.8°	31	Model nudged with ECMWF analysis for the year 2006	Kai Zhang Philip Stier Johann Feichter
NCAR-CAM3.5	GCM	2.5° × 1.9°	26	GCM-generated	Jean-Francois Lamarque
OsloCTM2	CTM	2.8° × 2.8°	60	ECMWF reanalysis from the Integrated Forecast System (IFS) model for year 2006	Gunnar Myhre Ragnhild B. Skeie Terje Berntsen
SPRINTARS	GCM	1.1° × 1.1°	56	NCEP/NCAR reanalysis (temperature and horizontal wind), nudged.	Toshihiko Takemura

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 2. Model description, aerosol information.

Model	S	BC	OC	BB	SOA	NO ₃	Aerosol microphysics	References for aerosol module
BCC	Y	Y	Y	Y	–	–	12 bin sizes for each aerosol with radii between 0.005–0.01, 0.01–0.02, 0.02–0.04, 0.04–0.08, 0.08–0.16, 0.16–0.32, 0.32–0.64, 0.64–1.28, 1.28–2.56, 2.56–5.12, 5.12–10.24, and 10.24–20.48 μm .	Zhang et al. (2012a)
CAM4-Oslo	Y	Y	Y	Y	–	–	Mass conc. of SO ₄ , BC, OM, sea-salt and dust in four size-classes are tagged according to production mechanism. Based on 44 sectional size bins and lognormal distributions at the point of emission, look-up tables yield physical properties of the processed aerosols.	Kirkevåg et al. (2012)
CAM5.1	Y	Y	Y	Y	Y	–	3 internally-mixed log-normal modes	Liu et al. (2012)
GEOS_Chem	Y	Y	Y	–	Y	Y	40 bins for secondary particles, 20 bins for sea salt, 15 bins for dust, 4 log-normal modes for BC and primary OC. Coating of primary particles by secondary species tracked. Extended SOA formation considering oxidation aging.	Yu and Luo (2009), Yu (2011), Ma et al. (2012)
GISS-MATRIX	Y	Y	Y	Y	–	Y	Aerosol microphysical scheme	Bauer et al. (2010; 2008)
GISS-modelE	Y	Y	Y	Y	Y	Y	Aerosol mass based scheme	Koch et al. (2007; 2006), Bauer et al. (2007), Tsigaridis et al. (2012)
GMI	Y	Y	Y	Y	–	Y	5 bin sizes for dust, 4 bin sizes for sea-salt, 3 bin size for nitrate and sulfate, all aerosols with log-normal size distributions.	Bian et al. (2009)
GOCART	Y	Y	Y	Y	Y	–	Parameterized with prescribed dry particle sizes: 8 bins for dust, 4 bins for sea salt, 1 bin for sulfate, BC, and OA, with log-normal distributions, particle growth parameterized as a function of RH	Chin et al. (2009; 2002; 2000), Ginoux et al. (2001)
IMPACT	Y	Y	Y	Y	Y	*	4 bin sizes for sea-salt and mineral dust, pure sulfate treated using 2 modes with predicted size and coagulation and condensation of SO ₄ with other aerosols explicitly resolved.	Wang and Penner (2009), Lin et al. (2012); Xu and Penner (2012)
INCA	Y	Y	Y	Y	–	Y	Soluble and insoluble aerosol treated separately, modal assumptions with log-normal size distributions. We distinguish between accumulation, coarse and super coarse modes.	Balkanski et al. (2004), Schulz (2007), Balkanski, (2011), Szopa et al. (2012)
HadGEM2	Y	Y	Y	Y	–	Y	Component aerosol mass transported in Aitken, accumulation, coarse, and dissolved modes. Size distributions assumed lognormal for interaction with radiation.	Bellouin et al. (2011)
ECHAM5-HAM	Y	Y	Y	Y	Y	–	Modal method, log-normal size distributions, 7 modes (4 soluble, 3 insoluble). Internal mixing is assumed for aerosol compositions considered in each mode, while external mixing is assumed among different aerosol modes.	Vignati et al. (2004), Stier et al. (2005), Zhang et al. (2012b)
NCAR-CAM3.5	Y	Y	Y	Y	–	Y	Bulk-aerosol model, except 4-bins for sea-salt and mineral dust	Lamarque et al. (2012)
OsloCTM2	Y	Y	Y	Y	Y	Y	8 bin sizes for sea-salt and mineral dust, aerosol mass scheme for other aerosols with log-normal size distributions in calculations of optical properties	Myhre et al. (2007; 2009), Skeie et al. (2011)
SPRINTARS	Y	Y	Y	Y	–	–	6 bins for dust, 4 bins for sea salt, 1 bin for sulfate, BC, and OA, with log-normal size distributions and particle growth as a function of relative humidity	Takemura et al. (2009; 2005)

Note: GOCART only includes SOA from biogenic sources (terpene oxidation)

* The NO₃ values for forcing from this model were simulated using the same model as the IMPACT model described here, but did not include the chemistry of formation of SOA and used the simplified NO_x chemistry described in Feng and Penner (2007). The resolution was 2.5 × 2.0.

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Table 3. Global mean anthropogenic value for all-sky and clear sky RF, normalized RF (NRF) with respect to AOD for clear sky, atmospheric absorption, atmospheric absorption divided by AAOD, AOD, anthropogenic fraction of AOD, AAOD, single scattering albedo (SSA), combined natural and anthropogenic change in SSA from PRE simulation to CTRL simulation, and present day cloud fraction (CLT).

Model	RF All-sky (W m^{-2})	RF Clear-sky (W m^{-2})	NRF Clear-sky (W m^{-2})	Atm.abs. (W m^{-2})	Atm.abs./AAOD (W m^{-2})	AOD (1)	AOD Ant.fr. (1)	AAOD (1)	SSA (1)	dSSA (1)	CLT (1)
BCC	-0.18	-0.75	-76.0	0.20	561	0.0099	0.138	0.0004	0.963	-0.0007	0.59
CAM4-Oslo	-0.08			1.75	479	0.0527	0.345	0.0037	0.931	-0.0148	0.54
CAM5.1	-0.016	-0.35	-23.6	0.69	470	0.0148	0.123	0.0015	0.901	-0.0064	0.64
GEOS_Chem	-0.49	-0.69	-22.0	0.72	400	0.0314		0.0018	0.943	-0.0106	0.59
GISS-MATRIX	-0.58	-0.79	-19.9			0.0398	0.229	0.0018	0.955	-0.0005	0.65
GISS-modelE	-0.32	-0.46	-20.9			0.0219	0.147	0.0020	0.907	-0.0096	0.65
GMI	-0.52	-0.91	-24.7	0.49	387	0.0368	0.271	0.0013	0.965	-0.0033	
GOCART	-0.36	-0.58	-21.8	0.73	432	0.0267	0.236	0.0017	0.937	0.0005	
HadGEM2	-0.31	-0.72	-27.2	0.61	429	0.0265	0.209	0.0014	0.947	-0.0073	0.55
IMPACT-Umich	-0.21	-1.01	-23.7	1.10	935	0.0428	0.325	0.0012	0.973	-0.0014	0.66
INCA	-0.36	-0.73	-17.4	0.95	723	0.0417	0.295	0.0013	0.968	-0.0046	0.47
ECHAM5-HAM	-0.15	-0.44	-17.8			0.0244	0.218	0.0016	0.936	-0.0101	0.63
NCAR-CAM3.5	-0.28	-0.74	-24.7	0.47	360	0.0298	0.277	0.0013	0.956		
OsloCTM2	-0.43	-1.18	-27.4	1.11	508	0.0432	0.252	0.0022	0.949	-0.0078	0.63
SPRINTARS	-0.14	-0.71	-27.4	0.85	685	0.0260	0.272	0.0012	0.952	-0.0071	0.60
Mean	-0.30	-0.72	-26.8	0.81	531	0.0312	0.238	0.0016	0.946	-0.0060	0.60
Median	-0.31	-0.73	-23.7	0.73	475	0.0298	0.244	0.0015	0.949	-0.0068	0.62
Stddev	0.17	0.22	14.5	0.40	170	0.0116	0.067	0.0007	0.021	0.0045	0.06

* For NCAR-CAM3 the sum of component forcings is used.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Table 4. Anthropogenic load, mass extinction coefficient (MEC), AOD, RF, normalized RF with respect to burden (NRFB), normalized RF with respect to AOD (NRFA) for sulphate.

Model	Load (mg m^{-2})	MEC ($\text{m}^2 \text{g}^{-1}$)	AOD (1)	RF (W m^{-2})	NRFB (W g^{-1})	NRFA (W m^{-2})
BCC	1.29	5.4	0.0069	-0.14	-108	-20.0
CAM4-Oslo	2.78	12.3	0.0342	-0.48	-173	-14.0
CAM5.1	1.69	5.6	0.0095	-0.18	-104	-18.4
GEOS_Chem	1.58	10.3	0.0163	-0.44	-277	-27.0
GISS-MATRIX	1.54			-0.30	-196	
GISS-modelE	1.03	38.6	0.0398	-0.32	-307	-8.0
GMI	2.14	12.0	0.0256	-0.42	-195	-16.3
GOCART	1.87	12.2	0.0228	-0.44	-238	-19.5
HadGEM2	1.59	8.9	0.0142	-0.31	-193	-21.7
IMPACT-Umich	1.42	10.3	0.0146	-0.16	-113	-11.0
INCA	2.26	12.5	0.0283	-0.41	-180	-14.4
ECHAM5-HAM	2.25	9.1	0.0204	-0.28	-125	-13.8
NCAR-CAM3.5	1.27	23.2	0.0295	-0.45	-354	-15.3
OsloCTM2	2.61	11.2	0.0293	-0.58	-223	-19.9
SPRINTARS	2.13	10.3	0.0220	-0.37	-172	-16.6
Mean	1.83	13.0	0.0224	-0.35	-197	-16.8
Median	1.69	10.8	0.0224	-0.37	-193	-16.5
Stddev	0.51	8.5	0.0095	0.13	74	4.8

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 5. Same as Table 4 for BC from FF and BF emissions.

Model	Load (mg m^{-2})	MEC ($\text{m}^2 \text{g}^{-1}$)	AOD (1)	RF (W m^{-2})	NRFB (W g^{-1})	NRFA (W m^{-2})
BCC	0.076	4.2	0.0003	0.05	650	155.1
CAM4-Oslo	0.21	8.2	0.0017	0.37	1763	216.0
CAM5.1	0.074	18.6	0.0014	0.20	2661	143.3
GEOS_Chem	0.12	8.9	0.0011	0.14	1121	126.0
GISS-MATRIX	0.075			0.19	2484	
GISS-modelE	0.16	13.8	0.0023	0.21	1253	90.9
GMI	0.14	12.0	0.0017	0.17	1208	100.4
GOCART	0.21	10.4	0.0021	0.18	874	84.3
HadGEM2	0.31	5.4	0.0016	0.19	612	114.1
IMPACT-Umich	0.09	14.0	0.0013	0.14	1467	104.6
INCA	0.15	9.5	0.0015	0.18	1160	122.5
ECHAM5-HAM	0.10	11.2	0.0011	0.14	1453	130.2
NCAR-CAM3.5	0.11			0.15	1364	
OsloCTM2	0.17	13.2	0.0022	0.38	2271	172.5
SPRINTARS	0.16	7.7	0.0012	0.21	1322	170.8
Mean	0.14	10.5	0.0015	0.19	1444	133.1
Median	0.14	10.4	0.0015	0.18	1322	126.0
Stddev	0.07	3.9	0.0006	0.08	615	37.7

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Table 6. Same as Table 4 for OA from FF and BF emissions.

Model	Load (mg m^{-2})	MEC ($\text{m}^2 \text{g}^{-1}$)	AOD (1)	RF (W m^{-2})	NRFB (W g^{-1})	NRFA (W m^{-2})
BCC	0.35	3.7	0.0013	−0.03	−97	−26.3
CAM4-Oslo	0.28	5.8	0.0017	−0.03	−118	−20.1
CAM5.1	0.44	6.5	0.0024	−0.02	−45	−8.3
GEOS_Chem	0.23	5.3	0.0012	−0.04	−164	−30.7
GISS-MATRIX	0.19			−0.02	−129	
GISS-modelE	0.46	6.1	0.0028	−0.03	−76	−12.4
GMI	0.30	6.6	0.0020	−0.06	−189	−28.5
GOCART	0.42	4.9	0.0021	−0.06	−144	−29.4
HadGEM2	0.24	7.0	0.0017	−0.04	−145	−20.6
IMPACT-Umich	0.20	14.1	0.0028	−0.03	−141	−10.0
INCA	0.62	7.5	0.0046	−0.05	−76	−10.1
ECHAM5-HAM	0.35	1.6	0.0006	−0.01	−41	−25.0
NCAR-CAM3.5	0.21			−0.01	−48	
OsloCTM2	0.45	6.7	0.0030	−0.08	−187	−28.0
SPRINTARS	0.22			−0.02	−102	
Mean	0.33	6.3	0.0022	−0.04	−113	−20.8
Median	0.30	6.3	0.0021	−0.03	−118	−22.8
Stddev	0.13	2.9	0.0010	0.02	49	8.5

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Table 7. Same as Table 4 for SOA.

Model	Load (mg m^{-2})	MEC ($\text{m}^2 \text{g}^{-1}$)	AOD (1)	RF (W m^{-2})	NRFB (W g^{-1})	NRFA (W m^{-2})
CAM5.1	0.27	5.9	0.0016	-0.01	-45	-7.5
GEOS_Chem	0.33	2.2	0.0007	-0.02	-70	-32.8
GISS-modelE	0.090	6.3	0.0006			
IMPACT-Umich	0.97	18.9	0.0184	-0.21	-218	-11.5
ECHAM5-HAM	0.15	10.9	0.0016	-0.02	-139	-12.8
OsloCTM2	0.36	6.9	0.0025	-0.07	-183	-26.4
Mean	0.36	8.5	0.0042	-0.07	-131	-18.2
Median	0.30	6.6	0.0016	-0.02	-139	-12.8
Stddev	0.32	5.8	0.0070	0.08	73	10.8

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 8. Same as Table 4 for nitrate.

Model	Load (mg m^{-2})	MEC ($\text{m}^2 \text{g}^{-1}$)	AOD (1)	RF (W m^{-2})	NRFB (W g^{-1})	NRFA (W m^{-2})
GEOS_Chem	0.82	10.3	0.0078	-0.21	-259	-27.2
GISS-MATRIX	0.44			-0.10	-240	
GISS-modelE	0.16	151.4	0.0246	-0.11	-684	-4.5
GMI	0.76	8.0	0.0061	-0.08	-103	-12.9
HadGEM2	0.44	11.8	0.0051	-0.11	-249	-21.1
IMPACT-Umich	0.78	11.2	0.0088	-0.12	-155	-13.8
INCA	0.44			-0.05	-110	
NCAR-CAM3.5	0.32	6.25	0.002	-0.03	-91	-14.5
OsloCTM2	0.16	11.3	0.0018	-0.03	-191	-16.9
Mean	0.48	30.0	0.0080	-0.09	-231	-15.8
Median	0.44	11.2	0.0061	-0.10	-191	-14.5
Stddev	0.26	53.6	0.0078	0.06	182	7.1

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Table 9. Same as Table 4 for combined OA and BC from BB emissions.

Model	Load (mg m^{-2})	MEC ($\text{m}^2 \text{g}^{-1}$)	AOD (1)	RF (W m^{-2})	NRFB (W g^{-1})	NRFA (W m^{-2})
BCC	0.46	3.9	0.0018	-0.03	-65	-16.8
CAM4-Oslo	2.96	5.4	0.0159	0.07	24	4.5
CAM5.1	0.24	7.0	0.0017	0.04	145	20.7
GEOS-Chem	1.91	2.2	0.0043	-0.05	-25	-11
GISS-modelE				-0.08		
GMI	0.21	7.1	0.0015	-0.06	-291	-40.9
GOCART				-0.02		
HadGEM2	0.48	8.0	0.0038	-0.07	-143	-17.8
IMPACT-Umich	0.88	6.4	0.0056	0.07	84	13.1
INCA	1.76	8.4	0.0147	-0.03	-16	-1.9
ECHAM5-HAM	0.07	22.6	0.0015	0.02	294	13.0
NCAR-CAM3.5	0.21			0.02	95	
OsloCTM2	0.72	6.1	0.0044	-0.02	-25	-4.1
SPRINTARS				0.00	0	0.0
Mean	0.90	7.7	0.0055	-0.01	6	-3.7
Median	0.48	6.7	0.0041	-0.02	-8	-1.9
Stddev	0.92	5.6	0.0054	0.05	146	17.5

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.




Fig. 1. Zonal mean top of the atmosphere short wave (TOA) albedo **(a)** and effective broadband surface albedo **(b)** shown for all the models. CERES TOA albedo data is shown together with the models.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.




Fig. 2. Zonal mean DAE RF for all-sky (a) and RF for clear sky (b).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.




Fig. 3. Anthropogenic AOD (a), anthropogenic AAOD (b), anthropogenic single scattering albedo (c), and change in single scattering albedo from pre-industrial to present conditions (d). All values are taken at 550 nm. Between 80° S and 30° S the anthropogenic AOD is extremely small for some of the models and anthropogenic single scattering albedo may reach unrealistic values and in such cases values have been removed from the figure.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.




Fig. 4. Radiative forcing from the six components, overlain with the (unmodified) model total forcing (yellow bars).

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.




Fig. 5. Model total RFs. Black bars show the bare modelled forcing, the colored bars show the forcing modified for untreated components (see text for details). The yellow bar shows the AeroCom mean of the total RF of DAE. Solid lines inside the boxes show the model mean, dashed lines show the median. The boxes indicate one standard deviation, while the whiskers indicate the max and min of the distribution. The yellow shaded bar shows the AeroCom mean when aerosol component adjustment is made for missing aerosol components.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.




Fig. 6. Correlation between anthropogenic absorption AOD and atmospheric absorption. Numbers show ratio AtmAbs/AAOD, the lines indicate the mean and one standard deviation of this ratio. $R^2 = 0.73$.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.




Fig. 8. Zonal mean SO₄ RF (a), burden (b), AOD at 550 nm (c), normalized RF with respect to AOD (NRF(A)) (d).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Fig. 9. Zonal mean BC RF (a), burden (b), AOD at 550 nm (c), NRF(A) (d).

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.




Fig. 10. Zonal mean OAFF RF (a), burden (b), AOD at 550 nm (c), NRF(A) (d).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.




Fig. 11. Zonal mean SOA RF (a), burden (b), AOD at 550 nm (c), NRF(A) (d).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.




Fig. 12. Zonal mean NO_3 RF (a), burden (b), AOD at 550 nm (c), NRF(A) (d).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.




Fig. 14. Aerosol forcing partial sensitivities for the AeroCom models. The partial sensitivities are calculated as $P_{x,n} = x_n / \langle x \rangle \cdot \langle \text{RF} \rangle$, where n is model, x is either burden, MEC, NRF (with respect to AOD) or RF, $\langle \rangle$ denote mean values. Black dotted line is the mean of the AeroCom models. (Some of the results from the GISS-models are removed due to unresolved issues). Results shown for sulphate, black carbon, organic carbon, biomass burning, SOA, and nitrate as given in the figure.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.




Fig. 15. Correlations between burdens (**a** and **d**), RF (**b** and **e**) and normalized forcing (**c** and **f**), for SO₄ vs. BCFF (**a**, **b**, and **c**) and OCFF vs. BCFF (**d**, **e**, and **f**).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

**Radiative forcing of
the direct aerosol
effect from AeroCom
Phase II simulations**

G. Myhre et al.




Fig. 16. Normalized PDFs of each aerosol component RF (dashed lines) based on the model spread shown above. The red line shows the mean of the modeled total aerosol RF, while the black line shows the spread when correcting for missing components as explained in the text.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Radiative forcing of the direct aerosol effect from AeroCom Phase II simulations

G. Myhre et al.




Fig. 17. Model mean RF per component (internal bars), modified from the study timeperiod of 1850–2000 to 1750–2010, based on numbers from Skeie et al. (2011). Details to be given in text.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)