Atmos. Chem. Phys. Discuss., 12, 24501–24530, 2012 www.atmos-chem-phys-discuss.net/12/24501/2012/ doi:10.5194/acpd-12-24501-2012 © Author(s) 2012. CC Attribution 3.0 License.



This discussion paper is/has been under review for the journal Atmospheric Chemistry and Physics (ACP). Please refer to the corresponding final paper in ACP if available.

The relative importance of impacts from climate change vs. emissions change on air pollution levels in the 21st century

G. B. Hedegaard^{1,2,*}, J. H. Christensen¹, and J. Brandt¹

¹Aarhus University, Department of Environmental Science, Roskilde, Denmark ²Danish Climate Center, Danish Meteorological Institute, Copenhagen, Denmark *now at: Center for Climate and Environmental Research, Lund University, Lund, Sweden

Received: 21 June 2012 – Accepted: 6 September 2012 – Published: 19 September 2012

Correspondence to: G. B. Hedegaard (gitte.brandt_hedegaard@cec.lu.se) and J. Brandt (jbr@dmu.dk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

	ACPD 12. 24501–24530. 2012			
Paner Di	Impacts of Climate vs. emission change on air pollution levels			
critecion	G. B. Hedegaard et al.			
Dung	Title	Title Page		
	Abstract	Introduction		
_	Conclusions	References		
	Tables	Figures		
	I.	►I.		
n D C R	•	•		
_	Back	Close		
	Full Sc	Full Screen / Esc		
	Printer-frie	Printer-friendly Version		
עס	Interactive	Interactive Discussion		
D				

Abstract

So far several studies have analysed the impacts of climate change on future air pollution levels. Significant changes due to impacts of climate change have been made clear. Nevertheless, these changes are not yet included in national, regional or global

- ⁵ air pollution reduction strategies. The changes in future air pollution levels are caused by both impacts from climate change and anthropogenic emission changes and the importance of these signals needs to be quantified and compared. In this study we use the Danish Eulerian Hemispheric Model (DEHM) driven on meteorological input data from the coupled Atmosphere-Ocean General Circulation Model ECHAM5/MPI-
- ¹⁰ OM and forced with the newly developed RCP4.5 emissions. The relative importance of the climate signal and the signal from changes in anthropogenic emissions on the future ozone, black carbon (BC), total particulate matter with a diameter below 2.5 μ m (total PM_{2.5} including BC, primary organic carbon (OC), mineral dust and secondary inorganic aerosols (SIA)) and total nitrogen (including NH_x + NO_y) has been deter-
- ¹⁵ mined. For ozone the impacts of anthropogenic emissions dominates though a climate penalty is found in the Arctic region and the Northwestern Europe where the signal from climate change dampens the effect from the projected emission reductions of anthropogenic ozone precursors. The investigated particles are even more dominated by the impacts from emission changes. For black carbon the emission signal dominates
- ²⁰ slightly at high latitudes increasing to be up to an order of magnitude larger close to the emission sources in temperate and subtropical areas. Including all particulate matter with a diameter below 2.5 μ m (total PM_{2.5}) enhances the dominance from emissions change. In contrast, total nitrogen (NH_x + NO_y) in parts of the Arctic and at low latitudes is dominated by impacts of climate change.



1 Introduction

All meteorological parameters do effect the chemistry and physics of the atmosphere either directly or indirectly through various chemical and physical interactions and feedback mechanisms. Since the last IPCC report (Pachauri and Reisinger, 2007) it had

- ⁵ become clear that climate change is already occurring and will continue in the future which means that the global, regional and local meteorological conditions will change in the future. Furthermore emissions will change due to population growth and technology evolution. In the early and mid 1970s, the first global pollutant problems were identified. The Arctic haze were (re-)discovered in the early 1970s (Wilkening, 2011) and after
- some years it became clear that it has its origin from Asian and European pollution. Since then, air pollution was no longer a local or regional problem but turned in to a global problem and national and international air pollution legislations were formulated and enforced. New legislations are formulated every year in order to prevent future atmospheric pollution levels to amplify or to clean up past and present pollution in the
- ¹⁵ purpose of returning to cleaner atmospheric conditions. Therefore large changes in both climate conditions and emission levels and distributions are to be expected in the future and these will together determine the future air pollution levels.

Future changes in emissions of air pollutants impacts the future air pollution levels both on a global, regional and local scale. Additionally, changes in the future climate
conditions have significant effects on the global, regional and local air pollution levels. Most recent studies have concentrated on the signal from climate change (Langner et al., 2005; Murazaki and Hess, 2006; Hedegaard et al., 2008, 2011) and only a few have compared the climate and the anthropogenic emission signals over a limited areas in the US (Tagaris et al., 2007; Wu et al., 2008b; Pye et al., 2009; Racherla and 25 Adams, 2009).

In this study we hypothesize that climate change can in some areas have significant impact on future air quality relative to the impacts from changes in future emissions. The signals from changes in climate conditions and emissions might cancel out, damp



or even amplify each other depending on the sign of the individual contributions. We aim to estimate the size and sign of the impact from changes in climate and changes in emissions relative to each other and relative to the total predicted change due to both drivers together. The entire Northern Hemisphere has been analysed. However,

it should be emphasized that nobody knows the future and that this study is to be considered as a sensitivity study and a first step in the direction of quantifying the relative importance of impacts from climate change vs. emission change in this century. In the following section the model setup and scientific method are described. In Sect. 4 the results of ozone, black carbon (BC), PM_{2.5}(including primary emitted mineral dust, black carbon (fresh and aged), organic carbon, and the secondary formed particles H₂SO₄, NO₃⁻, NH₄NO₃, NH₄HSO₄ and (NH₄)2SO₄) and total nitrogen (N_x + NO_y) are presented and discussed. The current model setup do not include Secondary Organic Aerosols (SOA). Finally the conclusions are summarised in Sect. 5.

2 Model setup and method

- ¹⁵ In this study the Danish Eulerian Hemispheric Model (DEHM), is driven by six-hourly meteorology input simulated by the coupled Atmosphere-Ocean General Circulation Model ECHAM5/MPI-OM and forced with the newly developed RCP4.5 emissions (Clarke et al., 2007; Smith and Wigley, 2006; Wise et al., 2009). The faith of 58 chemical species and 9 classes of particulate matter has been simulated for the 1990s and
- the 2090s decades in order to quantify the change air pollution levels due to climate chance and changes in emissions levels respectively. The current model version do include Black carbon (BC), primary emitted Organic Carbon (OC), mineral dust and Secondary Inorganic Aerosols (SIA) however Secondary Organic Aerosols (SOA) are not included in this model version. The performance of the total model system with ECHAM5/MPI-OM model coupled to the DEHM model system has been thoroughly
- tested in earlier studies (Hedegaard, 2007; Hedegaard et al., 2008). DEHM is an



Eulerian atmospheric chemistry transport model and a thorough model description can be found in Brandt et al. (2012) and references therein.

The coupled atmosphere-ocean model ECHAM5/MPI-OM consists of an Atmospheric General Circulation Model ECHAM5 (Roeckner et al., 2003, 2006) and the ocean-sea-ice model MPI-OM (Marsland et al., 2003). The specific simulation used in the climate model for this experiment is forced with the SRES A1B scenario (Nakicenovic et al., 2000) and details about the parametrization and aerosol effect for this particular model version can be found in May (2008). It should be noted that the forcing from emission A1B only applies to the projected meteorology. The anthropogenic emissions that feed into the chemical transport model DEHM is based on the newly developed RCP4.5 emission scenarios (Clarke et al., 2007; Smith and Wigley, 2006; Wise et al., 2009). The Representative Concentration Pathway (RCP) stabilizes at ra-

diative forcing of 4.5 W m^{-2} after year 2100. The emission data are global and have a resolution of $0.5^{\circ} \times 0.5^{\circ}$. One emission data set is provided for each decade, which in this case means that for the period 1990–1999, the emissions are represented by the RCP emissions for the year 2000 and for the period 2090–2099 we use the emission data set from year 2100. On the contrary the biogenic emission of isoprene is calcu-

lated dynamically in the model according to the GEIA natural VOC emission model (Guenther et al., 1995). Other naturally emitted VOCs like for example terpenes are not yet included in the model.

Several simulations has been carried out with different combinations of meteorology and emissions in order to study the relative importance of impacts from climate chance and impacts from changes in the anthropogenic emissions. The decade from 1990– 1999 (denoted met1990s) and from 2090–2099 (denoted met2090s) are investigated

²⁵ to give a first estimate of the changes during the 21st century. As a reference period the 1990s decade has been simulated with meteorology predicted by the ECHAM5/MPI-OM model for the same period (1990–1999) and constant 2000 emission. The reference simulation is from now on denoted x(met1990s, emis2000) where x is the concentration or deposition of a specific chemical specie like e.g. ozone.



The climate change signal (CS) relative to the reference period is given by:

 $CS = \frac{x(met2090s, emis2000) - x(met1990s, emis2000)}{x(met1990s, emis2000)}$

(1)

(2)

where *x* is a given parameter (like e.g. ozone concentration or nitrogen deposition) and *x* (met2090s,emis2000) represent the level of the given parameter *x* due to future meteorology for the period 2090–2099 and constant 2000-level emission derived from the RCP4.5 emission database. This means that in Eq. (1) above the signal from climate change is found by keeping the anthropogenic emissions constant at a present-day level and changing the meteorology.

For CS > 0: the given parameter increase due to climate change.

For CS = 0: the given parameter does not change due to climate change.

For CS < 0: the given parameter decrease due to climate change.

Similarly the signal from changes in emissions is identified by a simulation with present day meteorology 1990–1999 and scenario emissions for the year 2100, denoted x(met1990s,emis2100).

15 E

 $ES = \frac{x(met1990s, emis2100) - x(met1990s, emis2000)}{x(met1990s, emis2000)}$

The signal from emission change (ES) for a given parameter x is obtained by keeping the meteorology constant and force the simulation with future year-2100 level emissions based on the RCP4.5 emission scenario.

For ES > 0: the given parameter increase due to emission change.

²⁰ For ES = 0: the given parameter does not change due to emission change.

For ES < 0: the given parameter decrease due to emission change.



Finally the total signal (TOTS) or best guess of the future is based on future meteorology 2090–2099 and scenario predicted emissions for the year 2100 and this is denoted x(met2090s,emis2100).

 $TOTS = \frac{x(met2090s, emis2100) - x(met1990s, emis2000)}{x(met1990s, emis2000)}$

It should be noted that the any changes are non-linear for most species due to the non-linear nature of the atmospheric chemistry (except for the primary particles). E.g. CS + ES ≠ TOTS. The signal from climate change and the signal from emission change do not add up to the total signal from climate and emission change.

In order to quantify the size of the signals relative to one and another three fractions have been set up:

The climate signal relative to the total signal is given by:

10

15

20

$\frac{\text{CS}}{\text{TOTS}} = \frac{x(\text{met2090s}, \text{emis2000}) - x(\text{met1990s}, \text{emis2000})}{x(\text{met2090s}, \text{emis2100}) - x(\text{met1990s}, \text{emis2000})}$				
The emission signal relative to the total signal is given by:				
$\frac{\text{ES}}{\text{TOTO}} = \frac{x(\text{met1990s}, \text{emis2100}) - x(\text{met1990s}, \text{emis2000})}{(x + 10000)}$				
TOTS x (met2090s, emis2100) – x (met1990s, emis2000)				
The climate signal relative to the emission signal is given by:				

 $\frac{CS}{CS} = \frac{x(met2090s, emis2000) - x(met1990s, emis2000)}{x(met1990s, emis2000)}$

 $\overline{\text{ES}} = \frac{1}{x(\text{met1990s}, \text{emis2100}) - x(\text{met1990s}, \text{emis2000})}$

From Eqs. (4) and (5) the size of the climate and emission signal relative to the the total change including non-linear effects can be determined. Furthermore Eq. (6) give the size of the climate signal relative to the emission signal. Whether the two signals amplify or dampens each other can be seen from the climate signal (Eq. 1) and the emission signal (Eq. 2), respectively. This means;



(3)

(4)

(5)

(6)

For CS/ES = 1: the climate and emission signal is of equal size and sign and the sign can be determined from Eqs. (1) and (2).

For CS/ES > 1: the size of climate signal if larger than the size of the emission signal and both effects are either both positive (increasing) or both negative (decreasing) and therefore results in an amplified effect on a given concentration or deposition (x).

For CS/ES < 1: the emission signal dominates and either the climate signal or the emission signal is negative. The sign of the two signals can again be determined from Eqs. (1) and (2) (see e.g. Fig. 3a and b in the case of ozone concentration).

3 Meteorology by ECHAM5/MPI-OM

5

In the following, the output from the ECHAM5 climate simulation used to drive the DEHM model is described. The temperature, humidity and precipitation are direct output from ECHAM5 simulation whereas the mixing height is derived from an energy balance considerations and the radiation is calculated from the cloud cover (for details see Hedegaard, 2007). The ECHAM5 climate simulation used in this study where a part of the 4th IPCC Assessment Report (AR4) multi-model ensemble study. In the current simulation the global temperature is predicted to increase by 3.0 °C by the end of the 21st century and 4.3 °C by the end of the 22nd century, both relative to the period 1971–2000 (May, 2008). This increase is a little higher than the average value (2.7 °C and 3.4 °C, respectively) predicted by the multi-model ensemble following the SRES A1B scenario in the AR4 (Meehl et al., 2007). However, it is well within the standard deviation of the IPCC AR4 multi-model ensemble by the end of the 21st century.

The model setup is the same as in Hedegaard et al. (2011) and details about the temperature, humidity and global radiation can be found therein. The following illustrations

²⁵ of the meteorological output are limited to display the precipitation frequency and the mixing height of the 1990 and 2090 decades. In Fig. 1 the annual mean precipitation



frequency of the two decades considered together with the absolute change between these decadal mean values and the significance of these changes using a student's t-test (Spiegel, 1992).

The precipitation frequency is defined by a threshold value of 1 mm for precipitation occurring in a given six-hour interval and is defined as a fraction between 0 and 1, where 1 indicates precipitation in a given grid cell in a given six-hour time interval. The choice of 1 mm threshold value is based on the parameterizations of the in- and below-cloud scavenging in the DEHM model, for further details see Hedegaard (2007).

Figure 1 shows the (a) decadel mean precipitation frequency in 1990s, (b) in the 2090s, (c) the difference between the 2090s and the 1990s and finally in (d) the significance of the difference between the two decades according to the students t-test. The white colours indicates no significant change and the threshold value for significance is set to 10%. In general the precipitation frequency is projected to increase at high and low latitudes and decrease at mid-latitudes. Focusing on Europe the precipitation fre-

quency is projected to decrease significantly in the Southern Europe and oppositely an increase is projected in Scandinavia, Finland, Iceland, and Greenland. More generally, the precipitation frequency is projected to increase North of about 60° N and decrease significantly in the subtropical part of the Pacific and Atlantic Ocean, the Caribbean, Mexico and the Central South America and in Western Africa.

Figure 2 shows the mixing height in m. The mixing height is calculated from an energy balance equation for the internal boundary layer according to the methods described in Christensen (1997). Figure 2 is given similar to the precipitation frequency where (a) is shows the 1990s decadal mean mixing height, (b) the 2090s decadal mean, (c) the difference in m and (c) the significance of this difference. In South East-

ern Europe the mixing height is projected to increase in the range 50 m to above 100 m which gives a relative change of app. 20 %. Small increases (0–25 m) are found in general over Eurasia and the Arctic Ocean. In Mexico, the Caribbean and in Central South America the mixing height is also projected to increase in the range 50 m to above 100 m. In general the mixing height is projected to decrease over marine areas.



4 Results and discussion

In the following the results of ozone (O_3), Black Carbon (BC), and total $PM_{2.5}$ (including primary emitted mineral dust, black carbon (fresh and aged), organic carbon, and the secondary formed particles H_2SO_4 , NO_3^- , NH_4NO_3 , NH_4HSO_4 and $(NH_4)2SO_4$) and total nitrogen (sum of NO_v and NH_x) are displayed and discussed.

4.1 Ozone

5

10

Figure 3 illustrates the changes in ozone surface concentration due to impacts of climate change and changes in the anthropogenic emissions between the 1990s and 2090s. In Fig. 3a the signal from climate change is displayed. The change in ozone concentration in this figure is solely due to climate change and the anthropogenic emissions have been kept constant at a 2000 level. The ozone concentrations is projected to increase over the Arctic, the densely populated areas and the terrestrial tropics. Elsewhere the ozone concentration will decrease due to climate change alone.

The increase in Arctic is likely to be due to increased transport of ozone from the source areas at lower latitudes in combination with reduced amount of sea ice in the future. O₃ dry deposit more effectively to sea ice than to open water in the model and decreased sea ice in the 2090s results in increased ozone in the Arctic air masses. Further increased import of ozone from higher layers can also explain the projected increase in ozone concentration due to climate change (Hedegaard et al., 2011).

In densely populated areas and over the tropics the NO_x and/or the VOC level in general are high and the changed climate leads to enhanced productions of BVOCs and hence higher ozone levels over these areas. In the rest of the domain the effect from increased water vapour in the atmosphere enhances ozone destruction and this process is dominating in the areas with lower NO_x and VOC concentrations (for further details see discussion in Hedegaard et al., 2011).

Figure 3b illustrates the changes in the ozone concentration due to changes in the anthropogenic emission between the two decades 1990s and 2090s. In this simulation



the meteorology have been kept constant according to the 1990s meteorology in order to isolate the signal from changed anthropogenic emissions. In most of the domain the emission signal is opposing the signal from climate change. The model estimates a decrease in the current ozone concentration in the order of 20 % between the 1990s

- ⁵ decade and 2090s decade solely due to changes in the anthropogenic emissions. The tendency is different in parts of Northwestern Europe. In the Benelux countries and in the vicinity of this area the ozone concentrations are estimated to increase due to changes in the anthropogenic emissions. This is also the case in Africa south of Sahara.
- ¹⁰ The projected increase in ozone concentrations in the future in these areas can be explained by the applied emission inventory. The NO_x emissions are general prescribed to decrease in the future according to the RCP4.5 emission scenario (www.iiasa.ac.at/ web-apps/tnt/RcpDb). In the Benelux area the NO_x emissions are also projected to decrease, however this area differs from the rest of the densely populated areas in the
- ¹⁵ Northern Hemisphere. The largest density of NO_x emissions over the largest area is found in Benelux and the surrounding areas. The area is characterized by urban area chemistry (high NO_x area) and is at same time very large. The ozone present in urban area is used to convert the emitted NO to NO₂ which is called the ozone titration effect or urban deficit (Fowler et al., 2008). This means that lowering the emissions of NO
 ²⁰ will increase the amount of ozone which alters chemical regime. Moreover, European NO_x emissions are due to the regional legislations expected to be lowered significantly

more than other other high NO_x areas in the world.

In Africa a large increase in the future O_3 concentration is found (Fig. 3b) which is explained by a large increase in the anthropogenic NO_x emissions combined with general

high biogenic BVOC emissions in the tropics. About half of the global isoprene emissions originate from tropical broadleaf trees (Guenther et al., 2006) and the tropical meteorological conditions are highly conductive for isoprene emissions.

In Fig. 3c the combined effect on the future ozone concentration from both changes in anthropogenic emissions and changes in the climate is shown including any non-linear



effect from the ozone chemistry included in the model. The ozone concentration decrease due to impacts from changes in both the future anthropogenic emissions and climate, except over the terrestrial Tropics and Northwestern Europe (Benelux area).

Figure 3d shows the relative importance of the two individual signals. This is illus-

trated by the fraction: "Climate signal" divided by "Emission signal" (Eq. 6). Figure 3d reveals that the increase in surface ozone concentration in the terrestrial Tropics and Northwestern Europe are mainly due to the impacts of climate change. The climate signal dominates and is more than twice the size of the impact from changes in the anthropogenic emissions. Elsewhere the impacts from changes in anthropogenic emissions
 are dominating.

In the Arctic region a minor decrease in ozone concentration is projected due to a composite of two opposing signals. The impacts of climate change leads to a 5-10% increase (Fig. 3a) and the reduction in emission of ozone precursors implies a 5-10% decrease (Fig. 3b). The total signal is displayed in Fig. 3c and shows a minor overall decrease in the Arctic by the end of the 21st century. Figure 3d shows that the climate signal is a little weaker than the emission signal (0.75–1.00).

In general the opposition of the two signals (impacts from climate change and impacts of changes in anthropogenic emissions) means that to obtain a certain reduction target in the future, additional reductions must be made in order to compensate from the empering signal from climate shapes. This feature has in the literature have de-

²⁰ the opposing signal from climate change. This feature has in the literature been denoted "the climate penalty" (see e.g. Wu et al., 2008a).

4.2 Black carbon and total PM_{2.5}

15

25

In Fig. 4 the change in Black Carbon (BC) surface concentration due to changes in (a) climate, (b) anthropogenic emissions and (c) changes in both climate and anthropogenic emissions is shown. The latter is equal to the addition of (a) and (b), since BC is an inert tracer and do not react chemically with other species in the atmosphere. In Fig. 4d the climate signal is illustrated relative to emission signal.



The BC concentration is projected to decrease in the Arctic, in Scandinavia, over Eastern Europe and Russia and over large parts of the Pacific Ocean due to impacts of climate change (Fig. 4a). An increase is found nearly elsewhere. Since BC is an inert tracer in the model and the emissions are kept constant at a year 2000 level, only changes in the physical (e.g. meteorological) conditions can be the explanation of the projected changes.

5

10

In Fig. 5 the isolated effect from climate change on the BC (a) wet deposition, (b) dry deposition, (c) atmospheric concentrations and finally (d) the total deposition is shown. The plots display the relative change in decadal mean values between the 1990s and the 2090s. At high latitudes and over the North Eastern Pacific Ocean the atmospheric concentration of BC (Fig. 5c) is found to decrease significantly due to impacts of climate change.

In and South of the Mediterranean Sea, along the East Coast of North America and Asia and in general at low latitudes the atmospheric concentration of BC is found

- to increase due to climate change. In the Mediterranean area there is a close relation between decreased precipitation frequency (Fig. 1) and decreased wet deposition (Fig. 5a) and increased atmospheric concentrations of BC. Over the North American and Asian East Coast, on the contrary, the increased BC concentration in the air, can be explained by a decrease in the mixing height (Fig. 2). Over the tropical terrestrial
- ²⁰ areas (Brasil/Sahara) the precipitation frequency decreases and hence the wet deposition decreases which leads to increased atmospheric levels of BC.

Figure 4b shows the relative contribution from the impact of changes in anthropogenic emissions on the atmospheric BC distribution according to the RCP4.5 emission scenario. The contribution from emission change leads to a decrease in BC in

the majority of the domain, though there are some differences in the size of the signal. Since the meteorology is kept constant (1990s level) in this simulation, to isolate the impact from changes in the emissions, only change in emissions size and spatial distribution can explain the pattern of Fig. 4b.



The majority of BC emissions stems from the transportation, industry, residential and biomass burning sectors and BC is therefore mainly emitted in and near the terrestrial areas. In the future these emissions is prescribed by the RCP4.5 scenario to decline significantly. In contrast the emissions over the ocean are prescribed to increase. This

⁵ increase originates from increased shipping activities in the remote marine areas, see www.iiasa.ac.at/web-apps/tnt/RcpDb for emissions from the individual sectors. In the current model setup the emissions from the aviation sector have not been included.

In Fig. 4c the total signal on BC from changes in both emissions and climate conditions is shown and further the climate signal relative to the emission signal is given

- in Fig. 4d. In general the concentration level of BC are projected to be 30–60 % lower in the future. In the tropics a smaller decrease is found in the order 0–30 % and in small parts of the tropics even an increase is found. The distinguished area south of the Aleutian are strongly dominated by the signal from climate change. In this area the climate signal oppose the projected increase (orange color) due to changes in the
- anthropogenic emissions from ships. However, analysis of the absolute BC concentrations for the 1990s and the 2090s due to both climate change and emission change has shown that the distinct pattern South of the Aleutians is a results of small changes. In general it can be concluded that the changes due to impacts of climate change on an inert tracer like BC is small compared to the impacts from changes in the anthropogenic emissions which are absolutely depineding. The BC concentration will deprese due to
- ²⁰ emissions which are absolutely dominating. The BC concentration will decrease due to a general reduction in the future BC emissions.

The total $PM_{2.5}$ consists in the model of the sum of the following species: primary emitted mineral dust, black carbon (fresh and aged), organic carbon, and the secondary formed particles H_2SO_4 , NO_3^- , NH_4NO_3 , NH_4HSO_4 and $(NH_4)2SO_4$. Sec-

ondary formed organic aerosols (SOA) are not included in the current model setup. Figure 6 is similar to Fig. 4 and shows the relative impacts from (a) climate change, (b) emission change and (c) total change, (d) the climate signal relative to emission signal for total PM_{2.5}.



The impact of climate change alone (Fig. 6a) results in a decrease in the total $PM_{2.5}$ at high latitudes (orange and red colours) and in an small increase (0–15%) over a significant part of the Atlantic ocean and in subtropical and tropical areas in general. Though a small decrease (0–15%) is found over most of the subtropical Indian ocean

⁵ and Pacific ocean. In contrast the anthropogenic emissions acts to decrease the future total PM_{2.5} concentration significantly in most of the domain except over Central Africa where changes in anthropogenic emissions results in a small increase.

From Fig. 6d it can be seen that the impact from changes in the total $PM_{2.5}$ emissions dominate in the future. At high latitudes the overall signal from climate change and emissions change work in same direction and therefore amplify each other. In general

- ¹⁰ emissions change work in same direction and therefore amplify each other. In general the emission signal in future total PM_{2.5} concentration ranges from being twice the size of the climate signal to be an order of magnitude larger. In case of BC only a slight dominance of the emission signal can be found in the Arctic region increasing to be an order of magnitude higher close to and downstream of the emission sources at mid-15 latitudes. Further, the changes in total sulfate concentration have been analysed (not
- shown) and the pattern is very similar to the projection of BC.

4.3 Total N

Total N are defined by the sum of NO_y and NH_x . In Fig. 7 the individual contributions to the projected changes in total N concentration is shown. This figure is displayed similar

- to Figs. 3, 6 and 4 where the signal from (a) climate change, (b) signal from emission change and (c) total change and finally (d) the climate signal relative to emission signal is shown, respectively. Since total N contains both NH_x and NO_y the total N distribution is dependent on chemical reactions. Therefore the total signal displayed in Fig. 7c is like in the case of O_3 and total $PM_{2.5}$ different from the addition of Fig. 7a and b.
- ²⁵ Changing the climate alone (Fig. 7a leads to a decrease in the nitrogen concentration in the future north of ~ 50° N except over Europe where a small increase (0–10%) is found over Great Britain, Denmark, Southern Sweden and the North and Baltic sea



(Fig. 7a). At subtropical and tropical latitudes the impacts from climate change gives rise to an increase in the N concentration in the order 0-30 % (Fig. 7a).

In Fig. 7b the isolated effect of changed antropogenic emissions is shown. Over Asia (except Siberia), Africa, Central America and Southern US the impacts of changed emissions results in an increase above 30 % in the atmospheric nitrogen concentration. Elsewhere the total nitrogen concentration is projected to decrease due to impacts of emission change alone. The decrease is largest at high latitudes and is above 40 % in the Arctic, over Europe, Japan and coastal Areas of the US. The total changes,

including both impacts from climate and emission change, is shown in Fig. 7c. The
spatial pattern in Fig. 7c is very similar to the signal shown in Fig. 7b, except over
the Pacific ocean and South America. The impact from changed emission dominates
(Fig. 7b) and from Fig. 7d it can be seen that emission signal dominates in most areas
ranging from being twice the size to an order of magnitude higher. In a some regions the
impacts from climate change is dominating, however this is mainly where both signals
are very small (e.g. large part of the Tropics). In the terrestrial parts of the Arctic region
(Northern Canada, Greenland and Siberia) the impacts from climate change is twice

the size of the emission signal on future atmospheric total N concentration. Nitrogen deposition can have harmful effects on the marine and terrestrial ecosystems, hence the future nitrogen deposition has been analysed. The total (dry and wet)

- nitrogen deposition for the 1990s and 2090s decade is shown in Fig. 8 together with the difference (2090s–1990s) and the significance of the difference. Over the densely populated areas of Asia, in Africa and on the west coast of South America the future nitrogen deposition is projected to increase by more than 20% (Fig. 8c. Elsewhere the future total nitrogen deposition will decrease significantly. An exception is in the region
- ²⁵ of the Rocky Mountains and parts of China where an increase is projected similar to the tropical region. Further, the areas in the Arctic that from Fig. 7a are found to be controlled by the impacts of climate change are areas with extremely low nitrogen concentrations (not shown) and hence the changes due to climate change are also very low.



5 Summary and conclusions

10

So far only a few studies have concentrated on the relative importance of impacts from climate change and emission change and these studies has been limited to focus on smaller regions in the US. All three studies compares present day with small periods around 2050 and the results are therefore not comparable to this study. Nevertheless,

the overall conclusions are similar to the results found in this study.

In this study we used the Danish Eulerian Hemispheric Model (DEHM) driven on meteorological input from the coupled Atmosphere-Ocean General Circulation Model ECHAM5/MPI-OM and forced with the newly developed RCP4.5 emissions. The relative importance of the climate signal and the signal from changes in anthropogenic emissions on the future O_3 , BC, total $PM_{2.5}$ (Secondary Organic Aerosols (SOA) are not included in the model) and total N (sum of NO_v and NH_x) has been determined.

The changes in anthropogenic emissions dominate in general over the signal from climate change. However in some cases the signal from climate change are oppos-

ing the signal from the prescribed emission reductions, which implies that to obtain a certain reduction target in the future, additional reductions must be made in order to compensate from the opposing signal from climate change. This feature is known as "the climate penalty" (see e.g. Wu et al., 2008a).

Specifically, the ozone concentration over the Arctic area is found to decrease only a
 little in the future due to impact from both climate change and changes in the emission.
 Prescribed reduction of the ozone precursors in the source areas implies a significant decrease in the Arctic in the future. Oppositely do the impacts from climate result in a significant increase which is a little weaker than the decrease due to emissions reductions which minimize the overall effect to a minor decrease in the future Arctic ozone
 concentration.

In Northwestern Europe it was found that future NO_x reductions results in a rise in the ozone concentration, due to the extreme NO_x emission density in this area. This means that that future air pollution control policies have to account for the amplified



impact from climate change in order reach a specific reduction target. This implies that emissions of VOCs also needs to be considered as well in order to reduce both ozone and NO_x levels in the future in Northwestern Europe.

Compared to O₃ the investigated particles are even more dominated by the impacts from emission changes. For BC the emission signal dominates slightly at high latitudes increasing to be up to an order of magnitude larger close to the emission sources at temperate and subtropical areas. In the marine tropical areas the climate signal is twice the size of the impacts from changed emissions. In this region only very minor changes due to emission reductions is expected and this is combined with the general large uncertainties related to climate projections in the tropics.

The total $PM_{2.5}$ are similar to the BC concentration dominated by changes in emissions in most of the domain. The signal from changes in anthropogenic emissions ranges from being twice the size of the climate signal to be an order of magnitude larger than the climate signal for the total $PM_{2.5}$ concentration.

- ¹⁵ On the contrary the atmospheric total nitrogen concentration is in parts of the Arctic and at low latitudes dominated by impacts of climate change. In the terrestrial parts of the Arctic region (Northern Canada, Greenland and Siberia) the impacts from climate change is twice the size of the emission signal on future atmospheric total N concentration. However these are also regions with very low nitrogen concentrations and hence
- ²⁰ the changes are very small. In general, the impact from changed emissions dominate and is in some areas (e.g. over Europe) up to an order of magnitude higher than the signal from climate change.

The current work is to be considered as a sensitivity study. Nevertheless, it is the most comprehensive study so far, quantifying the impacts of climate change vs. an-

thropogenic emission change. Further the new emission inventory RCP4.5 has been used and the model domain covers the entire Northern Hemisphere. The current results indicate that impacts of climate change in general have less importance in the overall future concentration levels compared to anticipated changes in future anthropogenic emissions. Though, there are certain regions for the different chemical species



(e.g. ozone in the Arctic, north-western part Europe and Africa) where the impacts from climate change on future air pollution levels needs to be accounted for in order to reach a certain reduction level. On the other hand, the results also show that for certain areas, the future emission reductions might be less strict, if certain limit values have to be reached since the climate change impacts induces further decreases in the concen-

trations of particulate matter as well as in the deposition of nitrogen in specific regions.

Acknowledgements. This work was partly funded by the Centre for Energy, Environment and Health (CEEH), financed by The Danish Strategic Research Program on Sustainable Energy under contract no 2104-06-0027 (www.ceeh.dk) and by University of Copenhagen. Both institutions are acknowledged for their financial support. Furthermore Willy May from the Danish Climate Center, Danish meteorological Institute are acknowledge for providing us the ECHAM5 climate data.

References

5

15

20

Brandt, J., Silver, J. D., Frohn, L. M., Geels, C., Gross, A., Hansen, A. B., Hansen, K. M., Hedegaard, G. B., Skjøth, C. A., Villadsen, H., Zare, A., and Christensen, J. H.: An integrated model study for Europe and North America using the Danish Eulerian Hemispheric Model with focus on intercontinental transport of air pollution, Atmos. Environ., 53, 156–176, 2012. 24505

Christensen, J. H.: The Danish Eulerian Hemispheric Model - A three-dimensional air pollution model used for the Arctic, Atmos. Environ., 31, 4169–4191, 1997. 24509

- Clarke, L., Edmonds, J., Jacoby, H., Pitcher, H., Reilly, J., and Richels, R.: Synthesis and Assessment Product 2.1 by the U.S. Climate Change Science Program and the Subcommittee on Global Change Research, subreport 2.1A, Department of Energy, Office of Biological and Environmental Research, Washington DC, USA, 154 pp., 2007. 24504, 24505
- Fowler, D., Amann, M., Anderson, R., Ashmore, M., Cox, P., M, M. D., Derwent, D., Grennfelt, P., Hewitt, N., Hov, Ø., Jenkin, M., Kelly, F., Liss, P., Pilling, M., Pyle, J., Slingo, J., and Stefenson, D.: Ground-level ozone in the 21st century: future trends, impacts and policy implications, Science Policy Report 15/08, The Royal Society, pp. 148, 2008. 24511



- Guenther, A., Hewitt, C., Erickson, D., Fall, R., Geron, C., Graedel, T., Harley, P., Klinger, L., Lerdau, M., McKay, W., Pierce, T., Scholes, B., Steinbrecher, R., Tallamraju, R., Taylor, J., and Zimmerman, P.: A Global-Model of Natural Volatile Organic-Compound Emissions, J. Geophys. Res.-Atmos., 100, 8873–8892, 1995. 24505
- ⁵ Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), Atmos. Chem. Phys., 6, 3181–3210, doi:10.5194/acp-6-3181-2006, 2006. 24511

Hedegaard, G. B.: Impacts of Climate Change on Air Pollution Levels in the Northern Hemi-

- sphere, Technical report 240, National Environmental Research Institute, Aarhus University, Frederiksborgvej 399, P.O. Box 358, 4000 Roskilde, Denmark, www.dmu.dk, 2007. 24504, 24508, 24509
 - Hedegaard, G. B., Brandt, J., Christensen, J. H., Frohn, L. M., Geels, C., Hansen, K. M., and Stendel, M.: Impacts of climate change on air pollution levels in the Northern Hemi-
- sphere with special focus on Europe and the Arctic, Atmos. Chem. Phys., 8, 3337–3367, doi:10.5194/acp-8-3337-2008, 2008. 24503, 24504
 - Hedegaard, G. B., Gross, A., Christensen, J. H., May, W., Skov, H., Geels, C., Hansen, K. M., and Brandt, J.: Modelling the impacts of climate change on tropospheric ozone over three centuries, Atmos. Chem. Phys. Discuss., 11, 6805–6843, doi:10.5194/acpd-11-6805-2011, 2011. 24503, 24508, 24510
 - Langner, J., Bergstrøm, R., and Foltescu, V.: Impact of climate change on surface ozone and deposition of sulphur and nitrogen in Europe, Atmos. Environ., 39, 1129–1141, 2005. 24503
 Marsland, S. J., Haak, H., Jungclaus, J. H., Latif, M., and Roske, F.: The Max-Planck-Institute global ocean/sea ice model with orthogonal curvilinear coordinates, Ocean Modell., 5, 91–127. 2003. 24505

25 127, 2003. 24505 May W: Climatic changes ass

20

May, W.: Climatic changes associated with a global 2 degrees C-stabilization scenario simulated by the ECHAM5/MPI-OM coupled climate model, Clim. Dynam., 31, 283–313, 2008. 24505, 24508

Meehl, G. A., Stocker, T. F., Collins, W. D., Friedlingstein, P., Gaye, A. T., Gregory, J. M., Kitoh,

A., Knutti, R., Murphy, J. M., Noda, A., Raper, S. C. B., Watterson, I. G., Weaver, A. J., and Zhao, Z.-C.: Climate Change 2007: The Physical Science Basis, contribution of working group i to the fourth assessment report of the intergovernmental panel on climate change



24521

Global Climate Projections, pp. 747–846, Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2007. 24508

- Murazaki, K. and Hess, P.: How does climate change contribute to surface ozone change over the United States, J. Geophys. Res., 111, D05301, doi:10.1029/2005JD005873, 2006. 24503
- ⁵ Nakicenovic, N., Alcamo, J., Davis, G., de Vries, B., Fenhann, J., Gaffin, S., Gregory, K., Grübler, A., Jung, T., Kram, T., Rovere, E. L., Michaelis, L., Mori, S., Morita, T., Pepper, W., Pitcher, H., Price, L., Riahi, K., Roehrl, A., Rogner, H.-H., Sankovski, A., Schlesinger, M., Shukla, P., Smith, S., Swart, R., van Rooijen, S., Victor, N., and Dadi, Z.: Special Report on Emission Scenarios: a special report of Working Group III of the Intergovernmental Panel
- on Climate Change, Cambridge University Press, New York, NY (US), United States, 2000. 24505
 - Pachauri, R. K. and Reisinger: Climate Change 2007: Synthesis Report, Contribution of working group i, ii and iii to the fourth assessment report of the intergovernmental panel on climate change, IPCC, 2007. 24503
- Pye, H. O. T., Liao, H., Mickley, L. J., Jacob, D. J., and Henze, D. K.: Effects of changes in climate and emissions on future sulfate-nitrate-ammonium aerosol levels in the United States, J. Geophys. Res., 114, 1–18, doi:10.1029/2008JD010701, 2009. 24503
 - Racherla, P. N. and Adams, P. J.: U.S. ozone air quality under changing climate and antropogenic emissions, Environ Sci. Technol., 43, 571–577, 2009. 24503
- Roeckner, E., Bauml, G., Bonaventura, L., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kirchner, I., Kornblueh, L., Manzini, E., Rhodin, A., Schlese, U., Schulzweida, U., and Tompkins, A.: The atmospheric general circulation model ECHAM5, Part I, Report 249, Max-Planck-Institute für Meteorologie, Hamburg, Germany, 2003. 24505

Roeckner, E., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kornblueh, L. Manzini, E.,

- Schlese, U., and Schulzweida, U.: Sensitivity of simulated climate to horizontal and vertical resolution in the ECHAM5 atmosphere model, J. Climate, 19, 3771–3791, 2006. 24505
 Smith, S. and Wigley, T.: Multi-Gas Forcing Stabilization with the MiniCAM, Energy Journal, Special issue, 2006. 24504, 24505
 - Spiegel, M. R.: Theory and problems of statistics, Schaums Outline Series, McGraw-Hill, secound edn., 1992. 24509

30

Tagaris, E., Manomaiphiboon, K., Liao, K.-J., Leung, L. R., Woo, J.-H., He, S., Amar, P., and Russel, A. G.: Impacts of global climate change and emissions on regional ozone and



fine particulate matter concentrations over the United States, J. Geophys. Res., 12, 1–11, doi:10.1029/2006JD008262, 2007. 24503

- Wilkening, K.: Science and International Environmental Nonregimes: The Case of Arctic Haze, Review of Policy Research, 28, 125–148, 2011. 24503
- ⁵ Wise, M. A., Calvin, K. V., Thomson, A., Clarke, L. E., Bond-Lamberty, B., Sands, R. D., Smith, S. J., Janetos, A. C., and Edmonds, J. A.: Implications of Limiting CO2 Concentrations for Land Use and Energy, Science, 324, 1183–1186, 2009. 24504, 24505
 - Wu, S., Mickley, L. J., Jacob, D. J., Rind, D., and Streets, D. G.: Effects of 2000–2050 changes in climate and emissions on global tropospheric ozone and the policy-relevant background
- ozone in the United States, J. Geophys. Res., 113, D18312, doi:10.1029/2007JD009639, 2008a. 24512, 24517
 - Wu, S., Mickley, L. J., Leibensperger, E. M., Jacob, D. J., Rind, D., and Streets, D. G.: Effects of 2000–2050 global change on ozone air quality in the United States, J. Geophys. Res., 113, D06302, doi:10.1029/2007JD008917, 2008b. 24503

	ACPD			
]	12, 24501–24530, 2012			
-	Impacts o vs. emissio on air pollu	of Climate on change ition levels		
	G. B. Hedegaard et al.			
	Title Page			
	Abstract	Introduction		
,	Conclusions	References		
	Tables	Figures		
	I	▶1		
	•	•		
-	Back	Close		
	Full Screen / Esc			
	Printer-friendly Version			
	Interactive Discussion			

Jiscussion Pape

JISCUSSION Pape

Jiscussion Paper

Jiscussion Pape





Fig. 1. Annual mean precipitation frequency: Defined as a fraction between 0 and 1, the threshold value for precipitation in a given six-hour interval is 1 mm. Panel **(a)** shows the annual mean precipitation frequency during 1990s decade and **(b)** during the 2090s decade. In **(c)** the difference between the 2090s and the 1990s is shown and finally **(d)** illustrates the significance of the difference between the two decades, white colours indicates no significant change and the threshold value for significance is set to 10%.



(a) 1990s annual mean mixing height in m

(b) 2090s annual mean mixing height in m



(c) Difference in annual mean mixing height in m (2090s-1990s)



Fig. 2. Annual mean mixing height in m, displayed as in Fig. 1.

ACPD 12, 24501–24530, 201				
Impacts of Climate vs. emission change on air pollution level				
	Title Page			
DDr -	Abstract	Introduction		
	Conclusions	References		
000	Tables	Figures		
	I.	> 1		
	•	•		
	Back Close Full Screen / Esc			
000	Printer-frien	Printer-friendly Version		
	Interactive Discussion			



Fig. 3. O_3 : the signal from **(a)** future climate change, **(b)** future emission change, **(c)** the total change on the surface ozone concentration due to changes in both future climate and emissions and d) the climate signal relative to the emission signal. The climate signal is simulated with constant year 2000 emissions and ECHAM5 meteorology. The signal from changes in the anthropogenic emissions is simulated with projected RCP4.5 emissions and 1990s meteorology.





Fig. 4. Black Carbon (BC): as Fig. 3 but for black carbon.





Fig. 5. Black Carbon (BC): relative change due to impacts of climate change in % (2090s–1990s). (a) Wet deposition, (b) dry deposition, (c) concentration, and (d) total deposition.





Fig. 6. Total PM_{2.5}: as Fig. 3 but for total PM_{2.5}.





Fig. 7. Total nitrogen (N): as Fig. 3 but for total nitrogen N.





